NASA/ADS

ASASSN-16oh: A Nova Outburst with No Mass Ejection—A New Type of Supersoft X-Ray Source in Old Populations ()

Kato, Mariko (/search/?q=author%3A%22Kato%2C+Mariko%22); Saio, Hideyuki (/search/?q=author%3A%22Saio%2C+Hideyuki%22); Hachisu, Izumi (/search/?q=author%3A%22Hachisu%2C+Izumi%22)

ASASSN-16oh is a peculiar transient supersoft X-ray source without a mass-ejection signature in the field of the Small Magellanic Cloud. Maccarone et al. concluded that ASASSN-16oh is the first dwarf nova with supersoft X-ray that originated from an equatorial accretion belt on a white dwarf (WD). Hillman et al. proposed a thermonuclear runaway model that both the X-rays and V/I photons are emitted from the hot WD. We propose a nova model induced by a high rate of mass accretion during a dwarf nova outburst, I.e., the X-rays originate from the surface of the hydrogen-burning WD whereas the V/I photons are from the irradiated disk. Our model explains the main observational properties of ASASSN-16oh. We also obtained thermonuclear runaway models with no mass ejection for a wide range of parameters of the WD mass and mass-accretion rates including both natural and forced novae in low-metal environments of Z = 0.001 and Z = 0.0001. They are a new type of periodic supersoft X-ray sources with no mass ejection and also a bright transient in V/I bands if they have a large disk. We suggest that such objects are candidates for Type Ia supernova progenitors because its mass is increasing at a very high efficiency (~100%).

Publication:	The Astrophysical Journal
Pub Date:	March 2020
DOI:	10.3847/1538-4357/ab7996 (/link_gateway/2020ApJ89215K/doi:10.3847/1538-4357/ab7996) 🗹
arXiv:	arXiv:2002.10717 (/link_gateway/2020ApJ89215K/arXiv:2002.10717)
Bibcode:	2020ApJ89215K (/abs/2020ApJ89215K/abstract)
Keywords:	1127; 1284; 418; Astrophysics - Solar and Stellar Astrophysics; Astrophysics - High Energy Astrophysical Phenomena
E-Print:	references updated, 21 pages, 10 figures, published in ApJ, 892, 15 (2020)