



# Astronomy Today<sup>8e</sup>

CHAISSON McMILLAN

# Astronomy Today

*This page intentionally left blank*

---

# Astronomy Today

**Eric Chaisson**  
*Harvard University*

**Steve McMillan**  
*Drexel University*

**PEARSON**

Boston Columbus Indianapolis New York San Francisco Upper Saddle River  
Amsterdam Cape Town Dubai London Madrid Milan Munich Paris Montréal Toronto  
Delhi Mexico City São Paulo Sydney Hong Kong Seoul Singapore Taipei Tokyo

Publisher: Jim Smith  
Executive Editor: Nancy Whilton  
Editorial Manager: Laura Kenney  
Project Editor: Tema Goodwin  
Content Producers: Kate Brayton, Ziki Dekel  
Development Editor: Barbara Price  
Director of Marketing: Christy Lesko  
Marketing Manager: Will Moore  
Director of Product Management Services: Erin Gregg  
Team Lead, Program and Project Management: Corinne Benson  
Compositor: Cenveo® Publisher Services  
Production Service: Thistle Hill Publishing Services  
Illustrations: Rolin Graphics Inc.

Design Manager: Mark Ong  
Interior and Cover Design: Jeanne Calabrese  
Manufacturing Buyer: Jeffrey Sargent  
Specialist, Rights and Permissions: Joseph Croscup  
Image Permissions Coordinator: Maya Melunchuk  
Photo Research: Stefanie Ramsay  
Cover Printer: Lehigh-Phoenix  
Printer and Binder: R. R. Donnelley  
Cover Images:  
*Main Edition:* ESO/S. Guisard ([www.eso.org/~sguisard](http://www.eso.org/~sguisard))  
*Vol. 1: The Solar System:* NASA, JPL-Caltech, MSSS, Mastcam  
*Vol. 2: Stars and Galaxies:* ESO/F. Comeron

Copyright © 2014, 2011, 2008, 2005 Pearson Education, Inc., 1301 Sansome St., San Francisco, CA 94111. All rights reserved. Manufactured in the United States of America. This publication is protected by Copyright and permission should be obtained from the publisher prior to any prohibited reproduction, storage in a retrieval system, or transmission in any form or by any means, electronic, mechanical, photocopying, recording, or likewise. To obtain permission(s) to use material from this work, please submit a written request to Pearson Education, Inc., Permissions Department, 1900 E. Lake Ave., Glenview, IL 60025. For information regarding permissions, call (847) 486-2635.

Many of the designations used by manufacturers and sellers to distinguish their products are claimed as trademarks. Where those designations appear in this book, and the publisher was aware of a trademark claim, the designations have been printed in initial caps or all caps.

MasteringAstronomy® is a trademark, in the U.S. and/or other countries, of Pearson Education, Inc. or its affiliates.

#### Library of Congress Cataloging-in-Publication Data

Chaisson, Eric, author.

Astronomy today / Eric Chaisson, Harvard University, Steve McMillan, Drexel University. — Eighth edition.  
pages cm

Includes bibliographical references and index.

ISBN 978-0-321-90167-5 (student edition) — ISBN 978-0-321-90971-8 (volume 1)

— ISBN 978-0-321-90972-5 (volume 2) — ISBN 978-0-13-341279-6 (nasta)

I. Astronomy—Textbooks. I. McMillan, S. (Stephen), 1955– author. II. Title.

QB43.3.C48 2014

520—dc23

Proudly sourced and uploaded by [StormRG]

Kickass Torrents | TPB | ET | h33t

2013019295

ISBN 10 digit 0-321-90167-3; 13-digit 978-0-321-90167-5 (Student edition)

ISBN 10-digit 0-321-90971-2; 13-digit 978-0-321-90971-8 (Volume 1)

ISBN 10-digit 0-321-90972-0; 13-digit 978-0-321-90972-5 (Volume 2)

ISBN 10-digit 0-13-341279-2; 13-digit 978-0-13-341279-6 (NASTA)

**PEARSON**

[www.pearsonhighered.com](http://www.pearsonhighered.com)

1 2 3 4 5 6 7 8 9 10—VHC—19 18 17 16 15 14 13

# Brief Contents

## Part One: Astronomy and the Universe 2

- 1 Charting the Heavens: The Foundations of Astronomy 4
- 2 The Copernican Revolution: The Birth of Modern Science 32
- 3 Radiation: Information from the Cosmos 58
- 4 Spectroscopy: The Inner Workings of Atoms 78
- 5 Telescopes: The Tools of Astronomy 98

## Part Two: Our Planetary System 132

- 6 The Solar System: Comparative Planetology and Formation Models 134
- 7 Earth: Our Home in Space 160
- 8 The Moon and Mercury: Scorched and Battered Worlds 188
- 9 Venus: Earth's Sister Planet 216
- 10 Mars: A Near Miss for Life? 236
- 11 Jupiter: Giant of the Solar System 264
- 12 Saturn: Spectacular Rings and Mysterious Moons 290
- 13 Uranus and Neptune: The Outer Worlds of the Solar System 318
- 14 Solar System Debris: Keys to Our Origin 338
- 15 Exoplanets: Planetary Systems Beyond Our Own 366

## Part Three: Stars and Stellar Evolution 386

- 16 The Sun: Our Parent Star 388
- 17 The Stars: Giants, Dwarfs, and the Main Sequence 420
- 18 The Interstellar Medium: Gas and Dust among the Stars 448
- 19 Star Formation: A Traumatic Birth 468
- 20 Stellar Evolution: The Life and Death of a Star 494
- 21 Stellar Explosions: Novae, Supernovae, and the Formation of the Elements 520
- 22 Neutron Stars and Black Holes: Strange States of Matter 542

## Part Four: Galaxies and Cosmology 574

- 23 The Milky Way Galaxy: A Spiral in Space 576
- 24 Galaxies: Building Blocks of the Universe 606
- 25 Galaxies and Dark Matter: The Large-Scale Structure of the Cosmos 638
- 26 Cosmology: The Big Bang and the Fate of the Universe 666
- 27 The Early Universe: Toward the Beginning of Time 688
- 28 Life in the Universe: Are We Alone? 714

# Contents

About the Authors **xxi**

Preface **xxiii**

## Part One: Astronomy and the Universe **2**

### 1 Charting the Heavens

---

The Foundations of Astronomy **4**

- 1.1 Our Place in Space **6**
- 1.2 Scientific Theory and the Scientific Method **8**
- 1.3 The “Obvious” View **10**
- 1.4 Earth’s Orbital Motion **13**  
*More Precisely 1-1 Angular Measure 14*
- 1.5 The Motion of the Moon **18**
- 1.6 The Measurement of Distance **24**  
*More Precisely 1-2 Measuring Distances with Geometry 28*  
Chapter Review **29**

### 2 The Copernican Revolution

---

The Birth of Modern Science **32**

- 2.1 Ancient Astronomy **34**
- 2.2 The Geocentric Universe **36**
- 2.3 The Heliocentric Model of the Solar System **39**  
*Discovery 2-1 Foundations of the Copernican Revolution 40*
- 2.4 The Birth of Modern Astronomy **41**
- 2.5 The Laws of Planetary Motion **44**  
*More Precisely 2-1 Some Properties of Planetary Orbits 46*
- 2.6 The Dimensions of the Solar System **47**
- 2.7 Newton’s Laws **49**
- 2.8 Newtonian Mechanics **52**  
*More Precisely 2-2 Weighing the Sun 54*  
Chapter Review **56**

### 3 Radiation

---

Information from the Cosmos **58**

- 3.1 Information from the Skies **60**
- 3.2 Waves in What? **63**
- 3.3 Electromagnetic Spectrum **65**  
*Discovery 3-1 The Wave Nature of Radiation 67*
- 3.4 Thermal Radiation **68**  
*More Precisely 3-1 The Kelvin Temperature Scale 69*  
*More Precisely 3-2 More About the Radiation Laws 72*
- 3.5 The Doppler Effect **73**  
*More Precisely 3-3 Measuring Velocities with the Doppler Effect 75*  
Chapter Review **76**

## 4 Spectroscopy

---

### The Inner Workings of Atoms 78

- 4.1 Spectral Lines 80
- 4.2 Atoms and Radiation 84
  - More Precisely 4-1 The Hydrogen Atom 86
- 4.3 The Formation of Spectral Lines 87
  - Discovery 4-1 The Photoelectric Effect 88
- 4.4 Molecules 91
- 4.5 Spectral-Line Analysis 92
  - Chapter Review 95

## 5 Telescopes

---

### The Tools of Astronomy 98

- 5.1 Optical Telescopes 100
- 5.2 Telescope Size 105
- 5.3 Images and Detectors 109
- 5.4 High-Resolution Astronomy 111
- 5.5 Radio Astronomy 114
- 5.6 Interferometry 118
- 5.7 Space-Based Astronomy 121
  - Discovery 5-1 The ALMA Array 124
- 5.8 Full-Spectrum Coverage 128
  - Chapter Review 129

## Part Two: Our Planetary System 132

## 6 The Solar System

---

### Comparative Planetology and Formation Models 134

- 6.1 An Inventory of the Solar System 136
- 6.2 Measuring the Planets 138
- 6.3 The Overall Layout of the Solar System 139
- 6.4 Terrestrial and Jovian Planets 140
  - Discovery 6-1 Gravitational “Slingshots” 142
- 6.5 Interplanetary Matter 143
- 6.6 How Did the Solar System Form? 144
  - Discovery 6-2 Spacecraft Exploration of the Solar System 146
  - More Precisely 6-1 Angular Momentum 149
- 6.7 Jovian Planets and Planetary Debris 152
  - Chapter Review 156

## 7 Earth

---

### Our Home in Space 160

- 7.1 Overall Structure of Planet Earth 162
- 7.2 Earth’s Atmosphere 162
  - More Precisely 7-1 Why Is the Sky Blue? 165



[Discovery 7-1 The Greenhouse Effect and Global Warming](#) 167

- 7.3 Earth's Interior 168
  - [More Precisely 7-2 Radioactive Dating](#) 172
- 7.4 Surface Activity 173
- 7.5 Earth's Magnetosphere 180
- 7.6 The Tides 182
- Chapter Review 185

## 8 The Moon and Mercury

---

Scorched and Battered Worlds 188

- 8.1 Orbital Properties 190
- 8.2 Physical Properties 191
- 8.3 Surface Features on the Moon and Mercury 192
- 8.4 Rotation Rates 195
  - [More Precisely 8-1 Why Air Sticks Around](#) 196
  - [Discovery 8-1 Lunar Exploration](#) 198
- 8.5 Lunar Cratering and Surface Composition 201
- 8.6 The Surface of Mercury 206
- 8.7 Interiors 208
- 8.8 The Origin of the Moon 210
- 8.9 Evolutionary History of the Moon and Mercury 211
- Chapter Review 213

## 9 Venus

---

Earth's Sister Planet 216

- 9.1 Orbital Properties 218
- 9.2 Physical Properties 219
- 9.3 Long-Distance Observations of Venus 220
- 9.4 The Surface of Venus 221
- 9.5 The Atmosphere of Venus 228
- 9.6 Venus's Magnetic Field and Internal Structure 232
- Chapter Review 233

## 10 Mars

---

A Near Miss for Life? 236

- 10.1 Orbital Properties 238
- 10.2 Physical Properties 239
- 10.3 Long-Distance Observations of Mars 239
- 10.4 The Martian Surface 240
- 10.5 Water on Mars 244
  - [Discovery 10-1 Life on Mars?](#) 250
- 10.6 The Martian Atmosphere 256
- 10.7 Martian Internal Structure 259
- 10.8 The Moons of Mars 260
- Chapter Review 261

## 11 Jupiter

---

Giant of the Solar System 264

11.1 Orbital and Physical Properties 266

11.2 Jupiter Atmosphere 268

[Discovery 11-1 A Cometary Impact](#) 274

11.3 Internal Structure 274

[Discovery 11-2 Almost a Star?](#) 276

11.4 Jupiter's Magnetosphere 277

11.5 The Moons of Jupiter 279

11.6 Jupiter's Ring 287

Chapter Review 287

## 12 Saturn

---

Spectacular Rings and Mysterious Moons 290

12.1 Orbital and Physical Properties 292

12.2 Saturn's Atmosphere 293

12.3 Saturn's Interior and Magnetosphere 296

12.4 Saturn's Spectacular Ring System 298

12.5 The Moons of Saturn 304

[Discovery 12-1 Dancing Among Saturn's Moons](#) 306

Chapter Review 315

## 13 Uranus and Neptune

---

The Outer Worlds of the Solar System 318

13.1 The Discoveries of Uranus and Neptune 320

13.2 Orbital and Physical Properties 322

13.3 The Atmospheres of Uranus and Neptune 324

13.4 Magnetospheres and Internal Structure 326

13.5 The Moon Systems of Uranus and Neptune 328

13.6 The Rings of the Outermost Jovian Planets 332

Chapter Review 335

## 14 Solar System Debris

---

Keys to Our Origin 338

14.1 Asteroids 340

14.2 Comets 345

[Discovery 14-1 What Killed the Dinosaurs?](#) 350

14.3 Beyond Neptune 353

14.4 Meteoroids 358

Chapter Review 363

## 15 EXOPLANETS

---

Planetary Systems Beyond Our Own 366

15.1 Modeling Planet Formation 368

15.2 Solar System Regularities and Irregularities 369



- 15.3 Searching for Extrasolar Planets 370
- 15.4 Exoplanet Properties 373
  - Discovery 15-1 The Closest Exoplanet 376
- 15.5 Is Our Solar System Unusual? 379
  - Chapter Review 383

## Part Three: Stars and Stellar Evolution 386

### 16 The Sun

---

- Our Parent Star 388
- 16.1 Physical Properties of the Sun 390
- 16.2 The Solar Interior 392
  - Discovery 16-1 Eavesdropping on the Sun 395
- 16.3 The Sun's Atmosphere 397
- 16.4 Solar Magnetism 400
- 16.5 The Active Sun 405
  - Discovery 16-2 Solar–Terrestrial Relations 409
- 16.6 The Heart of the Sun 410
  - More Precisely 16-1 Fundamental Forces 413
- 16.7 Observations of Solar Neutrinos 414
  - More Precisely 16-2 Energy Generation in the Proton–Proton Chain 416
  - Chapter Review 417

### 17 The Stars

---

- Giants, Dwarfs, and the Main Sequence 420
- 17.1 The Solar Neighborhood 422
- 17.2 Luminosity and Apparent Brightness 425
- 17.3 Stellar Temperatures 428
  - More Precisely 17-1 More on the Magnitude Scale 430
- 17.4 Stellar Sizes 432
  - More Precisely 17-2 Estimating Stellar Radii 433
- 17.5 The Hertzsprung–Russell Diagram 434
- 17.6 Extending the Cosmic Distance Scale 437
- 17.7 Stellar Masses 440
  - More Precisely 17-3 Measuring Stellar Masses in Binary Stars 443
- 17.8 Mass and Other Stellar Properties 442
  - Chapter Review 445

### 18 The Interstellar Medium

---

- Gas and Dust Among The Stars 448
- 18.1 Interstellar Matter 450
- 18.2 Emission Nebulae 453
- 18.3 Dark Dust Clouds 459
- 18.4 21-Centimeter Radiation 462
- 18.5 Interstellar Molecules 463
  - Chapter Review 465

## 19 Star Formation

---

- A Traumatic Birth 468
- 19.1 Star-Forming Regions 470
  - More Precisely 19-1 Competition in Star Formation 471
- 19.2 The Formation of Stars Like the Sun 472
- 19.3 Stars of Other Masses 477
- 19.4 Observations of Cloud Fragments and Protostars 478
  - Discovery 19-1 Observations of Brown Dwarfs 479
- 19.5 Shock Waves and Star Formation 484
- 19.6 Star Clusters 486
  - Discovery 19-2 Eta Carinae 490
  - Chapter Review 491

## 20 Stellar Evolution

---

- The Life and Death of a Star 494
- 20.1 Leaving the Main Sequence 496
- 20.2 Evolution of a Sun-Like Star 496
- 20.3 The Death of a Low-Mass Star 502
  - Discovery 20-1 Learning Astronomy from History 508
- 20.4 Evolution of Stars More Massive than the Sun 509
  - Discovery 20-2 Mass Loss from Giant Stars 511
- 20.5 Observing Stellar Evolution in Star Clusters 512
- 20.6 Stellar Evolution in Binary Systems 515
  - Chapter Review 517

## 21 Stellar Explosions

---

- Novae, Supernovae, and the Formation of the Elements 520
- 21.1 Life after Death for White Dwarfs 522
- 21.2 The End of a High-Mass Star 524
- 21.3 Supernovae 526
- 21.4 The Formation of the Elements 530
  - Discovery 21-1 Supernova 1987A 532
- 21.5 The Cycle of Stellar Evolution 538
  - Chapter Review 539

## 22 Neutron Stars and Black Holes

---

- Strange States of Matter 542
- 22.1 Neutron Stars 544
- 22.2 Pulsars 545
- 22.3 Neutron-Star Binaries 548
- 22.4 Gamma-Ray Bursts 552
- 22.5 Black Holes 555
- 22.6 Einstein's Theories of Relativity 557
  - Discovery 22-1 Special Relativity 559
- 22.7 Space Travel Near Black Holes 561



- 22.8 Observational Evidence for Black Holes 564
  - More Precisely 22-1 Tests of General Relativity 566
  - Discovery 22-2 Gravity Waves: A New Window on the Universe 568
- Chapter Review 571

## Part Four: Galaxies and Cosmology 574

### 23 The Milky Way Galaxy

---

- A Spiral in Space 576
- 23.1 Our Parent Galaxy 578
- 23.2 Measuring the Milky Way 579
  - Discovery 23-1 Early “Computers” 584
- 23.3 Galactic Structure 586
- 23.4 The Formation of the Milky Way 589
- 23.5 Galactic Spiral Arms 591
  - Discovery 23-2 Density Waves 594
- 23.6 The Mass of the Milky Way Galaxy 595
- 23.7 The Galactic Center 599
- Chapter Review 603

### 24 Galaxies

---

- Building Blocks of the Universe 606
- 24.1 Hubble’s Galaxy Classification 608
- 24.2 The Distribution of Galaxies in Space 615
- 24.3 Hubble’s Law 619
  - More Precisely 24-1 Relativistic Redshifts and Look-Back Time 622
- 24.4 Active Galactic Nuclei 622
- 24.5 The Central Engine of an Active Galaxy 630
- Chapter Review 635

### 25 Galaxies and Dark Matter

---

- The Large-Scale Structure of the Cosmos 638
- 25.1 Dark Matter in the Universe 640
- 25.2 Galaxy Collisions 643
- 25.3 Galaxy Formation and Evolution 645
  - Discovery 25-1 The Sloan Digital Sky Survey 651
- 25.4 Black Holes in Galaxies 652
- 25.5 The Universe on Large Scales 656
- Chapter Review 663

### 26 Cosmology

---

- The Big Bang and the Fate of the Universe 666
- 26.1 The Universe on the Largest Scales 668
- 26.2 The Expanding Universe 670
- 26.3 The Fate of the Cosmos 673

- 26.4 The Geometry of Space 675
  - More Precisely 26-1 Curved Space 677
- 26.5 Will the Universe Expand Forever? 678
- 26.6 Dark Energy and Cosmology 680
  - Discovery 26-1 Einstein and the Cosmological Constant 681
- 26.7 The Cosmic Microwave Background 683
  - Chapter Review 685

## 27 The Early Universe

---

- Toward the Beginning of Time 688
- 27.1 Back to the Big Bang 690
- 27.2 Evolution of the Universe 693
  - More Precisely 27-1 More on Fundamental Forces 694
- 27.3 Formation of Nuclei and Atoms 697
- 27.4 The Inflationary Universe 700
- 27.5 Formation of Structure in the Universe 705
- 27.6 Cosmic Structure and the Microwave Background 707
  - Chapter Review 711

## 28 Life in the Universe

---

- Are We Alone? 714
- 28.1 Cosmic Evolution 716
  - Discovery 28-1 The Virus 717
- 28.2 Life in the Solar System 722
- 28.3 Intelligent Life in the Galaxy 724
- 28.4 The Search for Extraterrestrial Intelligence 729
  - Chapter Review 733

## Appendices

- Appendix 1 Scientific Notation **A-1**
- Appendix 2 Astronomical Measurement **A-2**
- Appendix 3 Tables **A-3**
- Glossary **G-1**
- Answers to Check Questions **AK-1**
- Answers to Self-Test Questions **AK-6**
- Photo Credits/Text Permissions **C-1**
- Index **I-1**
- Star Charts **S-1**

*This page intentionally left blank*

# Online Contents

## PART ONE: ASTRONOMY AND THE UNIVERSE 2

### Chapter 1 Charting the Heavens 4

- INTERACTIVE FIGURE Constellation Orion 10
- INTERACTIVE FIGURE Celestial Sphere 12
- INTERACTIVE FIGURE Northern Sky 12
- ANIMATION/VIDEO Summer Solstice 13
- INTERACTIVE FIGURE The Zodiac 15
- ANIMATION/VIDEO Winter Solstice 16
- ANIMATION/VIDEO The Earth's Seasons 16
- INTERACTIVE FIGURE Seasons 16
- ANIMATION/VIDEO The Equinoxes 17
- INTERACTIVE FIGURE Precession 18
- SELF-GUIDED TUTORIAL Phases of the Moon 19
- NARRATED FIGURE Lunar Phases 19
- INTERACTIVE FIGURE Sidereal Month 20
- INTERACTIVE FIGURE Lunar Eclipse 20
- INTERACTIVE FIGURE Types of Solar Eclipse 21
- ANIMATION/VIDEO Solar Eclipse in Indiana 21

### Chapter 2 The Copernican Revolution 32

- ANIMATION/VIDEO Retrograde Motion of Mars 37
- INTERACTIVE FIGURE Geocentric Model 38
- ANIMATION/VIDEO Geocentric Solar System 39
- ANIMATION/VIDEO Heliocentric Solar System 39
- INTERACTIVE FIGURE Retrograde Motion 41
- INTERACTIVE FIGURE Venus Phases 43
- INTERACTIVE FIGURE Ellipse 45
- INTERACTIVE FIGURE Kepler's Second Law 46

- ANIMATION/VIDEO Earth Captures a Temporary Moon 52
- INTERACTIVE FIGURE Orbits 53
- INTERACTIVE FIGURE Escape Speed 55

### Chapter 3 Radiation 58

- INTERACTIVE FIGURE Water Wave 61
- INTERACTIVE FIGURE Wave Properties 61
- ANIMATION/VIDEO Solar Eclipse Viewed in X-rays 66
- ANIMATION/VIDEO Multispectral View of Orion Nebula 66
- ANIMATION/VIDEO Earth Aurora in X-rays 66
- NARRATED INTERACTIVE FIGURE Electromagnetic Spectrum 66
- ANIMATION/VIDEO Fresnel Diffraction 67
- SELF-GUIDED TUTORIAL Continuous Spectra and Blackbody Radiation 70
- INTERACTIVE FIGURE Astronomical Thermometer 71
- SELF-GUIDED TUTORIAL Doppler Effect 73
- INTERACTIVE FIGURE Doppler Effect 74

### Chapter 4 Spectroscopy 78

- INTERACTIVE FIGURE Continuous and Emission Spectra 81
- SELF-GUIDED TUTORIAL Emission Spectra 82
- SELF-GUIDED TUTORIAL Absorption Spectra 82
- INTERACTIVE FIGURE Absorption Spectrum 82
- INTERACTIVE FIGURE Sodium Spectrum 83
- ANIMATION/VIDEO Classical Hydrogen Atom I 86
- ANIMATION/VIDEO Classical Hydrogen Atom II 86
- INTERACTIVE FIGURE Atomic Excitation 89

- ANIMATION/VIDEO Multispectral Views of the Orion Nebula 91
- INTERACTIVE FIGURE Doppler Shift 93

### Chapter 5 Telescopes 98

- SELF-GUIDED TUTORIAL The Optics of a Simple Lens 101
- SELF-GUIDED TUTORIAL Chromatic Aberration 102
- SELF-GUIDED TUTORIAL Reflecting Telescopes 103
- ANIMATION/VIDEO Hubble Space Telescope in Orbit 104
- ANIMATION/VIDEO Gemini Control Room 107
- INTERACTIVE FIGURE Resolving Power 108
- INTERACTIVE FIGURE Constructing an Image from Colored Filters 110
- ANIMATION/VIDEO Adaptive Optics 114
- ANIMATION/VIDEO Deployment of the James Webb Space Telescope 122
- ANIMATION/VIDEO Chandra Light and Data Paths 126
- NARRATED FIGURE Multiple Wavelengths 128

## PART TWO: OUR PLANETARY SYSTEM 132

### Chapter 6 The Solar System 134

- ANIMATION/VIDEO An Astronomical Ruler 139
- ANIMATION/VIDEO The Gas Giants 141
- ANIMATION/VIDEO Size and Scale of the Terrestrial Planets I & II 141
- INTERACTIVE FIGURE Gravitational Assist 142
- INTERACTIVE FIGURE Nebular Contraction 145
- ANIMATION/VIDEO Solar System Formation 152



- ANIMATION/VIDEO Protoplanetary Disk Destruction **153**
- ANIMATION/VIDEO Protoplanetary Disks in the Orion Nebula **153**
- INTERACTIVE FIGURE Jovian Condensation **153**

## Chapter 7 Earth 160

- ANIMATION/VIDEO Earth as Seen by *Galileo* **164**
- ANIMATION/VIDEO NEAR Earth Swingby **164**
- ANIMATION/VIDEO Ozone Hole Over the Antarctic **166**
- SELF-GUIDED TUTORIAL The Greenhouse Effect **166**
- INTERACTIVE FIGURE Greenhouse Effect **166**
- INTERACTIVE FIGURE Plate Drift **178**
- ANIMATION/VIDEO Northern and Southern Lights **182**
- INTERACTIVE FIGURE Solar and Lunar Tides **184**

## Chapter 8 The Moon and Mercury 188

- ANIMATION/VIDEO Transit of Mercury **191**
- ANIMATION/VIDEO Full Rotation of Moon **193**
- ANIMATION/VIDEO Lunar Flyby **193**
- INTERACTIVE FIGURE The Moon's Synchronous Rotation **195**
- ANIMATION/VIDEO First Step on the Moon **198**
- ANIMATION/VIDEO *Ranger* Spacecraft Descent to Moon **199**
- INTERACTIVE FIGURE Mercury's Rotation **200**
- INTERACTIVE FIGURE Meteoroid Impact **201**
- ANIMATION/VIDEO Protoplanetary Collision **209**
- NARRATED FIGURE Moon Formation **211**

## Chapter 9 Venus 216

- NARRATED FIGURE Venus's Brightness **218**
- SELF-GUIDED TUTORIAL Super-spaceship—Voyage to Venus **219**

- ANIMATION/VIDEO The Rotation of Venus **219**
- ANIMATION/VIDEO Transit of Venus **220**
- ANIMATION/VIDEO Topography of Venus **222**
- ANIMATION/VIDEO Flight Over Alpha Regio **225**
- ANIMATION/VIDEO Flight Over Sif Mons Volcano **225**

## Chapter 10 Mars 236

- ANIMATION/VIDEO *Hubble* View of Mars **240**
- INTERACTIVE FIGURE Mars Map **241**
- ANIMATION/VIDEO Flight Over Tharsis **242**
- ANIMATION/VIDEO Flight Over Mariner Valley **243**
- SELF-GUIDED TUTORIAL Comparative Planetology: Mars **245**
- ANIMATION/VIDEO Meteorites Ejected from Mars **251**
- ANIMATION/VIDEO *Hubble* View of Mars Polar Cap **252**
- ANIMATION/VIDEO Flight Over *Opportunity* at Gustav Crater **254**
- ANIMATION/VIDEO *Mars Rover* Landing **254**
- ANIMATION/VIDEO Flight Over Columbia Hills **254**
- SELF-GUIDED TUTORIAL Atmospheric Lifetimes **257**
- ANIMATION/VIDEO Martian Moons: Phobos & Deimos **260**

## Chapter 11 Jupiter 264

- SELF-GUIDED TUTORIAL Jupiter—Differential Rotation **267**
- ANIMATION/VIDEO Jupiter's Rotation **268**
- INTERACTIVE FIGURE Rotational Flattening **268**
- INTERACTIVE FIGURE Zonal Flow **270**
- ANIMATION/VIDEO *Galileo* Mission to Jupiter **272**
- ANIMATION/VIDEO Comet Impact with Jupiter **275**
- ANIMATION/VIDEO The Gas Giants II **276**

- ANIMATION/VIDEO Galilean Moons Transit Jupiter **279**
- INTERACTIVE FIGURE Galilean Moons **279**
- ANIMATION/VIDEO Io Cutaway **280**
- INTERACTIVE FIGURE Volcanoes on Io **282**
- ANIMATION/VIDEO *Galileo's* View of Europa **284**
- ANIMATION/VIDEO *Galileo's* View of Ganymede **285**
- ANIMATION/VIDEO Jupiter Icy Moons Orbiter Mission **286**

## Chapter 12 Saturn 290

- ANIMATION/VIDEO Saturn Cloud Rotation **295**
- ANIMATION/VIDEO Saturn Ring Plane Crossing **298**
- NARRATED FIGURE Roche Limit **299**
- INTERACTIVE FIGURE Saturn's Rings, Up Close **301**
- ANIMATION/VIDEO *Voyager* Ring Spokes **302**
- ANIMATION/VIDEO Saturn Satellite Transit **303**
- ANIMATION/VIDEO *Huygens* Landing on Titan **309**

## Chapter 13 Uranus and Neptune 318

- ANIMATION/VIDEO Neptune's Dark Spot **321**
- ANIMATION/VIDEO Rotation of Uranus **324**
- ANIMATION/VIDEO Rotation of Neptune **325**
- INTERACTIVE FIGURE Jovian Magnetic Fields **327**
- INTERACTIVE FIGURE Jovian Interiors **327**
- ANIMATION/VIDEO Geysers on Triton **331**

## Chapter 14 Solar System Debris 338

- NARRATED INTERACTIVE Inner Solar System **340**
- ANIMATION/VIDEO Orbiting Eros **341**
- ANIMATION/VIDEO NEAR Descent **341**

- ANIMATION/VIDEO NEAR Landing **341**
- ANIMATION/VIDEO Sun Grazing Comets **346**
- ANIMATION/VIDEO Anatomy of a Comet Part 1 **346**
- ANIMATION/VIDEO Comet Hale-Bopp Nucleus Animation **347**
- ANIMATION/VIDEO Anatomy of a Comet Part 2 **347**
- ANIMATION/VIDEO Deep Impact Simulation **352**
- INTERACTIVE FIGURE Comet Wild-2 **352**
- INTERACTIVE FIGURE Deep Impact **352**
- ANIMATION/VIDEO Orbits of Neptune and Pluto **354**
- INTERACTIVE FIGURE Neptune and Pluto **354**
- ANIMATION/VIDEO Hubble's View of Pluto **355**
- ANIMATION/VIDEO Historical Observations of Pluto **355**
- ANIMATION/VIDEO Asteroid/Comet Breakup **359**
- ANIMATION/VIDEO Delta Capricornid Meteor Near Orion **359**
- ANIMATION/VIDEO Daytime Passage of Meteor Fireball **359**

## Chapter 15 Exoplanets **366**

- ANIMATION/VIDEO Protoplanetary Disks in the Orion Nebula **369**
- ANIMATION/VIDEO Protoplanetary Disk Destruction **369**
- ANIMATION/VIDEO Evolution of Protoplanetary Disk **370**
- ANIMATION/VIDEO The Formation of the Solar System **370**
- INTERACTIVE FIGURE Planets Revealed **372**
- INTERACTIVE FIGURE An Extrasolar Transit **373**
- ANIMATION/VIDEO Hot Jupiter Extrasolar Planet Evaporating **380**
- ANIMATION/VIDEO Survey for Transiting Extrasolar Planets **380**
- INTERACTIVE FIGURE Jupiter-like Planet? **380**
- INTERACTIVE FIGURE Sinking Planet **380**

## PART THREE: STARS AND STELLAR EVOLUTION **386**

### Chapter 16 The Sun **388**

- SELF-GUIDED TUTORIAL Super-Spaceship—Voyage to the Sun **390**
- NARRATED FIGURE Stellar Balance **392**
- ANIMATION/VIDEO Solar Granulation **396**
- ANIMATION/VIDEO Solar Chromosphere **399**
- ANIMATION/VIDEO Sunspot **401**
- ANIMATION/VIDEO Solar Flare **407**
- ANIMATION/VIDEO Coronal Mass Ejections **407**
- INTERACTIVE FIGURE Solar Fusion **412**

### Chapter 17 The Stars **420**

- SELF-GUIDED TUTORIAL Stellar Parallax **422**
- ANIMATION/VIDEO The Inverse-Square Law **425**
- NARRATED FIGURE Inverse-Square Law **425**
- INTERACTIVE FIGURE Apparent Magnitude **427**
- INTERACTIVE FIGURE H-R Diagram of Well-Known Stars **435**
- INTERACTIVE FIGURE H-R Diagram of Nearby Stars **435**
- ANIMATION/VIDEO White Dwarfs in Globular Cluster **436**
- SELF-GUIDED TUTORIAL Hertzsprung–Russell Diagram **436**
- SELF-GUIDED TUTORIAL Binary Stars—Radial Velocity Curves **440**
- INTERACTIVE FIGURE Spectroscopic Binary **440**
- SELF-GUIDED TUTORIAL Eclipsing Binary Stars—Light Curves **441**
- INTERACTIVE FIGURE Eclipsing Binary **441**

### Chapter 18 The Interstellar Medium **448**

- ANIMATION/VIDEO Pillars Behind the Dust **451**
- ANIMATION/VIDEO Infrared View of Nebulae **451**
- NARRATED FIGURE Reddening **451**

- ANIMATION/VIDEO Orion Nebula Mosaic **453**
- ANIMATION/VIDEO The Tarantula Nebula **454**
- INTERACTIVE FIGURE Trifid Nebula **454**
- ANIMATION/VIDEO Gaseous Pillars of Star Birth, the Eagle Nebula **456**
- ANIMATION/VIDEO Horsehead Nebula **461**

### Chapter 19 Star Formation **468**

- ANIMATION/VIDEO Stellar Birth **475**
- INTERACTIVE FIGURE Newborn Star on the H-R Diagram **476**
- ANIMATION/VIDEO Binary Brown Dwarfs **479**
- INTERACTIVE FIGURE Orion Nebula, Up Close **481**
- ANIMATION/VIDEO Protostars **482**
- ANIMATION/VIDEO Herbig–Haro Objects **483**
- ANIMATION/VIDEO Bipolar Outflow **483**
- ANIMATION/VIDEO Triggered Star Formation **485**
- INTERACTIVE FIGURE Protostellar Collisions **489**
- INTERACTIVE FIGURE Young Stars in Orion **489**
- ANIMATION/VIDEO Carina Nebula **490**

### Chapter 20 Stellar Evolution **494**

- ANIMATION/VIDEO H-R Diagram Tracks Stellar Evolution **498**
- ANIMATION/VIDEO Red Giant Evolution **499**
- ANIMATION/VIDEO Death of the Sun Part 1 **502**
- INTERACTIVE FIGURE G-Type Star Evolution **502**
- ANIMATION/VIDEO Death of the Sun Part II **503**
- ANIMATION/VIDEO Helix Nebula Animation **504**
- ANIMATION/VIDEO Helix Nebula **504**
- ANIMATION/VIDEO Bi-Polar Planetary Nebula **504**
- INTERACTIVE FIGURE White Dwarf on the H-R Diagram **505**

- ANIMATION/VIDEO Helix Nebula White Dwarf **506**
- ANIMATION/VIDEO White Dwarf Cooling Sequence **506**
- SELF-GUIDED TUTORIAL Evolution of a 1-Solar-Mass Star **507**
- ANIMATION/VIDEO Light Echo **511**

## Chapter 21 Stellar Explosions **520**

- ANIMATION/VIDEO Recurrent Nova **523**
- INTERACTIVE FIGURE Heavy-Element Fusion **524**
- ANIMATION/VIDEO Structure of Supernova **526**
- ANIMATION/VIDEO Supernova Explosion **526**
- INTERACTIVE FIGURE Crab Supernova Remnant **529**
- ANIMATION/VIDEO Supernova Remnant in Cassiopeia **530**
- INTERACTIVE FIGURE Vela Supernova Remnant **530**
- ANIMATION/VIDEO Composition and Structure of the Ring Around Supernova 1987A **533**
- ANIMATION/VIDEO Shockwaves Hit the Ring of Supernova 1987A **533**
- INTERACTIVE FIGURE Stellar Recycling **538**

## Chapter 22 Neutron Stars and Black Holes **542**

- NARRATED FIGURE Pulsar Model **546**
- ANIMATION/VIDEO Pulsar in Crab Nebula **547**
- ANIMATION/VIDEO X-ray Binary Star **549**
- ANIMATION/VIDEO Colliding Binary Neutron Stars **554**
- SELF-GUIDED TUTORIAL Escape Speed and Black Hole Event Horizons **557**
- INTERACTIVE FIGURE Curved Space **560**
- ANIMATION/VIDEO Energy Released from a Black Hole? **563**

- INTERACTIVE FIGURE Gravitational Redshift **563**
- ANIMATION/VIDEO Black Hole and Companion Star **565**
- ANIMATION/VIDEO Black Hole Devours Neutron Star **565**
- INTERACTIVE FIGURE Stellar Black Hole **567**
- ANIMATION/VIDEO Supermassive Black Hole, Black Hole in the Center of M32 **569**
- ANIMATION/VIDEO Black Hole Accretion Disk and Jets **569**

## PART FOUR: GALAXIES AND COSMOLOGY **574**

### Chapter 23 The Milky Way Galaxy **576**

- ANIMATION/VIDEO Cepheid Variable Star in Distant Galaxy **581**
- NARRATED FIGURE Globular Cluster Distribution **585**
- INTERACTIVE FIGURE Stellar Populations in Our Galaxy **585**
- INTERACTIVE FIGURE Infrared View of the Milky Way **587**
- INTERACTIVE FIGURE Milky Way Spiral Structure **592**
- INTERACTIVE FIGURE Differential Galactic Rotation **593**
- INTERACTIVE FIGURE Spiral Density Waves **593**
- ANIMATION/VIDEO Rotating Globular Cluster **597**
- SELF-GUIDED TUTORIAL Gravitational Lensing **598**
- INTERACTIVE FIGURE Galactic Center **599**
- ANIMATION/VIDEO X-ray View of Galactic Core **600**
- ANIMATION/VIDEO Black Hole in the Center of the Milky Way? **601**

### Chapter 24 Galaxies **606**

- NARRATED FIGURE Galaxy Rotation **616**
- INTERACTIVE FIGURE Spacetime Diagram for an Extragalactic Supernova **623**

- ANIMATION/VIDEO Active Galaxy **624**
- ANIMATION/VIDEO M87 Jet **628**
- ANIMATION/VIDEO Eruption of a Supermassive Black Hole **628**
- INTERACTIVE FIGURE M87 Jet **628**

## Chapter 25 Galaxies and Dark Matter **638**

- INTERACTIVE FIGURE Rotation Curve for a Merry-Go-Round **640**
- INTERACTIVE FIGURE Galaxy Rotation Curves **640**
- ANIMATION/VIDEO Dark Matter **641**
- ANIMATION/VIDEO Galaxy Collision **644**
- ANIMATION/VIDEO Starburst Galaxy **644**
- INTERACTIVE FIGURE Starburst Galaxy **644**
- ANIMATION/VIDEO The Evolution of Galaxies **647**
- ANIMATION/VIDEO Hubble Deep Field Zoom I **647**
- ANIMATION/VIDEO Hubble Deep Field Zoom II **647**
- ANIMATION/VIDEO Galaxy Merger **650**
- NARRATED FIGURE Galaxy Evolution **655**
- ANIMATION/VIDEO Cluster Merger **656**
- INTERACTIVE FIGURE Absorption Line “Forest” **659**
- ANIMATION/VIDEO How a Gravitational Lens Works **660**
- INTERACTIVE FIGURE Gravitational Lens **660**
- ANIMATION/VIDEO Simulation of Gravitational Lens in Space **661**
- ANIMATION/VIDEO Dark Matter Collision **662**
- ANIMATION/VIDEO Bullet Cluster Collision **662**
- INTERACTIVE FIGURE Cluster Collision **662**

## Chapter 26 Cosmology 666

- ANIMATION/VIDEO Cosmic Structure **668**
- INTERACTIVE FIGURE The Expanding Raisin Cake (Universe) **672**
- INTERACTIVE FIGURE Receding Galaxies **672**
- INTERACTIVE FIGURE Cosmological Redshift **673**

## Chapter 27 The Early Universe 688

- ANIMATION/VIDEO The First Stars Reionize the Universe **696**
- INTERACTIVE FIGURE Creation of the Cosmic Microwave Background **699**
- ANIMATION/VIDEO Cosmic Structure **706**
- NARRATED FIGURE Structure Formation **706**
- INTERACTIVE FIGURE Early Structure **708**

## Chapter 28 Life in the Universe 714

- ANIMATION/VIDEO Icy Organics in Planet-Forming Disc **719**
- ANIMATION/VIDEO Earth's Biosphere in Action: Plankton Bloom **722**
- NARRATED FIGURE Drake Equation **725**
- ANIMATION/VIDEO Asteroid Impacting the Earth **728**

*This page intentionally left blank*

# About the Authors



**Eric Chaisson**

Eric holds a doctorate in astrophysics from Harvard University, where he spent 10 years on the faculty of Arts and Sciences. For more than two decades thereafter, he served on the senior science staff at the Space Telescope Science Institute and held various professorships at Johns Hopkins and Tufts universities. He is now back at Harvard, where he teaches and conducts research at the Harvard-Smithsonian Center for Astrophysics. Eric has written 12 books on astronomy and has published nearly 200 scientific papers in professional journals.



**Steve McMillan**

Steve holds a bachelor's and master's degree in mathematics from Cambridge University and a doctorate in astronomy from Harvard University. He held postdoctoral positions at the University of Illinois and Northwestern University, where he continued his research in theoretical astrophysics, star clusters, and high-performance computing. Steve is currently Distinguished Professor of Physics at Drexel University and a frequent visiting researcher at Princeton's Institute for Advanced Study and Leiden University. He has published more than 100 articles and scientific papers in professional journals.

*This page intentionally left blank*

# Preface

Astronomy is a science that thrives on new discoveries. Fueled by new technologies and novel theoretical insights, the study of the cosmos continues to change our understanding of the universe. We are pleased to have the opportunity to present in this book a representative sample of the known facts, evolving ideas, and frontier discoveries in astronomy today.

*Astronomy Today* has been written for students who have taken no previous college science courses and who will likely not major in physics or astronomy. It is intended for use in a one- or two-semester, nontechnical astronomy course. We present a broad view of astronomy, straightforwardly descriptive and without complex mathematics. The absence of sophisticated mathematics, however, in no way prevents discussion of important concepts. Rather, we rely on qualitative reasoning as well as analogies with objects and phenomena familiar to the student to explain the complexities of the subject without oversimplification. We have tried to communicate the excitement we feel about astronomy and to awaken students to the marvelous universe around us.

We are very gratified that the first seven editions of this text have been so well received by many in the astronomy education community. In using those earlier texts, many teachers and students have given us helpful feedback and constructive criticisms. From these, we have learned to communicate better both the fundamentals and the excitement of astronomy. Many improvements inspired by these comments have been incorporated into this new edition.

## Focus of the Eighth Edition

From the first edition, we have tried to meet the challenge of writing a book that is both accurate and approachable. To the student, astronomy sometimes seems like a long list of unfamiliar terms to be memorized and repeated. Many new terms and concepts will be introduced in this course, but we hope students will also learn and remember how science is done, how the universe works, and how things are connected. In the eighth edition, we have taken particular care to show how astronomers know what they know, and to highlight both the scientific principles underlying their work and the process used in discovery.

## New and Revised Material

Astronomy is a rapidly evolving field and, in the three years since the publication of the seventh edition of *Astronomy Today*, has seen many new discoveries covering the entire

spectrum of astronomical research. Almost every chapter in the eighth edition has been substantially updated with new information. Several chapters have also seen significant reorganization in order to streamline the overall presentation, strengthen our focus on the process of science, and reflect new understanding and emphases in contemporary astronomy.

In addition to updates throughout the text on the numbers and properties of the many astronomical objects, the many substantive changes include the following:

- A new *Discovery* box in Chapter 5 on the *ALMA* interferometric array.
- Significant revision in Chapter 5 of the discussion of infrared telescopes, including new coverage of *Herschel* and introduction of the *James Webb Space Telescope*.
- A new two-page box in Chapter 6 on planetary exploration.
- Incorporation and reorganization of the entire “standard” theory of solar system formation into Chapter 6, laying the groundwork for interpreting the planetary data presented in Part 2 and allowing Chapter 15 to focus on solar system details, irregularities, and exoplanets.
- Updated discussion in *Discovery 8-1* of *Chang’e*, *GRAIL*, and other recent lunar missions; new discussion of the *Prospector*, *LRO*, and *LCROSS* missions, with updated coverage of the search for lunar ice.
- Updated coverage in Chapter 8 of the lunar core and interior based on the latest *GRAIL* results.
- Updated discussion in Chapter 8 of surface features on Mercury, following the *Messenger* mission.
- Updated discussion in Chapter 8 of Mercury’s inner and outer core and magnetic field and formation, in light of new *Messenger* data.
- Updated discussion in Chapter 9 of *Venus Express* findings and status.
- Updated discussion in Chapter 10 of the collision hypothesis as the origin of the northern Martian lowlands.
- Reorganized and updated discussion in Chapter 10 of liquid water on the Martian surface.
- Updated discussion in Chapter 10 on the *Spirit*, *Opportunity*, and *Phoenix* landers; new material on the *Curiosity* lander and its findings.
- Revised discussion in Chapter 10 of the origin of the Martian moons.
- Updated coverage of cometary impacts in *Discovery 11-1*, indicating that such impacts are commonplace in the solar system.




## The Illustration Program

Visualization plays an important role in both the teaching and the practice of astronomy, and we continue to place strong emphasis on this aspect of our book. We have tried to combine aesthetic beauty with scientific accuracy in the artist's conceptions that adorn the text, and we have sought to present the best and latest imagery of a wide range of cosmic objects. Each illustration has been carefully crafted to enhance student learning; each is pedagogically sound and tied tightly to the nearby discussion of important scientific facts and ideas. This edition contains more than 100 revised figures that show the latest imagery and the results learned from them.


**Compound Art** It is rare that a single image, be it a photograph or an artist's conception, can capture all aspects of a complex subject. Wherever possible, multiple-part figures are used in an attempt to convey the greatest amount of information in the most vivid way:

- Visible images are often presented along with their counterparts captured at other wavelengths.
- Interpretive line drawings are often superimposed on or juxtaposed with real astronomical photographs, helping students to really “see” what the photographs reveal.
- Breakouts—often multiple ones—are used to zoom in from wide-field shots to close-ups so that detailed images can be understood in their larger context.

**Interactive Figures and Photos** Icons throughout the text direct students to dynamic, interactive versions of art and photos on MasteringAstronomy®. Using online applets, students can manipulate factors such as time, wavelength, scale, and perspective to increase their understanding of these figures.



**Interactive FIGURE 11.20 Volcanoes on Io** The main image shows a *Galileo* view of Io, whose surface is kept smooth and brightly colored by constant volcanism, revealed here as dark, circular features. The left inset shows an umbrella-like eruption of one of Io's volcanoes as *Galileo* flew past this fascinating moon in 1997; the plume measures about 150 km high and 300 km across. The right inset shows another volcano, this one face-on, where surface features here are resolved to just a few kilometers. (NASA)



**Interactive FIGURE 7.5 Greenhouse Effect** Sunlight that is not reflected by clouds reaches Earth's surface, warming it up. Infrared radiation reradiated from the surface is partially absorbed by carbon dioxide (and also water vapor, not shown here) in the atmosphere, causing the overall surface temperature to rise.

**MA** **► Narrated Figures (NEW)** Narrated Figures are brief videos that step students through complex figures from the text, expanding students' understanding of fundamental concepts in a presentation that includes narration, enhanced visuals, and one to two embedded questions, followed by short, one- to two-question Mastering activities that are graded. They mirror how an instructor might present a topic in class and can be assigned as homework, self-study, or as part of a pre-lecture program.


**Figure Annotations (REVISED)** The eighth edition incorporates the research-proven technique of strategically placing annotations (which always appear in blue type) within key pieces of art, fostering students' ability to read and interpret complex figures, focus on the most relevant information, and integrate written and visual knowledge.

**Full Spectrum Coverage and Spectrum Icons**

Astronomers exploit the full range of the electromagnetic spectrum to gather information about the cosmos. Throughout this book, images taken at radio, infrared, ultraviolet, X-ray, or gamma-ray wavelengths are used to supplement visible-light images. As it is sometimes difficult (even for a professional) to tell at a glance which images are visible-light photographs and which are false-color images created with other wavelengths, each photo in the text is accompanied by an icon that identifies the wavelength of electromagnetic radiation used to capture the image.

**MA** **► Narrated FIGURE 9.2 Venus's Brightness** Venus appears full when it is at its greatest distance from Earth, on the opposite side of the Sun from us (superior conjunction). As its distance decreases, less and less of its sunlit side becomes visible. When closest to Earth, it lies between us and the Sun (inferior conjunction), so we cannot see the sunlit side of the planet at all. Venus appears brightest when it is about 39° from the Sun. (Compare Figure 2.12.) (Insets: UC/Lick Observatory)

X-ray, or gamma-ray wavelengths are used to supplement visible-light images. As it is sometimes difficult (even for a professional) to tell at a glance which images are visible-light photographs and which are false-color images created with other wavelengths, each photo in the text is accompanied by an icon that identifies the wavelength of electromagnetic radiation used to capture the image.



- Revised discussion in Chapter 12 of storms on Saturn and new moons and features in Saturn's rings.
- Expanded coverage in Chapter 12 of *Cassini Solstice* observations of Titan and Enceladus.
- Updated discussion in Chapter 13 of Uranus's tilted spin axis and new imagery of weather patterns on Uranus and Neptune.
- New coverage in Chapter 14 of the *Dawn* mission to Vesta and Ceres.

- Updated coverage in Chapter 14 of Earth-crossing asteroids and asteroid near misses.
- Updated coverage in Chapter 14 of Pluto's moons and trans-Neptunian objects
- New *Discovery* box in Chapter 15 on the Alpha Centauri planetary system.
- Expanded coverage in Chapter 15 of exoplanet discoveries and properties and the *Kepler* candidates list.
- New discussion in Chapter 15 of Earths and super-Earths in the habitable zones of their parent stars.

- New coverage in Chapter 16 of the *Solar Dynamics Observatory* and its findings.
- Updated discussion in Chapter 19 of star cluster observations and formation.
- Revised discussion in Chapter 22 of gamma-ray bursts and hypernovae.
- Updated coverage in Chapter 23 of activity near the center of the Milky Way Galaxy.
- Significantly updated coverage in Chapter 25 of galaxies, including new discussion of inflow of gas from intergalactic space.
- Expanded discussion of tidal streams in the Milky Way halo.
- Significantly expanded coverage in Chapter 27 of baryon acoustic oscillations in the early universe and their connection to fluctuations in the microwave background.
- Updated discussion in Chapter 28 of the frequency of planetary systems and the numbers of habitable planets per system.
- Added 18 new Narrated Figure notations.
- Added helpful annotations so that now about half of the figures in the text employ this pedagogically useful tool.
- Added distance scales to many figures, helping students gain an understanding of the vastness of the universe.
- Replaced a number of older images for currency and clarity.
- Updated the art throughout the text.
- Added new table of contents for online material (Online Contents), which lists by chapter all the online assets the book delivers: Narrated Figures, Interactive Figures, Animation/Videos, and Self-Guided Tutorials.

## Other Pedagogical Features

As with many other parts of our text, instructors have helped guide us toward what is most helpful for effective student learning. With their assistance, we have revised both our in-chapter and end-of-chapter pedagogical apparatus to increase its utility to students.

**The Big Question** Our Sun will expand as it ages, and it is destined to balloon rapidly into a red giant as it begins running out of fuel in about 5 billion years. A burning question, often asked and then quickly dismissed as being too remote in time is, will the red-giant Sun expand enough to engulf Earth? No one is certain. We do know that the Sun is losing lots of matter, thereby lessening its gravitational pull. Perhaps that will allow Earth to recede eventually to a relatively safe orbit.

## Learning Outcomes

**(NEW)** Studies indicate that beginning students have trouble prioritizing textual material. For this reason, a few (typically five or six) well-defined Learning Outcomes are provided at the start of each chapter. These help students structure their reading of the chapter and then test their mastery of key concepts. The Learning Outcomes are numbered and keyed to

the items in the Chapter Summary, which in turn refer back to passages in the text. This highlighting of the most important aspects of the chapter helps students prioritize information and also aids in their review. The Learning Outcomes are organized and phrased in such a way as to make them objectively testable, affording students a means of gauging their own progress.

**The Big Picture (REVISED)** The Big Picture feature on every chapter opening spread encapsulates the overarching message that each chapter imparts, helping students see how chapter content is connected to a broad understanding of the universe.

**The Big Picture** Stars are everywhere in the nighttime sky. The naked eye can spot about 6000 of them, spread across 88 constellations. Millions more are visible even with binoculars or a small telescope. The total number of stars is impossible to count, and relatively few have been studied in detail. Yet, it is stars that tell us more about the fundamentals of astronomy than any other objects in the universe.

**The Big Question (NEW)** Each chapter now ends with a broad, open-ended query that is intended to ignite students' curiosity about the still-unanswered questions at the forefront of astronomical research. The Big Question builds on the material presented in the chapter and invites students to speculate on the larger scope of what they have just learned.

## Learning Outcomes

*Studying this chapter will enable you to*

- 1 Summarize the composition and physical properties of the interstellar medium.
- 2 Describe the characteristics of emission nebulae, and explain their significance in the life cycle of stars.
- 3 List the basic properties of dark interstellar clouds.
- 4 Specify the radio techniques used to probe the nature of interstellar matter.
- 5 Explain the nature and significance of interstellar molecules.

**Concept Checks** We incorporate into each chapter a number of “Concept Checks”—key questions that require the reader to reconsider some of the material just presented or attempt to place it into a broader context. Answers to these in-chapter questions are provided at the back of the book.

### CONCEPT Check


✓ Why do astronomers draw such a clear distinction between the inner and the outer planets?

**Process of Science Checks** Each chapter now also includes one or two “Process of Science Checks,” similar to the Concept Checks but aimed specifically at clarifying the questions of how science is done and how scientists reach the conclusions they do. Answers to these in-chapter questions are also provided at the back of the book.

### PROCESS OF SCIENCE Check

✓ In what sense are the comets we see *unrepresentative* of comets in general?

**Concept Links** In astronomy, as in many scientific disciplines, almost every topic seems to have some bearing on almost every other. In particular, the connection between the astronomical material and the physical principles set forth early in the text is crucial. Practically everything in Chapters 6–28 of this text rests on the foundation laid in the first five chapters. For example, it is important that students, when they encounter the discussion of high-redshift objects in Chapter 25, recall not only what they just learned about Hubble’s law in Chapter 24 but also refresh their memories, if necessary, about the inverse-square law (Chapter 17), stellar spectra (Chapter 4), and the Doppler shift (Chapter 3). Similarly, the discussions of the mass of binary-star components (Chapter 17) and of galactic rotation (Chapter 23) both depend on the discussion of Kepler’s and Newton’s laws in Chapter 2. Throughout, discussions of new astronomical objects and concepts rely heavily on comparison with topics introduced earlier in the text.

It is important to remind students of these links so that they recall the principles on which later discussions rest and, if necessary, review them. To this end, we have inserted “concept links” throughout the text—symbols that mark key intellectual bridges between material in different chapters. The links, denoted by the symbol  together with a section reference, signal that the topic under discussion is related in some significant way to ideas developed earlier and provide direction to material to review before proceeding.

**Key Terms** Like all subjects, astronomy has its own specialized vocabulary. To aid student learning, the most important astronomical terms are boldfaced at their first appearance in the text. Boldfaced Key Terms in the Chapter Summary are linked with the page number where the term

was defined. In addition, an expanded alphabetical glossary, defining each Key Term and locating its first use in the text, appears at the end of the book.

**H–R Diagrams and Acetate Overlays** All of the book’s H–R diagrams are drawn in a uniform format, using real data. In addition, a unique set of transparent acetate overlays dramatically demonstrates to students how the H–R diagram helps us to organize our information about the stars and track their evolutionary histories.

**More Precisely Boxes** These boxes provide more quantitative treatments of subjects discussed qualitatively in the text. Removing these more challenging topics from the main flow of the narrative and placing them within a separate modular element of the chapter design (so that they can be covered in class, assigned as supplementary material, or simply left as optional reading for those students who find them of interest) will allow instructors greater flexibility in setting the level of their coverage.

**Discovery Boxes** Exploring a wide variety of interesting supplementary topics, Discovery boxes provide the reader with insight into how scientific knowledge evolves and emphasizes the process of science.

**End-of-Chapter Questions, Problems, and Activities (NEW)** Many elements of the end-of-chapter material have seen substantial reorganization:

- Each chapter incorporates **Review and Discussion Questions**, which may be used for in-class review or for assignment. As with the Self-Test Questions, the material needed to answer Review Questions may be found within the chapter. The Discussion Questions explore particular topics more deeply, often asking for opinions, not just facts. As with all discussions, these questions usually have no single “correct” answer. Questions identified with a **POS** icon encourage students to explore the Process of Science, and each Learning Outcome is reflected in one of the Review and Discussion questions, marked by **LO**.
- Each chapter also contains **Conceptual Self-Test Questions** in a multiple-choice format, including select questions that are tied directly to a specific figure or diagram in the text, allowing students to assess their understanding of the chapter material. These questions are identified with a **VIS** icon. Answers to all these questions appear at the end of the book.
- The end-of-chapter material includes **Problems**, based on the chapter contents and requiring some numerical calculation. In many cases the problems are tied directly to quantitative statements made (but not worked out in detail) in the text. The solutions to the problems are not contained verbatim within the chapter, but the information necessary to solve them has been presented in the text. Answers to odd-numbered Problems appear at the end of the book.


- Also new to this edition, the end-of-chapter material now ends with collaborative and individual **Activities** relevant to the material presented in the text. These range from basic naked-eye and telescopic observing projects to opinion polls, surveys, group discussions, and astronomical research on the Web.

**Chapter Review Summaries** The Chapter Review Summaries, a primary review tool, are linked to the Learning Outcomes at the beginning of each chapter. Key Terms introduced in each chapter are listed again, in context and in boldface, along with key figures and page references to the text discussion.


### Chapter Review

SUMMARY


**1** The six main regions of Earth are (from inside to outside) a central metallic core (p. 162), which is surrounded by a thick rocky mantle (p. 162), topped with a thin crust (p. 162). The liquid oceans on our planet's surface make up the hydrosphere (p. 162). Above the surface is the atmosphere (p. 174), which is composed primarily of nitrogen and oxygen and thins rapidly with altitude. Surface winds and weather in the troposphere (p. 163), the lowest region of Earth's atmosphere, are caused by convection (p. 163), the process by which heat is moved from one place to another by the upwelling or downflow of a fluid, such as air or water. Higher above the atmosphere lies the magnetosphere (p. 162), where charged particles from the Sun are trapped by Earth's magnetic field.




by a liquid outer core (p. 169). The process by which heavy material sinks to the center of a planet and lighter material rises to the surface is called **differentiation** (p. 170). Earth's differentiation implies that our planet must have been at least partially molten in the past. One way in which this could have occurred is by the heat released during Earth's formation and subsequent bombardment by material from interplanetary space. Another possibility is the energy released by the decay of **radioactive** (p. 171) elements present in the material from which Earth formed.



**2** At high altitudes, in the ionosphere (p. 163), the atmosphere is kept ionized by the absorption of high-energy radiation and particles from the Sun. In the stratosphere (p. 163), just above the troposphere, lies the ozone layer (p. 164), where incoming solar ultraviolet radiation is absorbed. Both the ionosphere and the ozone layer help protect us from dangerous radiation from space. The greenhouse effect (p. 166) is the absorption and trapping of infrared radiation emitted by Earth's surface by atmospheric gases (primarily carbon dioxide and water vapor). It makes our planet's surface some 40 K warmer than would otherwise be the case. Earth's atmosphere was outgassed from our planet's interior by volcanoes and was then altered by solar radiation and, finally, by the emergence of life.




**4** Earth's surface is made up of about a dozen enormous slabs, or plates. The slow movement of these plates across the surface is called continental drift or **plate tectonics** (p. 174). Earthquakes, volcanism, and mountain building are associated with plate boundaries, where plates may collide, move apart, or rub against one another. The motion of the plates is thought to be driven by convection in Earth's mantle. The rocky upper layer of Earth that makes up the plates is the **lithosphere** (p. 174). The semisolid region in the upper mantle over which the plates slide is called the **asthenosphere** (p. 174). The constant recycling and transformation of crust material as plates separate, collide, and sink into the mantle is called the **rock cycle** (p. 179). Evidence for past plate motion can be found in the geographical fit of continents, in the fossil record, and in the ages and magnetism of surface rocks.



**3** We study Earth's interior by observing how seismic waves (p. 168), produced by earthquakes just below Earth's surface, travel through the mantle. We can also study the upper mantle by analyzing the material brought to the surface when a volcano erupts. Earth's center is dense and extremely hot. The planet's iron core consists of a solid **inner core** (p. 169) surrounded

**5** Earth's magnetic field extends far beyond the surface of our planet. Charged particles from the solar wind are trapped by Earth's magnetic field lines to form the **Van Allen belts** (p. 180) that surround our planet. When particles



## Instructor Resources



MasteringAstronomy is the most widely used and most advanced astronomy tutorial and assessment system in the world. By capturing the step-by-step work of students nationally, MasteringAstronomy has established an unparalleled database of learning challenges and patterns. Using this student data, a team of renowned astronomy education researchers has refined every activity and problem. The result is a library of activities of unique educational effectiveness and assessment accuracy. MasteringAstronomy provides students with two learning systems in one: a dynamic self-study area and the ability to participate in online assignments.

MasteringAstronomy provides instructors with a fast and effective way to assign uncompromising, wide-ranging online homework assignments of just the right difficulty and duration. The tutorials coach 90 percent of students to the correct answer with specific wrong-answer feedback. Powerful post-diagnostics allow instructors to assess the

progress of their class as a whole or to quickly identify an individual student's areas of difficulty. Tutorials built around text content and all the end-of-chapter problems from the text are available in MasteringAstronomy. A media-rich self-study area is included that students can use whether the instructor assigns homework or not.

**Instructor Guide** Revised by James Heath (Austin Community College), this online guide provides: sample syllabi and course schedules; an overview of each chapter; pedagogical tips; useful analogies; suggestions for classroom demonstrations; writing questions, selected readings, and answers/solutions to the end-of-chapter Review and Discussion Questions and Problems; and additional references and resources. ISBN 0-321-91021-4

**Test Bank** An extensive file of approximately 2800 test questions, newly compiled and revised for the eighth edition. The questions are organized and referenced by chapter section and by question type. The eighth edition Test Bank has been thoroughly revised and includes many new Multiple Choice and Essay questions for added conceptual emphasis. This Test Bank is available in both Microsoft® Word and TestGen® formats (see description of Instructor Resource DVD). ISBN 0-321-91008-7

### Instructor Resource Area in Mastering Astronomy

This instructor resource area resides in MasteringAstronomy and provides every electronic asset professors will need in and out of the classroom. The area not only contains an Instructor's Resource Manual, but also all text figures in jpeg and PowerPoint formats, including additional images, star charts, as well as the animations and videos from the MasteringAstronomy® Study Area. The area also contains TestGen®, an easy-to-use, fully networkable program for creating tests ranging from short quizzes to long exams. Questions from the Test Bank are supplied, and professors can use the Question Editor to modify existing questions or create new questions. It also contains chapter-by-chapter lecture outlines in PowerPoint and conceptual "clicker" questions in PowerPoint. It is available in both PC and Mac formats.

**Instructor Resource Center** The Pearson Instructor Resource Center contains everything found on the Instructor Resource Area in MasteringAstronomy and the Instructor DVD, above, with the exception of the text figures in jpeg and PowerPoint formats, which are too large to download.

**Instructor Resource DVD** This DVD contains every resource found in the Instructor Resource Area in MasteringAstronomy, and it provides virtually every electronic asset professors will need in and out of the classroom. The disc contain all text figures in jpeg and PowerPoint formats, as well as the animations and videos from the Mastering Astronomy® Study Area. The IR-DVD also contains TestGen®, an easy-to-use, fully networkable program for creating tests ranging

from short quizzes to long exams. Questions from the Test Bank are supplied, and professors can use the Question Editor to modify existing questions or create new questions. This disc set also contains chapter-by-chapter lecture outlines in PowerPoint and conceptual “clicker” questions in PowerPoint. ISBN 0-321-90974-7

### ***Learner-Centered Astronomy Teaching: Strategies for ASTRO 101***

Timothy F. Slater, *University of Wyoming*  
Jeffrey P. Adams, *Millersville University*

*Strategies for ASTRO 101* is a guide for instructors of the introductory astronomy course for nonscience majors. Written by two leaders in astronomy education research, this book details various techniques instructors can use to increase students' understanding and retention of astronomy topics, with an emphasis on making the lecture a forum for active student participation. Drawing from the large body of recent research to discover how students learn, this guide describes the application of multiple classroom-tested techniques to the task of teaching astronomy to predominantly nonscience students. ISBN 0-13-046630-1

### ***Peer Instruction for Astronomy***

Paul J. Green, *Harvard-Smithsonian Center for Astrophysics*

Peer instruction is a simple yet effective method for teaching science. Techniques of peer instruction for introductory physics were developed primarily at Harvard and have aroused interest and excitement in the physics education community. This approach involves students in the teaching process, making science more accessible to them. This book is an important vehicle for providing a large number of thought-provoking, conceptual short-answer questions aimed at a variety of class levels. While significant numbers of such questions have been published for use in physics, *Peer Instruction for Astronomy* provides the first such compilation for astronomy. ISBN 0-13-026310-9

## **Student Resources**

 **MasteringAstronomy**<sup>®</sup>  
www.masteringastronomy.com

This homework, tutorial, and assessment system is uniquely able to tutor each student individually by providing students with instantaneous feedback specific to their wrong answers, simpler subproblems upon request when they get stuck, and partial credit for their method(s) used. Students also have access to a self-study area that contains practice quizzes, self-guided tutorials, new narrated and interactive figures, animations, videos, and more.

**Pearson eText** is available through MasteringAstronomy, either automatically when MasteringAstronomy is packaged with new books, or available as a purchased upgrade online.

Allowing the students to access the text wherever they have access to the Internet, Pearson eText comprises the full text, including figures that can be enlarged for better viewing. Within Pearson eText students are also able to pop up definitions and terms to help with vocabulary and the reading of the material. Students also can take notes in Pearson eText using the annotation feature.

### ***Starry Night College™ Student Access Code Card, 7th Edition***

This best-selling planetarium software lets you escape the Milky Way and travel within 700 million light-years of space. View more than 16 million stars in stunningly realistic star fields. Zoom in on thousands of galaxies, nebulae, and star clusters. Move through 200,000 years of time to see key celestial events in a dynamic and ever-changing universe. Blast off from Earth and see the motions of the planets from a new perspective. Hailed for its breathtaking realism, powerful suite of features, and intuitive ease of use, *Starry Night College™* lives up to its reputation as astronomy software's brightest . . . night after night. ISBN 0-321-71295-1

### ***Starry Night College™ Activities & Observation and Research Projects***

This downloadable supplement contains activities for *Starry Night College* planetarium software by Erin O'Connor (Santa Barbara City College), as well as observation and research projects by Steve McMillan. It is downloadable free from the MasteringAstronomy Study Area and also from the Pearson *Starry Night College* download site. ISBN 0-321-75307-0

***SkyGazer 5.0 Student Access Code Card*** This access kit provides a one-time download of *SkyGazer 5.0* that combines exceptional planetarium software with informative pre-packaged tutorials. Based on the popular *Voyager* software, this access code card is available to be packaged at no additional charge with new copies of introductory astronomy textbooks. Along with the software, this access code card also enables users to download the *Astronomy Media Workbook* by Michael LoPresto. ISBN 0-321-76518-4

(Also available on CD-ROM. ISBN 0-321-89843-5)

***Sky and Telescope*** Based on the most popular amateur astronomy magazine, this special student supplement contains nine articles by Evan Skillman, each with a general overview and four question sets focused on the issues professors most want to address in this course: General Review, Process of Science, Scale of the Universe, and Our Place in the Universe. ISBN 0-321-70620-X

***Edmund Scientific Star and Planet Locator*** The famous rotating roadmap of the heavens shows the location of the stars, constellations, and planets relative to the horizon for the exact hour and date you determine. This eight-square star

chart was plotted by the late astronomer and cartographer George Lovi. The reverse side of the locator is packed with additional data on the planets, meteor showers, and bright stars. Included with each star chart is a 16-page, fully illustrated, pocket-size instruction booklet.  
 ISBN 0-13-140235-8

**Lecture-Tutorials for Introductory Astronomy, 3rd Edition**

Edward E. Prather, *University of Arizona*  
 Timothy F. Slater, *University of Wyoming*  
 Jeffrey P. Adams, *Millersville University*  
 Gina Brissenden, *University of Arizona*

Funded by the National Science Foundation, *Lecture-Tutorials for Introductory Astronomy* is designed to help make large-lecture-format courses more interactive. The third edition features six new tutorials on the Greenhouse Effect; Dark Matter; Making Sense of the Universe and Expansion; Hubble’s Law; Expansion, Lookback Times, and Distances; and The Big Bang. Each of the 44 Lecture-Tutorials is presented in a classroom-ready format that asks students to work in groups of two to three for between 10 and 15 minutes and requires no equipment. These lecture-tutorials challenge students with

a series of carefully designed questions that spark classroom discussion and engage students in critical reasoning.  
 ISBN 0-321-82046-0

**Observation Exercises in Astronomy** This workbook by Lauren Jones contains a series of astronomy exercises that integrate technology from planetarium software such as Stellarium, Starry Night College, WorldWide Telescope, and SkyGazer. Using these online products adds an interactive dimension to students’ learning.  
 ISBN: 0-321-63812-3

## Acknowledgments

Throughout the many drafts that have led to this book, we have relied on the critical analysis of many colleagues. Their suggestions ranged from the macroscopic issue of the book’s overall organization to the minutiae of the technical accuracy of each and every sentence. We have also benefited from much good advice and feedback from users of the first seven editions of the text. To these many helpful colleagues, we offer our sincerest thanks.

## Reviewers of the Eighth Edition

- |  |  |   |
|--|--|---|
| Brett Bochner<br><i>Hofstra University</i>             | Erika Gibb<br><i>University of Missouri, St. Louis</i>         | John Scalo<br><i>University of Texas, Austin</i>                    |
| James Brau<br><i>University of Oregon</i>              | James Higdon<br><i>Georgia Southern University</i>             | Trace Tessier<br><i>Central New Mexico Community College</i>        |
| Christina Cavalli<br><i>Austin Community College</i>   | Steve Kawaler<br><i>Iowa State University</i>                  | Robert K. Tyson<br><i>University of North Carolina at Charlotte</i> |
| Asif ud-Doula<br><i>Pennsylvania State University</i>  | Kristine Larsen<br><i>Central Connecticut State University</i> | Grant Wilson<br><i>University of Massachusetts, Amherst</i>         |
| Robert Egler<br><i>North Carolina State University</i> | George Nock<br><i>Northeast Mississippi Community College</i>  |   |
| David Ennis<br><i>The Ohio State University</i>        | Ron Olowin<br><i>Saint Mary’s College</i>                      |   |

## Reviewers of Previous Editions

- |  |   |   |   |
|--|---|---|---|
| Stephen G. Alexander<br><i>Miami University of Ohio</i>      | Peter A. Becker<br><i>George Mason University</i>         | Bruce Cragin<br><i>Richland College</i>                   | Michael N. Fanelli<br><i>University of North Texas</i>  |
| William Alexander<br><i>James Madison University</i>         | Timothy C. Beers<br><i>University of Evansville</i>       | Ed Coppola<br><i>Community College of Southern Nevada</i> | Richard Gelderman<br><i>Western Kentucky University</i> |
| Robert H. Allen<br><i>University of Wisconsin, La Crosse</i> | William J. Boardman<br><i>Birmingham Southern College</i> | David Currott<br><i>University of North Alabama</i>       | Harold A. Geller<br><i>George Mason University</i>      |
| Barlow H. Allen<br><i>University of Wisconsin, La Crosse</i> | Donald J. Bord<br><i>University of Michigan, Dearborn</i> | Norman Derby<br><i>Bennington College</i>                 | David Goldberg<br><i>Drexel University</i>              |
| Nadine G. Barlow<br><i>Northern Arizona University</i>       | Elizabeth P. Bozyan<br><i>University of Rhode Island</i>  | John Dykla<br><i>Loyola University, Chicago</i>           | Martin Goodson<br><i>Delta College</i>                  |
| Cecilia Barnbaum<br><i>Valdosta State University</i>         | Malcolm Cleaveland<br><i>University of Arkansas</i>       | Kimberly Engle<br><i>Drexel University</i>                | David G. Griffiths<br><i>Oregon State University</i>    |
|  | Anne Cowley<br><i>Arizona State University</i>            |   | Donald Gudehus<br><i>Georgia State University</i>       |

Thomasanna Hail <i>Parkland College</i>	Robert J. Leacock <i>University of Florida</i>	Andrew P. Odell <i>Northern Arizona University</i>	Harry L. Shipman <i>University of Delaware</i>
Clint D. Harper <i>Moorpark College</i>	Larry A. Lebofsky <i>University of Arizona</i>	Gregory W. Ojakangas <i>University of Minnesota, Duluth</i>	C. G. Pete Shugart <i>Memphis State University</i>
Marilynn Harper <i>Delaware County Community College</i>	Matthew Lister <i>Purdue University</i>	Ronald Olowin <i>Saint Mary's College of California</i>	Stephen J. Shulik <i>Clarion University</i>
Susan Hartley <i>University of Minnesota, Duluth</i>	M. A. Lohdi <i>Texas Tech University</i>	Robert S. Patterson <i>Southwest Missouri State University</i>	Tim Slater <i>University of Arizona</i>
Joseph Heafner <i>Catawaba Valley Community College</i>	Michael C. LoPresto <i>Henry Ford Community College</i>	Cynthia W. Peterson <i>University of Connecticut</i>	Don Sparks <i>Los Angeles Pierce College</i>
James Heath <i>Austin Community College</i>	Phillip Lu <i>Western Connecticut State University</i>	Lawrence Pinsky <i>University of Houston</i>	George Stanley, Jr. <i>San Antonio College</i>
Fred Hickok <i>Catonsville Community College</i>	Fred Marschak <i>Santa Barbara College</i>	Andreas Quirrenback <i>University of California, San Diego</i>	Maurice Stewart <i>Willamette University</i>
Lynn Higgs <i>University of Utah</i>	Matthew Malkan <i>University of California, Los Angeles</i>	Richard Rand <i>University of New Mexico</i>	Jack W. Sulentic <i>University of Alabama</i>
Darren L. Hitt <i>Loyola College, Maryland</i>	Steve Mellema <i>Gustavus Adolphus College</i>	James A. Roberts <i>University of North Texas</i>	Andrew Sustich <i>Arkansas State University</i>
F. Duane Ingram <i>Rock Valley College</i>	Chris Mihos <i>Case Western Reserve University</i>	Gerald Royce <i>Mary Washington College</i>	Donald Terndrup <i>The Ohio State University</i>
Steven D. Kawaler <i>Iowa State University</i>	Milan Mijic <i>California State University, Los Angeles</i>	Dwight Russell <i>Baylor University</i>	Craig Tyler <i>Fort Lewis College</i>
William Keel <i>University of Alabama</i>	Scott Miller <i>Pennsylvania State University</i>	Vicki Sarajedini <i>University of Florida</i>	Stephen R. Walton <i>California State University, Northridge</i>
Marvin Kemple <i>Indiana University-Purdue University, Indianapolis</i>	Mark Moldwin <i>University of California, Los Angeles</i>	Malcolm P. Savedoff <i>University of Rochester</i>	Peter A. Wehinger <i>University of Arizona</i>
Mario Klairc <i>Midlands Technical College</i>	Richard Nolthenius <i>Cabrillo College</i>	John Scalo <i>University of Texas at Austin</i>	Louis Winkler <i>Pennsylvania State University</i>
Kristine Larsen <i>Central Connecticut State University</i>	Edward Oberhofer <i>University of North Carolina, Charlotte</i>	John C. Schneider <i>Catonsville Community College</i>	Jie Zhang <i>George Mason University</i>
Andrew R. Lazarewicz <i>Boston College</i>		Larry Sessions <i>Metropolitan State College of Denver</i>	Robert Zimmerman <i>University of Oregon</i>

The publishing team at Pearson has assisted us at every step along the way in creating this text. Particular thanks go to Tema Goodwin, who managed with heroic fortitude the many conflicting variables and personalities that are a part of a multifaceted publication such as this. Executive editor Nancy Whilton steered this edition through all its phases, and development editor Barbara Price contributed her media expertise. Production managers Andrea Archer and Angela Urquhart of Thistle Hill Publishing Services have done an excellent job of tying together the threads of this very complex project, made all the more complex by the necessity of combining text, art, and electronic media into a coherent whole. Special thanks are in order to cover and interior designer Jeanne Calabrese for making the eighth

edition look spectacular and to Mark Ong for guiding the overall look of the book. We would also like to express our appreciation to Kate Brayton for updating and maintaining the media resources in the MasteringAstronomy® Study Area and to Christina Cavalli, author of the MasteringAstronomy Narrated Figures.

Finally, we would like to express our gratitude to renowned space artist Dana Berry for allowing us to use many of his beautiful renditions of astronomical scenes, and to Lola Judith Chaisson for assembling and drawing all the H–R diagrams (including the acetate overlays) for this edition.

**Eric Chaisson**  
**Steve McMillan**



*This page intentionally left blank*


# Astronomy Today



Galileo's sketch of Saturn

## PART ONE

# Astronomy and the Universe




Galileo's sketch of Orion

**It is often said** that we live in a golden age of astronomy. Yet the dawn of the 21st century is actually the second such period of rich discovery and rapid exploration. The first era of stunning scientific growth began in the late Renaissance. Foremost among the early architects of modern astronomy was the Italian scientist Galileo Galilei (1564–1642). By turning his telescope to the heavens, he changed radically and forever our view of the universe in which we live.

Although he did not invent the telescope, in 1610 Galileo was the first to record what he saw when he aimed a small (5-cm-diameter) lens at the sky. His findings created nothing less than a revolution in astronomy. Viewing for the first time dark blemishes on the Sun, rugged mountains on the Moon, and whole new worlds orbiting Jupiter, he demolished the Aristotelian notion that the heavens were perfect and unchanging. It was with the philosophers of the day, as much as with the theologians, that Galileo had trouble. In championing the scientific method, he used a tool to test his ideas, and what he found disagreed greatly with the leading thoughts and beliefs of the time.

Galileo's advance was simple yet profound: He used a telescope to focus, magnify, and study radiation reaching Earth from the heavens—in particular, light from the Sun, the Moon, and the planets. Light is the most familiar kind of radiation to humans on Earth, since it enables us to get around on the surface of our planet. But light also enables telescopes to see objects deep in space, allowing us to probe farther than the eye can alone. With his simple optical telescope, Galileo changed completely the way that the oldest science—astronomy—is pursued.


Among other “wondrous things” he found were star clusters along the Milky Way, moons and rings around the outer planets, and colorful nebulae unlike anything seen before. Some of Galileo's sketches are reproduced here (left side) and are compared with modern views at right.



Galileo Galilei



Galileo's sketch of the Pleiades



Saturn in the ultraviolet (STScI)

**Today**, we are again in the midst of another period of unsurpassed scientific achievement—a revolution in which modern astronomers are revealing the invisible universe as Galileo once spied the visible universe. We have learned how to detect, measure, and analyze invisible radiation streaming to us from dark objects in space. And once again our perceptions are changing.

Astronomy no longer evokes visions of plodding intellectuals peering through long telescope tubes. Nor does the cosmos any longer refer to that seemingly inactive, immutable domain seen visually when we gaze at the nighttime sky. Modern astronomers now decipher a more vibrant, changing universe—one in which stars emerge and perish much like living things, galaxies spew forth vast quantities of energy, and life itself is thought to be a natural consequence of the evolution of matter.

New discoveries are rapidly advancing our understanding of the universe, but they also raise new questions.

Astronomers will encounter many problems in the decades ahead, but this should neither dismay nor frustrate us, for it is precisely how science operates. Each discovery adds to our storehouse of information, generating a host of questions that lead in turn to more discoveries, and so on, causing an acceleration of basic knowledge.

Most notably, we are beginning to perceive the universe in all its multivariied ways. A single generation—not the generation of our parents and not that of our children, but our generation—has opened up the whole electromagnetic spectrum beyond visible light. And what we, too, have found are “wondrous things.”

Emerging largely from studies of the invisible universe, our view of the cosmos in its full splendor is one of many new scientific insights that we have recently been privileged to attain. Historians of the future may well regard our generation as the one that took a great leap forward, providing a whole new glimpse of our richly endowed universe. In all of history, there have been only two periods in which our perception of the universe has been so revolutionized within a single human lifetime.

The first occurred four centuries ago at the time of Galileo; the second is now under way.

Orion in the infrared (Caltech)

Pleiades in the optical (AURA)



# Charting the Heavens

## THE FOUNDATIONS OF ASTRONOMY

Nature offers no greater splendor than the starry sky on a clear, dark night. Silent and jeweled with the constellations of ancient myth and legend, the night sky has inspired wonder throughout the ages—a wonder that leads our imaginations far from the confines of Earth and the pace of the present day and out into the distant reaches of space and cosmic time itself.

Astronomy, born in response to that wonder, is built on two of the most basic traits of human nature: the *need to explore* and the *need to understand*. Through the interplay of curiosity, discovery, and analysis—the keys to exploration and understanding—people have sought answers to questions about the universe since the earliest times. Astronomy is the oldest of all the sciences, yet never has it been more exciting than it is today.

**The Big Picture** Our subject is science, and that means rich details and specific ideas. Even so, we also need to keep in mind a larger, general perspective. And when it comes to astronomy, there is perhaps no grander feature of the cosmos than stars—they're everywhere in the nighttime sky, like those seen in the photo opposite. Roughly as many stars reside in the observable universe as there are grains of sand in all the beaches of the world—about a hundred sextillion, or  $10^{23}$ .

**LEFT:** High overhead on a clear, dark night, we can see a rich band of stars known as the Milky Way—so-called for its resemblance to a milky band of countless stars. All these stars (and more) are part of a much larger system called the Milky Way Galaxy, of which our star, the Sun, is one member. This image shows the awesome splendor of the Milky Way shining above some of the big telescopes of the European Southern Observatory, a major astronomy facility high in the Chilean Andes. (ESO/Y. Beletsky)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Arrange the basic levels of structure in the universe in order of increasing size.
- 2 Distinguish among scientific theories, hypotheses, and observations, and describe how scientists combine observation, theory, and testing in their study of the universe.
- 3 Describe the celestial sphere, and tell how astronomers use constellations and angular measurement to locate objects in the sky.
- 4 Describe how and why the Sun and the stars appear to change their positions from night to night and from month to month.
- 5 Explain how Earth's axial tilt causes the seasons, and why the seasons change over time.
- 6 Account for the changing appearance of the Moon, and explain how the relative motions of Earth, the Sun, and the Moon lead to eclipses.
- 7 Give an example of how simple geometric reasoning can be used to measure the distances and sizes of otherwise inaccessible objects.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.

## 1.1 Our Place in Space

Of all the scientific insights attained to date, one stands out boldly: Earth is neither central nor special. We inhabit no unique place in the universe. Astronomical research, especially within the past few decades, strongly suggests that we live on what seems to be an ordinary rocky *planet* called Earth, one of eight known planets orbiting an average *star* called the Sun, a star near the edge of a huge collection of stars called the Milky Way Galaxy, which is one *galaxy* among billions of others spread throughout the observable universe. To begin to get a feel for the relationships among these very different objects, consult Figures 1.1 through 1.5.

We are connected to the most distant realms of space and time not only by our imaginations but also through a common cosmic heritage. Most of the chemical elements that make up our bodies (hydrogen, oxygen, carbon, and many more) were created billions of years ago in the hot centers of long-vanished stars. Their fuel supply spent, these giant stars died in huge explosions, scattering the elements created deep within their cores far and wide. Eventually, this matter collected into clouds of gas that slowly collapsed to give birth to new generations of stars. In this way, the Sun and its family of planets formed nearly 5 billion years ago. Everything on Earth embodies atoms from other parts of the universe and from a past far more remote than the beginning of human evolution. Elsewhere, other beings—perhaps with intelligence much greater than our own—may at this very moment be gazing in wonder at



◀ **FIGURE 1.1 Humans** We know our own size and scale well—adult humans are typically 1.5 meters tall. Earth in the next figure is about 10 million times bigger. (*J. Lodriguss*)



▲ **FIGURE 1.2 Earth** Earth is a planet, a mostly solid object, although it has some liquid in its oceans and core and gas in its atmosphere. In this view, the North and South American continents are clearly visible, though most of the scene shows Pacific waters. (*NASA*)



▲ **FIGURE 1.3 The Sun** The Sun is a star, a very hot ball of gas composed mainly of hydrogen and helium. Much bigger than Earth—more than 100 times larger in diameter—the Sun is held together by its own gravity. The dark blemishes are sunspots (see Chapter 16). (*AURA*)

their own night sky. Our own Sun may be nothing more than an insignificant point of light to them—if it is visible at all. Yet if such beings exist, they must share our cosmic origin.

Simply put, the **universe** is the totality of all space, time, matter, and energy. **Astronomy** is the study of the universe. It is a subject unlike any other, for it requires us to profoundly change our view of the cosmos and to consider matter on scales totally unfamiliar from everyday experience. Look again at the galaxy in Figure 1.4. It is a swarm of about a hundred billion stars—more stars than the number of people who have ever lived on Earth. The entire assemblage is spread across a vast expanse of space 100,000 **light-years** in diameter. Although it sounds like a

unit of time, a light-year is in fact the *distance* traveled by light in a year, at a speed of about 300,000 kilometers per second. Multiplying out, it follows that a light-year is equal to  $300,000 \text{ kilometers/second} \times 86,400 \text{ seconds/day} \times 365 \text{ days}$  or about 10 trillion kilometers, or roughly 6 trillion miles. Typical galactic systems are truly “astronomical” in size. For comparison, Earth’s roughly 13,000-km diameter is less than one-twentieth of a light-second.


The light-year is a unit introduced by astronomers to help them describe immense distances. We will encounter many such custom units in our studies. As discussed in more detail in Appendix 2, astronomers frequently augment the standard SI (Système Internationale) metric system with additional units tailored to the particular problem at hand.



▲ **FIGURE 1.4 Galaxy** A typical galaxy is a collection of a hundred billion stars, each separated by vast regions of nearly empty space. Our Sun is a rather undistinguished star near the edge of another such galaxy, called the Milky Way. (R. Gendler/Science Source)

A thousand (1000), a million (1,000,000), a billion (1,000,000,000), and even a trillion (1,000,000,000,000)—these words occur regularly in everyday speech. But let's take a moment to understand the magnitude of the numbers and appreciate the differences among them. One thousand is easy enough to understand: At the rate of one number per second, you could count to a thousand in 1000 seconds—about 16 minutes. However, if you wanted to count to a million, you would need more than 2 weeks of counting at the rate of one number per second, 16 hours per day (allowing 8 hours per day for sleep). To count from one to a billion at the same rate of one number per second and 16 hours per day would take nearly 50 years—the better part of an entire human lifetime.

In this book, we consider *distances* in space spanning not just billions of kilometers, but billions of light-years; *objects* containing not just trillions of atoms, but trillions of stars; and *time intervals* of not just billions of seconds or hours, but billions of years. You will need to become



▲ **FIGURE 1.5 Galaxy Cluster** This photograph shows a typical cluster of galaxies, spread across roughly a million light-years of space. Each galaxy contains hundreds of billions of stars, probably planets, and possibly living creatures. (NASA)

familiar—and comfortable—with such enormous numbers. A good way to begin is learning to recognize just how much larger than a thousand is a million, and how much larger still is a billion. Appendix 1 explains the convenient method used by scientists for writing and manipulating very large and very small numbers. If you are unfamiliar with this method, please read that appendix carefully—the *scientific notation* described there will be used consistently throughout our text, beginning in Chapter 2.

Lacking any understanding of the astronomical objects they observed, early skywatchers made up stories to explain them: The Sun was pulled across the heavens by a chariot drawn by winged horses, and patterns of stars traced heroes and animals placed in the sky by the gods. Today, of course, we have a radically different conception of the universe. The stars we see are distant, glowing orbs hundreds of times larger than our entire planet, and the patterns they form span hundreds of light-years. In this first chapter we present some basic methods used by astronomers to chart the space around us. We describe the slow progress of scientific knowledge, from chariots and gods to today's well-tested theories and physical laws, and explain why we now rely on science rather than on myth to help us explain the universe.




## 1.2 Scientific Theory and the Scientific Method

How have we come to know the universe around us and the cosmic perspective sketched in Figures 1.1–1.5? The earliest descriptions of the universe were based largely on imagination and mythology and made little attempt to explain the workings of the heavens in terms of earthly experience. However, some early scientists realized the importance of careful observation and testing to the formulation of their ideas. The success of their approach changed, slowly but surely, the way science was done and opened the door to a fuller understanding of nature. As the influence of logic and reasoned argument grew, the power of myth diminished. People began to inquire more critically about themselves and the universe. They realized that *thinking* about nature was no longer sufficient—*looking* at it was also necessary. Experiments and observations became a central part of the process of inquiry.

To be effective, a **theory**—the framework of ideas and assumptions used to explain some set of observations and make predictions about the real world—must be continually tested. Scientists accomplish this by using a theory to construct a **theoretical model** of a physical object (such as a planet or a star) or phenomenon (such as gravity or light) that accounts for its known properties. The model then makes further predictions about the object’s properties, or perhaps how it might behave or change under new circumstances. If experiments and observations favor those predictions, the theory can be further developed and refined. If not, the theory must be reformulated or rejected, no matter how appealing it originally seemed. This approach to investigation, combining thinking and doing—that is, theory and experiment—is known as the **scientific method**. The process, combining theoretical reasoning with experimental testing, is illustrated schematically in Figure 1.6. It lies at the heart of modern science, separating science from pseudoscience, fact from fiction.

The notion that theories must be tested and may be proven wrong sometimes leads people to dismiss their importance. We have all heard the expression, “Of course, it’s only a theory,” used to deride or dismiss an idea that someone finds unacceptable. Don’t be fooled! Gravity (see Section 2.7) is “only” a theory, but calculations based on it have guided human spacecraft throughout the solar system. Electromagnetism (Chapter 3) and quantum mechanics (Chapter 4) are theories, too, yet they form the foundation for technology. Facts about much of the universe are a dime a dozen. Theories are the intellectual “glue” that combine seemingly unrelated facts into a coherent and interconnected whole.


Notice that there is no end point to the process depicted in Figure 1.6. A theory can be invalidated by a single wrong prediction, but no amount of observation or



▲ **FIGURE 1.6 Scientific Method** Scientific theories evolve through a combination of observation, theoretical reasoning, and prediction, suggesting new observations. The process can begin at any point in the cycle, and it continues forever—or until the theory fails to explain an observation or makes a demonstrably false prediction.

experimentation can ever prove it “correct.” Theories simply become more and more widely accepted as their predictions are repeatedly confirmed. Modern scientific theories share several important defining characteristics:

- They must be *testable*—that is, they must admit the possibility that their underlying assumptions and their predictions can, in principle, be exposed to experimental verification. This feature separates science from, for example, religion, since, ultimately, divine revelations or scriptures cannot be challenged within a religious framework—we can’t design an experiment to “verify the mind of God.” Testability also distinguishes science from a pseudoscience such as astrology, whose underlying assumptions and predictions have been repeatedly tested and never verified, with no apparent impact on the views of those who continue to believe in it!
- They must continually be *tested*, and their consequences tested, too. This is the basic circle of scientific progress depicted in Figure 1.6.
- They should be *simple*. This is less a requirement than a practical outcome of centuries of scientific experience—the most successful theories tend to be the simplest ones that fit the facts. This viewpoint is often encapsulated in a principle known as *Occam’s razor*: If two competing theories both explain the facts and make the same predictions, then the simpler one is better. Put another way—“Keep it simple!” A good theory should be no more complex than is absolutely necessary.
- Finally, most scientists have the additional bias that a theory should in some sense be *elegant*. When a



▲ **FIGURE 1.7 A Lunar Eclipse** These photographs show Earth’s shadow (denoted by the dashed curve) sweeping across the Moon during a lunar eclipse. By observing this behavior, Aristotle reasoned that Earth was the cause of the shadow and concluded that Earth must be round. His theory has yet to be disproved. (G. Schneider)

clearly stated simple principle naturally ties together and explains several phenomena previously thought to be completely distinct, this is widely regarded as a strong point in favor of the new theory.

You may find it instructive to apply these criteria to the many physical theories—some old and well established, others much more recent and still developing—we will encounter throughout the text.

The birth of modern science is usually associated with the Renaissance, the historical period from the late 14th to the mid-17th century that saw a rebirth (*renaissance* in French) of artistic, literary, and scientific inquiry in European culture following the chaos of the Dark Ages. However, one of the first documented uses of the scientific method in an astronomical context was made by Aristotle (384–322 B.C.) some 17 centuries earlier. Aristotle is not normally remembered as a strong proponent of this approach—many of his best known ideas were based on pure thought, with no attempt at experimental test or verification. Nevertheless, his brilliance extended into many areas now thought of as modern science. He noted that, during a lunar eclipse (Section 1.6), Earth casts a curved shadow onto the surface of the Moon. Figure 1.7 shows a series of photographs taken during a recent lunar eclipse. Earth’s shadow, projected onto the Moon’s surface, is indeed slightly curved. This is what Aristotle must have seen and recorded so long ago.

Because the observed shadow seemed always to be an arc of the same circle, Aristotle theorized that Earth, the cause of the shadow, must be round. Don’t underestimate the scope of this apparently simple statement. Aristotle also had to reason that the dark region was indeed a shadow and that Earth was its cause—facts we regard as obvious today, but far from clear 25 centuries ago. On the basis of this *hypothesis*—one possible explanation of the observed facts—he then predicted that any and all future lunar eclipses would show Earth’s shadow to be curved, regardless of our planet’s orientation. That prediction has

been tested every time a lunar eclipse has occurred. It has yet to be proved wrong. Aristotle was not the first person to argue that Earth is round, but he was apparently the first to offer observational proof using this method.

This basic reasoning forms the basis of all modern scientific inquiry. Armed only with naked-eye observations of the sky (the telescope would not be invented for almost another 2000 years), Aristotle first made an observation. Next, he formulated a hypothesis to explain that observation. Then he tested the validity of his hypothesis by making predictions that could be confirmed or refuted by further observations. *Observation, theory, and testing*—these are the cornerstones of the scientific method, a technique whose power will be demonstrated again and again throughout our text.

Today, scientists throughout the world use an approach that relies heavily on testing ideas. They gather data, form a working hypothesis that explains the data, and then proceed to test the implications of the hypothesis using experiment and observation. Eventually, one or more “well-tested” hypotheses may be elevated to the stature of a physical law and come to form the basis of a theory of even broader applicability. The new predictions of the theory will in turn be tested, as scientific knowledge continues to grow. Experiment and observation are integral parts of the process of scientific inquiry. Untestable theories, or theories unsupported by experimental evidence, rarely gain any measure of acceptance in scientific circles. Used properly over a period of time, this rational, methodical approach enables us to arrive at conclusions that are mostly free of the personal bias and human values of any one scientist—it is designed to yield an objective view of the universe we inhabit.

### PROCESS OF SCIENCE Check

- ✓ Can a theory ever become a “fact,” scientifically speaking?

## 1.3 The “Obvious” View

To see how astronomers apply the scientific method to understand the universe around us, let’s start with some very basic observations. Our study of the cosmos, the modern science of astronomy, begins with looking at the night sky. The overall appearance of the sky is not so different now from what our ancestors would have seen hundreds or thousands of years ago, but our *interpretation* of what we see has changed immeasurably as the science of astronomy has evolved and grown.

### Constellations in the Sky

Between sunset and sunrise on a clear night, we can see about 3000 points of light. Including the view from the opposite side of Earth, nearly 6000 stars are visible to the unaided eye. A natural human tendency is to see patterns and relationships among objects even when no true connection exists, and people long ago connected the brightest stars into configurations called **constellations**, which ancient astronomers named after mythological beings, heroes, and animals—whatever was important to them. Figure 1.8 shows a constellation prominent in the nighttime sky from October through March: the hunter named Orion. Orion was a mythical Greek hero famed, among other things, for his amorous pursuit of the Pleiades, the seven daughters of the giant Atlas. According to Greek mythology, to protect the Pleiades from Orion,


the gods placed them among the stars, where Orion still stalks them across the sky. Many constellations have similarly fabulous connections with ancient lore.

Perhaps not surprisingly, the patterns have a strong cultural bias—ancient Chinese astronomers saw mythical figures different from those seen by the Greeks, the Babylonians, and the people of other cultures, even though they were all looking at the same stars in the night sky. Interestingly, different cultures often made the same basic *groupings* of stars, despite widely varying interpretations of what they saw. For example, the group of seven stars known in North America as “the Dipper” is called “the Wagon” or “the Plough” in western Europe. The ancient Greeks regarded these same stars as the tail of “the Great Bear,” the Egyptians saw them as the leg of an ox, the Siberians as a stag, and some Native Americans as a funeral procession.

Early astronomers had very practical reasons for studying the sky. Some constellations served as navigational guides. The star Polaris (part of the Little Dipper) indicates north, and the near constancy of its location in the sky, from hour to hour and night to night, has aided travelers for centuries. Other constellations served as primitive calendars to predict planting and harvesting seasons. For example, many cultures knew that the appearance of certain stars on the horizon just before daybreak signaled the beginning of spring and the end of winter.

In many societies, people came to believe that there were other benefits in tracing the regularly changing positions of heavenly bodies. The relative positions of stars and planets


This is a real photo of the Orion constellation . . .



. . . and this is a mapped interpretation, to exactly the same scale.



**Interactive FIGURE 1.8 Constellation Orion** (a) A photograph of the group of bright stars that make up the constellation Orion. (See the preface, p. xxv, for an explanation of the icon at the bottom, which simply indicates that this image was made in visible light.) (b) The stars are connected to show the pattern visualized by the Greeks: the outline of a hunter. The Greek letters serve to identify some of the brighter stars in the constellation (see also Figure 1.9). You can easily find Orion in the northern winter sky by identifying the line of three bright stars in the hunter’s “belt.” (*P. Sanz/Alamy*)



◀ **FIGURE 1.9 Orion in 3-D**

The true three-dimensional relationships among the most prominent stars in Orion. The distances were determined by the *Hipparcos* satellite in the 1990s. (See Chapter 17.)

at a person’s birth were carefully studied by *astrologers*, who used the data to make predictions about that person’s destiny. Thus, in a sense, astronomy and astrology arose from the same basic desire—to “see” into the future—and, indeed, for a long time they were indistinguishable from one another. Today, most people recognize that astrology is nothing more than an amusing diversion (although millions still study their horoscope in the newspaper every morning!). Nevertheless, the ancient astrological terminology—the names of the constellations and many terms used to describe the locations and motions of the planets—is still used throughout the astronomical world.

Generally speaking, as illustrated in Figure 1.9 for the case of Orion, the stars that make up any particular constellation are not actually close to one another in space, even by astronomical standards. They merely are bright enough to observe with the naked eye and happen to lie in roughly the same direction in the sky as seen from Earth. Still, the constellations provide a convenient means for astronomers to specify regions of the sky, much as geologists use continents or politicians use voting precincts to identify certain localities on planet Earth. Figure 1.10 shows how the conventionally defined constellations cover a portion of the sky in



◀ **FIGURE 1.10 Constellations**

**Near Orion** The region of the sky conventionally associated with the constellation Orion, together with some neighboring constellations (labeled in all capital letters). Some prominent stars are also labeled in lowercase letters. The 88 constellations span the entire sky, so that every astronomical object lies in precisely one of them.

the vicinity of Orion. In all, there are 88 constellations, most of them visible from North America at some time during the year.


## The Celestial Sphere

Over the course of a night, the constellations seem to move smoothly across the sky from east to west, but ancient skywatchers were well aware that the *relative* locations of stars remained unchanged as this nightly march took place.\* It was natural for those observers to conclude that the stars must be firmly attached to a **celestial sphere** surrounding Earth—a canopy of stars resembling an astronomical painting on a heavenly ceiling. Figure 1.11 shows how early astronomers pictured the stars as moving with this celestial sphere as it turned around a fixed, unmoving Earth. Figure 1.12 shows how all stars appear to move in circles around a point very close to the star Polaris (better known as the Pole Star or North Star). To the ancients, this point represented the axis around which the entire celestial sphere turned.

Today we recognize that the apparent motion of the stars is the result of the spin, or **rotation**, not of the celestial sphere, but of Earth. Polaris indicates the direction—due north—in which Earth’s rotation axis points. Even though we now know that the celestial sphere is an incorrect description of the heavens, we still use the idea as a convenient fiction that helps us visualize the positions of stars in the sky. The points where Earth’s axis intersects the celestial sphere are called the


\*We now know that stars do in fact move relative to one another, but this proper motion across the sky is too slow to be discerned with the naked eye (see Section 17.1).

Imagine yourself at the center of this sphere, looking out at the whole sky around you.



**Interactive FIGURE 1.11 Celestial Sphere** Planet Earth sits fixed at the hub of the celestial sphere. This is one of the simplest possible models of the universe, but it doesn't agree with the facts that astronomers now know about the universe.


**celestial poles.** In the Northern Hemisphere, the north celestial pole lies directly above Earth’s North Pole. The extension of Earth’s axis in the opposite direction defines the south celestial pole, directly above Earth’s South Pole. Midway between the north and south celestial poles lies



The duration of this exposure is about 5 hours, . . .

**Interactive FIGURE 1.12 Northern Sky** This time-lapse photograph of the northern sky shows how each star traces out a curved trail across the night sky. The concentric circles are centered near the North Star, Polaris. (AURA)

. . . since each star traces out approximately 20 percent of a circle.



the **celestial equator**, representing the intersection of Earth's equatorial plane with the celestial sphere. These parts of the celestial sphere are marked on Figure 1.11.

When discussing the locations of stars “on the sky,” astronomers naturally talk in terms of *angular* positions and separations. *More Precisely 1-1* presents some basic information on angular measure.

### CONCEPT Check

- ✓ Why do astronomers find it useful to retain the fiction of the celestial sphere to describe the sky? What vital piece of information about stars is lost when we talk about their locations “on” the sky?

## 1.4 Earth's Orbital Motion


### Day-to-Day Changes

We measure time by the Sun. Because the rhythm of day and night is central to our lives, it is not surprising that the period from one noon to the next, the 24-hour **solar day**, is our basic social time unit. The daily progress of the Sun and the other stars across the sky is known as *diurnal motion*. As we have just seen, it is a consequence of Earth's rotation. But the stars' positions in the sky do *not* repeat themselves exactly from one night to the next. Each night, the whole celestial sphere appears to be shifted a little relative to the horizon compared with the night before. The easiest way to confirm this difference is by noticing the stars that are visible just after sunset or just before dawn. You will find that they are in slightly different locations from those of the previous night. Because of this shift, a day measured by the stars—called a **sidereal day** after the Latin word *sidus*, meaning “star”—differs in length from a solar day. Evidently, there is more to the apparent motion of the heavens than simple rotation.

The reason for the difference between a solar day and a sidereal day is sketched in Figure 1.13. It is a result of the fact that Earth moves in two ways simultaneously: It rotates on its central axis while at the same time **revolving** around the Sun. Each time Earth rotates once on its axis, it also moves a small distance along its orbit about the Sun. Earth therefore has to rotate through slightly more than  $360^\circ$  (see *More Precisely 1-1*) for the Sun to return to the same apparent location in the sky. Thus, the interval of time between noon one day and noon the next (a solar day) is slightly greater than one true rotation period (one sidereal day). Our planet takes 365 days to orbit the Sun, so the additional angle is  $360^\circ/365 = 0.986^\circ$ . Because Earth, rotating at a rate of  $15^\circ$  per hour, takes about 3.9 minutes to rotate through this angle, the solar day is 3.9 minutes longer than the sidereal day (i.e., 1 sidereal day is roughly  $23^{\text{h}}56^{\text{m}}$  long).

## Seasonal Changes

Figure 1.14(a) illustrates the major stars visible from most locations in the United States on clear summer evenings. The brightest stars—Vega, Deneb, and Altair—form a conspicuous triangle high above the constellations Sagittarius and Capricornus, which are low on the southern horizon. In the winter sky, however, these stars are replaced as shown in Figure 1.14(b) by several other, well-known constellations, including Orion, Leo, and Gemini.



▲ **FIGURE 1.13 Solar and Sidereal Days** A sidereal day is Earth's true rotation period—the time taken for our planet to return to the same orientation in space relative to the distant stars. A solar day is the time from one noon to the next. The difference in length between the two is easily explained once we understand that Earth revolves around the Sun at the same time as it rotates on its axis. Frames (a) and (b) are 1 sidereal day apart. During that time, Earth rotates exactly once on its axis and also moves a little in its solar orbit—approximately  $1^\circ$ . Consequently, between noon at point A on one day and noon at the same point the next day, Earth actually rotates through about  $361^\circ$  (frame c), and the solar day exceeds the sidereal day by about 4 minutes. Note that the diagrams are not drawn to scale; the true  $1^\circ$  angle is in reality much smaller than shown here.

## MORE PRECISELY 1-1


### Angular Measure

Size and scale are often specified by measuring lengths and angles. The concept of length measurement is fairly intuitive to most of us. The concept of *angular measurement* may be less familiar, but it, too, can become second nature if you remember a few simple facts:

- A full circle contains 360 *degrees* ( $360^\circ$ ). Thus, the half-circle that stretches from horizon to horizon, passing directly overhead and spanning the portion of the sky visible to one person at any one time, contains  $180^\circ$ .
- Each  $1^\circ$  increment can be further subdivided into fractions of a degree, called *arc minutes*. There are 60 arc minutes (written  $60'$ ) in  $1^\circ$ . (The term “arc” is used to distinguish this angular unit from the unit of time.) Both the Sun and the Moon project an angular size of 30 arc minutes (half a degree) on the sky. Your little finger, held at arm’s length, has a similar angular size, covering about a  $40'$  slice of the  $180^\circ$  horizon-to-horizon arc.
- An arc minute can be divided into 60 *arc seconds* ( $60''$ ). Put another way, an arc minute is  $\frac{1}{60}$  of a degree, and an arc second is  $\frac{1}{60} \times \frac{1}{60} = \frac{1}{3600}$  of a degree. An arc second is an extremely small unit of angular measure—the angular size of a centimeter-sized object (a dime, say) at a distance of about 2 kilometers (a little over a mile).

The accompanying figure illustrates this subdivision of the circle into progressively smaller units.

Don’t be confused by the units used to measure angles. Arc minutes and arc seconds have nothing to do with the measurement of time, and degrees have nothing to do with temperature. Degrees, arc minutes, and arc seconds are simply ways to measure the size and position of objects in the universe.



The angular size of an object depends both on its actual size and on its distance from us. For example, the Moon at its present distance from Earth has an angular diameter of  $0.5^\circ$ , or  $30'$ . If the Moon were twice as far away, it would appear half as big— $15'$  across—even though its actual size would be the same. Thus, *angular size by itself is not enough to determine the actual diameter of an object—the distance to the object must also be known.* We return to this topic in more detail in *More Precisely 1-2*.


In the constellation Canis Major lies Sirius (the Dog Star), the brightest star in the sky. Year after year, the same stars and constellations return, each in its proper season. Every winter evening, Orion is high overhead; every summer, it is gone. (For more detailed maps of the sky at different seasons, consult the star charts at the end of the book.)

These regular seasonal changes occur because of Earth’s **revolution** around the Sun: Earth’s darkened hemisphere faces in a slightly different direction in space each evening. The change in direction is only about  $1^\circ$  per night (Figure 1.13)—too small to be easily noticed with the naked eye from one evening to the next, but clearly noticeable over the course of weeks and months, as illustrated in Figure 1.15. After 6 months, Earth has reached the opposite side of its orbit, and we face an entirely different group of stars and constellations at night. Because

of this motion, the Sun appears (to an observer on Earth) to move relative to the background stars over the course of a year. This apparent motion of the Sun on the sky traces out a path on the celestial sphere known as the **ecliptic**.

The 12 constellations through which the Sun passes as it moves along the ecliptic—that is, the constellations we would see looking in the direction of the Sun if they weren’t overwhelmed by the Sun’s light—had special significance for astrologers of old. These constellations are collectively known as the **zodiac**.


As illustrated in Figure 1.16, the ecliptic forms a great circle on the celestial sphere, inclined at an angle of  $23.5^\circ$  to the celestial equator. In reality, as illustrated in Figure 1.17, the plane of the ecliptic is *the plane of Earth’s orbit around the Sun*. Its tilt is a consequence of the *inclination* of our planet’s rotation axis to the plane of its orbit.



**▲ FIGURE 1.14 Typical Night Sky** (a) A typical summer sky above the United States. Some prominent stars (labeled in lowercase letters) and constellations (labeled in all capital letters) are shown. (b) A typical winter sky above the United States.


The point on the ecliptic where the Sun is at its northmost point above the celestial equator is known as the **summer solstice** (from the Latin words *sol*, meaning “sun,” and *stare*, “to stand”). As indicated in Figure 1.17, it represents the location in Earth’s orbit where our planet’s

North Pole comes closest to pointing in the direction of the Sun. This occurs on or near June 21—the exact date varies slightly from year to year because the actual length of a year is not a whole number of days. As Earth rotates, points north of the equator spend the greatest fraction of



**Interactive FIGURE 1.15 The Zodiac** The night side of Earth faces a different set of constellations at different times of the year. The 12 constellations named here make up the astrological zodiac. The arrows indicate the most prominent zodiacal constellations in the night sky at various times of the year. For example, in June, when the Sun is “in” Gemini, Sagittarius and Capricornus are visible at night.






◀ **FIGURE 1.16 Ecliptic** The seasons result from the changing height of the Sun above the celestial equator. At the summer solstice, the Sun is at its northernmost point on its path around the ecliptic; it is therefore highest in the sky, as seen from Earth's Northern Hemisphere, and the days are longest. The reverse is true at the winter solstice. At the vernal and autumnal equinoxes, when the Sun crosses the celestial equator, day and night are of equal length.

The tilt of Earth's rotation axis relative to the ecliptic is responsible for the **seasons** we experience—the marked difference in temperature between the hot summer and cold winter months. As illustrated in


Figure 1.17, two factors combine to cause this variation. First, there are more hours of daylight during the summer than in winter. To see why this is, look at the yellow lines on the surfaces of the drawings of Earth in the figure. (For definiteness, they correspond to a latitude of 45 degrees—roughly that of the Great Lakes or the south of France.) A much larger fraction of the line is sunlit in the summertime, and more daylight means more solar heating. Second, as illustrated in the insets in Figure 1.17, when the Sun is high in the sky in summer, rays of sunlight striking Earth's surface are more concentrated—spread out over a smaller area—than in winter. As a result, the Sun feels hotter. Therefore summer, when the Sun is highest above the horizon and the days are longest, is generally much warmer than winter, when the Sun is low and the days are short.

their time in sunlight on that date, so the summer solstice corresponds to the longest day of the year in the Northern Hemisphere and the shortest day in the Southern Hemisphere.

Six months later, the Sun is at its southernmost point below the celestial equator (Figure 1.16)—or, equivalently, the North Pole points farthest from the Sun (Figure 1.17). We have reached the **winter solstice** (December 21), the shortest day in Earth's Northern Hemisphere and the longest in the Southern Hemisphere.



▶ **Interactive FIGURE 1.17 Seasons** Earth's seasons result from the inclination of our planet's rotation axis with respect to its orbit plane. The summer solstice corresponds to the point on Earth's orbit where our planet's North Pole points most nearly toward the Sun. The opposite is true of the winter solstice. The vernal and autumnal equinoxes correspond to the points in Earth's orbit where our planet's axis is perpendicular to the line joining Earth and the Sun. The insets show how rays of sunlight striking the ground at an angle (e.g., during northern winter) are spread over a larger area than rays coming nearly straight down (e.g., during northern summer). As a result, the amount of solar heat delivered to a given area of Earth's surface is greatest when the Sun is high in the sky.



▲ **FIGURE 1.18 Earth's Orbit** Seen face on, Earth's orbit around the Sun is almost a perfect circle. The distance from Earth to the Sun varies only slightly over the course of a year and is *not* the cause of the seasonal temperature changes we experience on our planet.

A popular misconception is that the seasons have something to do with Earth's distance from the Sun. Figure 1.18 illustrates why this is *not* the case. It shows Earth's orbit “face on,” instead of almost edge-on, as in Figure 1.17. Notice that the orbit is almost perfectly circular, so the distance from Earth to the Sun varies very little (in fact, by only about 3 percent) over the course of a year—not nearly enough to explain the seasonal changes in temperature. What's more, Earth is actually *closest* to the Sun in early January, the dead of winter in the Northern Hemisphere, so distance from the Sun cannot be the main factor controlling our climate.

The two points where the ecliptic intersects the celestial equator (Figure 1.16)—that is, where Earth's rotation axis is perpendicular to the Earth-Sun line (Figure 1.17)—are known as **equinoxes**. On those dates, day and night are of equal duration. (The word *equinox* derives from the Latin for “equal night.”) In the fall (in the Northern Hemisphere), as the Sun crosses from the Northern into the Southern Hemisphere, we have the **autumnal equinox** (on September 21). The **vernal equinox** occurs in northern spring, on or near March 21, as the Sun crosses the celestial equator moving north. Because of its association with the end of winter and the start of a new growing season, the vernal equinox was particularly important to early astronomers and astrologers. It also plays an important role in human timekeeping: The interval of time from one vernal

equinox to the next—365.2422 mean solar days—is 1 **tropical year**.

## Long-Term Changes


Earth has many motions—it spins on its axis, it travels around the Sun, and it moves with the Sun through our Galaxy. We have just seen how some of these motions can account for the changing nighttime sky and the changing seasons. In fact, the situation is even more complicated. Like a spinning top that rotates rapidly on its own axis while that axis slowly revolves about the vertical, Earth's axis changes its *direction* over the course of time (although the angle between the axis and a line perpendicular to the plane of the ecliptic always remains close to 23.5°). Illustrated in Figure 1.19, this change is called **precession**. It is caused by torques (twisting forces) on Earth due to the gravitational pulls of the Moon and the Sun, which affect our planet in much the same way as the torque due to Earth's own gravity affects a top. During a complete cycle of precession—about 26,000 years—Earth's axis traces out a cone.

The time required for Earth to complete exactly one orbit around the Sun, relative to the stars, is called a **sidereal year**. One sidereal year is 365.256 mean solar days long—about 20 minutes longer than a tropical year. The reason for this slight difference is Earth's precession. Recall that the vernal equinox occurs when Earth's rotation axis is perpendicular to the line joining Earth and the Sun, and the Sun is crossing the celestial equator moving from south to north. In the absence of precession, this would occur exactly once per sidereal orbit, and the tropical and sidereal years would be identical. However, because of the slow precessional shift in the orientation of Earth's rotation axis, the instant when the axis is next perpendicular to the line from Earth to the Sun occurs slightly *sooner* than we would otherwise expect. Consequently, the vernal equinox drifts slowly westward (“backwards”) around the zodiac over the course of the precession cycle.

The tropical year is the year that our calendars measure. If our timekeeping were tied to the sidereal year, the seasons would slowly march around the calendar as Earth precessed—13,000 years from now, summer in the Northern Hemisphere would be at its height in late February! By using the tropical year, we ensure that July and August will always be (northern) summer months. However, in 13,000 years' time, Orion will be a summer constellation.

### CONCEPT Check

- ✓ In astronomical terms, what are *summer* and *winter*, and why do we see different constellations during those seasons?



**Interactive FIGURE 1.19 Precession** (a) Earth's axis currently points nearly toward the star **MA** Polaris. About 12,000 years from now—almost halfway through one cycle of precession—Earth's axis will point toward a star called Vega, which will then be the "North Star." Five thousand years ago, the North Star was a star named Thuban in the constellation Draco. (b) The yellow circle shows the precessional path of the north celestial pole among some prominent northern stars. Tick marks indicate intervals of a thousand years.

## 1.5 The Motion of the Moon

The Moon is our nearest neighbor in space. Apart from the Sun, it is the brightest object in the sky. Like the Sun, the Moon appears to move relative to the background stars. Unlike the Sun, however, the Moon really does revolve around Earth. It crosses the sky at a rate of about  $12^\circ$  per day, moving through an angular distance equal to its own diameter—30 arc minutes—in about an hour.


### Lunar Phases

The Moon's appearance undergoes a regular cycle of changes, or **phases**, taking roughly 29.5 days to complete. Figure 1.20 illustrates the appearance of the Moon at different times in this monthly cycle. Starting from the *new Moon*, which is all but invisible in the sky, the Moon appears to *wax* (or grow) a little each night and is visible as a growing *crescent* (photo 1 of Figure 1.20). One week after new Moon, half of the lunar disk can be seen (photo 2). This phase is known as a *quarter Moon*. During the next week, the Moon continues to wax, passing through the *gibbous* phase (photo 3) until, 2 weeks after new Moon, the *full Moon* (photo 4) is visible. During the next 2 weeks, the Moon *wanes* (or shrinks), passing in turn through the gibbous, quarter, crescent phases (photos 5–7) and eventually becoming new again.

The position of the Moon in the sky relative to the Sun, as seen from Earth, varies with lunar phase. For example, the full Moon rises in the east as the Sun sets in the west, while the first quarter Moon actually rises at noon, but may become visible only late in the day as the Sun's light fades and the Moon is already high in the sky. Some connections between the lunar phase and the rising and setting times of the Moon are indicated in Figure 1.20.

The Moon doesn't actually change its size and shape from night to night, of course. Its full circular disk is present at all times. Why, then, don't we always see a full Moon? The answer is that, unlike the Sun and the other stars, the Moon emits no light of its own. Instead, it shines by reflected sunlight. As illustrated in Figure 1.20, half of the Moon's surface is illuminated by the Sun at any instant. However, not all of the Moon's sunlit face can be seen because of the Moon's position with respect to Earth and the Sun. When the Moon is full, we see the entire "daylit" face because the Sun and the Moon are in opposite directions from Earth in the sky. In the case of a new Moon, the Moon and the Sun are in almost the same part of the sky, and the sunlit side of the Moon is oriented away from us. At new Moon, the Sun must be almost behind the Moon, from our perspective.

As the Moon revolves around Earth, our satellite's position in the sky changes with respect to the stars. In 1 **sidereal month** (27.3 days), the Moon completes one revolution and




**MA** **Narrated FIGURE 1.20 Lunar Phases** Because the Moon orbits Earth, the visible fraction of the lunar sunlit face varies from night to night, although the Moon always keeps the same face toward our planet. (Note the location of the small, straight arrows, which mark the same point on the lunar surface at each phase shown.) The complete cycle of lunar phases, shown here starting at the waxing crescent phase and following the Moon's orbit counterclockwise, takes 29.5 days to complete. Rising and setting times for some phases are also indicated. (UC/Lick Observatory)

returns to its starting point on the celestial sphere, having traced out a great circle in the sky. The time required for the Moon to complete a full cycle of phases, 1 **synodic month**, is a little longer—about 29.5 days. The synodic month is a little longer than the sidereal month for the same reason that a solar day is slightly longer than a sidereal day: Because of Earth's motion around the Sun, the Moon must complete slightly more than one full revolution to return to the same phase in its orbit (Figure 1.21).

## Eclipses

From time to time—but only at new or full Moon—the Sun and the Moon line up precisely as seen from Earth, and we observe the spectacular phenomenon known as an **eclipse**. When the Sun and the Moon are in exactly *opposite* directions, as seen from Earth, Earth's shadow sweeps across the Moon, temporarily blocking the Sun's light and darkening the Moon in a **lunar eclipse**, as illustrated in Figure 1.22.




**Interactive MA** **FIGURE 1.21 Sidereal Month** The difference between a *synodic* and a *sidereal* month stems from the motion of Earth relative to the Sun. Because Earth orbits the Sun in 365 days, in the 29.5 days from one new Moon to the next (1 synodic month), Earth moves through an angle of approximately 29°. Thus, the Moon must revolve more than 360° between new Moons. The sidereal month, which is the time taken for the Moon to revolve through exactly 360°, relative to the stars, is about 2 days shorter.

From Earth, we see the curved edge of Earth’s shadow cut into the face of the full Moon and slowly eat its way into the lunar disk. Usually, the alignment of the Sun, Earth, and Moon is imperfect, so the shadow never completely covers the Moon. Such an occurrence is known as a **partial lunar eclipse**. Occasionally, however, the entire lunar surface is obscured in a **total lunar eclipse**, such as that shown in the inset of Figure 1.22. Total lunar eclipses last only as long as is needed for the Moon to pass through Earth’s shadow—no more than about 100 minutes. During that time, the Moon often acquires an eerie, deep red coloration—the result of a small amount of sunlight reddened by Earth’s atmosphere (for the same reason that sunsets appeared—see *More Precisely 7-1*) and refracted (bent) onto the lunar surface, preventing the shadow from being completely black.

When the Moon and the Sun are in exactly the *same* direction, as seen from Earth, an even more awe-inspiring event occurs. The Moon passes directly in front of the Sun, briefly turning day into night in a **solar eclipse**. In a *total solar eclipse*, when the alignment is perfect, planets and some stars become visible in the daytime as the Sun’s light is reduced to nearly nothing. We can also see the Sun’s ghostly outer atmosphere, or *corona* (Figure 1.23).<sup>\*</sup> In a *partial solar eclipse*, the Moon’s path is slightly “off center,” and only a portion of the Sun’s face is covered. In either case, the sight of the Sun apparently being swallowed up by the black disk of the Moon is disconcerting even today. It must surely have inspired fear in early observers. Small wonder that the ability to predict such events was a highly prized skill.

Unlike a lunar eclipse, which is simultaneously visible from all locations on Earth’s night side, a total solar eclipse



<sup>\*</sup>Actually, although a total solar eclipse is undeniably a spectacular occurrence, the visibility of the corona is probably the most important astronomical aspect of such an event today. It enables us to study this otherwise hard-to-see part of our Sun (see Chapter 16).

**Interactive MA** **FIGURE 1.22 Lunar Eclipse** When the Moon passes through Earth’s shadow, we see a darkened, copper-colored Moon, as shown by the partial eclipse in the inset photograph. The red coloration is caused by sunlight deflected by Earth’s atmosphere onto the Moon’s surface. (Inset: G. Schneider)

*This is an actual photo of the eclipsed Moon, one of the great light shows visible to the naked eye.*




▲ **FIGURE 1.23 Total Solar Eclipse** During a total solar eclipse, the Sun's corona becomes visible as an irregularly shaped halo surrounding the blotted-out disk of the Sun. This was the August 1999 eclipse, as seen from the banks of the Danube River near Sofia, Bulgaria. (B. Angelov)

can be seen from only a small portion of Earth's daytime side. The Moon's shadow on Earth's surface is about 7000 kilometers wide—roughly twice the diameter of the Moon. Outside of that shadow, no eclipse is seen. However, within the central region of the shadow, called the **umbra**, the eclipse is total. Within the shadow, but outside the umbra, in the **penumbra**, the eclipse is partial, with less and less of the Sun obscured the farther one travels from the shadow's center.

The connections among the umbra, the penumbra, and the relative locations of Earth, Sun, and Moon are illustrated in Figure 1.24. The umbra is always very small. Even under the most favorable circumstances, its diameter never exceeds 270 kilometers. Because the shadow sweeps across Earth's surface at over 1700 kilometers per hour, the duration of a total eclipse at any given point on our planet can never exceed 7.5 minutes.

The Moon's orbit around Earth is not exactly circular. Thus, the Moon may be far enough from Earth at the moment of an eclipse that its disk fails to fully cover the disk of the Sun, even though their centers coincide. In that case, there is no region of totality—the umbra never reaches Earth, and a thin ring of sunlight can be seen surrounding the Moon. Such an occurrence, called an **annular eclipse**, is illustrated in Figure 1.24(c) and shown more clearly in Figure 1.25. Roughly half of all solar eclipses are annular.



▶ **Interactive FIGURE 1.24**  
**Types of Solar Eclipse**

MA (a) The Moon's shadow consists of two parts: the umbra, where no sunlight is seen, and the penumbra, where a portion of the Sun is visible. (b) If we are in the umbra, we see a total eclipse; in the penumbra, we see a partial eclipse. (c) If the Moon is too far from Earth at the moment of the eclipse, the umbra does not reach Earth and there is no region of totality; instead, an annular eclipse is seen. (Note that these figures are not drawn to scale.) (Insets: NOAA; G. Schneider)



▲ **FIGURE 1.25 Annular Solar Eclipse** During an annular solar eclipse, the Moon fails to completely hide the Sun, so a thin ring of light remains. No corona is seen in this case because even the small amount of the Sun still visible completely overwhelms the corona's faint glow. This was the December 1973 eclipse, as seen from Algiers. (The gray fuzzy areas at the top left and right are clouds in Earth's atmosphere.) (G. Schneider)

## Eclipse Seasons

Why isn't there a solar eclipse at every new Moon and a lunar eclipse at every full Moon? That is, why doesn't the Moon pass directly between Earth and the Sun once per orbit and directly through Earth's shadow 2 weeks later?

The answer is that the Moon's orbit is slightly inclined to the ecliptic (at an angle of  $5.2^\circ$ ), so the chance that a new (or full) Moon will occur just as the Moon happens to cross the plane of the ecliptic (with Earth, Moon, and Sun perfectly aligned) is quite low. Figure 1.26 illustrates some possible configurations of the three bodies. If the Moon happens to lie above or below the plane of the ecliptic when new (or full), a solar (or lunar) eclipse cannot occur. Such a configuration is termed *unfavorable* for producing an eclipse. In a *favorable* configuration, the Moon is new or full just as it crosses the plane of the ecliptic, and eclipses are seen. Unfavorable configurations are much more common, so eclipses are relatively rare events.

As indicated on Figure 1.26(b), the two points on the Moon's orbit where it crosses the plane of the ecliptic are known as the *nodes* of the orbit. The line joining the nodes, which is also the line of intersection of Earth's and the Moon's orbital planes, is known as the *line of nodes*. When the line of nodes is not directed toward the Sun, conditions are unfavorable for


eclipses. However, when the line of nodes briefly lies along the Earth–Sun line, eclipses are possible. These two periods, known as **eclipse seasons**, are the only times at which an eclipse *will* occur. Notice that there is no guarantee that an eclipse *will* occur. For a solar eclipse, we must have a new Moon during an eclipse season. Similarly, a lunar eclipse can occur only at full Moon during an eclipse season.

Because we know the orbits of Earth and the Moon to great accuracy, we can predict eclipses far into the future. Figure 1.27 shows the location and duration of all total eclipses of the Sun between 2010 and 2030. Note that the eclipse tracks run from west to east—just the opposite of more familiar phenomena such as sunrise and sunset, which are seen earlier by observers located farther east. The reason is that the Moon's shadow sweeps across Earth's surface faster than our planet rotates, so the eclipse actually *overtakes* observers on the ground.


The solar eclipses that we do see highlight a remarkable cosmic coincidence. Although the Sun is many times farther away from Earth than is the Moon, it is also much larger. In fact, the ratio of distances is almost exactly the same as the ratio of sizes, so the Sun and the Moon both have roughly the *same* angular diameter—about half a degree, seen from Earth. Thus, the Moon covers the face of the Sun almost exactly. If the Moon were larger, we would never see annular eclipses, and total eclipses would be much more common. If the Moon were a little smaller, we would see only annular eclipses.

The gravitational tug of the Sun causes the Moon's orbital orientation, and hence the direction of the line of nodes, to change slowly with time. As a result, the time between one orbital configuration with the line of nodes pointing at the Sun and the next (with the Moon crossing the ecliptic in the same sense in each case) is not exactly 1 year, but instead is 346.6 days—sometimes called 1 *eclipse year*. Thus, the eclipse seasons gradually progress backward through the calendar, occurring about 19 days earlier each year. For example, in 1999 the eclipse seasons were in February and August, and on August 11 much of Europe and southern Asia was treated to the last total eclipse of the millennium (Figure 1.23). By 2002, those seasons had drifted into December and June, and eclipses actually occurred on June 10 and December 4 of that year. By studying Figure 1.27, you can follow the progression of the eclipse seasons through the calendar.

The combination of the eclipse year and the Moon's synodic period leads to an interesting long-term cycle in solar (and lunar) eclipses. A simple calculation shows that 19 eclipse years is almost exactly 223 lunar months. Thus, every 6585 solar days (actually 18 years, 11.3 days) the "same" eclipse recurs, with Earth, the Moon, and the Sun in the same relative configuration. Several such repetitions are evident in Figure 1.27—see, for example, the similarly shaped July 11, 2010, and July 22, 2028, tracks. (Note that we must



(a)



(b)


**▲ FIGURE 1.26 Eclipse Geometry** (a) An eclipse occurs when Earth, Moon, and Sun are precisely aligned. If the Moon's orbital plane lay in exactly the plane of the ecliptic, this alignment would occur once a month. However, the Moon's orbit is inclined at about  $5^\circ$  to the ecliptic, so not all configurations are favorable for producing an eclipse. (b) For an eclipse to occur, the line of intersection of the two planes must lie along the Earth–Sun line. Thus, eclipses can occur just at specific times of the year. Only the umbra of each shadow is shown, for clarity (see Figure 1.24).

take leap years properly into account to get the dates right!) The roughly  $120^\circ$  offset in longitude corresponds to Earth's rotation in 0.3 day. This recurrence is called the *Saros cycle*. Well known to ancient astronomers, it undoubtedly was the key to their “mystical” ability to predict eclipses!

### CONCEPT Check

- ✓ What types of solar eclipses would you expect to see if Earth's distance from the Sun were to double? What if the distance became half its present value?





▲ **FIGURE 1.27 Eclipse Tracks** Regions of Earth that saw or will see total solar eclipses between the years 2010 and 2030. Each track represents the path of the Moon's umbra across Earth's surface during an eclipse. The width of the track depends upon the latitude on Earth and the distance from Earth to the Moon during the eclipse. High-latitude tracks are broader because sunlight strikes Earth's surface at an oblique angle near the poles (and also because of the projection of the map). The closer the Moon is to Earth during a total eclipse, the wider is the umbra (see Figure 1.24).

## 1.6 The Measurement of Distance


We have seen a little of how astronomers track and record the positions of the stars in the sky. But knowing the direction to an object is only part of the information needed to locate it in space. Before we can make a systematic study of the heavens, we must find a way of measuring *distances*, too. One distance-measurement method, called **triangulation**, is based on the principles of Euclidean geometry and finds widespread application today in both terrestrial and astronomical settings. Surveyors use these age-old geometric ideas to measure the distance to far-away objects indirectly. Triangulation forms the foundation of the family of distance-measurement techniques making up the **cosmic distance scale**.

### Triangulation and Parallax

Imagine trying to measure the distance to a tree on the other side of a river. The most direct method is to lay a tape across the river, but that's not the simplest way (nor, because of the current, may it even be possible). A smart surveyor would

make the measurement by visualizing an *imaginary* triangle (hence *triangulation*), sighting the tree on the far side of the river from two positions on the near side, as illustrated in Figure 1.28. The simplest possible triangle is a right triangle, in which one of the angles is exactly  $90^\circ$ , so it is usually convenient to set up one observation position directly opposite the object, as at point A. The surveyor then moves to another observation position at point B, noting the distance covered between points A and B. This distance is called the **baseline** of the imaginary triangle. Finally, the surveyor, standing at point B, sights toward the tree and notes the angle at point B between this line of sight and the baseline. Knowing the value of one side (AB) and two angles (the right angle at point A and the angle at point B) of the right triangle, the surveyor geometrically constructs the remaining sides and angles and establishes the distance from A to the tree.


To use triangulation to measure distances, a surveyor uses *trigonometry*, the mathematics of geometrical angles and distances. However, even knowing no trigonometry at all, we can still solve the problem by graphical means, as shown in Figure 1.29. Suppose that we pace off the baseline AB, measuring it to be 450 meters, and measure the angle between the



▲ **FIGURE 1.28 Triangulation** Surveyors often use simple geometry and trigonometry to estimate the distance to a faraway object by triangulation. By measuring the angles at A and B and the length of the baseline, the distance can be calculated without the need for direct measurement.

baseline and the line from B to the tree to be  $52^\circ$ , as illustrated in the figure. We can transfer the problem to paper by letting one box on our graph represent 25 meters on the ground. Drawing the line AB on paper and completing the other two sides of the triangle, at angles of  $90^\circ$  (at A) and  $52^\circ$  (at B), we measure the distance on paper from A to the tree to be 23 boxes—that is, 575 meters. We have solved the real problem by *modeling* it on paper. The point to remember here is this: Nothing more complex than basic geometry is needed to infer the distance, the size, and even the shape of an object that is too far away or inaccessible for direct measurement.

Obviously, for a fixed baseline the triangle becomes longer and narrower as the tree's distance from A increases. Narrow triangles cause problems, because it becomes hard to measure the angles at A and B with sufficient accuracy. The measurements can be made easier by "fattening" the triangle—that is, by lengthening the baseline—but there are limits on how long a baseline we can choose in astronomy. For example, consider an imaginary triangle extending from Earth to a nearby object in space, perhaps a neighboring planet. The triangle is now extremely long and narrow, even for a relatively nearby object (by cosmic standards). Figure 1.30(a) illustrates a case in which the longest baseline possible on Earth—Earth's diameter, measured from point A to point B—is used.




▲ **FIGURE 1.29 Geometric Scaling** Not even trigonometry is needed to estimate distances indirectly. Scaled estimates, like this one on a piece of graph paper, often suffice.

In principle, two observers could sight the planet from opposite sides of Earth, measuring the triangle's angles at A and B. However, in practice it is easier to measure the third angle of the imaginary triangle. The observers sight toward the planet, taking note of its position *relative to some distant stars* seen on the plane of the sky. The observer at point A sees the planet at apparent location A' relative to those stars, as indicated in Figure 1.30(a). The observer at B sees the planet at point B'. If each observer takes a photograph of the appropriate region of the sky, the planet will appear at slightly different places in the two images. The planet's position is slightly displaced, or shifted, relative to the field of distant background stars, as shown in Figure 1.30(b). The background stars themselves appear undisplaced because of their much greater distance from the observer.


This apparent displacement of a foreground object relative to the background as the observer's location changes is known as **parallax**. The size of the shift in Figure 1.30(b), measured as an angle on the celestial sphere, is the third, small angle in Figure 1.30(a). In astronomical contexts, the parallax is usually very small. For example, the parallax of a point on the Moon, viewed using a baseline equal to Earth's diameter, is about  $2^\circ$ ; the parallax of the planet Venus at closest approach (45 million kilometers), is just  $1'$  (see *More Precisely 1-2*).

The closer an object is to the observer, the larger is the parallax. Figure 1.31 illustrates how you can see this for yourself. Hold a pencil vertically in front of your nose and concentrate on some far-off object—a distant wall, perhaps.



**▲ FIGURE 1.30 Parallax** (a) An imaginary triangle extending from Earth to a nearby object in space. The group of stars at the top represents a background field of very distant stars. (b) Hypothetical photographs of the same star field showing the nearby object's apparent displacement, or shift, relative to the distant undisplaced stars.

Close one eye, and then open it while closing the other. You should see a large shift in the apparent position of the pencil projected onto the distant wall—a large parallax. In this example, one eye corresponds to point A, the other eye to point B, the distance between your eyeballs to the baseline, the pencil to the planet, and the distant wall to a remote field of stars. Now hold the pencil at arm's length, corresponding to a more distant object (but still not as far away as the even more distant stars). The apparent shift of the pencil will be less. You might even be able to verify that the apparent shift



**▲ FIGURE 1.31 Parallax Geometry** Parallax is inversely proportional to an object's distance. An object near your nose has a much larger parallax than an object held at arm's length.

is inversely proportional to the distance to the pencil. By moving the pencil farther away, we are narrowing the triangle and decreasing the parallax (and also making accurate measurement more difficult). If you were to paste the pencil to the wall, corresponding to the case where the object of interest is as far away as the background star field, blinking would produce no apparent shift of the pencil at all.

The amount of parallax is thus inversely proportional to an object's distance. Small parallax implies large distance, and large parallax implies small distance. Knowing the amount of parallax (as an angle) and the length of the baseline, we can easily derive the distance through triangulation. *More Precisely 1-2* explores the connection between angular measure and distance in more detail, showing how we can use elementary geometry to determine both the distances and the dimensions of far away objects.

Surveyors of the land use these simple geometric techniques to map out planet Earth. As surveyors of the sky, astronomers use the same basic principles to chart the universe.

## Sizing Up Planet Earth

Now that we have studied some of the tools available to astronomers, let's end the chapter with a classic example of how the scientific method, combined with the basic geometric techniques just described, enabled an early scientist to perform a calculation of truly “global” proportions.

In about 200 B.C., a Greek philosopher named Eratosthenes (276–194 B.C.) used simple geometric reasoning to calculate the size of our planet. He knew that at noon on the first day of summer observers in the city of Syene (now called Aswan) in Egypt saw the Sun pass directly overhead. This was evident from the fact that vertical objects cast no shadows and sunlight reached to the very bottoms of deep wells, as shown in the insets in Figure 1.32. However, at noon of the same day in Alexandria, a city 5000 *stadia* to the north, the Sun was seen to be displaced slightly from the vertical. (The *stadium* was a Greek unit of length, roughly equal to 0.16 km—the modern town of Aswan lies about 780 km, or 490 miles, south of Alexandria.) By measuring the length of the shadow of a vertical stick and applying elementary trigonometry, Eratosthenes determined the angular displacement of the Sun from the vertical at Alexandria to be 7.2°.

What could have caused this discrepancy between the two measurements? It was not the result of measurement error—the same results were obtained every time the observations were repeated. Instead, as illustrated in Figure 1.32, the explanation is simply that Earth's surface is not flat, but *curved*. Our planet is a sphere. Eratosthenes was not the first person to realize that Earth is spherical—the philosopher Aristotle had done that over 100 years earlier (see Section 1.2),—but he was apparently the first to build on this knowledge, combining geometry with direct measurement to infer the size of our planet. Here's how he did it.

Rays of light reaching Earth from a very distant object, such as the Sun, travel almost parallel to one another. Consequently, as shown in the figure, the angle measured at Alexandria between the Sun's rays and the vertical (i.e., the line joining Alexandria to the center of Earth) is equal to the angle between Syene and Alexandria, as seen from Earth's center. (For the sake of clarity, the angle has been exaggerated in the figure.) As discussed in *More Precisely 1-2*, the size of

this angle in turn is proportional to the fraction of Earth's circumference that lies between Syene and Alexandria:


$$\frac{7.2^\circ \text{ (angle between Syene and Alexandria)}}{360^\circ \text{ (circumference of a circle)}} = \frac{5000 \text{ stadia}}{\text{Earth's circumference}}$$

Earth's circumference is therefore  $50 \times 5000$ , or 250,000 stadia, or about 40,000 km, so Earth's radius is  $250,000/2\pi$  stadia, or 6366 km. The correct values for Earth's circumference and radius, now measured accurately by orbiting spacecraft, are 40,070 km and 6378 km, respectively.

Eratosthenes' reasoning was a remarkable accomplishment. More than 20 centuries ago, he estimated the circumference of Earth to within 1 percent accuracy, using only simple geometry and basic scientific reasoning. A person making measurements on only a small portion of Earth's surface was able to compute the size of the entire planet on the basis of observation and pure logic—an early triumph of the scientific method.

### CONCEPT Check

- ✓ Why is elementary geometry essential for measuring distances in astronomy?




► **FIGURE 1.32 Measuring Earth's Radius** The Sun's rays strike different parts of Earth's surface at different angles. The Greek philosopher Eratosthenes realized that the difference was due to Earth's curvature, enabling him to determine Earth's radius by using simple geometry.

## MORE PRECISELY 1-2

### Measuring Distances with Geometry

Simple geometrical reasoning forms the basis for almost every statement made in this book about size and scale in the universe. In a very real sense, our modern knowledge of the cosmos depends on the elementary mathematics of ancient Greece. Let's take a moment to look in a little more detail at how astronomers use geometry to measure the distances to, and sizes of, objects near and far.

We can convert baselines and parallaxes into distances, and vice versa, by using arguments made by the Greek geometer Euclid. The first figure represents Figure 1.30(a), but we have changed the scale and added the circle centered on the target planet and passing through our baseline on Earth:



To see how the planet's parallax relates to its distance, we note that the ratio of the baseline  $AB$  to the circumference of the large circle shown in the figure must be equal to the ratio of the parallax to one full revolution,  $360^\circ$ . Recall that the circumference of a circle is always  $2\pi$  times its radius (where  $\pi$ —the Greek letter “pi”—is approximately equal to 3.142). Applying this relation to the large circle in the figure, we find that

$$\frac{\text{baseline}}{2\pi \times \text{distance}} = \frac{\text{parallax}}{360^\circ},$$

from which it follows that

$$\text{parallax} = (360^\circ/2\pi) \times \frac{\text{baseline}}{\text{distance}}.$$

The angle  $360^\circ/2\pi \approx 57.3^\circ$  in the preceding equation is usually called 1 *radian*.


**EXAMPLE 1** The planet Venus lies roughly 45,000,000 km from Earth at closest approach. Two observers 13,000 km apart (i.e., at opposite ends of Earth's diameter) looking at the planet would measure a parallax of  $57.3^\circ \times (13,000 \text{ km}/45,000,000 \text{ km}) = 0.017^\circ = 1.0$  arc minutes, as stated in the text.

Alternatively, if we know the parallax (from direct measurement, such as the photographic technique described in Section 1.6), we can rearrange the above equation to tell us the distance to the planet:

$$\text{distance} = \text{baseline} \times \frac{57.3^\circ}{\text{parallax}}.$$

**EXAMPLE 2** Two observers 1000 km apart looking at the Moon might measure a parallax of 9.0 arc minutes—that is,  $0.15^\circ$ . It then follows that the distance to the Moon is  $1000 \text{ km} \times (57.3/0.15) \approx 380,000 \text{ km}$ . (More accurate measurements, based on laser ranging using equipment left on the lunar surface by *Apollo* astronauts, yield a mean distance of 384,000 km.)

Knowing the distance to an object, we can determine many other properties. For example, by measuring the object's *angular diameter*—the angle from one side of the object to the other as we view it in the sky—we can compute its size. The second figure illustrates the geometry involved:



Notice that this is basically the same diagram as the previous one, except that now the angle (the angular diameter) and distance are known, instead of the angle (the parallax) and baseline. Exactly the same reasoning as before then allows us to calculate the diameter. We have

$$\frac{\text{diameter}}{2\pi \times \text{distance}} = \frac{\text{angular diameter}}{360},$$

so

$$\text{diameter} = \text{distance} \times \frac{\text{angular diameter}}{57.3^\circ}.$$

**EXAMPLE 3** The Moon's angular diameter is measured to be about 31 arc minutes—a little over half a degree. From the preceding discussion, it follows that the Moon's actual diameter is  $380,000 \text{ km} \times (0.52^\circ/57.3^\circ) \approx 3450 \text{ km}$ . A more precise measurement gives 3476 km.


Study the foregoing reasoning carefully. We will use these simple arguments, in various forms, many times throughout this text.

**The Big Question** Take another look at the spectacular photo at the beginning of this chapter. Contemplate for a moment all those stars—about 100,000,000,000 in our Galaxy alone—of which our Sun is just one. We cannot help wondering: Are there planets around some of those stars, and perhaps intelligent beings on some of those planets? One of the grandest of all unsolved questions in astronomy concerns life on other worlds. No one yet knows the answer, but we will return to this fascinating topic in the last chapter.


# Chapter Review

## SUMMARY


**1** The **universe** (p. 6) is the totality of all space, time, matter, and energy. **Astronomy** (p. 6) is the study of the universe. In order of increasing size, the basic constituents of the cosmos are planets, stars, galaxies, galaxy clusters, and the universe itself. They differ enormously in scale—a factor of a billion billion from planet Earth to the entire observable universe.



**2** The **scientific method** (p. 8) is a methodical approach employed by scientists to explore the universe around us in an objective manner. A **theory** (p. 8) is a framework of ideas and assumptions used to explain some set of observations and construct **theoretical models** (p. 8) that make predictions about the real world. These predictions in turn are amenable to further observational testing. In this way, the theory expands and science advances.



**3** Early observers grouped the thousands of stars visible to the naked eye into patterns called **constellations** (p. 10), which they imagined were attached to a vast **celestial sphere** (p. 12) centered on Earth. Constellations have no physical significance, but are still used to label regions of the sky. The points where Earth's axis of rotation intersects the celestial sphere are called the north and




south **celestial poles** (p. 12). The line where Earth's equatorial plane cuts the celestial sphere is the **celestial equator** (p. 13).


**4** The nightly motion of the stars across the sky is the result of Earth's **rotation** (p. 12) on its axis. The time from one noon to the next is called a **solar day** (p. 13). The time between successive risings of any given star is 1 **sidereal day** (p. 13). Because of Earth's **revolution** (p. 14) around the Sun, we see different stars at night at different times of the year, and the Sun appears to move relative to the stars. The Sun's apparent yearly path around the celestial sphere (or the plane of Earth's orbit around the Sun) is called the **ecliptic** (p. 14).



**5** We experience **seasons** (p. 16) because Earth's rotation axis is inclined to the ecliptic plane. At the **summer solstice** (p. 15), the Sun is highest in the sky and the length of the day is greatest. At the **winter solstice** (p. 16), the Sun is lowest and the day is shortest. At the **vernal** (p. 17) and **autumnal equinoxes** (p. 17), Earth's axis of rotation is perpendicular to the line joining Earth to the Sun, so day and night are of equal length. Because of **precession** (p. 17), the slow "wobble" of Earth's axis due to the influence of the Moon, the orientation of Earth's axis changes slowly over time. As a result, the particular constellations visible during any given season change over the course of thousands of years.




**6** The Moon emits no light of its own, but instead shines by reflected sunlight. As the Moon orbits Earth, we see **lunar phases** (p. 18) as the amount of the Moon's sunlit face visible to us varies. A **lunar eclipse** (p. 19) occurs when the Moon enters Earth's shadow.



A **solar eclipse** (p. 20) occurs when the Moon passes between Earth and the Sun. An eclipse may be **total** (p. 21) if the body in question (Moon or Sun) is completely obscured, or **partial** (p. 21) if only a portion of the surface is affected. If the Moon happens to be too far from Earth for its disk to completely hide the Sun, an **annular eclipse** (p. 21) occurs. Because the Moon's orbit around Earth is slightly inclined with respect to the ecliptic, solar and lunar eclipses are relatively rare events.

**7** Astronomers use **triangulation** (p. 24) to measure the distances to planets and stars, forming the foundation of the

**cosmic distance scale** (p. 24), the family of distance-measurement techniques used to chart the universe. **Parallax** (p. 25) is the apparent motion of a foreground object relative to a distant background as the observer's position changes. The larger the **baseline** (p. 24)—the distance between the two observation points—the greater is the parallax. The same basic geometric reasoning is used to determine the sizes of objects whose distances are known.



**MasteringAstronomy**® For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- LO1** Compare the size of Earth with that of the Sun, the Milky Way Galaxy, and the entire universe.
- What does an astronomer mean by “the universe”?
- LO2 POS** What is the scientific method, and how does science differ from religion?
- LO3** What is a constellation? Why are constellations useful for mapping the sky?
- Why does the Sun rise in the east and set in the west each day? Does the Moon also rise in the east and set in the west?
- LO4** How and why does a day measured with respect to the Sun differ from a day measured with respect to the stars?
- Why do we see different stars at different times of the year?
- LO5** Why are there seasons on Earth?
- What is precession, and what causes it?
- If one complete hemisphere of the Moon is always lit by the sun, why do we see different phases of the Moon?
- LO6** What causes a lunar eclipse? A solar eclipse? Why aren't there lunar and solar eclipses every month?
- POS** Do you think an observer on another planet in the solar system might see eclipses? Why or why not?
- What is parallax? Give an everyday example.
- Why is it necessary to have a long baseline when using triangulation to measure the distances to objects in space?
- LO7** What two pieces of information are needed to determine the diameter of a faraway object?

## Conceptual Self-Test: Multiple Choice

- If Earth rotated twice as fast as it currently does, but its motion around the Sun stayed the same, then (a) the night would be twice as long; (b) the night would be half as long; (c) the year would be half as long; (d) the length of the day would be unchanged.
- A long, thin cloud that stretched from directly overhead to the western horizon would have an angular size of (a) 45°; (b) 90°; (c) 180°; (d) 360°.
- VIS** According to Figure 1.15 (“The Zodiac”), in January the Sun is in the constellation (a) Cancer; (b) Gemini; (c) Leo; (d) Aquarius.
- If Earth orbited the Sun in 9 months instead of 12, then, compared with a sidereal day, a solar day would be (a) longer; (b) shorter; (c) unchanged.
- When a thin crescent of the Moon is visible just before sunrise, the Moon is in its (a) waxing phase; (b) new phase; (c) waning phase; (d) quarter phase.
- If the Moon's orbit were a little larger, solar eclipses would be (a) more likely to be annular; (b) more likely to be total; (c) more frequent; (d) unchanged in appearance.
- If the Moon orbited Earth twice as fast but in the same orbit the frequency of solar eclipses would (a) double; (b) be cut in half; (c) stay the same.
- VIS** In Figure 1.28 (“Triangulation”), using a longer baseline would result in (a) a less accurate distance to the tree; (b) a more accurate distance to the tree; (c) a smaller angle at point B; (d) a greater distance across the river.
- VIS** In Figure 1.32 (“Parallax”), a smaller Earth would result in (a) a smaller parallax angle; (b) a shorter distance measured to the object; (c) a larger apparent displacement; (d) stars appearing closer together.
- Today, distances to stars are measured by (a) bouncing radar signals; (b) reflected laser beams; (c) travel time by spacecraft; (d) geometry.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

- In 1 second, light leaving Los Angeles reaches approximately as far as (a) San Francisco, about 500 km; (b) London, roughly 10,000 km; (c) the Moon, 384,000 km; (d) Venus, 45,000,000 km from Earth at closest approach; or (e) the nearest star, about 4 light-years from Earth. Which is correct?
- (a) Write the following numbers in scientific notation (see Appendix 1 if you are unfamiliar with this notation): 1000; 0.000001; 1001; 1,000,000,000,000,000; 123,000; 0.000456. (b) Write the following numbers in “normal” numerical form:  $3.16 \times 10^7$ ;  $2.998 \times 10^5$ ;  $6.67 \times 10^{-11}$ ;  $2 \times 10^0$ . (c) Calculate:  $(2 \times 10^3) + 10^{-2}$ ;  $(1.99 \times 10^{30}) / (5.98 \times 10^{24})$ ;  $(3.16 \times 10^7) \times (2.998 \times 10^5)$ .
- The vernal equinox is now just entering the constellation Aquarius (see Figure 1.15). In what constellation will it lie in A.D. 10,000?
- Relative to the stars, through how many degrees, arc minutes, or arc seconds does the Moon move in (a) 1 hour of time; (b) 1 minute; (c) 1 second? How long does it take for the Moon to move a distance equal to its own diameter?
- At what distance is an object if its parallax, as measured from either end of a 1000-km baseline, is (a)  $1^\circ$ ; (b)  $1'$ ; (c)  $1''$ ?
- Given that the angular size of Venus is  $55''$  when the planet is 45,000,000 km from Earth, calculate Venus’s diameter (in kilometers).
- The Moon lies 384,000 km from Earth, and the Sun lies 150,000,000 km away. If both have the same angular size as seen from Earth, how many times larger than the Moon is the Sun?
- Estimate the angular diameter of your thumb, held at arm’s length.

## Activities

### Collaborative

1. Measure the nightly and monthly motion of the Moon. On a clear night, sketch a  $10^\circ$ -wide patch of the sky containing the Moon, with the Moon initially toward the west side of the patch. (See Individual Activity 2 below for how to estimate angles on the sky.) Repeat the observation of the same collection of stars every hour over the course of a night. You will see that the Moon’s position relative to the stars changes noticeably even in a few hours. What is the Moon’s angular speed (in degrees per hour)? Now observe the Moon at the *same* time each night over the course of a month. Sketch its appearance and note its position on the sky each night. Can you interpret its changing phase in terms of the relative positions of Earth, the Sun, and the Moon? (See Figure 1.20.)

### Individual

1. Find the star Polaris, also known as the North Star, in the evening sky. Identify any separate pattern of stars in the same general vicinity of the sky. Wait several hours, at least until after midnight, and then locate Polaris again. Has Polaris moved? What has happened to the nearby pattern of stars? Why?
2. Hold your little finger out at arm’s length. Can you cover the disk of the Moon? The Moon projects an angular size of  $30'$  (half a degree); your finger should more than cover it. You can use this fact to make some basic sky measurements. As a simple rule, your little finger at arm’s length is about  $1^\circ$  across, your middle three fingers are about  $4^\circ$  across, and your clenched fist is about  $10^\circ$  across. If the constellation Orion is visible, use this information to estimate the angular size of Orion’s belt and the angular distance between Betelgeuse and Rigel. Compare your findings with Figure 1.8(a).





# The Copernican Revolution

# 2

## THE BIRTH OF MODERN SCIENCE

Living in the Space Age, we have become accustomed to the modern view of our place in the universe. Images of our planet taken from space leave little doubt that Earth is round, and no one seriously questions the idea that we orbit the Sun. Yet there was a time, not so long ago, when some of our ancestors maintained that Earth was flat and lay at the center of all things.

Our view of the universe—and of ourselves—has undergone a radical transformation since those early days. Earth has become a planet like many others, and humankind has been torn from its throne at the center of the cosmos and relegated to a rather unremarkable position on the periphery of the Milky Way Galaxy. But we have been amply compensated for our loss of prominence: We have gained a wealth of scientific knowledge in the process. The story of how all this came about is the story of the rise of the scientific method and the genesis of modern astronomy.

**The Big Picture** Exploration is at the heart of the modern scientific method used by all scientists around the world. Ideas must be tested against what is observed in nature, and those ideas that fail the test are discarded. In this way, astronomers progressively generate, not “truth,” but better and better approximations of reality.

**LEFT:** In this colorized piece of historical artwork, a young Nicholas Copernicus is observing a lunar eclipse in Rome in the year 1500. He is not actually using a telescope, which would not be invented for another century; rather, he is looking along a transit device that improved naked-eye estimates of angular sizes. Perhaps more than anyone else, this Polish astronomer began the revolution that overthrew more than a thousand years of philosophical thinking that claimed Earth to be the immovable center of all things. (S. Terry; engraving from the 1875 edition of *Vies des Savants Illustres*)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Describe how some ancient civilizations attempted to explain the heavens in terms of Earth-centered models of the universe.
- 2 Explain how the observed motions of the planets led to our modern view of a Sun-centered solar system.
- 3 Describe the major contributions of Galileo and Kepler to our understanding of the solar system.
- 4 State Kepler’s laws of planetary motion.
- 5 Explain how astronomers have measured the true size of the solar system.
- 6 State Newton’s laws of motion and universal gravitation and explain how they account for Kepler’s laws.
- 7 Explain how the law of gravitation enables us to measure the masses of astronomical bodies.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.

## 2.1 Ancient Astronomy

Many ancient cultures took a keen interest in the changing nighttime sky. The records and artifacts that have survived until the present make that abundantly clear. But unlike today, the major driving force behind the development of astronomy in those early societies was probably neither scientific nor religious. Instead, it was decidedly practical and down to earth. Seafarers needed to navigate their vessels, and farmers had to know when to plant their crops. In a real sense, then, human survival depended on knowledge of the heavens. The ability to predict accurately the arrival of the seasons, as well as other astronomical events, was undoubtedly a highly prized, perhaps jealously guarded, skill.

In Chapter 1, we saw that the human mind's ability to perceive patterns in the stars led to the "invention" of constellations as a convenient means of labeling regions of the celestial sphere. ∞ (Sec. 1.3) The realization that these patterns returned to the night sky at the same time each year met the need for a practical means of tracking the seasons. Widely separated cultures all over the world built elaborate structures to serve, at least in part, as primitive calendars. Often the keepers of the secrets of the sky enshrined their knowledge in myth and ritual, and these astronomical sites were also used for religious ceremonies.

Perhaps the best-known such site is *Stonehenge*, located on Salisbury Plain in England, and shown in Figure 2.1. This ancient stone circle, which today is one of the most popular tourist attractions in Britain, dates from the Stone Age. Researchers think it was an early astronomical observatory

of sorts—not in the modern sense of the term (a place for making new observations and discoveries pertaining to the heavens)—but rather a kind of three-dimensional calendar or almanac, enabling its builders and their descendants to identify important dates by means of specific celestial events. Its construction apparently spanned a period of about 17 centuries, beginning around 2800 B.C. Additions and modifications continued to about 1100 B.C., indicating its ongoing importance to the Stone Age and, later, Bronze Age people who built, maintained, and used Stonehenge. The largest stones shown in Figure 2.1 weigh up to 50 tons and were transported from quarries many miles away.


Many of the stones are aligned so that they point toward important astronomical events. For example, the line joining the center of the inner circle to the so-called heel stone, set off some distance from the rest of the structure, points in the direction of the rising Sun on the summer solstice. Other alignments are related to the rising and setting of the Sun and the Moon at other times of the year. The accurate alignments (within a degree or so) of the stones of Stonehenge were first noted in the 18th century, but it was only relatively recently—in the second half of the 20th century, in fact—that the scientific community began to credit Stone Age technology with the ability to carry out such a precise feat of engineering. Although some of Stonehenge's purposes remain uncertain and controversial, the site's function as an astronomical almanac seems well established. Although Stonehenge is the most impressive and the best preserved, other stone circles, found all over Europe, are thought to have performed similar functions.



◀ **FIGURE 2.1 Stonehenge** This remarkable site in the south of England was probably constructed as a primitive calendar or almanac. The inset shows sunrise at Stonehenge at the summer solstice. As seen from the center of the stone circle, the Sun rose directly over the "heel stone" on the longest day of the year. (S. Pitamitz/Superstock; inset D. Numuk/All Canada Photos/Superstock)



(a)



(c)

**▲ FIGURE 2.2 Observatories in the Americas** (a) The Big Horn Medicine Wheel in Wyoming, built by the Plains Indians, has spokes and other features that roughly align with risings and settings of the Sun and other stars. (b) The Caracol temple in Mexico, built by the Mayan civilization, has some windows that seem to align with astronomical events, suggesting that at least part of Caracol's function may have kept track of the seasons and the heavens. (c) This thin streak of light and shadow, created by the Sun's rays playing off the cliffs in Chaco Canyon of America's Southwest, aligns exactly with a carved rock pattern at noon on the summer solstice—almost certainly an intentional sign for astronomical or agricultural purposes. (G. Gerster; H. Lapahie Jr; F1online Digitale Bildagentur)

Many North American cultures were interested in the heavens. The Big Horn Medicine Wheel in Wyoming (Figure 2.2a) is similar to Stonehenge in design—and, perhaps, intent—although it is somewhat simpler in execution. Some researchers have identified alignments between the Medicine Wheel's spokes and the rising and setting Sun at solstices and equinoxes, and with some bright stars, suggesting that its builders—the Plains Indians—had much more than a passing familiarity with the changing nighttime sky. Other experts disagree, however, arguing that



(b)

the alignments are quite inaccurate and consistent with pure chance and that the Medicine Wheel's purpose was more likely symbolic, rather than practical. A similar controversy swirls around the Caracol temple (Figure 2.2b) in the famous Mayan city of Chitzen Itza, built around A.D. 1000 on Mexico's Yucatán peninsula. Was it an observatory, as some suggest, perhaps tied to human sacrifices when Venus appeared in the morning or evening sky? Or are the claimed alignments of its windows just wishful thinking and the temple's purpose simply religious, rather than astronomical?

Experts do seem to agree—for now, at least—that the Sun Dagger (Figure 2.2c), in Chaco Canyon, New Mexico, is a genuine astronomical calendar. It is constructed so that the sliver of light passes precisely through the center of the carved stone spiral at noon on the summer solstice. Numerous similar sites have been found throughout the American Southwest.

The ancient Chinese also observed the heavens. Their astrology attached particular importance to “omens” such as comets and “guest stars”—stars that appeared suddenly in the sky and then slowly faded away—and they kept careful and extensive records of such events. Twentieth-century astronomers still turn to the Chinese records to obtain observational data recorded during the Dark Ages (roughly from the 5th to the 10th century A.D.), when turmoil in Europe



▲ **FIGURE 2.3 Persian Astronomers at Work** During the Dark Ages, much scientific information was preserved and new discoveries were made by astronomers in the Islamic world, as depicted in this illustration from a 16th-century manuscript. (Bridgeman Art Library)

largely halted the progress of Western science. Perhaps the best-known guest star was one that appeared in A.D. 1054 and was visible in the daytime sky for many months. We now know that the event was actually a *supernova*: the explosion of a giant star, which scattered most of its mass into space (see Chapter 21). It left behind a remnant that is still detectable today, nine centuries later. The Chinese data are a prime source of historical information for supernova research.

A vital link between the astronomy of ancient Greece and that of medieval Europe was provided by astronomers in the Muslim world (see Figure 2.3). For six centuries, from the depths of the Dark Ages to the beginning of the Renaissance, Islamic astronomy flourished and grew, preserving and augmenting the knowledge of the Greeks. Its influence on modern astronomy is subtle, but widespread. Many of the mathematical techniques involved in trigonometry were developed by Islamic astronomers in response to practical problems,

such as determining the precise dates of holy days or the direction of Mecca from any given location on Earth. Astronomical terms such as *zenith* and *azimuth* and the names of many stars—for example, Rigel, Betelgeuse, and Vega—all bear witness to this extended period of Muslim scholarship.

Astronomy is not the property of any one culture, civilization, or era. The same ideas, the same tools, and even the same misconceptions have been invented and reinvented by human societies all over the world in response to the same basic driving forces. Astronomy came into being because people knew that there was a practical benefit in being able to predict the positions of the stars, but its roots go much deeper than that. The need to understand where we came from and how we fit into the cosmos is an integral part of human nature.

## 2.2 The Geocentric Universe


The Greeks of antiquity, and undoubtedly civilizations before them, built models of the universe. The study of the workings of the universe on the largest scales is called *cosmology*. Today, cosmology entails looking at the universe on scales so large that even entire galaxies can be regarded as mere points of light scattered throughout space. To the Greeks, however, the universe was basically the *solar system*—the Sun, Earth, and Moon, and the planets known at that time. The stars beyond were surely part of the universe, but they were considered to be fixed, unchanging beacons on the celestial sphere. The Greeks did not consider the Sun, the Moon, and the planets to be part of this mammoth celestial dome, however. Those objects had patterns of behavior that set them apart.

### Observations of the Planets

Greek astronomers observed that over the course of a night, the stars slid smoothly across the sky. Over the course of a month, the Moon moved smoothly and steadily along its path on the sky relative to the stars, passing through its familiar cycle of phases. Over the course of a year, the Sun progressed along the ecliptic at an almost constant rate, varying little in brightness from day to day. In short, the behavior of both Sun and Moon seemed fairly simple and orderly. But ancient astronomers were also aware of five other bodies in the sky—the planets Mercury, Venus, Mars, Jupiter, and Saturn—whose behavior was not so easy to grasp. Their motions ultimately led to the downfall of an entire theory of the solar system and to a fundamental change in humankind's view of the universe.

To the naked eye (or even through a telescope), planets do not behave in as regular and predictable a fashion as the Sun, Moon, and stars. They vary in brightness, and they don't maintain a fixed position in the sky. Unlike the Sun and Moon, the planets seem to wander around the celestial

Motions of the planets relative to the stars produce continuous streaks on a planetarium "sky."




Observed planet motions can be complicated because each planet travels with a different speed around the Sun.

sphere—indeed, the word *planet* derives from the Greek word *planetes*, meaning “wanderer.” Planets never stray far from the ecliptic and generally traverse the celestial sphere from west to east, like the Sun. However, they seem to speed up and slow down during their journeys, and at times they even appear to loop back and forth relative to the stars, as shown in Figure 2.4. In other words, there are periods when a planet’s eastward motion (relative to the stars) stops, and the planet appears to move westward in the sky for a month or two before reversing direction again and continuing on its eastward journey. Motion in the eastward sense is usually referred to as *direct*, or *prograde*, motion; the backward (westward) loops are known as **retrograde motion**.

Ancient astronomers knew well that the periods of retrograde motion were closely correlated with other planetary properties, such as apparent brightness and position in the sky. Figure 2.5 (a modern view of the

► **FIGURE 2.5 Inferior and Superior Orbits** Diagram of Earth’s orbit and two other possible planetary orbits. An “inferior” orbit lies between Earth’s orbit and the Sun. Mercury and Venus move in such orbits. A “superior” orbit (such as the orbit of Mars, Jupiter, or Saturn) lies outside that of Earth. The points noted on the orbits indicate times when a planet appears to come close to the Sun (conjunction) or is diametrically opposite the Sun on the celestial sphere (opposition).



◀ **FIGURE 2.4 Planetary Motion** Most of the time, planets move from west to east relative to the background stars. Occasionally—roughly once per year—they change direction and temporarily undergo retrograde motion (east to west) before looping back. The main illustration shows an actual retrograde loop in the motion of the planet Mars. The inset depicts the movements of several planets over the course of several years, as reproduced on the inside dome of a planetarium. (*Boston Museum of Science*)

solar system, note!) shows three schematic planetary orbits and defines some time-honored astronomical terminology describing a planet’s location relative to Earth and the Sun. Mercury and Venus are referred to as *inferior* (“lower”) planets because their orbits lie between Earth and the Sun. Mars, Jupiter, and Saturn, whose orbits lie outside Earth’s, are known as *superior* (“higher”) planets. For early astronomers, the key observations of planetary orbits were the following:

- An inferior planet never strays too far from the Sun, as seen from Earth. As illustrated in the inset to Figure 2.5, because its path on the celestial sphere is close to the ecliptic, an inferior planet makes two *conjunctions* (or close approaches) with the Sun during each orbit. (It doesn’t actually come close to the Sun, of course. Conjunction is simply the occasion when the planet and the Sun are in the same direction in the sky.) At *inferior conjunction*, the planet is closest to Earth and moves past the Sun from east to west—that is, in the retrograde

sense. At *superior conjunction*, the planet is farthest from Earth and passes the Sun in the opposite (prograde) direction.

- Seen from Earth, the superior planets are not “tied” to the Sun as the inferior planets are. The superior planets make one prograde conjunction with the Sun during each trip around the celestial sphere. However, they exhibit retrograde motion (Figure 2.4) when they are at *opposition*, diametrically opposite the Sun on the celestial sphere.
- The superior planets are brightest at opposition, during retrograde motion. By contrast, the inferior planets are brightest a few weeks before and after inferior conjunction.


The challenge facing astronomers—then as now—was to find a solar system model that could explain all the existing observations and that could also make testable and reliable predictions of future planetary motions. ∞ (Sec. 1.2)

Ancient astronomers correctly reasoned that the changing brightness of a planet in the night sky is related to the planet’s distance from Earth. Like the Moon, the planets produce no light of their own. Instead, they shine by reflected sunlight and, generally speaking, appear brightest when closest to us. Looking at Figure 2.5, you may already be able to discern the basic reasons for some of the planetary properties just listed; we’ll return to the “modern” explanation in the next section. However, as we now discuss, the ancients took a very different path in their attempts to explain planetary motion.

## A Theoretical Model

The earliest models of the solar system followed the teachings of the Greek philosopher Aristotle (384–322 B.C.) and were **geocentric**, meaning that Earth lay at the center of the universe and all other bodies moved around it. ∞ (Sec. 1.3) The celestial sphere, shown in Figures 1.11 and 1.16, illustrates the basic geocentric view. These models employed what Aristotle, and Plato before him, had taught was the perfect form: the circle. The simplest possible description—uniform motion around a circle with Earth at its center—provided a fairly good approximation to the orbits of the Sun and the Moon, but it could not account for the observed variations in planetary brightness or the retrograde motion of the planets. A more complex model was needed to describe these heavenly “wanderers.”

In the first step toward this new model, each planet was taken to move uniformly around a small circle, called an **epicycle**, whose *center* moved uniformly around Earth on a second and larger circle, known as a **deferent** (Figure 2.6). The motion was now composed of two separate circular orbits, creating the possibility that, at some times, the planet’s apparent motion could be retrograde. Also, the distance from the planet to Earth would vary, accounting for changes




**Interactive FIGURE 2.6 Geocentric Model** In the geocentric model of the solar system, the observed motions of the planets made it impossible to assume that they moved on simple circular paths around Earth. Instead, each planet was thought to follow a small circular orbit (the epicycle) about an imaginary point that itself traveled in a large, circular orbit (the deferent) about Earth.

in brightness. By tinkering with the relative sizes of the epicycle and deferent, with the planet’s speed on the epicycle, and with the epicycle’s speed along the deferent, early astronomers were able to bring this “epicyclic” motion into fairly good agreement with the observed paths of the planets in the sky. Moreover, the model had good predictive power, at least to the accuracy of observations at the time.

However, as the number and the quality of observations increased, it became clear that the simple epicyclic model was not perfect. Small corrections had to be introduced to bring it into line with new observations. The center of the deferents had to be shifted slightly from Earth’s center, and the motion of the epicycles had to be imagined uniform with respect to yet another point in space, not Earth. Furthermore, in order to explain the motions of the inferior planets, the model simply had to assume that the deferents of Mercury and Venus were, for some (unknown) reason, tied to that of the Sun. Similar assumptions also applied to the superior planets, to ensure that their retrograde motion occurred at opposition.

Around A.D. 140, a Greek astronomer named Ptolemy constructed perhaps the most complete geocentric model of all time. Illustrated in simplified form in Figure 2.7, it explained remarkably well the observed paths of the five planets then known, as well as the paths of the Sun and the Moon. However, to achieve its explanatory and predictive power, the full **Ptolemaic model** required a series of no fewer than 80 distinct circles. To account for the paths of the



▲ **FIGURE 2.7 Ptolemaic Model** The basic features, drawn roughly to scale, of Ptolemy's geocentric model of the inner solar system, a model that enjoyed widespread popularity prior to the Renaissance. Only the five planets visible to the naked eye and hence known to the ancients—Mercury, Venus, Mars, Jupiter, and Saturn—are shown. To avoid confusion, partial paths (dashed) of only two planets—Venus and Jupiter—are drawn here.

Sun, Moon, and all eight planets (and their moons) that we know today would require a vastly more complicated set. Nevertheless, Ptolemy's comprehensive text on the topic, *Syntaxis* (better known today by its Arabic name, *Almagest*, “the greatest”), provided the intellectual framework for all discussion of the universe for well over a thousand years.

## Evaluating the Geocentric Model

Today, our scientific training leads us to seek simplicity, because, in the physical sciences, simplicity has so often proved to be an indicator of truth. We would regard the intricacy of a model as complicated as the Ptolemaic system as a clear sign of a fundamentally flawed theory. ∞ (Sec. 1.2) Why was the Ptolemaic model so complex? With the benefit of hindsight, we now recognize that its major error lay in its assumption of a geocentric universe. This misconception was compounded by the insistence on uniform circular motion, whose basis was largely philosophical, rather than scientific, in nature.

Actually, history records that some ancient Greek astronomers reasoned differently about the motions of heavenly bodies. Foremost among them was Aristarchus of Samos (310–230 B.C.), who proposed that all the planets, including Earth, revolve around the Sun and, furthermore, that Earth

rotates on its axis once each day. This combined revolution and rotation, he argued, would create an *apparent* motion of the sky—a simple idea that is familiar to anyone who has ridden on a merry-go-round and watched the landscape appear to move past in the opposite direction. However, Aristarchus's description of the heavens, though essentially correct, did not gain widespread acceptance during his lifetime. Aristotle's influence was too strong, his followers too numerous, and his writings too comprehensive. The geocentric model went largely unchallenged until the 16th century A.D.

The Aristotelian school did present some simple and (at the time) compelling arguments in favor of their views. First, of course, Earth doesn't *feel* as if it's moving—and if it were moving, wouldn't there be a strong wind as the planet revolves at high speed around the Sun? Also, considering that the vantage point from which we view the stars changes over the course of a year, why don't we see stellar parallax?

∞ (Sec. 1.4)

Nowadays we might dismiss the first points as merely naive, but the last is a valid argument and the reasoning essentially sound. Indeed, we now know that there *is* stellar parallax as Earth orbits the Sun. However, because the stars are so distant, it amounts to less than 1 arc second ( $1''$ ), even for the closest stars. Early astronomers simply would not have noticed it. (In fact, stellar parallax was conclusively measured only in the middle of the 19th century.)

We will encounter many other instances in astronomy wherein correct reasoning led to the wrong conclusions because it relied on inadequate data. Even when the scientific method is properly applied and theoretical predictions are tested against reality, a theory can be only as good as the observations on which it is based. ∞ (Sec. 1.2)

## 2.3 The Heliocentric Model of the Solar System

The Ptolemaic picture of the universe survived, more or less intact, for almost 14 centuries, until a 16th-century Polish cleric, Nicolaus Copernicus (Figure 2.8), rediscovered Aristarchus's **heliocentric** (Sun-centered) model and showed how, in its harmony and organization, it provided a more natural explanation of the observed facts than did the tangled geocentric cosmology. Copernicus asserted that Earth spins on its axis and, like the other planets, orbits the Sun. Only the Moon, he said, orbits Earth. As we will see, not only does this model explain the observed daily and seasonal changes in the heavens, but it also naturally accounts for retrograde motion and variations in brightness of the planets. ∞ (Sec. 1.4)

The critical realization that Earth is not at the center of the universe is now known as the **Copernican revolution**. The seven crucial statements that form its foundation are summarized in *Discovery 2-1*.



## DISCOVERY 2-1

## Foundations of the Copernican Revolution

The following seven points are essentially Copernicus's own words, with the italicized material providing additional explanation:


1. The celestial spheres do not have just one common center. *Specifically, Earth is not at the center of everything.*
2. The center of Earth is not the center of the universe, but is instead only the center of gravity and of the lunar orbit.
3. All the spheres revolve around the Sun. *By spheres, Copernicus meant the planets.*
4. The ratio of Earth's distance from the Sun to the height of the firmament is so much smaller than the ratio of Earth's radius to the distance to the Sun that the distance to the Sun is imperceptible compared with the height of the firmament.
5. The motions appearing in the firmament are not its motions, but those of Earth. Earth performs a daily rotation around its fixed poles, while the firmament remains immobile as the highest heaven. *Because the stars are so far away, any apparent motion we see in them is the result of Earth's rotation.*
6. The motions of the Sun are not its motions, but the motion of Earth. *Similarly, the Sun's apparent daily and yearly motion are actually due to the various motions of Earth.*
7. What appears to us as retrograde and forward motion of the planets is not their own, but that of Earth. *The heliocentric picture provides a natural explanation for retrograde planetary motion, again as a consequence of Earth's motion.*

Figure 2.9 shows how the Copernican view explains the varying brightness of a planet (in this case, Mars), its observed looping motions, and the fact that the retrograde motion of a superior planet occurs at opposition. If we suppose that Earth moves faster than Mars, then every so often Earth “overtakes” that planet. Mars will then appear to move backward in the sky, in much the same way as a car we overtake on the highway seems to slip backward relative to us. Replace Mars by Earth and Earth by Venus, and you should also be able to extend the explanation to the inferior

planets. (To complete the story with a full explanation of their apparent brightnesses, however, you'll have to wait until Section 9.1!) Notice that, in the Copernican picture, the planet's looping motions are only apparent. In the Ptolemaic view, they are real.

Copernicus's major motivation for introducing the heliocentric model was simplicity. Even so, he was still influenced by Greek thinking and clung to the idea of circles to model the planets' motions. To bring his theory into agreement with observations of the night sky, he was forced to retain the idea of epicyclic motion, although with the deferent centered on the Sun rather than on Earth and with smaller epicycles than in the Ptolemaic picture. Thus, he retained unnecessary complexity and actually gained little in predictive power over the geocentric model. The heliocentric model did rectify some small discrepancies and inconsistencies in the Ptolemaic system, but for Copernicus, the primary attraction of heliocentricity was its simplicity—its being “more pleasing to the mind.” His theory was more something he *felt* than he could *prove*. To the present day, scientists still are guided by simplicity, symmetry, and beauty in modeling all aspects of the universe.


Despite the support of some observational data, neither his fellow scholars nor the general public easily accepted Copernicus's model. For the learned, heliocentricity went against the grain of much previous thinking and violated many of the religious teachings of the time, largely because it relegated Earth to a noncentral and undistinguished place within the solar system and the universe. And Copernicus's work had little impact on the general populace of his time, at least in part because it was published in Latin (the standard language of academic discourse at the time), which most people could not read. Only long after Copernicus's death, when others—notably Galileo Galilei—popularized his ideas, did the Roman Catholic Church take them seriously enough to



▲ FIGURE 2.8 Nicolaus Copernicus (1473–1543). (E. Lessing/Art Resource, NY)

### Interactive **FIGURE 2.9** **Retrograde Motion**

**MA** The Copernican model of the solar system explains both the varying brightnesses of the planets and the phenomenon of retrograde motion. Here, for example, when Earth and Mars are relatively close to one another in their respective orbits (as at position 6), Mars seems brighter. When they are farther apart (as at position 1), Mars seems dimmer. Also, because the (light blue) line of sight from Earth to Mars changes as the two planets orbit the Sun, Mars appears to loop back and forth in retrograde motion. Follow the lines in numerical order, and note how the line of sight moves backward relative to the stars between locations 5 and 7. That's because Earth, on the inside track, moves faster in its orbit than does Mars. The white curves are actual planetary orbits; the red curve is Mars's motion as seen from Earth.



both banning them. Copernicus's writings on the heliocentric universe were placed on the Church's *Index of Prohibited Books* in 1616, 73 years after they were first published. They remained there until the end of the 18th century.

#### CONCEPT Check


- ✓ How do the geocentric and heliocentric models of the solar system differ in their explanations of planetary retrograde motion?

## 2.4 The Birth of Modern Astronomy

In the century following the death of Copernicus and the publication of his theory of the solar system, two scientists—Galileo Galilei and Johannes Kepler—made indelible imprints on the study of astronomy. Contemporaries, they were aware of each other's work and corresponded from time to time about their theories. Each achieved fame for his discoveries and made great strides in popularizing the Copernican viewpoint, yet in their approaches to astronomy they were as different as night and day.

### Galileo's Historic Observations

Galileo Galilei (Figure 2.10) was an Italian mathematician and philosopher. By his willingness to perform experiments to test his ideas—a rather radical approach in those days—and by embracing the brand-new technology of the telescope, he revolutionized the way in which science was done, so much so that he is now widely regarded as the father of experimental science.




▲ **FIGURE 2.10** Galileo Galilei (1564–1642). (Art Resource, NY)

The telescope was invented in Holland in the early 17th century. Hearing of the invention (but without having seen one), Galileo built a telescope for himself in 1609 and aimed it at the sky. What he saw conflicted greatly with the philosophy of Aristotle and provided much new data to support the ideas of Copernicus.\*

Using his telescope, Galileo discovered that the Moon had mountains, valleys, and craters—terrain in many ways

\*In fact, Galileo had already abandoned Aristotle in favor of Copernicus, although he had not published his opinions at the time he began his telescopic observations.



▲ **FIGURE 2.11 Galilean Moons** The four Galilean moons of Jupiter, as sketched by Galileo in his notebook on 7 nights between January 7 and 15, 1610. More of Galileo's remarkable sketches of Saturn, star clusters, and the Orion constellation can be seen on the first page of Part 1. (From *Sidereus Nuncius*)

reminiscent of that on Earth. Looking at the Sun (something that should *never* be done directly and that may have eventually blinded Galileo), he found imperfections—dark blemishes now known as *sunspots*. These observations ran directly counter to the orthodox wisdom of the day. By noting the changing appearance of sunspots from day to day, Galileo inferred that the Sun *rotates*, approximately once per month, around an axis roughly perpendicular to the ecliptic plane.

Galileo also saw four small points of light, invisible to the naked eye, orbiting the planet Jupiter and realized that they were moons. Figure 2.11 shows some sketches of these moons, taken from Galileo's notes. To Galileo, the fact that another planet had moons provided the strongest support for the Copernican model. Clearly, Earth was not the center of all things. He also found that Venus varied in apparent size and showed a complete cycle of phases, like those of our Moon (Figure 2.12), findings that could be explained only by the planet's motion around the Sun. These observations were more strong evidence that Earth is not the center of all things and that at least one planet orbited the Sun. For more of Galileo's sketches, with comparisons to modern photographs, see the Part 1 Opener on p. 1.


In 1610, Galileo published a book called *Sidereus Nuncius* (*The Starry Messenger*), detailing his observational findings

and his controversial conclusions supporting the Copernican theory. In reporting and interpreting the wondrous observations made with his new telescope, Galileo was directly challenging both the scientific orthodoxy and the religious dogma of his day. He was (literally) playing with fire—he must certainly have been aware that only a few years earlier, in 1600, the astronomer Giordano Bruno had been burned at the stake in Rome, in part for his heretical teaching that Earth orbited the Sun. However, by all accounts, Galileo delighted in publicly ridiculing and irritating his Aristotelian colleagues. In 1616 his ideas were judged heretical, Copernicus's works were banned by the Roman Catholic Church, and Galileo was instructed to abandon his astronomical pursuits.



But Galileo would not desist. In 1632 he raised the stakes by publishing *Dialogue Concerning the Two Chief World Systems*, which compared the Ptolemaic and Copernican models. The book presented a discussion among three people, one of them a dull-witted Aristotelian whose views (which were in fact the stated opinions of the then Pope, Urban VIII) time and again were roundly defeated by the arguments of one of his two companions, an articulate proponent of the heliocentric system. To make the book accessible to a wide popular audience, Galileo wrote it in Italian rather than Latin. These actions brought Galileo into direct conflict with the authority of the Church. Eventually, the Inquisition forced him, under threat of torture, to retract his claim that Earth orbits the Sun, and he was placed under house arrest in 1633. He remained imprisoned for the rest of his life. Not until 1992 did the Church publicly forgive Galileo's "crimes." But the damage to the orthodox view of the universe was done, and the Copernican genie was out of the bottle once and for all.

## Ascendancy of the Copernican System

Although Renaissance scholars were correct, they could not *prove* that our planetary system is centered on the Sun or even that Earth moves through space. The observational consequences of Earth's orbital motion were just too small for the technology of the day to detect. Direct evidence of Earth's motion was obtained only in 1728, when English astronomer James Bradley discovered the **aberration of starlight**—a slight (roughly 20") shift in the observed direction to a star, caused by Earth's motion perpendicular to the line of sight, much as rain drops falling vertically leave slanted tracks on the passenger window of a moving car or train. Bradley's observation was the first proof that Earth revolves around the Sun; subsequent observations of many stars in many different directions have repeatedly confirmed the effect. Additional proof of Earth's orbital motion came in 1838, with the first unambiguous determination of stellar parallax (see Figure 1.30) by German astronomer Friedrich Bessel. ∞ (Sec. 1.6)



(a) Sun-centered model



(b) Ptolemy's model

**Interactive FIGURE 2.12 Venus Phases** Both the Ptolemaic and the Copernican models of the solar system predict that Venus should show phases as it moves in its orbit. (a) In the Copernican picture, when Venus is directly between Earth and the Sun, its unlit side faces us and the planet is invisible to us. As Venus moves in its orbit, progressively more of its illuminated face is visible from Earth. Note the connection between the orbital phase and the apparent size of the planet: Venus seems much larger in its crescent phase than when it is full because it is much closer to us during its crescent phase. This is the behavior actually observed. The insets at bottom left and right are actual photographs of Venus taken at two of its crescent phases. (Courtesy N. Mex. St. Univ.) (b) The Ptolemaic model (see also Figure 2.7) is unable to account for these observations. In particular, the full phase of the planet cannot be explained. Seen from Earth, Venus reaches only a "fat crescent", yet never a full phase, then begins to wane as it nears the Sun. (Both these views are from a sideways perspective; from overhead, both orbits are very nearly circular, as shown in Figure 2.18.)

Following those early measurements, support for the heliocentric solar system has grown steadily, as astronomers have subjected the theory to more and more sophisticated observational tests, culminating in the interplanetary expeditions of our unmanned space probes of the 1960s, 1970s, and 1980s. Today, the evidence is overwhelming. The development and eventual acceptance of the heliocentric model were milestones in human thinking. This removal of Earth from any position of great cosmic significance is generally known, even today, as the *Copernican principle*. It has become a cornerstone of modern astrophysics.

The Copernican revolution is a prime example of how the scientific method, though affected at any given time by the subjective whims, human biases, and even sheer luck of researchers, does ultimately lead to a definite degree of objectivity. ∞ (Sec. 1.2) Over time, many groups of scientists checking, confirming, and refining experimental

tests can neutralize the subjective attitudes of individuals. Usually, one generation of scientists can bring sufficient objectivity to bear on a problem, although some especially revolutionary concepts are so swamped by tradition, religion, and politics that more time is necessary. In the case of heliocentricity, objective confirmation was not obtained until about three centuries after Copernicus published his work and more than 2000 years after Aristarchus had proposed the concept. Nonetheless, objectivity *did in fact* eventually prevail, and our knowledge of the universe has expanded immeasurably as a result.

### PROCESS OF SCIENCE Check


- ✓ In terms of the Scientific Method presented in Chapter 1, what were the principal advantages of the heliocentric theory over the geocentric model?

## 2.5 The Laws of Planetary Motion

At about the same time as Galileo was becoming famous—or notorious—for his pioneering telescopic observations and outspoken promotion of the Copernican system, Johannes Kepler (Figure 2.13), a German mathematician and astronomer, was developing the laws of planetary motion that now bear his name. Galileo was in many ways the first “modern” observer. He used emerging technology, in the form of the telescope, to achieve new insights into the universe. In contrast, Kepler was a pure theorist. His groundbreaking work that so clarified our knowledge of planetary motion was based almost entirely on the observations of others, principally an extensive collection of data compiled by Tycho Brahe (1546–1601), Kepler’s employer and arguably one of the greatest observational astronomers that has ever lived.

### Brahe’s Complex Data

Tycho, as he is often called, was both an eccentric aristocrat and a skillful observer. Born in Denmark, he was educated at some of the best universities in Europe, where he studied astrology, alchemy, and medicine. Most of his observations, which predated the invention of the telescope by several decades, were made at his own observatory, named *Uraniborg*, in Denmark (Figure 2.14). There, using instruments of his own design, Tycho maintained meticulous and accurate records of the stars, planets, and other noteworthy celestial



▲ **FIGURE 2.13** Johannes Kepler (1571–1630). (E. Lessing/Art Resource, NY)




▲ **FIGURE 2.14** Tycho Brahe The astronomer in his observatory Uraniborg, on the island of Hveen in Denmark. Brahe’s observations of the positions of stars and planets on the sky were the most accurate and complete set of naked-eye measurements ever made. (Newberry Library/Superstock)

events, including a comet and a supernova (see Chapter 21), the appearance of which helped convince him that the Aristotelian view of the universe could not be correct.

In 1597, having fallen out of favor with the Danish court, Tycho moved to Prague as Imperial Mathematician of the Holy Roman Empire. Prague happens to be fairly close to Graz, in Austria, where Kepler lived and worked. Kepler joined Tycho in Prague in 1600 and was put to work trying to find a theory that could explain Brahe’s planetary data. When Tycho died a year later, Kepler inherited not only his position, but also his most priceless possession: the accumulated observations of the planets, spanning several decades. These observations, though made with the naked eye, were nevertheless of very high quality. In most cases, his measured positions of stars and planets were accurate to within about 1′. Kepler set to work seeking a unifying principle to explain in detail the motions of the planets, without the need for epicycles. The effort was to occupy much of the remaining 29 years of his life.

Kepler had already accepted the heliocentric picture of the solar system. His goal was to find a simple and elegant description of planetary motion within the Copernican framework that fit Brahe's complex mass of detailed observations. In the end, he found it necessary to abandon Copernicus's simple idea of circular planetary orbits. However, even greater simplicity emerged as a result. After long years of studying Brahe's planetary data, and after many false starts and blind alleys, Kepler developed the laws that now bear his name.

Kepler determined the shape of each planet's orbit by triangulation—not from different points on Earth, but from different points on Earth's orbit, using observations made at many different times of the year.  (Sec. 1.6) By using a portion of Earth's orbit as a baseline for his triangle, Kepler was able to measure the relative sizes of the other planetary orbits. Noting where the planets were on successive nights, he found the speeds at which the planets move. We do not know how many geometric shapes Kepler tried for the orbits before he hit upon the correct one. His difficult task was made even more complicated because he had to determine Earth's own orbit, too. Nevertheless, he eventually succeeded in summarizing the motions of all the known planets, including Earth, in just three laws: the **laws of planetary motion**.


## Kepler's Simple Laws


Kepler's first law of planetary motion has to do with the *shapes* of the planetary orbits:

The orbital paths of the planets are elliptical (*not* circular), with the Sun at one focus.

An **ellipse** is simply a flattened circle. Figure 2.15 illustrates a means of constructing an ellipse with a piece of string and two thumbtacks. Each point at which the string is pinned is called a **focus** (plural: *foci*) of the ellipse. The long axis of the ellipse, containing the two foci, is known as the *major axis*. Half the length of this long axis is referred to as the **semimajor axis**, a measure of the ellipse's size. A circle is a special case in which the two foci happen to coincide; its semimajor axis is simply its radius.

The **eccentricity** of an ellipse is simply a measure of how flattened it is. Technically, eccentricity is defined as the ratio of the distance between the foci to the length of the major axis, but the most important thing to remember here is that an eccentricity of zero corresponds to no flattening—a perfect circle—whereas an eccentricity of one means that the circle has been squashed all the way down to a straight line. Note that, while the Sun resides at one focus of the elliptical orbit, the other focus is empty and has no particular physical significance. (However, we can still figure out where it is, because the two



 **Interactive FIGURE 2.15 Ellipse** An ellipse can be drawn with the aid of a string, a pencil, and two thumbtacks.

foci are symmetrically placed about the center, along the major axis.)

The length of the semimajor axis and the eccentricity are all we need to describe the size and shape of a planet's orbital path (see *More Precisely 2-1*). In fact, no planet's elliptical orbit is nearly as elongated as the one shown in Figure 2.15. With one exception (the orbit of Mercury), planetary orbits in our solar system have such small eccentricities that our eyes would have trouble distinguishing them from true circles. Only because the orbits are so nearly circular were the Ptolemaic and Copernican models able to come as close as they did to describing reality.

Kepler's substitution of elliptical for circular orbits was no small advance. It amounted to abandoning an aesthetic bias—the Aristotelian belief in the perfection of the circle—that had governed astronomy since Greek antiquity. Even Galileo Galilei, not known for his conservatism in scholarly matters, clung to the idea of circular motion and never accepted the notion that the planets move in elliptical paths.

The second law, illustrated in Figure 2.16, addresses the *speed* at which a planet traverses different parts of its orbit:


An imaginary line connecting the Sun to any planet sweeps out equal areas of the ellipse in equal intervals of time.

While orbiting the Sun, a planet traces the arcs labeled A, B, and C in the figure in equal times. Notice, however, that the

## MORE PRECISELY 2-1

### Some Properties of Planetary Orbits


Two numbers—*semimajor axis* and *eccentricity*—are all that are needed to describe the size and shape of a planet's orbital path. From them, we can derive many other useful quantities. Two of the most important are the planet's *perihelion* (its point of closest approach to the Sun) and its *aphelion* (point of greatest distance from the Sun). From the definitions presented in the text, it follows that if the planet's orbit has semimajor axis  $a$  and eccentricity  $e$ , the planet's perihelion is at a distance  $a(1 - e)$  from the Sun, while its aphelion is at  $a(1 + e)$ . These points and distances are illustrated in the following figure:



**EXAMPLE 1** We can locate the other focus of the ellipse in the diagram and hence determine the eccentricity from the definition in the text quite simply. The second focus is placed symmetrically along the major axis, at the point marked with an "X." With a ruler, measure (1) the length of the major axis and (2) the distance between the two foci. Dividing the second distance by the first, you should find an eccentricity of  $3.4 \text{ cm}/6.8 \text{ cm} = 0.5$ . Alternatively, we could use the formula given for the perihelion. Measure the perihelion distance to be  $a(1 - e) = 1.7 \text{ cm}$ . Dividing this by  $a = 3.4 \text{ cm}$ , we obtain  $1 - e = 0.5$ , so once again,  $e = 0.5$ .

**EXAMPLE 2** A (hypothetical) planet with a semimajor axis of 400 million km and an eccentricity of 0.5 (i.e., with an orbit as shown in the diagram) would range between  $400 \times (1 - 0.5) = 200$  million km and  $400 \times (1 + 0.5) = 600$  million km from the Sun over the course of one complete orbit. With  $e = 0.9$ , the range in distances would be 40 to 760 million km, and so on.

No planet has an orbital eccentricity as large as 0.5—the planet with the most eccentric orbit is Mercury, with  $e = 0.206$  (see Table 2.1). However, many meteoroids and all comets (see Chapter 14) have eccentricities considerably greater than that. In fact, most comets visible from Earth have eccentricities very close to  $e = 1$ . Their highly elongated orbits approach within a few astronomical units of the Sun at perihelion, yet these tiny frozen worlds spend most of their time far beyond the orbit of Pluto.



**Interactive** **FIGURE 2.16 Kepler's Second Law** A line joining a planet to the Sun sweeps out equal areas in equal intervals of time. The three shaded areas A, B, and C are equal. Any object traveling along the elliptical path would take the same amount of time to cover the distance indicated by the three red arrows. Planets move faster when closer to the Sun.

distance traveled by the planet along arc C is greater than the distance traveled along arc A or arc B. Because the time is the same and the distance is different, the speed must vary. When a planet is close to the Sun, as in sector C, it moves much faster than when farther away, as in sector A.

By taking into account the relative speeds and positions of the planets in their elliptical orbits about the Sun, Kepler's first two laws explained the variations in planetary brightness and some observed peculiar nonuniform motions that could not be accommodated within the assumption of circular motion, even with the inclusion of epicycles. Gone at last were the circles within circles that rolled across the sky. Kepler's modification of the Copernican theory to allow the possibility of elliptical orbits both greatly simplified the model of the solar system and at the same time provided much greater predictive accuracy than had previously been possible. Note, too, that these laws are not restricted to planets. They apply to *any* orbiting object. Spy satellites, for example, move very rapidly as they swoop close to Earth's surface, not because they are

TABLE 2.1 Some Solar System Dimensions

Planet	Orbital Semimajor Axis, $a$ (AU)	Orbital Period, $P$ (years)	Orbital Eccentricity, $e$	$P^2/a^3$
Mercury	0.387	0.241	0.206	1.002
Venus	0.723	0.615	0.007	1.001
Earth	1.000	1.000	0.017	1.000
Mars	1.524	1.881	0.093	1.000
Jupiter	5.203	11.86	0.048	0.999
Saturn	9.537	29.42	0.054	0.998
Uranus	19.19	83.75	0.047	0.993
Neptune	30.07	163.7	0.009	0.986

propelled with powerful onboard rockets, but because their highly eccentric orbits are governed by Kepler's laws.

Kepler published his first two laws in 1609, stating that he had proved them only for the orbit of Mars. Ten years later, he extended them to all the then-known planets (Mercury, Venus, Earth, Mars, Jupiter, and Saturn) and added a third law relating the size of a planet's orbit to its sidereal orbital **period**—the time needed for the planet to complete one circuit around the Sun:

The square of a planet's orbital period is proportional to the cube of its semimajor axis.

This law becomes particularly simple when we choose the (Earth sidereal) year as our unit of time and the *astronomical unit* as our unit of length. One **astronomical unit** (AU) is the semimajor axis of Earth's orbit around the Sun—essentially the average distance between Earth and the Sun. Like the light-year, the astronomical unit is custom-made for the vast distances encountered in astronomy. Using these units for time and distance, we can rewrite Kepler's third law for any planet as

$$P^2 \text{ (in Earth years)} = a^3 \text{ (in astronomical units)},$$

where  $P$  is the planet's sidereal orbital period and  $a$  is the length of its semimajor axis. The law implies that a planet's "year"  $P$  increases more rapidly than does the size of its orbit,  $a$ . For example, Earth, with an orbital semimajor axis of 1 AU, has an orbital period of 1 Earth year. The planet Venus, orbiting at a distance of roughly 0.7 AU, takes only 0.6 Earth year—about 225 days—to complete one circuit. By contrast, Saturn, almost 10 AU from the Sun, takes considerably more than 10 Earth years—in fact, nearly 30 years—to orbit the Sun just once.

Table 2.1 presents basic data describing the orbits of the eight planets now known. Renaissance astronomers knew these properties for the innermost six planets and used them to construct the currently accepted heliocentric model of

the solar system. The second column presents each planet's orbital semimajor axis, measured in astronomical units; the third column gives the orbital period, in Earth years. The fourth column lists the planets' orbital eccentricities. For purposes of verifying Kepler's third law, the fifth column lists the ratio  $P^2/a^3$ . As we have just seen, the third law implies that this number should always be unity in the units used in the table.

The main points to be grasped from Table 2.1 are these: (1) With the exception of Mercury, the planets' orbits are nearly circular (i.e., their eccentricities are close to zero) and (2) the farther a planet is from the Sun, the greater is its orbital period, in agreement with Kepler's third law to within the accuracy of the numbers in the table. (The small, but significant, deviations of  $P^2/a^3$  from unity in the cases of Uranus and Neptune are caused by the gravitational attraction between those two planets; see Chapter 13.) Most important, note that Kepler's laws are obeyed by *all* the known planets, *not just by the six on which he based his conclusions*.

The laws developed by Kepler were far more than mere fits to existing data. They also made definite, testable predictions about the future locations of the planets. Those predictions have been borne out to high accuracy every time they have been tested by observation—the hallmark of any credible scientific theory. ∞ (Sec. 1.2)

### CONCEPT Check


✓ Why is it significant that Kepler's laws also apply to Uranus and Neptune?

## 2.6 The Dimensions of the Solar System

Kepler's laws allow us to construct a scale model of the solar system, with the correct shapes and *relative* sizes of all the planetary orbits, but they do not tell us the *actual* size of any orbit. We can express the distance to each planet only in terms of the distance from Earth to the Sun. Why is this? Because Kepler's triangulation measurements all used a portion of Earth's orbit as a baseline, distances could be expressed only relative to the size of that orbit, which was itself not determined by the method. Thus, our model of the solar system would be like a road map of the United States showing the *relative* positions of cities and towns, but lacking the all-important scale marker indicating distances in kilometers or miles. For example, we would know that Kansas City is about three times more distant from New York than it is from Chicago, but we would not know the actual mileage between any two points on the map.

If we could somehow determine the value of the astronomical unit—in kilometers, say—we would be able to add the vital scale marker to our map of the solar system and compute the precise distances between the Sun and each of






▲ **FIGURE 2.17 Solar Transit** The transit of Mercury across the face of the Sun. Such transits happen only about once per decade because Mercury's orbit does not quite coincide with the plane of the ecliptic. Transits of Venus are even rarer, occurring only about twice per century. The most recent took place in 2006. (*P. Jones*)

the planets. We might propose using triangulation to measure the distance from Earth to the Sun directly. However, we would find it impossible to measure the Sun's parallax using Earth's diameter as a baseline. The Sun is too bright, too big, and too fuzzy for us to distinguish any apparent displacement relative to a field of distant stars. To measure the Sun's distance from Earth, we must resort to some other method.

Before the middle of the 20th century, the most accurate measurements of the astronomical unit were made by using triangulation on the planets Mercury and Venus during their rare *transits* of the Sun—that is, during the brief periods when those planets passed directly between the Sun and Earth (as shown for the case of Mercury in Figure 2.17). Because the time at which a transit occurs can be measured with great precision, astronomers can use this information to make accurate measurements of a planet's position in the sky. They can then employ simple geometry to compute the distance to the planet by combining observations made from different locations on Earth, as discussed earlier in Chapter 1. ∞ (Sec. 1.6) For example, the parallax of Venus at closest approach to Earth, as seen from two diametrically opposite points on Earth (separated by about 13,000 km), is about  $1'$  ( $1/60^\circ$ )—at the limit of naked-eye capabilities, but easily measurable telescopically. Using the second formula presented in *More Precisely 1-2*, we find that this parallax represents a distance of  $13,000 \text{ km} \times 57.3^\circ/(1/60^\circ)$ , or approximately 45,000,000 km.

Knowing the distance to Venus, we can compute the magnitude of the astronomical unit. Figure 2.18 is an idealized diagram of the Sun–Earth–Venus orbital geometry. The planetary orbits are drawn as circles here, but in reality they are slight ellipses. This is a subtle difference, and



▲ **FIGURE 2.18 Astronomical Unit** The wavy blue lines represent the paths along which radar signals are transmitted toward Venus and received back at Earth when Venus is at its minimum distance from Earth. Because the radius of Earth's orbit is 1 AU and that of Venus is about 0.7 AU, the one-way distance covered by the signal is 0.3 AU. Thus, we can calibrate the astronomical unit in kilometers.

we can correct for it using detailed knowledge of orbital motions. Assuming for the sake of simplicity that the orbits are perfect circles, we see from the figure that the distance from Earth to Venus at closest approach is approximately 0.3 AU. Knowing that 0.3 AU is 45,000,000 km makes determining 1 AU straightforward—the answer is  $45,000,000/0.3$ , or 150,000,000 km.

The modern method for deriving the absolute scale (that is, the scale expressed in kilometers, rather than just relative to Earth's orbit) of the solar system uses radar rather than triangulation. The word **radar** is an acronym for **radio detection and ranging**. In this technique, radio waves are transmitted toward an astronomical body, such as a planet. (We cannot use radar ranging to measure the distance to the Sun directly, because radio signals are absorbed at the solar surface and are not reflected to Earth.) The returning echo indicates the body's direction and range, or distance, in absolute terms—that is, in kilometers rather than in astronomical units. Multiplying the 300-second round-trip travel time of the radar signal (the time elapsed between transmission of the signal and reception of the echo) by the speed of light (300,000 km/s, which is also the speed of radio waves), we obtain twice the distance to the target planet.

Venus, whose orbit periodically brings it closest to Earth, is the most common target for radar ranging. The round-trip travel time (for example, at closest approach, as indicated by the wavy lines in Figure 2.18) can be measured with high

precision—in fact, well enough to determine the planet's distance to an accuracy of about 1 km. In this way, the astronomical unit is now known to be 149,597,870 km. We will use the rounded-off value of  $1.5 \times 10^8$  km in this text.

Having determined the value of the astronomical unit, we can reexpress the sizes of the other planetary orbits in terms of more familiar units, such as miles or kilometers. The entire scale of the solar system can then be calibrated to high precision.

### CONCEPT Check

✓ Why don't Kepler's laws tell us the value of the astronomical unit?

## 2.7 Newton's Laws

Kepler's three laws, which so simplified the solar system, were discovered *empirically*. In other words, they resulted solely from the analysis of observational data and were not derived from any theory or mathematical model. Indeed, Kepler did not have any appreciation of the physics underlying his laws. Nor did Copernicus understand *why* his heliocentric model of the solar system worked. Even Galileo, often called the father of modern physics, failed to understand why the planets orbit the Sun (although Galileo's work laid vital groundwork for Newton's theories).

What prevents the planets from flying off into space or from falling into the Sun? What causes them to revolve about the Sun, apparently endlessly? To be sure, the motions of the planets obey Kepler's three laws, but only by considering something more fundamental than those laws can we really understand planetary motion. The heliocentric system was secured when, in the 17th century, the British mathematician Isaac Newton (Figure 2.19) developed a deeper understanding of the way *all* objects move and interact with one another.

### The Laws of Motion

Isaac Newton was born in Lincolnshire, England, on Christmas Day in 1642, the year Galileo died. Newton studied at Trinity College of Cambridge University, but when the bubonic plague reached Cambridge in 1665, he returned to the relative safety of his home for 2 years. During that time he made probably the most famous of his discoveries, the law of gravity (although it is but one of the many major scientific advances for which Newton was responsible). However, either because he regarded the theory as incomplete or possibly because he was afraid that he would be attacked or plagiarized by his colleagues, he did not tell anyone of his monumental achievement for almost 20 years. It was not until 1684, when Newton was discussing the leading astronomical



▲ FIGURE 2.19 Isaac Newton (1642–1727). (S. Terry)


problem of the day—*Why* do the planets move according to Kepler's laws?—with Edmund Halley (of Halley's comet fame) that he astounded his companion by revealing that he had solved the problem in its entirety nearly two decades before!

Prompted by Halley, Newton published his theories in perhaps the most influential physics book ever written: *Philosophiae Naturalis Principia Mathematica* (*The Mathematical Principles of Natural Philosophy*—what we would today call “science”), usually known simply as Newton's *Principia*. The ideas expressed in that work form the basis for what is now known as **Newtonian mechanics**. Three basic laws of motion, the law of gravity, and a little calculus (which Newton also developed) are sufficient to explain and quantify virtually all the complex dynamic behavior we see on Earth and throughout the universe.

Figure 2.20 illustrates Newton's first law of motion:

Every body continues in a state of rest or in a state of uniform motion in a straight line, unless it is compelled to change that state by a force acting on it.

The first law simply states that a moving object will move forever in a straight line, unless some external **force**—a push or a pull—changes its speed or direction of motion. For example, the object might glance off a brick wall or be hit with a baseball bat; in either case, a force changes the original motion of the object. Another example of a force, well known to most of us, is **weight**—the force (commonly



▲ **FIGURE 2.20 Newton's First Law** An object at rest will remain at rest (a) until some force acts on it. When a force (represented by the red arrow) does act (b), the object will remain in that state of uniform motion until another force acts on it. When a second force (green arrow) acts in a direction different from the first force (c), the object changes its direction of motion.

measured in pounds in the United States) with which gravity pulls you toward Earth's center.

The tendency of an object to keep moving at the same speed and in the same direction unless acted upon by a force is known as **inertia**. Newton's first law implies that it requires no force to maintain motion in a straight line with constant speed. This contrasts sharply with the view of Aristotle, who maintained (incorrectly) that the natural state of an object was to be *at rest*—most probably an opinion based on Aristotle's observations of the effect of friction. In our discussion, we will neglect friction—the force that slows balls rolling along the ground, blocks sliding across tabletops, and baseballs moving through the air. In any case, it is not an issue for the planets because there is no appreciable friction in outer space—there is no air or any other matter to impede a planet's motion. The fallacy in Aristotle's argument was first realized and exposed by Galileo, who conceived of the notion of inertia long before Newton formalized it into a law.

A familiar measure of an object's inertia is its **mass**—loosely speaking, the total amount of matter the object contains. The greater an object's mass, the more inertia it has, and the greater is the force needed to change its state of motion.

Newton's first law describes motion in a straight line with constant speed—that is, motion with constant **velocity**. An object's velocity includes both its speed (in miles per hour or meters per second, say) *and* its direction in space (up, down, northwest, and so on). In everyday speech, we tend to use the terms “speed” and “velocity” more or less interchangeably, but we must realize that they are actually different quantities and that Newton's laws are always stated in terms of the latter. As a specific illustration of the difference, consider a rock tied to a string, moving at a constant rate in a circle as you whirl it around your head. The rock's *speed* is constant, but its *direction* of motion, and hence its velocity, is continually changing. Thus, according to

Newton's first law, a force must be acting. That force is the tension you feel in the string. In a moment, we'll see reasoning similar to this applied to the problem of planetary motion.

The rate of change of the velocity of an object—speeding up, slowing down, or simply changing direction—is called the object's **acceleration** and is the subject of Newton's second law, which states that the acceleration of an object is directly proportional to the applied force and inversely proportional to its mass:

When a force  $F$  acts on a body of mass  $m$ , it produces in it an acceleration  $a$  equal to the force divided by the mass. Thus,  $a = F/m$ , or  $F = ma$ .

Hence, the greater the force acting on the object or the smaller the mass of the object, the greater is the acceleration of the object. If two objects are pulled with the same force, the more massive one will accelerate less; if two identical objects are pulled with different forces, the one acted on by the greater force will accelerate more.

Acceleration is the rate of change of velocity, so its units are velocity units per unit of time, such as meters per second *per second* (usually written as  $\text{m/s}^2$ ). In honor of Newton, the SI unit of force is named after him. By definition, 1 newton (N) is the force required to cause a mass of 1 kilogram to accelerate at a rate of 1 meter per second every second ( $1 \text{ m/s}^2$ ). One newton is approximately 0.22 pound.

At Earth's surface, the force of gravity produces a downward acceleration of approximately  $9.8 \text{ m/s}^2$  on *all* bodies, regardless of mass. According to Newton's second law, this means that your weight (in newtons) is directly proportional to your mass (in kilograms). We will return to this very important point later.

Finally, Newton's third law simply tells us that forces cannot occur in isolation:

To every action, there is an equal and opposite reaction.

In other words, if body  $A$  exerts a force on body  $B$ , then body  $B$  necessarily exerts a force on body  $A$  that is equal in magnitude, but oppositely directed. For example, when a baseball player hits a home run, Newton's third law says that the bat and the ball exert equal and opposite forces on one another during the instant they are in contact. According to the second law, the ball subsequently moves away much faster than the bat because the *mass* of the ball is much less than the combined mass of the bat plus batter (whose body absorbs much of the reaction force), so the ball's *acceleration* is much greater.


Only in extreme circumstances—when speeds approach the speed of light—do Newton's laws break down, and this

fact was not realized until the 20th century, when Albert Einstein's theories of relativity once again revolutionized our view of the universe (see Chapter 22). Most of the time, however, Newtonian mechanics provides an excellent description of the motions of planets, stars, and galaxies through the cosmos.

## Gravity

Forces may act *instantaneously* or *continuously*. The force from the baseball bat that hits the home run can reasonably be thought of as being instantaneous. A good example of a continuous force is the one that prevents the baseball from zooming off into space—**gravity**, the phenomenon that started Newton on the path to the discovery of his laws. Newton hypothesized that any object having mass always exerts an attractive **gravitational force** on all other massive objects. The more massive an object, the stronger is its gravitational pull.

Consider a baseball thrown upward from Earth's surface, as illustrated in Figure 2.21. In accordance with Newton's first law, the downward force of Earth's gravity continuously modifies the baseball's velocity, slowing the initial upward motion and eventually causing the ball to fall back to the ground. Of course, the baseball, having some mass of its own, also exerts a gravitational pull on Earth. By Newton's third law, this force is equal and opposite to the weight of the ball (the force with which Earth attracts it). But, by Newton's second law, Earth has a much greater effect on the light



▲ **FIGURE 2.21 Gravity** A ball thrown up from the surface of a massive object, such as a planet, is pulled continuously downward (arrows) by the gravity of that planet—and, conversely, the gravity of the ball continuously pulls the planet (although very, very little).

baseball than the baseball has on the much more massive Earth. The ball and Earth act upon each other with the same gravitational force, but Earth's acceleration is much smaller.

Now consider the trajectory of the same baseball batted from the surface of the Moon. The pull of gravity is about one-sixth as great on the Moon as on Earth, so the baseball's velocity changes more slowly—a typical home run in a ballpark on Earth would travel nearly half a mile on the Moon. Less massive than Earth, the Moon has less gravitational influence on the baseball. The magnitude of the gravitational force, then, depends on the *masses* of the attracting bodies. In fact, the force is *directly proportional* to the product of the two masses.

Studying the motions of the planets uncovers a second aspect of the gravitational force. At locations equidistant from the Sun's center, the gravitational force has the same strength and is always directed toward the Sun. Furthermore, detailed calculation of the planets' accelerations as they orbit the Sun reveals that the strength of the Sun's gravitational pull decreases in proportion to the *square* of the distance from the Sun. The force of gravity is said to obey an **inverse-square law**. As shown in Figure 2.22, inverse-square forces decrease rapidly with distance from their source. For example, tripling the distance makes the force  $3^2 = 9$  times weaker, whereas multiplying the distance by five results in a force that is  $5^2 = 25$  times weaker. Despite this rapid decrease, the force never quite reaches zero. The gravitational pull of an object having some mass can never be completely extinguished.

We can combine the preceding statements about mass and distance to form a law of gravity that dictates the way in which *all* massive objects (i.e., objects having some mass) attract one another:

Every particle of matter in the universe attracts every other particle with a force that is directly proportional to the product of the masses of the particles and inversely proportional to the square of the distance between their centers.



As a proportionality, the law of gravity may be written as

$$\text{gravitational force} \propto \frac{\text{mass of object 1} \times \text{mass of object 2}}{\text{distance}^2}$$

(The symbol  $\propto$  here means “is proportional to.”) The rule for computing the force  $F$  between two bodies of masses  $m_1$  and  $m_2$ , separated by a distance  $r$ , is usually written more compactly as

$$F = \frac{Gm_1m_2}{r^2}.$$

The quantity  $G$  is known as the *gravitational constant*, or, often, simply as Newton's constant. It is one of the fundamental constants of the universe. The value of  $G$  has been measured in extremely delicate laboratory experiments as  $6.67 \times 10^{-11}$  newton meter<sup>2</sup>/kilogram<sup>2</sup> ( $\text{N} \cdot \text{m}^2/\text{kg}^2$ ).



**▲ FIGURE 2.22 Gravitational Force** (a) The gravitational force between two bodies is proportional to the mass of each and is inversely proportional to the square of the distance between them. (b) Inverse-square forces rapidly weaken with distance from their source, never quite reaching zero, no matter how far away.

## 2.8 Newtonian Mechanics

Newton's three laws of motion and the law of gravitation provide a solid theoretical foundation upon which we can base a deeper understanding of planetary orbits, the laws of planetary motion, and many other important aspects of orbital motion. With the development of Newtonian mechanics, the transition from geocentric lore to heliocentric fact was complete.

### Planetary Motion

The mutual gravitational attraction between the Sun and the planets, as expressed by Newton's law of gravity, is responsible for the observed planetary orbits. As depicted in Figure 2.23, this gravitational force continuously pulls each planet toward


the Sun, deflecting its forward motion into a curved orbital path. Because the Sun is much more massive than any of the planets, it dominates the interaction. We might say that the Sun “controls” the planets, not the other way around.

The Sun–planet interaction sketched here is analogous to our earlier example of the rock whirling on a string. The Sun's gravitational pull is your hand and the string, and the planet is the rock at the end of that string. The tension in the string provides the force necessary for the rock to move in a circular path. If you were suddenly to release the string—which would be like eliminating the Sun's gravity—the rock would fly away along a tangent to the circle, in accordance with Newton's first law.

In the solar system, at this very moment, Earth is moving under the combined influence of gravity and inertia. The net result is a stable orbit, despite our continuous rapid motion through space. (In fact, Earth orbits the Sun at a speed of about 30 km/s, or approximately 70,000 mph. You can verify this for yourself by calculating how fast Earth must move to complete a circle of radius 1 AU—and hence of circumference  $2\pi$  AU, or 940 million km—in 1 year, or  $3.2 \times 10^7$  seconds. The answer is  $9.4 \times 10^8$  km /  $3.2 \times 10^7$  s, or 29.4 km/s.) *More Precisely 2-2* describes how astronomers can use Newtonian mechanics and the law of gravity to quantify planetary motion and measure the masses of Earth, the Sun, and many other astronomical objects by studying the orbits of objects near them.

### Kepler's Laws Reconsidered

Newton's laws of motion and law of universal gravitation provide a theoretical explanation for Kepler's empirical laws of planetary motion. Kepler's three laws follow directly from




**▲ FIGURE 2.23 Solar Gravity** The Sun's inward pull of gravity on a planet competes with the planet's tendency to continue moving in a straight line. These two effects combine, causing the planet to move smoothly along an intermediate path, which continuously “falls around” the Sun. This unending tug-of-war between the Sun's gravity and the planet's inertia results in a stable orbit.


Newtonian mechanics, as solutions of the equations describing the motion of a body moving in response to an inverse-square force. However, just as Kepler modified the Copernican model by introducing ellipses rather than circles, so, too, did Newton make corrections to Kepler's first and third laws. It turns out that a planet does not orbit the exact center of the Sun. Instead, both the planet and the Sun orbit their common **center of mass**—the “average” position of all the matter making up the two bodies (Figure 2.24). Because the Sun and the planet are acted upon by equal and opposite gravitational forces (by Newton's third law), the Sun must also move (by Newton's first law), driven by the gravitational influence of the planet. The Sun, however, is so much more massive than any planet that the center of mass of the planet–Sun system is very close to the center of the Sun, which is why Kepler's laws are so accurate. Thus, Kepler's first law becomes

The orbit of a planet around the Sun is an ellipse, with the *center of mass of the planet–Sun system* at one focus.

As shown in Figure 2.24, the center of mass of two objects of comparable mass does not lie within either object. For identical masses orbiting one another (Figure 2.25a), the orbits are identical ellipses, with a common focus located midway between the two objects. For unequal masses (as in Figure 2.25b), the elliptical orbits still share a focus, and both have the same eccentricity, but the more massive object moves more slowly and on a tighter orbit. (Note that Kepler's second law, as stated earlier, continues to apply without modification to each orbit separately, but the *rates* at which the two orbits sweep out areas are different.)



(a) Equal masses Center of mass



(b) Unequal masses Center of mass


**▲ FIGURE 2.24 Center of Mass** (a) The center of mass of two bodies of equal mass lies midway between them. (b) As the mass of one body increases, the center of mass moves toward it. Experienced seesawers know that when both sides are balanced, the center of mass is at the pivot point.

The change to Kepler's third law is also small in the case of a planet orbiting the Sun, but very important in other circumstances, such as the orbital motion of two stars that are gravitationally bound to each other. Following through the mathematics of Newton's theory, we find that the true relationship between the semimajor axis  $a$  (measured in astronomical units) of the planet's orbit relative to the Sun and its orbital period  $P$  (in Earth years) is

$$P^2 \text{ (in Earth years)} = \frac{a^3 \text{ (in astronomical units)}}{M_{\text{total}} \text{ (in solar units)}}$$

where  $M_{\text{total}}$  is the *combined* mass of the two objects. Notice that Newton's restatement of Kepler's third law preserves the proportionality between  $P^2$  and  $a^3$ , but now the proportionality includes  $M_{\text{total}}$ , so it is *not* quite the same for all the

*The resulting orbits for mutually gravitating bodies depend on their masses and distances from each other.*



**MA** **Interactive FIGURE 2.25 Orbits** (a) The orbits of two bodies (stars, for example) with equal masses, under the influence of their mutual gravity, are identical ellipses with a common focus. The pairs of numbers (e.g., the two 2s in each orbit) indicate the positions of the two bodies at three different times. (Note that a line joining the bodies at any give time always passes through the common focus.) (b) The orbits of two bodies, one twice as massive as the other, are again elliptical and with the same eccentricity, but according to Newton's laws, the more massive body moves more slowly and in a smaller orbit. (c) In the case of an extremely small planet orbiting the massive Sun, the common focus of the two orbits could be inside the Sun.

## MORE PRECISELY 2-2

### Weighing the Sun

We can use Newtonian mechanics to calculate some useful formulae relating the properties of planetary orbits to the mass of the Sun. Again for simplicity, let's assume that the orbits are circular (not a bad approximation in most cases, and Newton's laws easily extend to cover the more general case of eccentric orbits). Consider a planet of mass  $m$  moving at speed  $v$  in an orbit of radius  $r$  around the Sun, of mass  $M$ . Even though the planet's *speed* is constant, the *direction* of its motion is not, so the planet's velocity is changing—it is accelerating. In fact, the planet's acceleration is

$$a = \frac{v^2}{r},$$

so, by Newton's second law, the force required to keep the planet in orbit is

$$F = ma = \frac{mv^2}{r}.$$

Setting this equation equal to the gravitational force due to the Sun, we obtain

$$\frac{mv^2}{r} = \frac{GmM}{r^2},$$

so the speed of the planet in the circular orbit is

$$v = \sqrt{\frac{GM}{r}}.$$

planets. The Sun's mass is so great, however, that the differences in  $M_{\text{total}}$  among the various combinations of the Sun and the other planets are almost unnoticeable, so Kepler's third law, as originally stated, is a very good approximation. This modified form of Kepler's third law is true in all circumstances, inside or outside the solar system.

### PROCESS OF SCIENCE Check

✓ In what ways did Newtonian mechanics supersede Kepler's laws as a model of the solar system?

### Escaping Forever

The law of gravity that describes the orbits of planets around the Sun applies equally well to natural moons and artificial satellites orbiting any planet. All our Earth-orbiting, human-made satellites move along paths governed by a combination of the inward pull of Earth's gravity and the forward motion gained during the rocket launch. If the rocket initially imparts enough speed to the satellite, it can go into orbit. Satellites not given enough speed at launch, by accident or design (e.g., intercontinental ballistic

missiles), fail to achieve orbit and fall back to Earth (see Figure 2.26).

$$M = \frac{rv^2}{G}$$

Now let's turn the problem around. Because we have measured  $G$  in the laboratory on Earth, and because we know the length of a year and the size of the astronomical unit, we can use Newtonian mechanics to *weigh* the Sun—that is, find its mass by measuring its gravitational influence on another body (in this case, Earth). Rearranging the last equation to read

and substituting the known values of  $v = 30$  km/s,  $r = 1$  AU =  $1.5 \times 10^{11}$  m, and  $G = 6.7 \times 10^{-11}$  Nm<sup>2</sup>/kg<sup>2</sup>, we calculate the mass of the Sun to be  $2.0 \times 10^{30}$  kg—an enormous mass by terrestrial standards.


**EXAMPLE** Similarly, knowing the distance from Earth to the Moon ( $r = 384,000$  km) and the length of the (sidereal) month ( $P = 27.3$  days), we can calculate the Moon's orbital speed to be  $v = 2\pi r/P = 1.02$  km/s, and hence, using the preceding formula, measure Earth's mass to be  $6.0 \times 10^{24}$  kg.

In fact, this is basically how *all* masses are measured in astronomy. Because we can't just go out and attach a scale to an astronomical object when we need to know its mass, we must look for its gravitational influence on something else. This principle applies to planets, stars, galaxies, and even clusters of galaxies—very different objects, but all subject to the same physical laws.

Some space vehicles, such as the robot probes that visit other planets, attain enough speed to escape our planet's gravity and move away from Earth forever. This speed, known as the **escape speed**, is about 41% greater (actually,  $\sqrt{2} = 1.414 \dots$  times greater) than the speed of an object traveling in a circular orbit at any given radius.\* At less than the escape speed, the old adage "What goes up must come down" (or at least stay in orbit) still applies. At more than the escape speed, however, a spacecraft will leave Earth for good. Planets, stars, galaxies—all gravitating bodies—have escape speeds. No matter how massive the body, gravity decreases with distance. As a result, the escape speed diminishes with increasing separation. The farther we go from Earth (or any gravitating body), the easier it becomes to escape.

The speed of a satellite in a circular orbit just above Earth's atmosphere is 7.9 km/s (roughly 18,000 mph). The satellite would have to travel at 11.2 km/s (about 25,000 mph) to escape from Earth altogether. If an object exceeds the escape speed, its motion is said to be **unbound**, and the orbit is no longer

\*In terms of the formula presented in More Precisely 2-2, the escape speed is given by  $v_{\text{escape}} = \sqrt{2GM/r}$ .



**Interactive FIGURE 2.26 Escape Speed** The effect of launch speed on the trajectory of a satellite. With too low a speed at point A, the satellite will simply fall back to Earth. Given enough speed, however, the satellite will go into orbit—it “falls around Earth.” As the initial speed at point A is increased, the orbit will become more and more elongated. When the initial speed exceeds the escape speed, the satellite will become unbound from Earth and will escape along a hyperbolic trajectory.

an ellipse. In fact, the path of the spacecraft relative to Earth is a related geometric figure called a *hyperbola*. If we simply change the word *ellipse* to *hyperbola*, the modified version of Kepler’s first law still applies, as does Kepler’s second law. (Kepler’s third law does not extend to unbound orbits because it doesn’t make sense to talk about a period in those cases.)

## The Circle of Scientific Progress

The progression from the complex Ptolemaic model of the universe to the elegant simplicity of Newton’s laws is a case

study in the scientific method. [↻ \(Sec. 1.2\)](#) Copernicus made a radical conceptual leap away from the Ptolemaic view, gaining much in insight but little in predictive power. Kepler made critical changes to the Copernican picture and gained both accuracy and predictive power but still fell short of a true physical explanation of planetary motion within the solar system, or of orbital motion in general. Eventually, Newton showed how all known planetary motion could be explained in detail by the application of four, simple, fundamental laws—the three laws of motion and the law of gravity. The process was slow, with many starts and stops and a few wrong turns, but it worked!

In a sense, then, the development of Newton’s laws and their application to planetary motion represented the end of the first “loop” around the schematic diagram presented in Figure 1.6. The long-standing practical and conceptual questions raised by ancient observations of retrograde motion were finally resolved, and new predictions, themselves amenable to observational testing, became possible. And the laws are still being tested today. Every time a comet appears in the night sky right on schedule, or a spacecraft reaches the end of a billion-kilometer journey within meters of its target and seconds of the predicted arrival time, our confidence in Newtonian mechanics is further strengthened.

But unlike the essentially descriptive models of Ptolemy, Copernicus, and Kepler, Newtonian mechanics is not limited to the motions of planets, or even to events occurring within our own solar system. They apply to moons, comets, spacecraft, stars, and even the most distant galaxies, extending the range of our scientific inquiries across the observable universe—as well as apples falling to the ground.

### CONCEPT Check

- ✓ Explain, in terms of Newton’s laws of motion and gravity, why planets orbit the Sun.


**The Big Question** The concept of gravity is well developed, thanks to Isaac Newton’s efforts in the 17th century; it works well for small masses and small velocities, thus for almost every application on or near Earth. But, as we shall see later in this book, in the 20th century Albert Einstein overthrew the idea of gravity with a better one—relativity theory, which deals with fast-moving, often massive objects in curved space. Even so, now in the 21st century, relativity is coming up short, especially when studying exotic objects such as black holes. Who will take the next great leap in understanding and what will the new concept be? No one knows the answers, but this is the way the scientific method works, by constantly refining our ideas about the nature of the universe.




# Chapter Review

## SUMMARY


**1** **Geocentric** (p. 38) models of the universe had the Sun, the Moon, and the planets all orbiting Earth. The most successful of these was the **Ptolemaic model** (p. 38). Planets sometimes appear to temporarily reverse their motion relative to the stars and later resume their normal “forward” course. This is called **retrograde motion** (p. 37). Geocentric models explained retrograde motion as a real backward motion of a planet as it moved along its epicyclic path around Earth.




**2** The **heliocentric** (p. 39) view of the solar system, due to Aristarchus and later Copernicus, holds that Earth, like all the planets, orbits the Sun. This model naturally explains both retrograde motion as Earth overtakes other planets in its orbit and the observed brightness variations of the planets. The widespread realization during the Renaissance that the solar system is Sun centered, and not Earth centered, is known as the **Copernican revolution** (p. 39).



**3** Galileo’s telescopic observations of the Moon, the Sun, Venus, and Jupiter played a crucial role in supporting and strengthening the Copernican picture of the solar system. Johannes Kepler improved on Copernicus’s model by condensing the observational data of Tycho Brahe into three **laws of planetary motion** (p. 45).




**4** Kepler’s Laws state: (1) Planetary orbits are **ellipses** (p. 45) with the Sun at one **focus** (p. 45); (2) a planet moves faster as its orbit takes it closer to the Sun; (3) the




**semimajor axis** (p. 45) of the orbit is related in a simple way to the planet’s orbital **period** (p. 47). Most planetary orbits differ only slightly from perfect circles.

**5** The average distance from Earth to the Sun is one **astronomical unit** (p. 47), today most accurately determined by bouncing **radar** (p. 48) signals off the planet Venus. Once this distance is known, the distances to all other planets can be inferred from Kepler’s laws.



**6** To change the body’s velocity, a force (p. 48) must be applied. The rate of change of velocity, called **acceleration** (p. 50), is equal to the applied force divided by the body’s **mass** (p. 50). When bodies interact, the forces between them are always equal and opposite to one another. To explain Kepler’s laws, Newton postulated that **gravity** (p. 51) attracts the planets to the Sun. Every object having any mass exerts a **gravitational force** (p. 51) on all other objects. The strength of this force decreases with distance according to an **inverse-square law** (p. 51).



**7** For one object to escape from the gravitational pull of another, its speed must exceed the **escape speed** (p. 54) of that other object. By determining the gravitational force needed to keep one body orbiting another, Newton’s laws allow astronomers to measure the masses of distant objects.



**MasteringAstronomy**® For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information.

**LO** connects to the introduction’s numbered Learning Outcomes.

## Review and Discussion

- LO1** What contributions to astronomy were made by Chinese and Islamic astronomers during the Dark Ages in Europe?
- Briefly describe the geocentric model of the universe.
- POS** The benefit of our current knowledge lets us see flaws in the Ptolemaic model of the universe. What is its basic flaw?
- LO2** What was the great contribution of Copernicus to our knowledge of the solar system? How was his model still flawed?
- What is a theory? Can a theory ever be proved to be true?
- What is the Copernican principle?
- LO3 POS** What discoveries of Galileo helped confirm the views of Copernicus, and how did they do so?
- LO4** Briefly describe Kepler’s three laws of planetary motion.
- How did Tycho Brahe contribute to Kepler’s laws?
- LO5** If radio waves cannot be reflected from the Sun, how can radar be used to find the distance from Earth to the Sun?
- POS** What does it mean to say that Kepler’s laws are empirical?
- LO6** What are Newton’s laws of motion and gravity?
- Why do we say that a baseball falls toward Earth, and not Earth toward the baseball?
- LO7** In what sense is the Moon falling toward Earth? How can we use this fact to measure Earth’s mass?
- What is the meaning of the term *escape speed*?

## Conceptual Self-Test: Multiple Choice

- Planets near opposition (a) rise in the east; (b) rise in the west; (c) do not rise or set; (d) have larger deferents.
- A major flaw in Copernicus's model was that it still had (a) the Sun at the center; (b) Earth at the center; (c) retrograde loops; (d) circular orbits.
- VIS** As shown in Figure 2.12 ("Venus Phases"), Galileo's observations of Venus demonstrated that Venus must be (a) orbiting Earth; (b) orbiting the Sun; (c) about the same diameter as Earth; (d) similar to the Moon.
- An accurate sketch of Jupiter's orbit around the Sun would show (a) the Sun far off center; (b) an oval twice as long as it is wide; (c) a nearly perfect circle; (d) phases.
- A calculation of how long it takes a planet to orbit the Sun would be most closely related to Kepler's (a) first law of orbital shapes; (b) second law of orbital speeds; (c) third law of planetary distances; (d) first law of inertia.
- An asteroid with an orbit lying entirely inside Earth's (a) has an orbital semimajor axis of less than 1 AU; (b) has a longer orbital period than Earth's; (c) moves more slowly than Earth; (d) must have a highly eccentric orbit.
- If Earth's orbit around the Sun were twice as large as it is now, the orbit would take (a) less than twice as long; (b) two times longer; (c) more than two times longer to traverse.
- VIS** Figure 2.21 ("Gravity"), showing the motion of a ball near Earth's surface, depicts how gravity (a) increases with altitude; (b) causes the ball to accelerate downward; (c) causes the ball to accelerate upward; (d) has no effect on the ball.
- If the Sun *and its mass* were suddenly to disappear, Earth would (a) continue in its current orbit; (b) suddenly change its orbital speed; (c) fly off into space; (d) stop spinning.
- VIS** Figure 2.25(b) ("Orbits") shows the orbits of two stars of unequal masses. If one star has twice the mass of the other, then the more massive star (a) moves more slowly than; (b) moves more rapidly than; (c) has half the gravity of; (d) has twice the eccentricity of the less massive star.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

- Tycho Brahe's observations of the stars and planets were accurate to about 1 arc minute (1'). To what distance does this angle correspond at the distance of (a) the Moon; (b) the Sun; and (c) Saturn (at closest approach)?
- Halley's comet has a perihelion distance of 0.6 AU and an orbital period of 76 years. What is its greatest distance from the Sun?
- A spacecraft has an orbit that just grazes Earth's orbit at aphelion and just grazes Venus's orbit at perihelion. Assuming that Earth and Venus are in the right places at the right times, how long will the spacecraft take to travel from Earth to Venus?
- How long would a radar signal take to complete a round-trip between Earth and Mars when the two planets are 0.7 AU apart?
- What is the maximum possible parallax of Mercury during a solar transit, as seen from either end of a 3000-km baseline on Earth?
- The acceleration due to gravity at Earth's surface is  $9.80 \text{ m/s}^2$ . What is the acceleration at altitudes of (a) 100 km; (b) 1000 km; (c) 10,000 km? Earth's radius is 6400 km.
- The Moon's mass is  $7.4 \times 10^{22} \text{ kg}$  and its radius is 1700 km. What is the speed of a spacecraft moving in a circular orbit just above the lunar surface? What is the escape speed from the Moon?
- Use Newton's law of gravity to calculate the force of gravity between you and Earth. Convert your answer, which will be in newtons, to pounds, using the conversion 4.45 N equals 1 pound (1 lb). What do you normally call this force?

## Activities

### Collaborative

- Select what you believe to be Galileo's single most important astronomical observation, say why you think it was the most important, and explain, using sketches, what he observed.
- Use a small telescope to replicate Galileo's observations of Jupiter's four largest moons. Note the moons' brightnesses and their locations with respect to Jupiter. If you watch over a period of several nights, draw what you see; you'll notice that these moons change their positions as they orbit the giant planet.

### Individual

- Draw an ellipse (see Figure 1.11). You'll need two pins, a piece of string, and a pencil. Tie the string in a loop and place it around the pins. Place the pencil inside the loop and run it around the inside of the string, holding the loop taut. The two pins will be at the foci of the ellipse. How does the shape of the ellipse change as you vary the distance between the pins?



# Radiation

## INFORMATION FROM THE COSMOS

Astronomical objects are more than just things of beauty in the night sky. Planets, stars, and galaxies are of vital significance if we are to fully understand our place in the larger scheme of things in the universe. Each object is a source of information about the material aspects of our universe—its state of motion, its temperature, its chemical composition, and even its past history.

Much of this information comes to us in the form of light. When we look at the stars, the light we see actually began its journey to Earth decades, centuries—even millennia—ago. The faint rays from the most distant galaxies have taken billions of years to reach us. The stars and galaxies in the night sky show us the far away and the long ago. In this chapter, we begin our study of how astronomers extract information from the light emitted by astronomical objects. These basic concepts of radiation are central to modern astronomy.

**The Big Picture** Our human eyes actually see only a small part of the universe—we literally see optical, or visible, light. There is a much bigger picture to be sensed beyond visible light—invisible radiation such as heat or radio waves or X-rays. Many different kinds of radiation are constantly traveling through space, and detailed study of this wider range of visible and invisible information is the main way that astronomers study stars and other distant objects well beyond Earth.

**LEFT:** In about 5 billion years, the Sun will begin running out of fuel. Its hydrogen gas will become depleted at its center, causing the bulk of our old and decrepit star to slowly dissipate into space. We can actually watch such an amazing event by observing other stars that are now dying. This stunning image captures the Helix Nebula about 650 light-years away—but not in visible light. Rather, it shows invisible radiation emitted by the former star. Infrared radiation (mostly yellow in this image) captured by the *Spitzer Space Telescope* and ultraviolet radiation (mostly bluish) by the *Galaxy Evolution Explorer* satellite enable close inspection of the remarkable process of stellar death. (NASA, Caltech)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Outline the basic properties of wave motion.
- 2 Tell how electromagnetic radiation transfers energy and information through interstellar space.
- 3 Describe the major regions of the electromagnetic spectrum and explain how Earth's atmosphere affects our ability to make astronomical observations at different wavelengths.
- 4 Explain what is meant by the term "blackbody radiation" and describe the basic properties of such radiation.
- 5 Tell how we can determine the temperature of an object by observing the radiation it emits.
- 6 Describe how the relative motion between a source of radiation and its observer can change the perceived wavelength of the radiation, and explain the importance of this phenomenon to astronomy.

### MasteringAstronomy®


Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.

## 3.1 Information from the Skies

Figure 3.1 shows a galaxy in the constellation Andromeda. On a dark, clear night, far from cities or other sources of light, the Andromeda galaxy, as it is generally called, can be seen with the naked eye as a faint, fuzzy patch on the sky, comparable in diameter to the full Moon. Yet the fact that it is visible from Earth belies this galaxy's enormous distance from us: It lies roughly 2.5 million light-years away.

An object at such a distance is truly inaccessible in any realistic human sense. Even if a space probe could miraculously travel at the speed of light, it would need 2.5 million years to reach this galaxy and 2.5 million more to return with its findings. Considering that civilization has existed on Earth for less than 10,000 years, and its prospects for the next 10,000 are far from certain, even this unattainable technological feat would not provide us with a practical means of exploring other galaxies. Even the farthest reaches of our own galaxy, “only” a few tens of thousands of light-years distant, are effectively off limits to visitors from Earth, at least for the foreseeable future.

Given the practical impossibility of traveling to such remote parts of the universe, how do astronomers know anything about objects far from Earth? How do we obtain detailed information about planets, stars, or galaxies too distant for a personal visit or any kind of controlled experiment? The answer is that we use the laws of physics, as we know them here on Earth, to interpret the **electromagnetic radiation** emitted by those objects.



▲ **FIGURE 3.1 Andromeda Galaxy** The pancake-shaped Andromeda Galaxy lies about 2.5 million light-years away and contains a few hundred billion stars. (R. Gendler)

## Light and Radiation

*Radiation* is any way in which energy is transmitted through space from one point to another without the need for any physical connection between the two locations. The term *electromagnetic* just means that the energy is carried in the form of rapidly fluctuating *electric* and *magnetic* fields (to be discussed in more detail later in Section 3.2). Virtually all we know about the universe beyond Earth's atmosphere has been gleaned from painstaking analysis of electromagnetic radiation received from afar. Our understanding depends completely on our ability to decipher this steady stream of data reaching us from space.

How bright are the stars (or galaxies, or planets), and how hot? What are their masses? How rapidly do they spin, and what is their motion through space? What are they made of, and in what proportion? The list of questions is long, but one fact is clear: Electromagnetic theory is vital to providing the answers—without it, we would have no way of testing our models of the cosmos, and the modern science of astronomy simply would not exist. ∞ (Sec. 1.2)

**Visible light** is the particular type of electromagnetic radiation to which our human eyes happen to be sensitive. As light enters our eye, small chemical reactions triggered by the incoming energy send electrical impulses to the brain, producing the sensation of sight. But modern instruments (see Chapter 5) can also detect many forms of *invisible* electromagnetic radiation, which goes completely unnoticed by our eyes. **Radio**, **infrared**, and **ultraviolet** waves, as well as **X-rays** and **gamma rays**, all fall into this category.

Note that, despite the different names, the words *light*, *rays*, *radiation*, and *waves* all really refer to the same thing. The names are just historical accidents, reflecting the fact that it took many years for scientists to realize that these apparently very different types of radiation are in reality one and the same physical phenomenon. Throughout this text, we will use the general terms *light* and *electromagnetic radiation* more or less interchangeably.

## Wave Motion

Despite the early confusion still reflected in current terminology, scientists now know that all types of electromagnetic radiation travel through space in the form of **waves**. To understand the behavior of light, then, we must know a little about wave motion.

Simply stated, a wave is a way in which energy is transferred from place to place without the physical movement of mate-

rial from one location to another. In wave motion, the energy is carried by a *disturbance* of some sort. This disturbance, whatever its nature, occurs in a distinctive repeating pattern. Ripples on the surface of a pond, sound waves in air, and electromagnetic waves in space, despite their many obvious differences, all share this basic defining property.

Imagine a twig floating in a pond (Figure 3.2). A pebble thrown into the pond at some distance from the twig disturbs the surface of the water, setting it into up-and-down motion. This disturbance will move outward from the point of impact in the form of waves. When the waves reach the twig, some of the pebble's energy will be imparted to it, causing the twig to bob up and down. In this way, both energy and *information*—the fact that the pebble entered the water—are transferred from the place where the pebble landed to the location of the twig. We could tell that a pebble (or, at least, some object) had entered the water just by observing the twig. With a little additional physics, we could even estimate the pebble's energy.


A wave is not a physical object. No water traveled from the point of impact of the pebble to the twig—at any location on the surface, the water surface simply moved up and down as the wave passed. What, then, *did* move across the surface of the pond? As illustrated in the figure, the answer is that the wave was the *pattern* of up-and-down motion. This pattern was transmitted from one point to the next as the disturbance moved across the water.

Figure 3.3 shows how wave properties are quantified and illustrates some standard terminology. The wave's **period** is the number of seconds needed for the wave to repeat itself at any given point in space. The **wavelength** is the number of meters needed for the wave to repeat itself at a given moment in time. It can be measured as the distance

Waves ripple out from where a pebble hit the water . . .

. . . to where a twig is floating.


This insert shows a series of "snapshots" of the pond surface as the wave passes by.



**Interactive FIGURE 3.2 Water Wave** The passage of a wave across a pond causes the surface of the water to bob up and down, but there is no movement of water from one part of the pond to another.

between two adjacent wave *crests*, two adjacent wave *troughs*, or any other two similar points on adjacent wave cycles (e.g., the points marked  $\times$  in the figure). A wave moves a distance equal to one wavelength in one period. The maximum departure of the wave from the undisturbed state—still air, say, or a flat pond surface—is called the wave's **amplitude**.

The number of wave crests passing any given point per unit time is called the wave's **frequency**. If a wave of a given wavelength moves at high speed, then many crests pass per second and the frequency is high. Conversely, if



**Interactive FIGURE 3.3 Wave Properties** A typical wave has a direction of motion, wavelength, and amplitude. In one wave period, the entire pattern shown here moves one wavelength to the right.

the same wave moves slowly, then its frequency will be low. The frequency of a wave is just the reciprocal of the wave's period; that is,

$$\text{frequency} = \frac{1}{\text{period}}$$

Frequency is expressed in units of inverse time (that is, 1/second, or cycles per second), called hertz (Hz) in honor of the 19th-century German scientist Heinrich Hertz, who studied the properties of radio waves. Thus, a wave with a period of 5 seconds (5 s) has a frequency of (1/5) cycles/s = 0.2 Hz, meaning that one wave crest passes a given point in space every 5 seconds.

Because a wave travels one wavelength in one period, it follows that the *wave velocity* is simply equal to the wavelength divided by the period:

$$\text{velocity} = \frac{\text{wavelength}}{\text{period}}$$

Since the period is the reciprocal of the frequency, we can equivalently (and more commonly) write this relationship as

$$\text{velocity} = \text{wavelength} \times \text{frequency}.$$

Thus, if the wave in our earlier example had a wavelength of 0.5 m, its velocity would be (0.5 m)/(5 s), or (0.5 m)  $\times$  (0.2 Hz) = 0.1 m/s. In the case of electromagnetic radiation, the velocity is the speed of light. Notice that wavelength and wave frequency are *inversely* related—doubling one halves the other.


## Components of Visible Light

White light is a mixture of colors, which we conventionally divide into six major hues: red, orange, yellow, green, blue, and violet. As shown in Figure 3.4, we can separate a beam of white light into a rainbow of these basic colors—called a *spectrum* (plural, *spectra*)—by passing it through a prism. This experiment was first reported by Isaac Newton over 300 years ago. In principle, the original beam of white light could be recovered by passing the spectrum through a second prism to recombine the colored beams.

What determines the color of a beam of light? The answer is its frequency (or alternatively, its wavelength). We see different colors because our eyes react differently to electromagnetic waves of different frequencies. A prism splits a beam of light up into separate colors because light rays of different frequencies are bent, or *refracted*, slightly differently as they pass through the prism—red light the least, violet light the most. Red light has a frequency of roughly  $4.3 \times 10^{14}$  Hz, corresponding to a wavelength of about  $7.0 \times 10^{-7}$  m. Violet light, at the other end of the visible range, has nearly double the frequency— $7.5 \times 10^{14}$  Hz—and (since the speed of light is always the same) just over half the wavelength— $4.0 \times 10^{-7}$  m. The other colors we see have frequencies and wavelengths intermediate between these two extremes, spanning the entire *visible spectrum* shown in Figure 3.4. Radiation outside this range is invisible to human eyes.

Scientists often use a unit called the *nanometer* (nm) in describing the wavelength of light (see Appendix 2). There are  $10^9$  nanometers in 1 meter. An older unit called

▼ **FIGURE 3.4 Visible Spectrum** When passed through a prism, white light splits into its component colors, spanning red to violet in the visible part of the electromagnetic spectrum. The slit narrows the beam of radiation. The “rainbow” of colors projected on the screen is just a series of different-colored images of the slit.



the *angstrom* ( $1 \text{ \AA} = 10^{-10} \text{ m} = 0.1 \text{ nm}$ ) is also widely used. (The unit is named after the 19th-century Swedish physicist Anders Ångström—pronounced “ong-strem.”) However, in SI units, the nanometer is preferred. Thus, the visible spectrum covers the range of wavelengths from 400 nm to 700 nm ( $4000 \text{ \AA}$  to  $7000 \text{ \AA}$ ). The radiation to which our eyes are most sensitive has a wavelength near the middle of this range, at about 550 nm ( $5500 \text{ \AA}$ ), in the yellow-green region of the spectrum. It is no coincidence that this wavelength falls within the range of wavelengths at which the Sun emits most of its electromagnetic energy—our eyes have evolved to take greatest advantage of the available light.


## 3.2 Waves in What?

Waves of radiation differ fundamentally from water waves, sound waves, or any other waves that travel through a material medium. Radiation needs *no* such medium. When light travels from a distant galaxy, or from any other cosmic object, it moves through the virtual vacuum of space. Sound waves, by contrast, cannot do this, despite what you have probably heard in almost every sci-fi movie ever made! If we were to remove all the air from a room, conversation would be impossible (even with suitable breathing apparatus to keep our test subjects alive!) because sound waves cannot exist without air or some other physical medium to support them. Communication by flashlight or radio, however, would be entirely feasible.

The ability of light to travel through empty space was once a great mystery. The idea that light, or any other kind of radiation, could move as a wave through nothing at all seemed to violate common sense, yet it is now a cornerstone of modern physics.

### Interactions Between Charged Particles


To understand more about the nature of light, consider for a moment an *electrically charged* particle, such as an **electron** or a **proton**. Like mass, electrical charge is a fundamental property of matter. Electrons and protons are elementary particles—“building blocks” of atoms and all matter—that carry the basic unit of charge. Electrons are said to carry a *negative* charge, whereas protons carry an equal and opposite *positive* charge.

Just as a massive object exerts a gravitational force on every other massive body, an electrically charged particle exerts an *electrical* force on every other charged particle in the universe.  (Sec. 2.7) The buildup of electrical charge (a net excess of positive over negative, or vice versa) is what causes “static cling” on your clothes when you take them out of a hot clothes dryer; it also causes the shock you sometimes feel when you touch a metal door frame on a particularly dry day.

Unlike the gravitational force, which is always attractive, electrical forces can be either attractive or repulsive. As illustrated in Figure 3.5(a), particles with *like* charges (i.e., both negative or both positive—for example, two electrons or two protons) repel one another. Particles with *unlike* charges (i.e., having opposite signs—an electron and a proton, say) attract.

How is the electrical force transmitted through space? Extending outward in all directions from any charged particle is an **electric field**, which determines the electrical force exerted by the particle on all other charged particles in the universe (Figure 3.5b). The strength of the electric field, like the strength of the gravitational field, decreases with increasing distance from the charge according to an inverse-square law. By means of the electric field, the particle’s presence is “felt” by all other charged particles, near and far.

Now, suppose our particle begins to vibrate, perhaps because it becomes heated or collides with some other particle.



**▲ FIGURE 3.5 Charged Particles** (a) Particles carrying like electrical charges repel one another, whereas particles with unlike charges attract. (b) A charged particle is surrounded by an electric field, which determines the particle’s influence on other charged particles. We represent the field by a series of field lines. (c) If a charged particle begins to vibrate, its electric field changes.



Its changing position causes its associated electric field to change, and this changing field in turn causes the electrical force exerted on other charges to vary (Figure 3.5c). If we measure the change in the force on these other charges, we learn about our original particle. Thus, *information about the particle's state of motion is transmitted through space via a changing electric field*. This disturbance in the particle's electric field travels through space as a wave.


## Electromagnetic Waves

The laws of physics tell us that a **magnetic field** must accompany every changing electric field. Magnetic fields govern the influence of *magnetized* objects on one another, much as electric fields govern interactions among charged particles. The fact that a compass needle always points to magnetic north is the result of the interaction between the magnetized needle and Earth's magnetic field (Figure 3.6). Magnetic fields also exert forces on *moving* electric charges (i.e., electric currents)—electric meters and motors rely on this basic fact. Conversely, moving charges *create* magnetic fields (electromagnets are a familiar example). In short, electric and magnetic fields are inextricably linked to one another: A change in either one necessarily creates the other.

Thus, as illustrated in Figure 3.7, the disturbance produced by the moving charge in Figure 3.5(c) actually consists of vibrating electric *and* magnetic fields, moving together through space. Furthermore, as shown in the diagram, these fields are always oriented *perpendicular* to one another and to the direction in which the wave is traveling. The fields do not exist as independent entities; rather, they are different aspects of a single physical phenomenon: **electromagnetism**. Together, they constitute an *electromagnetic wave* that carries energy and information from one part of the universe to another.


Now consider a real cosmic object—a star, say. When some of its charged contents move around, their electric fields change, and we can detect that change. The resulting electromagnetic ripples propagate (travel) outward as waves through space, requiring no material medium in which to move. Small charged particles, either in our eyes or in our experimental equipment, eventually respond to the electromagnetic field changes by vibrating in tune with the radiation that is received. This response is how we detect the radiation—how we see.

How *quickly* is one charge influenced by the change in the electromagnetic field when another charge begins to move? This is an important question because it is equivalent to asking how fast an electromagnetic wave travels. Does it propagate at some measurable speed, or is it instantaneous? Both theory and experiment tell us that all electromagnetic waves move at a



▲ **FIGURE 3.6 Magnetism** Earth's magnetic field interacts with a magnetic compass needle, causing the needle to become aligned with the field—that is, to point toward Earth's north (magnetic) pole. The north magnetic pole currently lies at latitude 80° N, longitude 107° W, some 1140 km from the geographic North Pole.

very specific speed—the **speed of light** (always denoted by the letter  $c$ ). Its exact value is 299,792.458 km/s in a vacuum (and somewhat less in material substances such as air or water). We will round this value off to  $c = 3.00 \times 10^5$  km/s, an extremely high speed. In the time needed to snap your fingers (about a tenth of a second), light can travel three-quarters of the way around our planet! If the currently known laws of physics are



▲ **FIGURE 3.7 Electromagnetic Wave** Electric and magnetic fields vibrate perpendicularly to each other. Together they form an electromagnetic wave that moves through space at the speed of light in the direction perpendicular to both the electric and the magnetic fields comprising it.


correct, then the speed of light is the fastest speed possible (see *More Precisely 22-1*).

The speed of light is very large, but it is still *finite*. That is, light does not travel instantaneously from place to place. This fact has some interesting consequences for our study of distant objects. It takes time—often lots of time—for light to travel through space. The light we see from the nearest large galaxy—the Andromeda Galaxy, shown in Figure 3.1—left that object about 2.5 million years ago, around the time our first human ancestors appeared on planet Earth. We can know nothing about this galaxy as it exists today. For all we know, it may no longer even exist! Only our descendants, 2.5 million years into the future, will know whether it exists now. So as we study objects in the cosmos, remember that the light we see left those objects long ago. We can never observe the universe as it is—only as it was.

## The Wave Theory of Radiation

The description presented in this chapter of light and other forms of radiation as electromagnetic waves traveling through space is known as the **wave theory of radiation**. It is a spectacularly successful scientific theory, full of explanatory and predictive power and deep insight into the complex interplay between light and matter—a cornerstone of modern physics.

Two centuries ago, however, the wave theory stood on much less solid scientific ground. Before about 1800, scientists were divided in their opinions about the nature of light. Some thought that light was a wave phenomenon (although at the time, electromagnetism was unknown), whereas others maintained that light was in reality a stream of particles that moved in straight lines. Given the experimental apparatus available at the time, neither camp could find conclusive evidence to disprove the other's theory. *Discovery 3-1* discusses some wave properties that are of particular importance to modern astronomers and describes how their detection in experiments using visible light early in the 19th century tilted the balance of scientific opinion in favor of the wave theory.

But that's not the end of the story. The wave theory, like all good scientific theories, can and must continually be tested by experiment and observation.  (Sec. 1.2) Around the turn of the 20th century, physicists made a series of discoveries about the behavior of radiation and matter on very small (atomic) scales that simply could not be explained by the “classical” wave theory just described. Changes had to be made. As we will see in Chapter 4, the modern theory of radiation is actually a hybrid of the once-rival wave and particle views, combining key elements of each in a unified and—for now—undisputed whole.

### PROCESS OF SCIENCE Check

- ✓ Describe the scientific reasoning leading to the conclusion that light is an electromagnetic wave.

## 3.3 Electromagnetic Spectrum

Figure 3.8 plots the entire range of electromagnetic radiation, illustrating the relationships among the different types of electromagnetic radiation listed earlier. Notice that the only characteristic distinguishing one from another is wavelength, or frequency. To the low-frequency, long-wavelength side of visible light lie *radio* and *infrared* radiation. Radio frequencies include radar, microwave radiation, and the familiar AM, FM, and TV bands. We perceive infrared radiation as heat. At higher frequencies (shorter wavelengths) are the domains of *ultraviolet*, *X-ray*, and *gamma-ray* radiation. Ultraviolet radiation, lying just beyond the violet end of the visible spectrum, is responsible for suntans and sunburns. The shorter-wavelength X-rays are perhaps best known for their ability to penetrate human tissue and reveal the state of our insides without resorting to surgery. Gamma rays are the shortest-wavelength radiation. They are often associated with radioactivity and are invariably damaging to living cells they encounter.


### The Spectrum of Radiation

All these spectral regions, including the visible spectrum, collectively make up the **electromagnetic spectrum**. Remember that, despite their greatly differing wavelengths and the different roles they play in everyday life on Earth, all are basically the same phenomenon, and all move at the same speed—the speed of light,  $c$ .

Figure 3.8 is worth studying carefully, as it contains a great deal of information. Note that wave frequency (in hertz) increases from left to right, and wavelength (in meters) increases from right to left. Scientists often disagree on the “correct” way to display wavelengths and frequencies in diagrams of this type. In picturing wavelengths and frequencies, this book consistently adheres to the convention that frequency increases toward the *right*.

Notice also that the wavelength and frequency scales in Figure 3.8 do not increase by equal increments of 10. Instead, successive values marked on the horizontal axis differ by *factors of 10*—each is 10 times greater than its neighbor. This type of scale, called a *logarithmic* scale, is often used in science to condense a large range of some quantity into a manageable size. Had we used a linear scale for the wavelength range shown in the figure, it would have been many light-years long! Throughout the text, we will often find it convenient to use a logarithmic scale to compress a wide range of some quantity onto a single, easy-to-view plot.

Figure 3.8 shows that wavelengths extend from the size of mountains (radio radiation) to the size of an atomic nucleus (gamma-ray radiation). The box at the upper right emphasizes how small the visible portion of the electromagnetic spectrum is. Most objects in the universe emit large amounts of invisible radiation. Indeed, many of them emit only a tiny fraction of their total energy in the visible range.



**Narrated Interactive** **FIGURE 3.8 Electromagnetic Spectrum** The entire electromagnetic

spectrum, running from long-wavelength, low-frequency radio waves to short-wavelength, high-frequency gamma rays.

A wealth of extra knowledge can be gained by studying the invisible regions of the electromagnetic spectrum.

To remind you of this important fact and to identify the region of the electromagnetic spectrum in which a particular observation was made, we have attached the following spectrum icon—an idealized version of the wavelength scale in Figure 3.8—to every astronomical image presented in this text (Figure 3.8). Hence, we can tell at a glance from the highlighted “V” that, for example, Figure 3.1 (p. 60) is an image made with the use of visible light, whereas the first image in

Figure 3.11 (p. 71) was captured in the radio (“R”) part of the spectrum. Chapter 5 discusses in more detail how astronomers actually make such observations, using telescopes and sensitive detectors tailored to different electromagnetic waves.


## Atmospheric Opacity

Only a small fraction of the radiation produced by astronomical objects actually reaches Earth’s surface, because of the **opacity** of our planet’s atmosphere. Opacity is the extent

## DISCOVERY 3-1

## The Wave Nature of Radiation


Until the early 19th century, debate raged in scientific circles regarding the true nature of light. On the one hand, the particle, or *corpuscular*, theory, proposed by Isaac Newton, held that light consisted of tiny particles moving in straight lines. Different colors were presumed to correspond to different particles. On the other hand, the *wave* theory, championed by the 17th-century Dutch astronomer Christian Huygens, viewed light as a wave phenomenon, in which color was determined by frequency, or wavelength. During the first few decades of the 19th century, growing experimental evidence that light displayed two key wave properties—*diffraction* and *interference*—argued strongly in favor of the wave theory.



**Diffraction** is the deflection, or “bending,” of a wave as it passes a corner or moves through a narrow gap. We might expect that light passing through a sharp-edged hole in a barrier would produce a sharp shadow, especially if radiation were composed of rays or particles moving in perfectly straight lines. As depicted in the first figure, however, closer inspection reveals that the shadow actually has a “fuzzy” edge, as shown in the photograph at the right—the diffraction pattern produced by a small circular opening.

We are not normally aware of such effects in everyday life, because diffraction is generally very small for visible light. For any wave, the amount of diffraction is proportional to the ratio of the wavelength to the width of the gap. The longer the wavelength or the smaller the gap, the greater is the angle through which the wave is diffracted. Thus, visible light, with its extremely short wavelengths, shows perceptible diffraction only when passing through very narrow openings. (The effect is much more noticeable for sound waves. No one thinks twice about our ability to hear people, even when they are around a corner and out of our line of sight.)

**Interference** is the ability of two or more waves to reinforce or diminish each other. The second figure shows two sets of waves moving through the same region of space. The waves are positioned so that their crests and troughs exactly



coincide. In the upper frames, the waves have the same wavelength, but the green one has twice the amplitude in the opposite direction to the orange one. The net effect is that the two wave motions interfere with each other, resulting in the wave at the right. This phenomenon is known as *destructive interference*. When the waves reinforce each other instead, as in the lower frames, the effect is called *constructive interference*.

As with diffraction, interference between waves of visible light is not noticeable in everyday experience, but it is readily measured in the laboratory. The final photograph shows the characteristic *interference pattern* that results when two identical light sources are placed side by side. The light and dark bands are formed by constructive and destructive interference of the beams from the two sources. This classic experiment, first performed by English physicist Thomas Young around 1805, was instrumental in establishing the wave nature of radiation.



Both diffraction and interference are predicted by the wave theory of light. The particle theory did not predict them; in fact, it predicted that they should *not* occur. By the 1830s, experimenters had reported clear measurements of both phenomena, convincing most scientists that the wave theory was the proper description of electromagnetic radiation. It would be almost a century before the particle description of radiation would resurface, but in a radically different form, as we will see in Chapter 4.

to which radiation is blocked by the material through which it is passing—in this case, air. The more opaque an object is, the less radiation gets through it: Opacity is just the opposite of transparency. Earth’s atmospheric opacity is plotted along the wavelength and frequency scales at the bottom of Figure 3.8. The extent of shading is proportional to the opacity. Where the shading is greatest (such as at the X-ray or “far” infrared regions of the spectrum), no radiation can get in or out. Where there is no shading at all (in the optical and part of the radio domain), the atmosphere is almost completely transparent. In some parts of the spectrum (e.g., the microwave band and much of the infrared portion), Earth’s atmosphere is partly transparent, meaning that some, but not all, incoming radiation makes it to the surface.

The effect of atmospheric opacity is that there are only a few *spectral windows* at well-defined locations in the electromagnetic spectrum where Earth’s atmosphere is transparent. In much of the radio domain and in the visible portions of the spectrum, the opacity is low, and we can study the universe at those wavelengths from ground level. In parts of the infrared range, the atmosphere is partially transparent, so we can make certain infrared observations from the ground. Moving to the tops of mountains, above as much of the atmosphere as possible, improves observations. In the rest of the spectrum, however, the atmosphere is opaque: Ultraviolet, X-ray, and gamma-ray observations can be made only from above the atmosphere, from orbiting satellites.

What causes opacity to vary along the spectrum? Certain atmospheric gases absorb radiation very efficiently at some wavelengths. For example, water vapor ( $\text{H}_2\text{O}$ ) and oxygen ( $\text{O}_2$ ) absorb radio waves having wavelengths less than about a centimeter, whereas water vapor and carbon dioxide ( $\text{CO}_2$ ) are strong absorbers of infrared radiation. Ultraviolet, X-ray, and gamma-ray radiation are completely blocked by the *ozone* ( $\text{O}_3$ ) *layer* high in Earth’s atmosphere (see Section 7.2). A passing, but unpredictable, source of atmospheric opacity in the visible part of the spectrum is the blockage of light by atmospheric clouds.

In addition, the interaction between the Sun’s ultraviolet radiation and the upper atmosphere produces a thin, electrically conducting layer at an altitude of about 100 km. The *ionosphere*, as this layer is known, reflects long-wavelength radio waves (wavelengths greater than about 10 m) as well as a mirror reflects visible light. In this way, extraterrestrial waves are kept out, and terrestrial waves—such as those produced by AM radio stations—are kept in. (That’s why it is possible to transmit some radio frequencies beyond the horizon—the broadcast waves bounce off the ionosphere.)

### CONCEPT Check

✓ In what sense are radio waves, visible light, and X-rays one and the same phenomenon?

## 3.4 Thermal Radiation

All macroscopic objects—fires, ice cubes, people, stars—emit radiation at all times, regardless of their size, shape, or chemical composition. They radiate mainly because the microscopic charged particles they are made up of are in constantly varying random motion, and whenever charges interact (“collide”) and change their state of motion, electromagnetic radiation is emitted. The **temperature** of an object is a direct measure of the amount of microscopic motion within it (see *More Precisely 3-1*). The hotter the object—that is, the higher its temperature—the faster its component particles move, the more violent are their collisions, and the more energy they radiate.

### The Blackbody Spectrum

*Intensity* is a term often used to specify the amount or strength of radiation at any point in space. Like frequency and wavelength, intensity is a basic property of radiation. No natural object emits all its radiation at just one frequency. Instead, because particles collide at many different speeds—some gently, others more violently—the energy is generally spread out over a range of frequencies. By studying how the intensity of this radiation is distributed across the electromagnetic spectrum, we can learn much about the object’s properties.

Figure 3.9 sketches the distribution of radiation emitted by an object. The curve peaks at a single, well-defined frequency and falls off to lesser values above and below that frequency. Note that the curve is not shaped like a symmetrical bell that declines evenly on either side of the peak. Instead, the intensity falls off more slowly from the peak to lower frequencies than it does on the high-frequency side. This overall shape is characteristic of the thermal radiation emitted by *any* object, regardless of its size, shape, composition, or temperature.

The curve drawn in Figure 3.9(a) is the radiation-distribution curve for a mathematical idealization known as a *blackbody*—an object that absorbs all radiation falling on it. In a steady state, a blackbody must reemit the same amount of energy it absorbs. The **blackbody curve** shown in the figure describes the distribution of that reemitted radiation. (The curve is also known as the *Planck curve*, after Max Planck, the German physicist whose mathematical analysis of such thermal emission in 1900 played a key role in the development of modern physics.)

No real object absorbs and radiates as a perfect blackbody. For example, the Sun’s actual curve of emission is shown in Figure 3.9(b). However, in many cases, the blackbody curve is a good approximation to reality, and the properties of blackbodies provide important insights into the behavior of real objects.


## MORE PRECISELY 3-1

### The Kelvin Temperature Scale

The atoms and molecules that make up any piece of matter are in constant random motion. This motion represents a form of energy known as *thermal energy*, or, more commonly, *heat*. The quantity we call *temperature* is a direct measure of an object's internal motion: The higher the object's temperature, the faster, on average, is the random motion of its constituent particles. Note that the two concepts, though obviously related, are different. The temperature of a piece of matter specifies the *average* thermal energy of the particles it contains.

Our familiar Fahrenheit temperature scale, like the archaic English system in which length is measured in feet and weight in pounds, is of somewhat dubious value. In fact, the “degree Fahrenheit” is now a peculiarity of American society. Most of the world uses the Celsius scale of temperature measurement (also called the centigrade scale). In the Celsius system, water freezes at 0 degrees (0°C) and boils at 100 degrees (100°C), as illustrated in the accompanying figure.

There are, of course, temperatures below the freezing point of water. In principle, temperatures can reach as low as  $-273.15^{\circ}\text{C}$  (although we know of no matter anywhere in the universe that is actually that cold). Known as *absolute zero*, this is the temperature at which, theoretically, all thermal atomic and molecular motion ceases. Since no object can have a temperature below that value, scientists find it convenient to use a temperature scale that takes absolute zero as its starting point. This scale is called the *Kelvin scale*, in honor of the 19th-century British physicist Lord Kelvin. Since it starts at absolute zero, the Kelvin scale differs from




the Celsius scale by  $273.15^{\circ}$ . In this book, we round off the decimal places and simply use

$$\text{kelvins} = \text{degrees Celsius} + 273.$$

Thus,

- all thermal motion ceases at 0 kelvins (0 K).
- water freezes at 273 kelvins (273 K).
- water boils at 373 kelvins (373 K).

Note that the unit is *kelvins*, or *K*, but not *degrees kelvin* or  $^{\circ}\text{K}$ . (Occasionally, the term *degrees absolute* is used instead.)




**▲ FIGURE 3.9 Blackbody Curves, Ideal vs. Reality** The blackbody, or Planck, curve represents the spread of the intensity of radiation emitted all across the electromagnetic spectrum. Note the contrast between the clean, “textbook” case (a) with a real graph (dashed) of the Sun’s emission (b). Absorption in the atmospheres of the Sun and Earth causes the difference.

## The Radiation Laws

The blackbody curve shifts toward higher frequencies (shorter wavelengths) and greater intensities as an object's temperature increases. Even so, the *shape* of the curve remains the same. This shifting of radiation's peak frequency with temperature is familiar to us all: Very hot glowing objects, such as toaster filaments or stars, emit visible light. Cooler objects, such as warm rocks, household radiators, or people, produce invisible radiation—warm to the touch, but not glowing hot to the eye. These latter objects emit most of their radiation in the lower frequency infrared part of the electromagnetic spectrum (Figure 3.8).

Imagine a piece of metal placed in a hot furnace. At first, the metal becomes warm, although its visual appearance doesn't change. As it heats up, the metal begins to glow dull red, then orange, brilliant yellow, and finally white. How do we explain this phenomenon? As illustrated in Figure 3.10, when the metal is at room temperature (300 K—see *More Precisely 3-1* for a discussion of the Kelvin temperature scale), it emits only invisible infrared radiation. As the metal becomes hotter, the peak of its blackbody curve shifts toward higher frequencies. At 1000 K, for instance, most of the emitted radiation is still infrared, but now there is also a small amount of visible (dull red) radiation being emitted. (Note that the high-frequency portion of the 1000 K curve just overlaps the visible region of the graph.)



▲ **FIGURE 3.10 Multiple Blackbody Curves** As an object is heated, the radiation it emits peaks at higher and higher frequencies. Shown here are curves corresponding to temperatures of 300 K (room temperature), 1000 K (beginning to glow dull red), 4000 K (red hot), and 7000 K (white hot).

As the temperature continues to rise, the peak of the metal's blackbody curve moves through the visible spectrum, from red (the 4000 K curve) through yellow. Eventually, the metal becomes white hot because, when its blackbody curve peaks in the blue or violet part of the spectrum (the 7000 K curve), the low-frequency tail of the curve extends through the entire visible spectrum (to the left in the figure), meaning that substantial amounts of green, yellow, orange, and red light are also emitted. Together, all these colors combine to produce white.

From studies of the precise form of the blackbody curve, we obtain a very simple connection between the wavelength at which most radiation is emitted and the absolute temperature (i.e., the temperature measured in kelvins) of the emitting object:

$$\text{wavelength of peak emission} \propto \frac{1}{\text{temperature}}$$

(Recall that the symbol “ $\propto$ ” here just means “is proportional to.”) This relationship is called **Wien's law**, after Wilhelm Wien, the German scientist who formulated it in 1897.


Simply put, Wien's law tells us that the hotter the object, the bluer is its radiation. For example, an object with a temperature of 6000 K emits most of its energy in the visible part of the spectrum, with a peak wavelength of 480 nm. At 600 K, the object's emission would peak at a wavelength of 4800 nm, well into the infrared portion of the spectrum. At a temperature of 60,000 K, the peak would move all the way through the visible spectrum to a wavelength of 48 nm, in the ultraviolet range (see Figure 3.11).

It is also a matter of everyday experience that, as the temperature of an object increases, the *total* amount of energy it radiates (summed over all frequencies) increases rapidly. For example, the heat given off by an electric heater increases very sharply as it warms up and begins to emit visible light. Careful experimentation leads to the conclusion that the total amount of energy radiated per unit time is actually proportional to the fourth power of the object's temperature:

$$\text{total energy emission} \propto \text{temperature}^4.$$

This relation is called **Stefan's law**, after the 19th-century Austrian physicist Josef Stefan. From the form of Stefan's law, we can see that the energy emitted by a body rises dramatically as its temperature increases. Doubling the temperature causes the total energy radiated to increase by a factor of  $2^4 = 16$ ; tripling the temperature increases the emission by  $3^4 = 81$ , and so on.

The radiation laws are presented in more detail in *More Precisely 3-2*.



**Interactive FIGURE 3.11**  
**Astronomical Thermometer**

**MA** Comparison of blackbody curves of four cosmic objects.  
 (a) A cool, dark galactic gas cloud called Barnard 68. At a temperature of 60 K, it emits mostly radio radiation, shown here as overlaid contours. (b) A dim, young star (shown white in the inset photograph) called Herbig-Haro 46. The star's atmosphere, at 600 K, radiates mainly in the infrared. (c) The Sun's surface, at approximately 6000 K, is brightest in the visible region of the electromagnetic spectrum. (d) Some very hot, bright stars in a cluster called Messier 2, as observed by an orbiting space telescope above Earth's atmosphere. At a temperature of 60,000 K, these stars radiate strongly in the ultraviolet. (ESO; AURA; SST; GALEX)



## MORE PRECISELY 3-2

### More About the Radiation Laws

As mentioned in Section 3.4, *Wien's law* relates the temperature  $T$  of an object to the wavelength  $\lambda_{\max}$  at which the object emits the most radiation. (The Greek letter  $\lambda$ —lambda—is conventionally used to denote wavelength.) Mathematically, if we measure  $T$  in kelvins and  $\lambda_{\max}$  in millimeters, we can determine the constant of proportionality in the relation presented in the text, to find that

$$\lambda_{\max} = \frac{2.9 \text{ mm}}{T}.$$

We could also convert Wien's law into an equivalent statement about frequency  $f$ , using the relation  $f = c/\lambda$  (see Section 3.1), where  $c$  is the speed of light, but the law is most commonly stated in terms of wavelength and is probably easier to remember that way.

**EXAMPLE 1** For a blackbody with the same temperature  $T$  as the surface of the Sun ( $\approx 6000$  K), the wavelength of maximum intensity is  $\lambda_{\max} = (2.9/6000)$  mm, or 480 nm, corresponding to the yellow-green part of the visible spectrum. A cooler star with a temperature of  $T = 3000$  K has a peak wavelength of  $\lambda_{\max} = (2.9/3000)$  mm  $\approx 970$  nm, just beyond the red end of the visible spectrum, in the near infrared. The blackbody curve of a hotter star with a temperature of 12,000 K peaks at 242 nm, in the near ultraviolet, and so on.

In fact, this application—looking at the spectrum and determining where it peaks—is an important way of estimating the temperature of planets, stars, and other objects across the universe. We will use it extensively throughout this text.

We can also give *Stefan's law* a more precise mathematical formulation. With  $T$  measured in kelvins, the total amount of energy emitted per square meter of the body's surface per second (a quantity known as the *energy flux*  $F$ ) is given by

$$F = \sigma T^4.$$

Energy per  
unit area
Constant
Temperature  
to the fourth  
power

This equation is usually referred to as the *Stefan-Boltzmann* equation. Stefan's student, Ludwig Boltzmann, was an Austrian physicist who played a central role in the development of the laws of thermodynamics during the late 19th and early 20th centuries. The constant  $\sigma$  (the Greek letter sigma) is known as the Stefan-Boltzmann constant.

The SI unit of energy is the *joule* (J). Probably more familiar is the closely related unit called the *watt* (W), which measures power—the *rate* at which energy is emitted or expended by an object. One watt is the emission of 1 J per second. For example, a 100-W lightbulb emits energy (mostly in the form of infrared and visible light) at a rate of 100 J/s. In SI units, the Stefan-Boltzmann constant has the value  $\sigma = 5.67 \times 10^{-8} \text{ W/m}^2 \cdot \text{K}^4$ .

**EXAMPLE 2** Notice just how rapidly the energy flux increases with increasing temperature. A piece of metal in a furnace, when at a temperature of  $T = 3000$  K, radiates energy at a rate of  $\sigma T^4 \times (1 \text{ cm})^2 = 5.67 \times 10^{-8} \text{ W/m}^2 \cdot \text{K}^4 \times (3000 \text{ K})^4 \times (0.01 \text{ m})^2 = 460 \text{ W}$  for every square centimeter of its surface area. Doubling this temperature to 6000 K (so that the metal becomes yellow hot, by Wien's law), the surface temperature of the Sun, increases the energy emitted by a factor of 16 (four “doublings”) to 7.3 kilowatts (7300 W) per square centimeter.

Finally, note that Stefan's law relates to energy emitted *per unit area*. The flame of a blowtorch is considerably hotter than a bonfire, but the bonfire emits far more energy *in total* because it is much larger. Thus, in computing the total energy emitted from a hot object, both the object's temperature *and* its surface area must be taken into account. This fact is of great importance in determining the “energy budget” of planets and stars, as we will see in later chapters.

## Astronomical Applications

No known natural terrestrial objects reach temperatures high enough to emit very high frequency radiation. Only human-made thermonuclear explosions are hot enough for their spectra to peak in the X-ray or gamma-ray range. (Most human inventions that produce short-wavelength, high-frequency radiation, such as X-ray machines, are designed to emit only a specific range of wavelengths and do not operate at high temperatures. They are said to produce a *nonthermal* spectrum of radiation.) Many extraterrestrial

objects, however, do emit large amounts of ultraviolet, X-ray, and even gamma-ray radiation.

Astronomers often use blackbody curves as thermometers to determine the temperatures of distant objects. For example, an examination of the solar spectrum indicates the temperature of the Sun's surface. Observations of the radiation from the Sun at many frequencies yield a curve shaped somewhat like that shown in Figure 3.9. The Sun's curve peaks in the visible part of the electromagnetic spectrum; the Sun also emits a lot of infrared and a little ultraviolet radiation. Using Wien's law, we find that the temperature of the

Sun's surface is approximately 6000 K. (A more precise measurement, applying Wien's law to the blackbody curve that best fits the solar spectrum, yields a temperature of 5800 K.)

Other cosmic objects have surfaces very much cooler or hotter than the Sun's, emitting most of their radiation in invisible parts of the spectrum. For example, the relatively cool surface of a very young star may measure 600 K and emit mostly infrared radiation. Cooler still is the interstellar gas cloud from which the star formed; at a temperature of 60 K, such a cloud emits mainly long-wavelength radiation in the radio and infrared parts of the spectrum. The brightest stars, by contrast, have surface temperatures as high as 60,000 K and hence emit mostly ultraviolet radiation (see Figure 3.11).


### CONCEPT Check

- ✓ Describe, in terms of the radiation laws, how and why the appearance of an incandescent lightbulb changes as you turn a dimmer switch to increase its brightness from "off" to "maximum."

## 3.5 The Doppler Effect

Imagine a rocket ship launched from Earth with enough fuel to allow it to accelerate to speeds approaching that of light. As the ship's speed increased, a remarkable thing would happen (Figure 3.12). Passengers would notice that the light from the star system toward which they were traveling seemed to be getting *bluer*. In fact, *all* stars in front of the ship would appear bluer than normal, and the greater the ship's speed, the greater the color change would be. Furthermore, stars behind the vessel would seem *redder* than normal, but stars to either side would be unchanged in appearance. As the spacecraft slowed down and came to rest relative to Earth, all stars would resume their usual appearance.

The travelers would have to conclude that the stars had changed their colors, not because of any real change in their physical properties, but because of the spacecraft's own *motion*. This motion-induced change in the observed frequency of a wave is known as the **Doppler effect**, in honor of Christian Doppler, the 19th-century Austrian physicist who first explained it in 1842. This phenomenon is not restricted to electromagnetic radiation and fast-moving spacecraft. Waiting at a railroad crossing for an express train to pass, most of us have had the experience of hearing the pitch of a train whistle change from high shrill (high frequency, short wavelength) to low blare (low frequency, long wavelength) as the train approaches and then recedes. The explanation is basically the same. Applied to cosmic sources of electromagnetic radiation, the Doppler effect has become one of the most important measurement techniques in all of modern astronomy. Here's how it works.




▲ **FIGURE 3.12 High-Speed Observers** Observers in a fast-moving spacecraft see the stars ahead of them bluer than normal, while those behind are reddened. The stars have not actually changed their properties—the color changes result from the observers' motion relative to the stars.

Imagine a wave moving from the place where it is created toward an observer who is not moving with respect to the source of the wave, as shown in Figure 3.13(a). By noting the distances between successive crests, the observer can determine the wavelength of the emitted wave. Now suppose that not just the wave, but the *source* of the wave, also is moving. As illustrated in Figure 3.13(b), because the source moves between the times of emission of one crest and the next, successive crests in the direction of motion of the source will be seen to be *closer together* than normal, whereas crests behind the source will be more widely spaced. An observer in front of the source will therefore measure a *shorter* wavelength than normal, whereas one behind will see a *longer* wavelength. The numbers indicate (a) successive crests emitted by the source and (b) the location of the source at the instant each crest was emitted.

The greater the relative speed between source and observer, the greater is the observed shift. If the other velocities involved are not too large compared with the wave speed—less than a few percent, say—we can write down a particularly simple formula for what the observer sees. In terms of the net velocity of *recession* between source and observer, the apparent wavelength and frequency (measured by the observer) are related to the true quantities (emitted by the source) as follows:

$$\frac{\text{apparent wavelength}}{\text{true wavelength}} = \frac{\text{true frequency}}{\text{apparent frequency}}$$

$$= 1 + \frac{\text{recession velocity}}{\text{wave speed}}$$



**Interactive FIGURE 3.13**  
**Doppler Effect** (a) Wave motion from a source toward an observer at rest with respect to the source. As seen by the observer, the source is not moving, so the wave crests are just concentric spheres (shown here as circles). (b) Waves from a moving source tend to “pile up” in the direction of motion and be “stretched out” on the other side. As a result, an observer situated in front of the source measures a shorter-than-normal wavelength—a *blueshift*—while an observer behind the source sees a *redshift*. In this diagram, the source is shown in motion. However, the same general statements hold whenever there is any relative motion between source and observer, allowing astronomers to probe the motions of distant objects.

The recession velocity measures the rate at which the distance between the source and the observer is changing. A positive recession velocity means that the two are moving apart; a negative velocity means that they are approaching.

The wave speed is the speed of light,  $c$ , in the case of electromagnetic radiation. For most of this text, the assumption that the recession velocity is small compared to the speed of light will be a good one. Only when we discuss the properties of black holes (Chapter 22) and the structure of the universe on the largest scales (Chapters 25 and 26) will we have to reconsider this formula.

Note that in Figure 3.13 the *source* is shown in motion (as in our train analogy), whereas in our earlier spaceship example (Figure 3.12) the *observers* were in motion. For electromagnetic radiation, the result is the same in either case—only the *relative* motion between source and observer matters. Note also that only motion along the line joining source and observer—known as *radial* motion—appears in the foregoing equation. Motion that is *transverse* (perpendicular) to the line of sight has no significant effect.\* Notice, incidentally, that the Doppler effect depends only on the *relative motion* between source and observer; it does not depend on the *distance* between them in any way.

A wave measured by an observer situated in front of a moving source is said to be *blueshifted*, because blue light has a shorter wavelength than red light. Similarly, an observer situated behind the source will measure a longer-than-normal wavelength—the radiation is said to be *redshifted*. This terminology is used even for invisible radiation, for which “red” and “blue” have no meaning. Any shift toward shorter wavelengths is called a blueshift, and any shift toward longer wavelengths is called a redshift. For example, ultraviolet radiation might be blueshifted into the X-ray part of the spectrum or redshifted into the visible; infrared radiation could be redshifted into the microwave range, and so on. *More Precisely 3-3* describes how the Doppler effect is used in practice to measure velocities in astronomy.

\*In fact, Einstein’s theory of relativity (see Chapter 22) implies that when the transverse velocity is comparable to the speed of light, a change in wavelength, called the transverse Doppler shift, does occur. For most terrestrial and astronomical applications, however, this shift is negligibly small, and we will ignore it here.


## MORE PRECISELY 3-3

### Measuring Velocities with the Doppler Effect

Because the speed of light,  $c$ , is so large—300,000 km/s—the Doppler effect is extremely small for everyday terrestrial velocities. For example, consider a source receding from the observer at Earth’s orbital speed of 30 km/s, a velocity much greater than any encountered in day-to-day life. Using the formula in the text, we find that the shift in wavelength of a beam of blue light would be just

$$\begin{aligned} \frac{\text{change in wavelength}}{\text{true wavelength}} &= \frac{\text{recession velocity}}{\text{wave speed}} \\ &= \frac{30 \text{ km/s}}{300,000 \text{ km/s}} = 0.01 \text{ percent.} \end{aligned}$$

That is, the wavelength would lengthen from 400 nm to 400.04 nm—a very small change indeed, and one that the human eye cannot distinguish. However, it *is* easily detectable with modern instruments.



In practice, it is hard to measure the Doppler shift of an entire blackbody curve, simply because it is spread over many wavelengths, making small shifts hard to determine with any accuracy. However, if the radiation is more narrowly defined and takes up just a narrow “sliver” of the spectrum, then precise measurements of Doppler effect *can* be made. We will see in the next chapter that in many circumstances this is precisely what does happen, making the

Astronomers can use the Doppler effect to find the line-of-sight speed of any cosmic object simply by measuring the extent to which its light is redshifted or blueshifted. To see how, let’s use a simple example.

**EXAMPLE** Suppose that the beam of blue light just mentioned is observed to have a wavelength of 401 nm, instead of the 400 nm with which it was emitted. (Let’s defer until the next chapter the question of *how* an observer might know the wavelength of the emitted light.) Using the earlier equation, rewritten as

$$\begin{aligned} \frac{\text{recession velocity}}{\text{wave speed, } c} &= \frac{\text{change in wavelength}}{\text{true wavelength}} \\ &= \frac{1 \text{ nm}}{400 \text{ nm}} = 0.0025, \end{aligned}$$

the observer could calculate the source’s recession velocity to be 0.0025 times 300,000 km/s, or 750 km/s.

The basic reasoning is simple, but very powerful. The motions of nearby stars and distant galaxies—even the expansion of the universe itself—have all been measured in this way.

Motorists stopped for speeding on the highway have experienced another, much more down-to-earth application: As illustrated in the accompanying figure, police radar measures speed by means of the Doppler effect, as do the radar guns used to clock the velocity of a pitcher’s fastball or a tennis player’s serve. As shown in the illustration, the reflected radiation (blue crests) from the oncoming car is shifted to shorter wavelengths by an amount proportional to the car’s speed. Gotcha!

Doppler effect one of the observational astronomer’s most powerful tools.

#### CONCEPT Check

- ✓ Astronomers observe two stars orbiting one another. How might the Doppler effect be useful in determining the masses of the stars?

**The Big Question** The speed of light plays a central role throughout all of physics and astronomy. But a nagging question that often comes up is, Can anything travel faster than light? Contrary to popular belief, Einstein’s theory of relativity, which we will study later, does not prohibit objects from traveling any faster. And there is evidence that the early universe did in fact expand faster than light. Yet no one knows for sure if light velocity is the ultimate speed limit in the universe, so experiments will continue to test this very fundamental concept.

# Chapter Review

## SUMMARY

### 1 Electromagnetic radiation

(p. 60) travels through space

in the form of a **wave** (p. 60).

A wave is characterized by its

**period** (p. 61), the length


of time taken for one complete cycle; its **wavelength** (p. 61),

the distance between successive wave crests; and its **amplitude**

(p. 61), which measures the size of the disturbance associated

with the wave. A wave's **frequency** (p. 61) is the number of

wave crests that pass a given point in one second.



### 2 Any electrically charged object

is surrounded by an **electric field**

(p. 63) that determines the force the

object exerts on other charged objects.

When a charged particle moves, information

about its motion is transmitted

via the particle's changing **electric and**


**magnetic fields** (pp. 63, 64). The

information travels at the **speed of light**

(p. 64) as an electromagnetic wave. **Diffraction** (p. 67) and

**interference** (p. 67) are properties of radiation that mark it as

a wave phenomenon.



### 3 The color of visible light is simply

a measure of its wavelength—red light

has a longer wavelength than blue

light. The entire **electromagnetic**


**spectrum** (p. 65) consists of (in

order of increasing frequency) **radio**

**waves, infrared radiation, visible**

**light, ultraviolet radiation, X-rays,**

and **gamma rays** (p. 60). Only radio



waves, some infrared wavelengths, and visible light can penetrate the atmosphere and reach the ground from space.

### 4 The **temperature** (p. 68) of

an object is a measure of the speed with


which its constituent particles

move. The intensity of radiation

emitted by an object has a character-

istic distribution, called a **blackbody**

**curve** (p. 68), which depends only on the object's temperature.



### 5 Wien's law (p. 70) tells us that

the wavelength at which the object

radiates most of its energy is inversely

proportional to its temperature.

Measuring that peak wavelength tells


us the object's temperature. **Stefan's**

**law** (p. 70) states that the total

amount of energy radiated is pro-

portional to the fourth power of the

temperature.



### 6 Our perception of the

wavelength of a beam of light

can be altered by the source's

velocity relative to us. This

motion-induced change in the

observed frequency of a wave

is called the **Doppler effect**

(p. 73). Any net motion of the

source away from the observer

causes a redshift—a shift to lower

frequencies—in the received


beam. Motion toward the observer

causes a blueshift. The extent

of the shift is directly proportional

to the source's radial velocity

relative to the observer.



**MasteringAstronomy**® For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information.

**LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

1. What is a wave?
2. **LO1** What is the relationship between wavelength, wave frequency, and wave velocity?
3. What is diffraction, and how does it relate to wave motion?
4. **POS** What's so special about  $c$ ?
5. Name the colors that make up white light. What is it about these colors that causes us to perceive them differently?
6. What effect does a positive charge have on a nearby negatively charged particle?
7. **LO2** Describe the way in which light leaves a star, travels through the vacuum of space, and finally is seen by someone on Earth.
8. Why is light referred to as an electromagnetic wave?
9. What do radio waves, infrared radiation, visible light, ultraviolet radiation, X-rays, and gamma rays have in common? How do they differ?
10. **LO3** In what parts of the electromagnetic spectrum is the atmosphere transparent enough for ground-based astronomy?

11. **LO4** What is a blackbody? What are the main characteristics of the radiation it emits?
12. **POS** What does Wien's law reveal about stars in the sky?
13. **LOS** In terms of its blackbody curve, describe what happens as a red-hot glowing coal cools.
14. **LOG** How do astronomers use the Doppler effect to determine the velocities of astronomical objects?
15. **POS** If Earth were completely blanketed with clouds and we couldn't see the sky, could we learn about the realm beyond the clouds? What forms of radiation might be received?

## Conceptual Self-Test: Multiple Choice

1. Compared with ultraviolet radiation, infrared radiation has a greater (a) wavelength; (b) amplitude; (c) frequency; (d) energy.
2. Compared with red light, blue wavelengths of visible light travel (a) faster; (b) slower; (c) at the same speed.
3. An electron that collides with an atom will (a) cease to have an electric field; (b) produce an electromagnetic wave; (c) change its electric charge; (d) become magnetized.
4. **VIS** According to Figure 3.8 ("Electromagnetic Spectrum"), the wavelength of green light is about the size of (a) an atom; (b) a bacterium; (c) a fingernail; (d) a skyscraper.
5. An X-ray telescope located in Antarctica would not work well because of (a) the extreme cold; (b) the ozone hole; (c) continuous daylight; (d) Earth's atmosphere.
6. **VIS** In Figure 3.11 ("Multiple Blackbody Curves"), an object at 1000 K emits mostly (a) infrared light; (b) red light; (c) multiple green light; (d) blue light.
7. According to Wien's law, the hottest stars also have (a) the longest peak wavelength; (b) the shortest peak wavelength; (c) maximum emission in the infrared region of the spectrum; (d) the largest diameters.
8. Stefan's law says that if the Sun's temperature were to double, its energy emission would (a) become half its present value; (b) double; (c) increase four times; (d) increase 16 times.
9. A star much cooler than the Sun would appear (a) red; (b) blue; (c) smaller; (d) larger.
10. The blackbody curve of a star moving toward Earth would have its peak shifted (a) to a higher intensity; (b) toward higher energies; (c) toward longer wavelengths; (d) to a lower intensity.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

1. ● A sound wave moving through water has a frequency of 256 Hz and a wavelength of 5.77 m. What is the speed of the wave?
2. ● What is the wavelength of a 100-MHz ("FM 100") radio signal?
3. ●● Estimate the total amount of energy you radiate to your surroundings.
4. ● Normal human body temperature is about 37°C. What is this temperature in kelvins? What is the peak wavelength emitted by a person with this temperature? In what part of the spectrum does this lie?
5. ● The Sun has a temperature of 5800 K, and its blackbody emission peaks at a wavelength of approximately 500 nm. At what wavelength does a protostar with a temperature of 1000 K radiate most strongly?
6. ● Two otherwise identical bodies have temperatures of 300 K and 1500 K, respectively. Which one radiates more energy, and by what factor does its emission exceed the emission of the other?
7. ● At what velocity and in what direction would a spacecraft have to be moving for a radio station on Earth transmitting at 100 MHz to be picked up by a radio tuned to 99.9 MHz?
8. ● Radiation from the nearby star Alpha Centauri is observed to be reduced in wavelength (after correction for Earth's orbital motion) by a factor of 0.999933. What is the recession velocity of Alpha Centauri relative to the Sun?

## Activities


### Collaborative

1. Stand near (but not too near!) a train track or busy highway and wait for a train or traffic to pass by. Can you notice the Doppler effect in the pitch of the engine noise or whistle blowing? How does the sound frequency depend on the train's (a) speed and (b) motion toward or away from you? Divide your group into two. One subgroup should time the train's motion and hence calculate approximately its speed. The other (consisting of the more musically inclined!) should

estimate the perceived frequency change of the whistle when the train is moving first toward you and then away from you.

### Individual

1. Locate the constellation Orion. Its two brightest stars are Betelgeuse and Rigel. Which is hotter? How can you tell? Which of the other stars scattered across the night sky are hot, and which are cool?



# 4

# Spectroscopy

## THE INNER WORKINGS OF ATOMS

The wave description of radiation allowed 19th-century astronomers to begin deciphering information reaching Earth from the cosmos in the form of visible and invisible light. However, early in the 20th century, it became clear that the wave theory of electromagnetic phenomena was incomplete—some aspects of light simply could not be explained in purely wave terms.

When radiation interacts with matter on atomic scales, it does so not as a continuous wave, but in a jerky, discontinuous way—in fact, as a particle. With this discovery, scientists quickly realized that atoms, too, must behave in a discontinuous way, and the stage was set for a scientific revolution—*quantum mechanics*—that has affected virtually every area of modern life.

**The Big Picture** Spectroscopy is an observational method used by astronomers, along with telescopes and theoretical insight, to infer the nature of matter by the way it emits and absorbs radiation. This powerful technique can not only reveal the chemical composition of distant stars, but also provide much information about the origin, evolution, and destiny of stars throughout the universe. Spectroscopy is an indispensable foundation of modern astrophysics.

**LEFT:** The beautiful visible spectrum of the star Procyon is shown here from red to blue, interrupted by hundreds of dark lines caused by the absorption of light in the hot star's cooler atmosphere. The whole spectrum is normally 6 meters (20 feet) across, but in order to display it on a single page the full spectrum is cut into dozens of horizontal segments and stacked vertically. (NOAO/AURA)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Describe the characteristics of continuous, emission, and absorption spectra and the conditions under which each is produced.
- 2 Explain the relation between emission and absorption lines and say what we can learn from those lines.
- 3 Specify the basic components of the atom and describe our modern conception of its structure.
- 4 Outline the observations that led scientists to conclude that light has particle as well as wave properties.
- 5 Explain how electron transitions within atoms produce unique emission and absorption features in the spectra of those atoms.
- 6 Describe the general features of spectra produced by molecules.
- 7 List and explain the kinds of information that can be obtained by analyzing the spectra of astronomical objects.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.




## 4.1 Spectral Lines

In Chapter 3, we saw something of how astronomers can analyze electromagnetic radiation received from space to obtain information about distant objects. A vital step in this process is the formation of a *spectrum*—a splitting of the incoming radiation into its component wavelengths. But in reality, *no* cosmic object emits a perfect blackbody spectrum like those discussed earlier. ∞ (Sec 3.4) All spectra deviate from this idealized form—some by only a little, others by a lot. Far from invalidating our earlier studies, however, these deviations contain a wealth of detailed information about physical conditions within the source of the radiation. Because spectra are so important, let's examine how astronomers obtain and interpret them.

Radiation can be analyzed with an instrument known as a **spectroscope**. In its most basic form, this device consists of an opaque barrier with a slit in it (to define a beam of light), a prism (to split the beam into its component colors), and an eyepiece or screen (to allow the user to view the resulting spectrum). Figure 4.1 shows such an arrangement. The research instruments called *spectrographs*, or *spectrometers*, used by professional astronomers are rather more complex, consisting of a telescope (to capture the radiation), a dispersing device (to spread the radiation out into a spectrum), and a detector (to record the result). Despite their greater sophistication, their basic operation is conceptually similar to the simple spectroscope shown in the figure.

In many large instruments, the prism is replaced by a device called a *diffraction grating*, consisting of a sheet of transparent material with numerous closely spaced parallel lines ruled on it. The spacing between the lines is typically a few microns ( $10^{-6}$  m), comparable to the wavelength of visible light. The spaces act as many tiny openings, and light is diffracted as it passes through the grating (or is reflected from it, depending on the design of the device). ∞ (Discovery 3-1) Because different wavelengths of electromagnetic radiation are diffracted by different amounts on encountering the grating,



the effect is to split a beam of light into its component colors. You are probably more familiar with diffraction gratings than you think—the “rainbow” of colors seen in light reflected from a compact disk is the result of precisely this process.

### Emission Lines

The spectra we encountered in Chapter 3 are examples of **continuous spectra**. A lightbulb, for example, emits radiation of all wavelengths (mostly in the visible range), with an intensity distribution that is well described by the blackbody curve corresponding to the bulb's temperature. ∞ (Sec. 3.4) Viewed through a spectroscope, the spectrum of the light from the bulb would show the familiar rainbow of colors, from red to violet, without interruption, as presented in Figure 4.2(a).

Not all spectra are continuous, however. For instance, if we took a glass jar containing pure hydrogen gas and passed an electrical discharge through it (a little like a lightning bolt arcing through Earth's atmosphere), the gas would begin to glow—that is, it would emit radiation. If we were to examine that radiation with our spectroscope, we would find that its spectrum consists of only a few bright lines on an otherwise dark background, quite unlike the continuous spectrum described for the incandescent lightbulb. Figure 4.2(b) shows the experimental arrangement and its result schematically. (A more detailed rendering of the spectrum of hydrogen appears in the top panel of Figure 4.3.) Note that the light produced by the hydrogen in this experiment does *not* consist of all possible colors, but instead includes only a few narrow, well-defined **emission lines**—thin “slices” of the continuous spectrum. The black background represents all the wavelengths *not* emitted by hydrogen.

After some experimentation, we would also find that, although we could alter the *intensity* of the lines—for example, by changing the amount of hydrogen in the jar or the strength of the electrical discharge—we could not alter their *color* (in other words, their frequency or wavelength). The pattern of spectral emission lines shown is a property of the element hydrogen. Whenever we perform this experiment, the same characteristic colors result.

By the early 19th century, scientists had carried out similar experiments on many different gases. By vaporizing solids and liquids in a flame, they extended their inquiries to include materials that are not normally



◀ **FIGURE 4.1 Spectroscope** A simple spectroscope allows a narrow beam of light to pass through a thin slit and then into a prism where the light is split into its component colors. A lens then focuses the light into a sharp image that is either projected onto a screen, as shown here, or analyzed as it strikes a detector.

**Interactive** **FIGURE 4.2 Continuous and Emission Spectra**

When passed through a slit and split up by a prism, light from a source of continuous radiation (a) gives rise to the familiar rainbow of colors. By contrast, the light from excited hydrogen gas (b) consists of a series of distinct bright spectral lines called emission lines. (The focusing lenses have been omitted for clarity—see Section 5.1.)

**MA**

found in the gaseous state. Sometimes the pattern of lines was fairly simple, and sometimes it was complex, but it was always *unique* to that element. Even though the origin of the lines was not understood, researchers quickly realized that the lines provided a one-of-a-kind “fingerprint” of the substance under investigation. They could detect the presence of a particular atom or molecule (a group of atoms held together by chemical bonds—see Section 4.4) solely through the study of the light it emitted. Scientists have accumulated extensive catalogs of the specific wavelengths at which many different hot gases emit radiation. The particular pattern of light emitted by a gas of a given chemical composition is known as the **emission spectrum** of the gas. The emission spectra of some common substances are shown in Figure 4.3.



**▲ FIGURE 4.3 Elemental Emission** The emission spectra of some well-known elements. In accordance with the convention adopted throughout this text, frequency increases to the right. Note that wavelengths shorter than approximately 400 nm, shown here in shades of purple, are actually in the ultraviolet part of the spectrum and thus invisible to the human eye. (Wabash Instrument Corp.)

## Absorption Lines


When sunlight is split by a prism, at first glance it appears to produce a continuous spectrum. However, closer scrutiny with a spectroscope shows that the solar spectrum is interrupted vertically by a large number of narrow dark lines, as shown in Figure 4.4. We now know that many of these lines represent wavelengths of light that have been removed (absorbed) by gases present either in the outer layers of the Sun or in Earth's atmosphere. These gaps in the spectrum are called **absorption lines**.

The English astronomer William Wollaston first noticed the solar absorption lines in 1802. They were studied in greater detail about 10 years later by the German physicist Joseph von Fraunhofer, who measured and cataloged over 600 of them. They are now referred to


collectively as *Fraunhofer lines*. Although the Sun is by far the easiest star to study, and so has the most extensive set of observed absorption lines, similar lines are known to exist in the spectra of all stars.

At around the same time as the solar absorption lines were discovered, scientists found that such lines could also be produced in the laboratory by passing a beam of light from a source that produces a continuous spectrum through a cool gas, as shown in Figure 4.5. The scientists quickly observed an intriguing connection between emission and absorption lines: The absorption lines associated with a given gas occur at precisely the *same* wavelengths as the emission lines produced when the gas is heated.

As an example, consider the element sodium, whose emission spectrum appears in Figure 4.6. When heated to high temperatures, a sample of sodium vapor emits visible light strongly at just two wavelengths—589.6 nm and 589.0 nm—lying in the yellow part of the spectrum. When a continuous




◀ **FIGURE 4.4 Solar Spectrum** The Sun's visible spectrum shows hundreds of vertical dark absorption lines superimposed on a bright continuous spectrum. This high-resolution spectrum is displayed in a series of 48 horizontal strips stacked vertically; each strip covers a small portion of the entire spectrum from left to right. The scale extends from long wavelengths (red) at the upper left to short wavelengths (blue) at the lower right. See also the chapter-opening spectrum on page 78. (AURA)



### Interactive FIGURE 4.5 Absorption Spectrum

◀ **MA** cool gas is placed between a source of continuous radiation (such as a hot lightbulb) and a detector/screen, the resulting color spectrum is crossed by a series of dark absorption lines. These lines are formed when the intervening cool gas absorbs certain wavelengths (colors) from the original beam of light. The absorption lines appear at precisely the same wavelengths as the emission lines that would be produced if the gas were heated to high temperatures (see Figure 4.2). (b) An everyday analogy for any of these line spectra is a supermarket bar code that uniquely determines the type and cost of a product.



**Interactive FIGURE 4.6 Sodium Spectrum** (a) The characteristic emission lines of sodium appear as two bright lines in the yellow part of the spectrum. (b) The absorption spectrum of sodium shows two dark lines at exactly the same wavelengths as the bright lines in sodium's emission spectrum.

spectrum is passed through some relatively cool sodium vapor, two sharp, dark absorption lines appear at precisely the same wavelengths. The emission and absorption spectra of sodium are compared in Figure 4.6, clearly showing the relation between emission and absorption features.

## Kirchhoff's Laws

The analysis of the ways in which matter emits and absorbs radiation is called **spectroscopy**. One early spectroscopist, the German physicist Gustav Kirchhoff, summarized the observed relationships among the three types of spectra—continuous, emission line, and absorption line—in 1859. He formulated three spectroscopic rules, now known as **Kirchhoff's laws**, governing the formation of spectra:

1. A luminous solid or liquid, or a sufficiently dense gas, emits light of all wavelengths and so produces a *continuous spectrum* of radiation.
2. A low-density, hot gas emits light whose spectrum consists of a series of bright *emission lines* that are characteristic of the chemical composition of the gas.
3. A cool, thin gas absorbs certain wavelengths from a continuous spectrum, leaving dark *absorption lines* in their place, superimposed on the continuous spectrum. Once again, these lines are characteristic of the composition of the intervening gas—they occur at precisely the same wavelengths as the emission lines produced by that gas at higher temperatures.

Figure 4.7 illustrates Kirchhoff's laws and the relationship between absorption and emission lines. Viewed directly, the light source, a hot solid (the filament of the bulb), has a continuous (blackbody) spectrum. When the light source is viewed through a cloud of cool hydrogen gas, a series of dark absorption lines appear, superimposed on the spectrum at wavelengths characteristic of hydrogen. The lines appear because the light at those wavelengths is absorbed by the hydrogen. As we will see later in this chapter, the absorbed energy is subsequently reradiated into space—but in all directions, not just the original direction of the beam. Consequently, when the cloud is viewed from the side against an otherwise dark background, a series of

faint emission lines is seen. These lines contain the energy lost by the forward beam. If the gas was heated to incandescence, it would produce stronger emission lines at precisely the same wavelengths.

## Identifying Starlight

By the late 19th century, spectroscopists had developed a formidable arsenal of techniques for interpreting the radiation received from space. Once astronomers knew that spectral lines were indicators of chemical composition, they set about identifying the observed lines in the solar spectrum. Almost all the lines in light from extra-terrestrial sources could be attributed to known elements. For example, many of the Fraunhofer lines in sunlight are associated with the element iron, a fact first recognized by Kirchhoff and coworker Robert Bunsen (of Bunsen burner fame) in 1859. However, some unfamiliar lines also appeared in the solar spectrum. In 1868, astronomers realized that those lines must correspond to a previously unknown element. It was given the name helium, after the Greek word *helios*, meaning “Sun.” Not until 1895, almost three decades after its detection in sunlight, was helium discovered on Earth! (A laboratory spectrum of helium is included in Figure 4.3.)


Yet, for all the information that 19th-century astronomers could extract from observations of stellar spectra, they still lacked a theory explaining how the spectra themselves arose. Despite their sophisticated spectroscopic equipment, they knew scarcely any more about the physics of stars than did Galileo or Newton. To understand how spectroscopy can be used to extract detailed information about astronomical objects from the light they emit, we must delve more deeply into the processes that produce line spectra.

### CONCEPT Check

- ✓ What are absorption and emission lines, and what do they tell us about the composition of the gas producing them?

## ► FIGURE 4.7 Kirchhoff's

**Laws** A source of continuous radiation, here represented by a lightbulb, is used to illustrate Kirchhoff's laws of spectroscopy. (a) The unimpeded beam shows the familiar continuous spectrum of colors. (b) When the source is viewed through a cloud of hydrogen gas, a series of dark hydrogen absorption lines appears in the continuous spectrum. These lines are formed when the gas absorbs some of the bulb's radiation and reemits it in random directions. Because most of the reemitted radiation does not go through the slit, the effect is to remove the absorbed radiation from the light that reaches the screen at the left. (c) When the gas is viewed from the side, a fainter hydrogen emission spectrum is seen, consisting of reemitted radiation. The absorption lines in (b) and the emission lines in (c) have the same wavelengths.



## 4.2 Atoms and Radiation

By the start of the 20th century, physicists had accumulated substantial evidence that light sometimes behaves in a manner that cannot be explained by the wave theory. As we have just seen, the production of absorption and emission lines involves only certain very specific frequencies or wavelengths of light. This would not be expected if light behaved like a continuous wave and matter always obeyed the laws of Newtonian mechanics. Other experiments conducted around the same time strengthened the conclusion that the notion of radiation as a wave was incomplete. It became clear that when light interacts with matter on very small scales, it does so not in a continuous way, but in a discontinuous, “stepwise” manner. The challenge was to find an explanation for this unexpected behavior. The eventual solution revolutionized our view of nature and now forms the foundation for all of physics and astronomy—indeed, for virtually all modern science.

### Atomic Structure

To explain the formation of emission and absorption lines, we must understand not just the nature of light, but also the structure of **atoms**—the microscopic building blocks from which all matter is constructed. Let's start with the simplest atom of all: hydrogen. A hydrogen atom consists of an electron with a negative electrical charge orbiting a proton carrying a positive charge. The proton forms the central **nucleus** (plural: nuclei) of the atom. The hydrogen atom as a whole is electrically neutral. The equal and opposite charges of the proton and the orbiting electron produce an electrical attraction that binds them together within the atom.

How does this picture of the hydrogen atom relate to the characteristic emission and absorption lines associated with hydrogen gas? If an atom absorbs some energy in the form

of radiation, that energy must cause some internal change. Similarly, if the atom emits energy, that energy must come from somewhere within the atom. It is reasonable (and correct) to suppose that the energy absorbed or emitted by the atom is associated with changes in the motion of the orbiting electron.

The first theory of the atom to provide an explanation of hydrogen's observed spectral lines was set forth by the Danish physicist Niels Bohr in 1912. Now known simply as the *Bohr model* of the atom, its essential features are as follows:

1. There is a state of lowest energy—the **ground state**—which represents the “normal” condition of the electron as it orbits the nucleus.
2. There is a maximum energy that the electron can have and still be part of the atom. Once the electron acquires more than that maximum energy, it is no longer bound to the nucleus, and the atom is said to be **ionized**; an atom missing one or more of its electrons is called an *ion*.
3. Most important (and also least intuitive), between those two energy levels, the electron can exist only in certain sharply defined energy states, often referred to as **orbitals**.


This description of the atom contrasts sharply with the predictions of Newtonian mechanics, which would permit orbits with *any* energy, not just at certain specific values. ∞ (Sec. 2.8) In the atomic realm, such discontinuous behavior is the norm. In the jargon of the field, the orbital

energies are said to be **quantized**. The rules of **quantum mechanics**, the branch of physics governing the behavior of atoms and subatomic particles, are far removed from everyday experience.


In Bohr's original model, each electron orbital was pictured as having a specific radius, much like a planetary orbit in the solar system, as shown in Figure 4.8. However, the modern view is not so simple. Although each orbital *does* have a precise energy, the orbits are not sharply defined, as indicated in the figure. Rather, the electron is now envisioned as being smeared out in an "electron cloud" surrounding the nucleus, as illustrated in Figure 4.9. We cannot tell "where" the electron is—we can only speak of the *probability* of finding it in a certain location within the cloud. It is common to speak of the average distance from the cloud to the nucleus as the "radius" of the electron's orbit. When a hydrogen atom is in its ground state, the radius of the orbit is about 0.05 nm (0.5 Å). As the orbital energy increases, the radius increases, too.

For the sake of clarity in the diagrams that follow, we will represent electron orbitals in this chapter as solid lines. (See *More Precisely 4-1* for a more detailed rendition of hydrogen's energy levels.) However, you should always bear in mind that Figure 4.9 is a more accurate depiction of reality.

Atoms do not always remain in their ground state. An atom is said to be in an **excited state** when an electron occupies an orbital at a greater-than-normal distance from its parent nucleus. An atom in such an excited state has a greater-than-normal amount of energy. The excited state with the lowest energy (that is, the state closest in energy to the ground state) is called the *first excited state*, that with the second-lowest energy is the *second excited state*, and so on. An atom can become excited in one of two ways: by absorbing some energy from a source of electromagnetic radiation or by colliding with some other particle—another atom, for example. However, the electron cannot stay in a higher orbital forever; the ground state is the only level where it can remain indefinitely. After about  $10^{-8}$  s, an excited atom returns to its ground state.



▲ **FIGURE 4.8 Classical Atom** An early-20th-century conception of the hydrogen atom—the Bohr model—pictured its electron orbiting the central proton in a well-defined orbit, much like a planet orbiting the Sun. Two electron orbitals of different energies are shown: (a) the ground state and (b) an excited state.



▲ **FIGURE 4.9 Modern Atom** The modern interpretation of the hydrogen atom visualizes the electron as a "cloud" surrounding the nucleus. The same two energy states are shown as in Figure 4.8.

## Radiation as Particles

Because electrons can exist only in orbitals having specific energies, atoms can absorb only specific amounts of energy as their electrons are boosted into excited states. Likewise, atoms can emit only specific amounts of energy as their electrons fall back to lower energy states. Thus, the amount of light energy absorbed or emitted in these processes *must correspond precisely to the energy difference between two orbitals*. The atom's quantized energy levels require that light be absorbed and emitted in the form of distinct "packets" of electromagnetic radiation, each carrying a specific amount of energy. We call these packets **photons**. A photon is, in effect, a "particle" of electromagnetic radiation.

The idea that light sometimes behaves not as a continuous wave, but as a stream of particles, was proposed by Albert Einstein in 1905 to explain a number of experimental results (especially the *photoelectric effect*—see *Discovery 4-1*) then puzzling physicists. Furthermore, Einstein was able to quantify the relationship between the two aspects of light's double nature. He found that the energy carried by a photon had to be proportional to the *frequency* of the radiation:

$$\text{photon energy} \propto \text{radiation frequency.}$$

For example, a "deep red" photon having a frequency of  $4 \times 10^{14}$  Hz (or a wavelength of approximately 750 nm) has half the energy of a violet photon of frequency of  $8 \times 10^{14}$  Hz (wavelength = 375 nm) and 500 times the energy of an  $8 \times 10^{11}$  Hz (wavelength = 375  $\mu$ m) microwave photon.

The constant of proportionality in the preceding relation is now known as *Planck's constant*, in honor of the German physicist Max Planck, who determined its numerical value. It is always denoted by the symbol  $h$ , and the equation relating the photon energy  $E$  to the radiation frequency  $f$  is usually written

$$E = hf.$$

# MORE PRECISELY 4-1

## The Hydrogen Atom

By observing the emission spectrum of hydrogen and using the connection between photon energy and color first suggested by Einstein (Section 4.2), Niels Bohr determined early in the 20th century what the energy differences between the various energy levels must be. Using that information, he was then able to infer the actual energies of the excited states of hydrogen.

A unit of energy often used in atomic physics is the *electron volt* (eV). (The name actually has a rather technical definition: the amount of energy gained by an electron when it accelerates through an electric potential of 1 volt. For our purposes, however, it is just a convenient quantity of energy.) One electron volt (1 eV) is equal to  $1.60 \times 10^{-19}$  J (joule)—roughly half the energy carried by a single photon of red light. The minimum amount of energy needed to ionize hydrogen from its ground state is 13.6 eV. Bohr numbered the energy levels of hydrogen, with level 1 the ground state, level 2 the first excited state, and so on. He found that, by assigning zero energy to the ground state, the energy of any state (the  $n$ -th, say) could then be written as follows:

$$E_n = 13.6 \left( 1 - \frac{1}{n^2} \right) \text{eV}.$$

Thus, the ground state ( $n = 1$ ) has energy  $E_1 = 0$  eV, the first excited state ( $n = 2$ ) has energy  $E_2 = 13.6 \times (1 - 1/4)$  eV = 10.2 eV, the second excited state has energy  $E_3 = 13.6 \times (1 - 1/9)$  eV = 12.1 eV, and so on. There are infinitely many excited states between the ground state and the energy at which the atom is ionized, crowding closer and closer together as  $n$  increases and  $E_n$  approaches 13.6 eV.

**EXAMPLE** Using Bohr's formula for the energy of each electron orbital, we can reverse his reasoning and calculate the energy associated with a transition between any two given states. To boost an electron from the first excited state to the second, an atom must be supplied with  $E_3 - E_2 = 12.1$  eV - 10.2 eV = 1.9 eV of energy, or  $3.0 \times 10^{-19}$  J. Now, from the formula  $E = hf$  presented in the text, we find that this energy corresponds to a photon with a frequency of  $4.6 \times 10^{14}$  Hz, having a wavelength of 656 nm, and lying in the red portion of the spectrum. (A more precise calculation gives the value 656.3 nm reported in the text.)


The accompanying diagram summarizes the structure of the hydrogen atom. The increasing energy levels are depicted as a series of circles of increasing radius. The electronic transitions between these levels (indicated by arrows) are conventionally grouped into families, named after their discoverers, that define the terminology used to identify specific spectral lines. (Note that the spacings of the energy levels are not drawn to scale here to provide room for all labels on the diagram. In reality, the circles should become more and more closely spaced as we move outward.)

Transitions starting from or ending at the ground state (level 1) form the *Lyman series*, named after American

spectroscopist Theodore Lyman, who discovered these lines in 1914. The first is *Lyman alpha* (Lya), corresponding to the transition between the first excited state (level 2) and the ground state. The energy difference is 10.2 eV, and the Lya photon has a wavelength of 121.6 nm (1216 Å). The Lyβ (beta) transition, between level 3 (the second excited state) and the ground state, corresponds to an energy change of 12.10 eV and a photon of wavelength 102.6 nm (1026 Å). Lyγ (gamma) corresponds to a jump from level 4 to level 1, and so on. All Lyman-series energies lie in the ultraviolet region of the spectrum.

The next series of lines, the *Balmer series*, involves transitions down to (or up from) level 2, the first excited state. The series is named after the Swiss mathematician Johann Balmer, who didn't discover these lines (they were well known to spectroscopists early in the 19th century), but who published a mathematical formula for their wavelengths in 1885. All the Balmer series lines lie in or close to the visible portion of the electromagnetic spectrum.

Because they form the most easily observable part of the hydrogen spectrum and were the first to be discovered, the Balmer lines are often referred to simply as the *Hydrogen series*, denoted by the letter *H*. As with the Lyman series, the individual transitions are labeled with Greek letters. An Hα photon (level 3 to level 2) has a wavelength of 656.3 nm, in the red part of the visible spectrum; Hβ (level 4 to level 2) has a wavelength of 486.1 nm (green); Hγ (level 5 to level 2) has a wavelength of 434.1 nm (blue); and so on. We will use these designations (especially Hα and Hβ) frequently in later chapters. The most energetic Balmer series photons have energies that place them just beyond the blue end of the visible spectrum, in the near ultraviolet.



A few of the transitions making up the Lyman and Balmer (Hydrogen) series are marked on the figure. There are infinitely many other families of lines, all lying in the infrared and radio regions of the spectrum, but astronomically, the Lyman and Balmer sequences are the most important.

Like the gravitational constant  $G$  and the speed of light,  $c$ , Planck's constant is one of the fundamental physical constants of the universe.

In SI units, the value of Planck's constant is a very small number:  $h = 6.63 \times 10^{-34}$  joule seconds ( $\text{J} \cdot \text{s}$ ). Consequently, the energy of a single photon is tiny. Even a very high frequency gamma ray (the most energetic type of electromagnetic radiation) with a frequency of  $10^{22}$  Hz has an energy of just  $(6.63 \times 10^{-34}) \times 10^{22} \approx 7 \times 10^{-12}$  J—about the same energy carried by a flying gnat. Nevertheless, this energy is more than enough to damage a living cell. The basic reason that gamma rays are so much more dangerous to life than visible light is that each gamma-ray photon typically carries millions, if not billions, of times more energy than a photon of visible radiation.

The equivalence between the energy and frequency (or inverse wavelength) of a photon completes the connection between atomic structure and atomic spectra. Atoms absorb and emit radiation at characteristic wavelengths determined by their own particular internal structure. Because this structure is *unique* to each element, the colors of the absorbed and emitted photons—that is, the spectral lines we observe—are characteristic of that element *and only that element*. The spectrum we see is thus a unique identifier of the atom involved.

Many people find it confusing that light can behave in two such different ways. To be truthful, modern physicists don't yet fully understand *why* nature displays this wave-particle duality. Nevertheless, there is irrefutable experimental evidence for both of these aspects of radiation. Environmental conditions ultimately determine which description—wave or stream of particles—better fits the behavior of electromagnetic radiation in a particular instance. As a general rule of thumb, in the macroscopic realm of everyday experience, radiation is more usefully described as a wave, whereas in the microscopic domain of atoms, it is best characterized as a series of particles.

### PROCESS OF SCIENCE Check

- ✓ Describe the scientific reasoning leading to the conclusion that light behaves both as a wave and a particle.

## 4.3 Formation of Spectral Lines

With quantum mechanics as our guide to the internal structure of atoms, we can now explain quantitatively the spectral lines we see. Let's start with hydrogen, the simplest element, then move on to more complex systems.

### The Spectrum of Hydrogen

The full spectrum of hydrogen consists of many lines, spread across much of the electromagnetic spectrum, from

ultraviolet to radio; we focus here on just a few of those lines. The energy levels and spectrum of hydrogen are discussed in more detail in *More Precisely 4-1*.

Figure 4.10 illustrates schematically the absorption and emission of photons by a hydrogen atom. Figure 4.10(a) shows the atom absorbing a photon and making a transition from the ground state to the first excited state. It then emits a photon of precisely the same energy and drops back to the ground state. The energy difference between the two states corresponds to an ultraviolet photon of wavelength 121.6 nm (1216 Å).

Absorption may also boost an electron into an excited state higher than the first excited state. Figure 4.10(b) depicts the absorption of a more energetic (higher frequency, shorter wavelength) ultraviolet photon, one with a wavelength of 102.6 nm (1026 Å). The absorption of this photon causes the atom to jump to the *second* excited state. As before, the atom returns rapidly to the ground state, but this time, because there are two states lying below the excited state, the atom can do so in one of two possible ways:

1. It can proceed directly back to the ground state, in the process emitting an ultraviolet photon identical to the one that excited the atom in the first place.
2. Alternatively, the electron can *cascade* down, one orbital at a time. If this occurs, the atom will emit *two* photons: one with an energy equal to the difference between the second and first excited states and the other with an energy equal to the difference between the first excited state and the ground state.

Either possibility can occur, with roughly equal probability. The second step of the cascade process produces a 121.6-nm ultraviolet photon, just as in Figure 4.10(a). However, the first transition—the one from the second to the first excited state—produces a photon with a wavelength of 656.3 nm (6563 Å), which is in the visible part of the spectrum. This photon is seen as red light. An individual atom—if one could be isolated—would emit a momentary red flash. This is the origin of the red line in the hydrogen spectrum shown in Figure 4.3.

The absorption of additional energy can boost the electron to even higher orbitals within the atom. As the excited electron cascades back down to the ground state, the atom may emit many photons, each with a different energy and hence a different wavelength, and the resulting spectrum shows many spectral lines. In a sample of heated hydrogen gas, at any instant atomic collisions ensure that atoms are found in many different excited states. The complete emission spectrum therefore consists of wavelengths corresponding to all possible transitions between those states and states of lower energy.

In the case of hydrogen, all transitions ending at the ground state produce ultraviolet photons. However, downward transitions ending at the *first* excited state give rise to spectral lines in or near the visible portion of the




## DISCOVERY 4-1

## The Photoelectric Effect

Einstein developed his breakthrough insight into the nature of radiation partly as a means of explaining a puzzling experimental result known as the **photoelectric effect**. This effect can be demonstrated by shining a beam of light on a metal surface (as shown in the accompanying figure). When high-frequency ultraviolet light is used, bursts of electrons are dislodged from the surface by the beam, much as when one billiard ball hits another, knocking it off the table. However, the speed with which the particles are ejected from the metal is found to depend only on the *color* of the light, and not on its intensity. For lower-frequency light—blue, say—an electron detector still records bursts of electrons, but now their speeds, and hence their *energies*, are less. For even lower frequencies—red or infrared light—*no* electrons are kicked out of the metal surface at all.

These results are difficult to reconcile with a wave model of light, which would predict that the energies of the ejected electrons should increase steadily with increasing intensity at any frequency. Instead, the detector shows an abrupt cutoff in ejected electrons as the frequency of the incoming radiation drops below a certain level. Einstein realized that the only way to explain the cutoff, and the increase in electron speed with frequency above the cutoff, was to envision radiation as traveling as “bullets,” or particles, or *photons*. Furthermore, to account for the experimental findings, the energy of any photon had to be proportional to the *frequency* of the radiation. Low-frequency, long-wavelength photons carry less energy than high-frequency, short-wavelength ones.

If we also suppose that some minimum amount of energy is needed just to “unplug” the electrons from the metal, then we can see why no electrons are emitted below some critical frequency: The photons associated with red light



in the diagram just don't carry enough energy. Above the critical frequency, photons do have enough energy to dislodge the electrons. Moreover, any energy they possess above the necessary minimum is imparted to the electrons as *kinetic energy*, the energy of motion. Thus, as the frequency of the radiation increases, so, too, does the photon's energy and hence the speed of the electrons that they liberate from the metal.

The realization and acceptance of the fact that light can behave both as a wave and as a particle is another example of the scientific method at work. Despite the enormous success of the wave theory of radiation in the 19th century, the experimental evidence led 20th-century scientists to the inevitable conclusion that the theory was incomplete—it had to be modified to allow for the fact that light sometimes acts like a particle. Although Einstein is perhaps best known today for his theories of relativity, in fact his 1919 Nobel prize was for his work on the photoelectric effect. In addition to bringing about the birth of a whole new branch of physics—the field of quantum mechanics—Einstein's explanation of the photoelectric effect radically changed the way physicists view light and all other forms of radiation.


electromagnetic spectrum (Figure 4.3; *More Precisely 4-1*). Other transitions ending in higher states generally give rise to infrared and radio spectral lines.

The inset in Figure 4.10 shows an astronomical object whose red coloration is the result of precisely the process mentioned in step 2 on p. 87. As ultraviolet photons from a young, hot star pass through the surrounding cool hydrogen gas out of which the star recently formed, some photons are absorbed by the gas, boosting its atoms into excited states or ionizing

them completely. The 656.3-nm red glow characteristic of excited hydrogen gas results as the atoms cascade back to their ground states. The phenomenon is called *fluorescence*.

## Kirchhoff's Laws Explained

Let's reconsider our earlier discussion of emission and absorption lines in terms of the model just presented. In Figure 4.7, a beam of continuous radiation shines



**Interactive FIGURE 4.10 Atomic Excitation** (a) Absorption of an ultraviolet (UV) photon (left) by a hydrogen atom causes the momentary excitation of the atom into its first excited state (center). After about  $10^{-8}$  s, the atom returns to its ground state (right), in the process emitting a photon having exactly the same energy as the original photon. (b) Absorption of a higher-energy UV photon may boost the atom into a higher excited state, from which there are several possible paths back to the ground state. At the top, the electron falls immediately back to the ground state, emitting a photon identical to the one it absorbed. At the bottom, the electron initially falls into the first excited state, producing visible radiation of wavelength 656.3 nm—the characteristic ( $H\alpha$ ) red glow of excited hydrogen. (That's why many nebulae, like the one in the inset, glow red.) Subsequently, the atom emits another photon (having the same energy as in part (a)) as it falls back to the ground state. (Inset: NASA)

through a cloud of hydrogen gas. The beam contains photons of all energies, but most don't interact with the gas—the gas can absorb only those photons having just the right energy to cause a change in an electron's orbit from one state to another. All other photons in the beam—with energies that cannot produce a transition—do not interact with the gas at all, but pass through it unhindered. Photons having the right energies are absorbed, excite the gas, and are removed from the beam. This is the cause of the dark absorption lines in the spectrum of Figure 4.7(b). The lines are direct indicators of the energy differences between orbitals in the atoms making up the gas.

The excited atoms rapidly return to their original states, each emitting one or more photons in the process.

Most of these reemitted photons leave at angles that do *not* take them through the slit and onto our detector. A second detector looking at the cloud from the side would record the reemitted energy as an emission spectrum, as illustrated in Figure 4.7(c). An astronomical example is the *emission nebula* shown in the inset to Figure 4.10. Like the absorption spectrum, the emission spectrum is characteristic of the gas, not of the original beam. The type of spectrum we see depends on our chance location with respect to both the source and the intervening cloud.

Figure 4.7(a) shows a *continuous* spectrum, in which emitted photons escape from the bulb without further interaction with matter. Actually, the situation in a dense source of radiation (a thick gas cloud or in a liquid or solid body) is more complex. There, a photon is likely to interact with

atoms, free electrons, and ions in the body many times before finally escaping, exchanging some energy with the matter at each encounter. The net result is that the emitted radiation displays a continuous spectrum, in accordance with Kirchhoff's first law. The spectrum is approximately that of a blackbody with the same temperature as the source.

## More Complex Spectra


All hydrogen atoms have basically the same structure—a single electron orbiting a single proton—but, of course, there are many other kinds of atoms, each kind having a unique internal structure. The number of protons in the nucleus of an atom determines the **element** that it represents. Just as all hydrogen atoms have a single proton, all oxygen atoms have 8 protons, all iron atoms have 26 protons, and so on.

The next simplest element after hydrogen is helium. The central nucleus of the most common form of helium is made up of two protons and two **neutrons** (another kind of elementary particle having a mass slightly larger than that of a proton, but having no electrical charge). Two electrons orbit this nucleus. As with hydrogen and all other atoms, the “normal” condition for helium is to be electrically neutral, with the negative charge of the orbiting electrons exactly canceling the positive charge of the nucleus (Figure 4.11a).

More complex atoms contain more protons (and neutrons) in the nucleus and have correspondingly more orbiting electrons. For example, an atom of carbon, shown in Figure 4.11(b), consists of six electrons orbiting a nucleus containing six protons and six neutrons. As we progress to heavier and heavier elements, the number of orbiting electrons increases, and the number of possible electron transitions rises rapidly. The result is that very complicated spectra can be produced. The complexity of atomic spectra generally reflects the complexity of the atoms themselves. A good example is the element iron, which contributes nearly 800 of the Fraunhofer absorption lines seen in the solar spectrum (Figure 4.4).

Atoms of a single element such as iron can yield many lines for two main reasons. First, the 26 electrons of a normal iron atom can make an enormous number of different transitions among available energy levels. Second, many iron atoms are *ionized*, with some of their 26 electrons stripped away. The removal of electrons alters an atom's electromagnetic structure, and the energy levels of ionized iron are quite different from those of

► **FIGURE 4.11 Helium and Carbon** (a) A helium atom in its ground state has two electrons within its lowest-energy orbital, around a nucleus containing two protons and two neutrons. (b) A carbon atom in its ground state has six electrons orbiting around a six-proton, six-neutron nucleus—two of the electrons in an inner orbital and the other four at a greater distance from the center.



*Remember, the clean orbitals shown here and in other atomic diagrams are really more like fuzzy “clouds” of electron energy levels, as shown in Figure 4.9.*


neutral iron. Each new level of ionization introduces a whole new set of spectral lines. Besides iron, many other elements, also in different stages of excitation and ionization, absorb photons at visible wavelengths. When we observe the entire Sun, all these atoms and ions absorb simultaneously, yielding the rich spectrum we see.

The power of spectroscopy is most apparent when a cloud contains many different gases mixed together, because it enables us to study one kind of atom or ion to the exclusion of all others simply by focusing on specific wavelengths of radiation. By identifying the superimposed absorption and emission spectra of many different atoms, we can determine the cloud's composition (and much more—see Section 4.4). Figure 4.12 shows an actual spectrum observed from a real cosmic object. As in Figure 4.10, the characteristic red glow of this emission nebula comes from the H $\alpha$  transition in hydrogen, the nebula's main constituent.

Spectral lines occur throughout the entire electromagnetic spectrum. Usually, electron transitions among the lowest orbitals of the lightest elements, such as hydrogen and helium, produce visible and ultraviolet spectral lines. Transitions among very highly excited states of hydrogen and other elements can produce spectral lines in the infrared and radio parts of the electromagnetic spectrum. Conditions on Earth make it all but impossible to detect these radio and infrared features in the laboratory, but they are routinely observed by radio and infrared telescopes (see Chapter 5) in radiation coming from space. Electron transitions among lower energy levels in heavier, more complex elements produce X-ray spectral lines, which have been observed in the laboratory. Some have also been observed in stars and other cosmic objects.

## CONCEPT Check

- ✓ How does the structure of an atom determine the atom's emission and absorption spectra?



▲ **FIGURE 4.12 Emission Nebula** The visible spectrum of hot gases in the Omega Nebula (M17). (The word *nebula* means “gas cloud”—one of many places in our Galaxy where new stars are forming today.) Shining by the light of several very hot stars, the gas in the nebula produces a complex spectrum of bright and dark lines (bottom). That same spectrum can also be displayed, as shown here, as a white graph of intensity versus frequency, spanning the spectrum from red to blue. (Adapted from ESO)

## 4.4 Molecules


A **molecule** is a tightly bound group of atoms held together by interactions among their orbiting electrons—interactions that we call *chemical bonds*. Much like atoms, molecules can exist only in certain well-defined energy states, and again like atoms, molecules produce characteristic emission or absorption spectral lines when they make a transition from one state to another. Because molecules are more complex than individual atoms, the rules of molecular physics are also more complex. Nevertheless, as with atomic spectral lines, painstaking experimental work over many decades has determined the precise frequencies (or wavelengths) at which millions of molecules emit and absorb radiation.

In addition to the lines resulting from electron transitions, molecular lines result from two other kinds of change not possible in atoms: Molecules can *rotate*, and they can *vibrate*. Figure 4.13 illustrates these basic molecular motions. Molecules rotate and vibrate in specific ways. Just as with atomic states, only certain spins and vibrations are allowed by the rules of molecular physics. When a molecule *changes* its rotational or vibrational state, a photon is emitted or absorbed. Spectral lines characteristic of the specific kind of molecule result. Like their atomic counterparts, these lines are unique molecular fingerprints, enabling researchers to


identify and study one kind of molecule to the exclusion of all others. As a rule of thumb,

- *electron transitions* within molecules produce visible and ultraviolet spectral lines (the largest energy changes).
- changes in molecular *vibration* produce infrared spectral lines.
- changes in molecular *rotation* produce spectral lines in the radio part of the electromagnetic spectrum (the smallest energy changes).

Molecular lines usually bear little resemblance to the spectral lines associated with their component atoms. For example, Figure 4.14(a) shows the emission spectrum of the simplest molecule known: molecular hydrogen. Notice how different it is from the spectrum of atomic hydrogen shown in part (b) of the figure.



▲ **FIGURE 4.13 Molecular Emission** Molecules can change in three ways while emitting or absorbing electromagnetic radiation. The colors and wavelengths of the emitted photons represent the relative energies involved. Sketched here is the molecule carbon monoxide (CO) undergoing (a) a change in which an electron in the outermost orbital of the oxygen atom drops to a lower energy state (emitting a photon of shortest wavelength, in the visible or ultraviolet range), (b) a change in vibrational state (of intermediate wavelength, in the infrared), and (c) a change in rotational state (of longest wavelength, in the radio range).



▲ **FIGURE 4.14 Hydrogen Spectra** The emission spectrum of molecular hydrogen (a) differs greatly from that of simpler atomic hydrogen (b). (*Bausch & Lomb, Inc.*)

### CONCEPT Check

✓ What kinds of internal changes within a molecule can cause radiation to be emitted or absorbed?

## 4.5 Spectral-Line Analysis

Astronomers apply the laws of spectroscopy in analyzing radiation from beyond Earth. A nearby star or a distant galaxy takes the place of the lightbulb in our previous examples. An interstellar cloud or a stellar (or even planetary) atmosphere plays the role of the intervening cool gas, and a spectrograph attached to a telescope replaces our simple prism and detector. We began our study of electromagnetic radiation by stating that virtually all we know about planets, stars, and galaxies is gleaned from studies of the light we receive from them, and we have presented some of the ways in which that knowledge is obtained. Here, we describe a few of the ways in which the properties of emitters and absorbers can be determined by careful analysis of radiation received on (or near) Earth. We will encounter other important examples as our study of the cosmos unfolds.

### A Spectroscopic Thermometer

In the hot interior of a star, atoms are fully ionized. Electrons travel freely through the gas, unbound to any nucleus, and the spectrum of radiation is continuous. However, near the relatively cool stellar surface, some atoms retain a few, or even most, of their orbital electrons. As noted earlier, astronomers can determine the star's chemical composition by matching the spectral lines they see with the laboratory spectra of known atoms, ions, and molecules.

The strength of a spectral line (brightness or darkness, depending on whether the line is seen in emission or absorption) depends on the number of atoms giving rise to it. The more atoms there are to emit or absorb photons of the appropriate frequency, the stronger the line. But the strength


of line also depends critically on the *temperature* of the gas containing the atoms, because temperature determines how many atoms at any instant are in the right orbital to undergo any particular transition. Simply put, at low temperatures, only low-lying energy states tend to be populated, and transitions into and out of those states dominate the spectrum. At higher temperatures, more atoms are in excited states, and some may be ionized, radically changing the character of the possible transitions and hence the spectrum we see.

Spectroscopists have developed mathematical formulas that relate the number of emitted or absorbed photons to the energy levels of the atoms involved and the temperature of the gas. Once an object's spectrum is measured, astronomers can interpret it by matching the observed intensities of the spectral lines with those predicted by the formulas. In this way, astronomers can refine their measurements of both the composition *and* the temperature of the gas producing the lines. These temperature measurements are generally much more accurate than crude estimates based on the radiation laws and the assumption of blackbody emission. ∞ (Sec. 3.4) In Chapter 17, we will see how these ideas are put to use in the classification and interpretation of stellar spectra.

### Measurement of Radial Velocity

The Doppler effect—the apparent shift in the frequency of a wave due to the motion of the source relative to the observer—is a classical phenomenon common to all waves. ∞ (Sec. 3.5) However, by far its most important astronomical application comes when it is combined with observations of atomic and molecular spectral lines.

The spectra of many atoms, ions, and molecules are well known from laboratory measurements. Often, however, a familiar pattern of lines appears, but the lines are displaced from their usual locations. In other words, as illustrated in Figure 4.15, a set of spectral lines may be recognized as belonging to a particular element, except that the lines are all offset—blueshifted or redshifted—by the same fractional amount from their normal wavelengths. These shifts are due



**Interactive FIGURE 4.15 Doppler Shift** The Doppler effect shifts the entire spectrum of a moving object to higher or lower frequencies. The spectrum at top shows the redshift of the hydrogen lines from an object moving at a speed of 300 km/s away from the observer. The amount of the shift (0.1 percent here) tells us the object's recession velocity—0.001  $c$ . The spectrum at bottom shows the blueshift of the same set of lines from an object approaching us at 600 km/s. The shift is twice as large (0.2 percent), because the speed has doubled, and in the opposite sense because the direction has reversed.

to the Doppler effect, and they allow astronomers to measure how fast the source of the radiation is moving along the line of sight from the observer (the *radial velocity* of the source).

For example, in Figure 4.15, the 486.1-nm  $H\beta$  line of hydrogen in the spectrum of a distant galaxy is received on Earth at a wavelength of 485.1 nm—blueshifted to a slightly shorter wavelength. (Remember, we know that it is the  $H\beta$  line because *all* the hydrogen lines are observed to have the same fractional shift—the characteristic line pattern identifies the spectrum as that of hydrogen.) We can compute the galaxy's line-of-sight velocity relative to Earth by using the Doppler equation presented in Section 3.5. The calculation is essentially the same as that presented in *More Precisely 3-3*: The change in wavelength (apparent minus true) is 485.1 nm – 486.1 nm = –1.0 nm (the negative sign simply indicating that the wavelength has decreased). It then follows that the recession velocity is

$$\frac{-1.0 \text{ nm}}{486.1 \text{ nm}} \times c = -620 \text{ km/s.}$$

In other words, the galaxy is *approaching* us (this is the meaning of the negative sign) at a speed of 620 km/s.


This book will have a lot to say about the motions of planets, stars, and galaxies throughout the universe. Just bear in mind that almost all of that information is derived from telescopic observations of Doppler-shifted spectral lines in many different parts of the electromagnetic spectrum.

## Line Broadening


The structure of the lines themselves reveals still more information. At first glance the emission lines shown earlier may seem uniformly bright, but more careful study shows that this is in fact not the case. As illustrated in Figure 4.16, the brightness of a line is greatest at the center and falls off toward

either side. Earlier, we stressed that photons are emitted and absorbed at very precise energies, or frequencies. Why, then, aren't spectral lines extremely narrow, occurring only at specific wavelengths? This *line broadening* is not the result of some inadequacy of our experimental apparatus; rather, it is caused by the *environment* in which the emission or absorption occurs—the physical state of the gas or star in which the line is formed. For definiteness, we have drawn Figure 4.16 and subsequent figures to refer to emission lines, but realize that the ideas apply equally well to absorption features.

Several processes can broaden spectral lines. The most important involve the Doppler effect. Imagine a hot cloud of gas containing individual atoms in random thermal motion in every possible direction, as illustrated in Figure 4.17(a). If an atom happens to be moving away from us as it emits



**▲ FIGURE 4.16 Line Profile** By tracing the changing brightness across a typical emission line (a) and expanding the scale, we obtain a graph of the line's intensity versus its frequency (b).



▲ **FIGURE 4.17 Thermal Broadening** Atoms moving randomly (a) produce broadened spectral lines (b) as their individual redshifted and blueshifted emission lines merge in our detector. The hotter the gas, the greater is the degree of thermal broadening.


a photon, that photon is redshifted by the Doppler effect—we do not record it at the precise wavelength predicted by atomic physics, but rather at a slightly longer wavelength. The extent of this redshift is proportional to the atom’s instantaneous velocity away from the detector. Similarly, if the atom is moving toward us at the instant of emission, its light is blueshifted. In short, because of thermal motion within the gas, emission and absorption lines are observed at frequencies slightly different from those we would expect if all atoms in the cloud were motionless.

Most atoms in a typical cloud have small thermal velocities, so in most cases the line is Doppler shifted just a little. Only a few atoms have large shifts. As a result, the center of a spectral line is much more pronounced than its “wings,” producing a bell-shaped spectral feature like that shown in Figure 4.17(b). Thus, even if all atoms emitted and absorbed photons at only one precise wavelength, the effect of

their thermal motion would be to smear the line out over a range of wavelengths. The hotter the gas, the larger the spread of Doppler motions and the greater the width of the line. ∞ (*More Precisely 3-1*) By measuring a line’s width, astronomers can estimate the average speed of the particles and hence the temperature of the gas producing it.

Other processes, such as *rotation* and *turbulence*, can produce similar effects. Consider an astronomical object (a star or a gas cloud) that is spinning about some axis as sketched in Figure 4.18 or that has some other internal motion, such as turbulent eddies or vortices on many scales. Photons emitted from regions that happen to be moving toward us are blueshifted by the Doppler effect; photons emitted from regions moving away from us are redshifted. Often the object under study is so small or far away that our equipment cannot distinguish, or *resolve*, different parts from one another—all the emitted light is blended together in our detector. In that case, the result is a net broadening of the observed spectral lines. The more rapid the internal motion, the more broadening we see. Note that this broadening has *nothing* to do with the temperature of the gas producing the lines and is generally superimposed on the thermal broadening just discussed.

Still other broadening mechanisms do not depend on the Doppler effect at all. For example, if electrons are moving between orbitals while their parent atom is colliding with another atom, the energy of the emitted or absorbed photons changes slightly, blurring the spectral lines. This mechanism, which occurs most often in dense gases where



▲ **FIGURE 4.18 Rotational Broadening** Rotation of a star can cause spectral line broadening. Since most stars are unresolved—that is, they are so distant that we cannot distinguish one part of the star from another—light rays from all parts of the star merge to produce broadened lines.

collisions are most frequent, is usually referred to as *collisional broadening*. The amount of broadening increases as the density of the emitting or absorbing gas rises.

Finally, *magnetic fields* can also broaden spectral lines by a process called the *Zeeman effect*. The electrons and nuclei within atoms behave as tiny spinning magnets, and the basic emission and absorption rules of atomic physics change slightly whenever atoms are immersed in a magnetic field, as is the case in many stars to greater or lesser extents. The result is a slight splitting of a spectral line, which then blurs into an overall line broadening. Generally, the stronger the magnetic field, the more pronounced is the broadening.

## The Message of Starlight

Given sufficiently sensitive equipment, there is almost no end to the wealth of data that can be obtained from starlight. Table 4.1 lists some basic measurable properties of an incoming beam of radiation and indicates what sort of information can be obtained from them.

It is important to realize, however, that deciphering the extent to which each of the factors just described influences a spectrum can be a very difficult task. Typically, the spectra of many elements are superimposed on one another, and often several competing physical effects are occurring simultaneously, each modifying the spectrum in its own way. Further analysis is generally required to disentangle them. For example, if we know the temperature of the emitting gas (perhaps by comparing intensities of different spectral lines,

**TABLE 4.1** Spectral Information Derived from Starlight

Observed Spectral Characteristic	Information Provided
Peak frequency or wavelength (continuous spectra only)	Temperature (Wien's law)
Lines present	Composition, temperature
Line intensities	Composition, temperature
Line width	Temperature, turbulence, rotation speed, density, magnetic field
Doppler shift	Line-of-sight velocity

as discussed earlier), then we can calculate how much of the broadening is due to thermal motion and therefore how much is due to the other mechanisms just described. In addition, it is often possible to distinguish between the various broadening mechanisms by studying the detailed *shapes* of the lines.

The challenge facing astronomers is to decode spectral-line profiles to obtain meaningful information about the sources of the lines. In the next chapter, we will discuss some of the means by which astronomers obtain the raw data they need in their quest to understand the cosmos.

### CONCEPT Check


- ✓ Why is it so important for astronomers to analyze spectral lines in detail?

**The Big Question** Atoms constitute the basic essence of normal matter—the kind of matter from which stars, planets, and we ourselves are built. In turn, all atoms are made of smaller elementary particles, including protons, neutrons, and electrons. But even these particles are not fundamental, for physicists know that protons and neutrons are made of quarks. We are left wondering, Are quarks made of entities even smaller?

# Chapter Review


## SUMMARY

**1** A **spectroscope** (p. 80) is a device for splitting a beam of radiation into its component frequencies for detailed study. Many hot objects emit a **continuous spectrum** (p. 80) of radiation, containing light of all wavelengths. A hot gas may instead produce an **emission spectrum** (p. 81), consisting of only a few well-defined **emission lines** (p. 80) of specific frequencies, or colors. Passing a continuous beam of radiation through cool gas will produce



**absorption lines** (p. 82) at precisely the same frequencies as would be present in the gas's emission spectrum.


**2** **Kirchhoff's laws** (p. 83) describe the relationships among these different types of spectra. The emission and absorption lines produced by each element are unique—they provide a “fingerprint” of that element. The study of the spectral






lines produced by different substances is called **spectroscopy** (p. 83). Spectroscopic studies of the Fraunhofer lines in the solar spectrum yield detailed information about the Sun's composition.


**3 Atoms** (p. 84) are made up of negatively charged electrons orbiting a positively charged **nucleus** (p. 84) consisting of positively charged protons and electrically neutral **neutrons** (p. 90). The number of protons in the nucleus determines the particular **element** (p. 90) the atom represents. In the **Bohr model** (p. 84), a hydrogen atom has a minimum energy **ground state** (p. 84) representing its “normal” condition. When the electron has a higher than normal energy, the atom is in an **excited state** (p. 85). For any given atom, only certain, well-defined energies are possible. In the modern view, the electron is envisaged as being spread out in a “cloud” around the nucleus, but still with a sharply defined energy.




**4** Electromagnetic radiation exhibits both wave and particle properties. Particles of radiation are called **photons** (p. 85). In order to explain the **photoelectric effect** (p. 88), Einstein found that the energy of a photon must be directly proportional to the photon's frequency.




**5** As electrons move between energy levels within an atom, the difference in energy between the states is emitted or absorbed in the form of photons. Because the energy levels have definite energies, the photons also have definite energies, and hence colors, that are characteristic of the type of atom involved.



**6 Molecules** (p. 91) are groups of two or more atoms bound together by electromagnetic forces. Like atoms, molecules exist in energy states that obey rules similar to those governing the internal structure of atoms. When a molecule makes a transition between energy states, it emits or absorbs a characteristic spectrum of radiation that identifies it uniquely.



**7** Astronomers apply the laws of spectroscopy to analyze radiation from beyond Earth. Several physical mechanisms can broaden spectral lines. The most important is the Doppler effect, which occurs because stars are hot, or rotating, or turbulent, so their atoms are in constant motion.



**MasteringAstronomy**® For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- LO1** What is an absorption spectrum? An emission spectrum? How are they related?
- Describe the basic components of a simple spectroscope.
- LO2** What is spectroscopy? How can spectroscopy be used to infer the composition and temperature of a star?
- Why are gamma rays generally harmful to life-forms, but radio waves generally harmless?
- In the particle description of light, what is color?
- LO3** What is an atom? In what ways does the Bohr model of atomic structure differ from the modern view?
- Give a brief description of a hydrogen atom.
- LO4** What does it mean to say that a physical quantity is quantized? Why do we think light is quantized?
- What is the normal condition for atoms? What is an excited atom? What are orbitals?
- LO5** Why do excited atoms absorb and reemit radiation at characteristic frequencies?
- POS** How are absorption and emission lines produced in a stellar spectrum? What information might absorption lines in the spectrum of a star reveal about a cloud of cool gas lying between us and the star?
- Why might spectral lines of an element in a star's spectrum be weak, even though that element is abundant in the star?
- LO6** How do molecules produce spectral lines unrelated to the movement of electrons between energy levels?
- How can the Doppler effect broaden a spectral line?
- LO7 POS** List three properties of a star that can be determined from observations of its spectrum.

## Conceptual Self-Test: Multiple Choice

- Compared with a ground-based spectrum, the spectrum of a star observed from above Earth's atmosphere would show (a) no absorption lines; (b) fewer emission lines; (c) fewer absorption lines; (d) many more absorption lines.
- The visible spectrum of sunlight reflected from Saturn's cold moon Titan would be expected to be (a) continuous; (b) an emission spectrum; (c) an absorption spectrum.
- VIS** Figure 4.3 ("Elemental Emission") shows the emission spectrum of neon gas. If the temperature of the gas were increased, we would observe (a) fewer red lines and more blue lines; (b) even more red lines; (c) some faint absorption features; (d) no significant change.
- Compared with a star having many blue absorption lines, a star with many red and blue absorption lines must be (a) cooler; (b) of different composition; (c) moving away from the observer; (d) moving away from the other star.
- An atom that has been ionized (a) has equal numbers of protons and electrons; (b) has more protons than electrons; (c) is radioactive; (d) is electrically neutral.
- VIS** In Figure 4.10 ("Atomic Excitation"), compared with an electron transition from the first excited state to the ground state, a transition from the third excited state to the second excited state emits a photon of (a) greater energy; (b) lower energy; (c) identical energy.
- Compared with a complex atom like neon, a simple atom such as hydrogen has (a) more excited states; (b) fewer excited states; (c) the same number of excited states.
- Compared with cooler stars, the hottest stars have absorption lines that are (a) thin and distinct; (b) broad and fuzzy; (c) identical to the lines in the cooler stars.
- Compared with slowly rotating stars, the fastest spinning stars have absorption lines that are (a) thin and distinct; (b) broad and fuzzy; (c) identical to the lines in the slowly rotating stars.
- Astronomers analyze starlight to determine a star's (a) temperature; (b) composition; (c) motion; (d) all of the above.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

- What is the energy (in electron volts—see *More Precisely 4-1*) of a 450-nm blue photon? A 200-nm ultraviolet photon?
- What is the energy (in electron volts) of a 100-GHz (1 gigahertz =  $10^9$  Hz) microwave photon?
- What are the wavelengths of (a) a 2-eV red photon, (b) an 0.1-eV infrared photon, and (c) a 5000-eV (5-keV) X-ray?
- How many times more energy has a 1-nm gamma ray than a 10-MHz radio photon?
- List all the spectral lines of hydrogen that lie in the visible range (taken to run from 400 to 700 nm in wavelength).
- How many different photons (i.e., photons of different frequencies) can be emitted as a hydrogen atom in the third excited state falls back, directly or indirectly, to the ground state? What are their wavelengths?
- A distant galaxy is receding from Earth with a radial velocity of 3000 km/s. At what wavelength would its Ly $\alpha$  line be received by a detector above Earth's atmosphere?
- In a demonstration of the photoelectric effect, suppose that a minimum energy of  $5 \times 10^{-19}$  J (3.1 eV) is required to dislodge an electron from a metal surface. What is the minimum frequency (and longest wavelength) of radiation for which the detector registers a response?


## Activities

### Collaborative

- Find a spectrum of the Sun that also has a wavelength scale on it. Google is a good place to start. Select some absorption lines and determine their wavelengths by interpolation. Now, try to identify the element that produced these lines. Use a reference of lines such as Moore's "A Multiplet Table of Astrophysical Interest," available on the NASA Astrophysics Data System. Work with the darkest lines before trying the fainter ones. How many elements can you find?

### Individual

- Obtain a handheld spectroscope, available from your school science lab or online. In the shade, point the spectroscope at a white cloud or white piece of paper that is in direct sunlight. Look for the absorption lines in the Sun's spectrum. Note their wavelength from the scale inside the spectroscope. How many of the lines can you identify by comparing your list with the Fraunhofer lines given in many astronomy reference books, or on Wikipedia?



# Telescopes

## THE TOOLS OF ASTRONOMY

At its heart, astronomy is an observational science. More often than not, observations of cosmic phenomena precede any clear theoretical understanding of their nature. As a result, our detecting instruments—our telescopes—have evolved to observe as broad a range of wavelengths as possible.

Until the middle of the 20th century, telescopes were limited to collecting visible light. Since then, technological advances have expanded our view of the universe to all regions of the electromagnetic spectrum. Some telescopes are sited on Earth, whereas others must be placed in space. However they are built and wherever they operate, telescopes are devices whose basic purpose is to collect electromagnetic radiation and deliver it to a detector for detailed study.

**The Big Picture** Telescopes are time machines, and astronomers, in a sense, are historians. Their detectors enhance our senses, enabling us to look far out in space—hence far back in time. Some of the biggest telescopes allow us to explore objects much farther than would be possible with our unaided eyes and to perceive radiation at wavelengths far beyond human vision. Almost everything in this book would be unknown without telescopes.

**LEFT:** Astronomers like to think big, and really big telescopes are now on the drawing board. This artist's conception for the European Southern Observatory shows ELT—the Extremely Large Telescope. With a mirror diameter of nearly 40 m, ELT would combine unrivaled light-gathering power with the ability to examine cosmic objects with unprecedented detail. This largest telescope in the world will be built on Cerro Armazones, a 3000-m mountaintop in Chile's Atacama Desert. (ESO/L. Calçada)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Sketch how optical telescopes work, and specify the advantages of reflecting telescopes over refractors.
- 2 Explain why larger telescopes gather more light and can make more detailed images.
- 3 Outline the purpose of some of the detectors used in astronomical telescopes.
- 4 Describe how Earth's atmosphere limits astronomical observations, and how astronomers overcome these limitations.
- 5 List some relative advantages and disadvantages of radio and optical astronomy.
- 6 Explain how interferometry is used to improve astronomical observations.
- 7 Describe the design of infrared, ultraviolet, and high-energy telescopes, and explain why some telescopes must be placed in space.
- 8 Say why it is important to make astronomical observations at many different wavelengths across the electromagnetic spectrum.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.

## 5.1 Optical Telescopes

In essence, a **telescope** is a “light bucket” whose primary function is to capture as many photons as possible from a given region of the sky and concentrate them into a focused beam for analysis. Much like a water bucket that collects only the rain falling into it, a telescope intercepts only that radiation falling onto it.


*Optical* telescopes are designed specifically to collect the wavelengths that are visible to the human eye. These telescopes have a long history, stretching back to the days of Galileo in the early 17th century, and for most of the past four centuries astronomers have built their instruments primarily for use in the narrow, visible, portion of the electromagnetic spectrum. ∞ (Sec. 3.3) Optical telescopes are probably also the best-known type of astronomical hardware, so it is fitting that we begin our study with them.

Although the various telescope designs presented in this section all come to us from optical astronomy, the discussion applies equally well to many instruments designed to capture invisible radiation, particularly in the infrared and ultraviolet regimes. Many large ground-based optical facilities are also used extensively for infrared work.\* Indeed, many ground-based observatories have recently been constructed with infrared observing as their principal function.

\*Recall from Chapter 3 that, while Earth’s atmosphere effectively blocks all ultraviolet, and most infrared, radiation, there remain several fairly broad spectral windows through which ground-based infrared observations can be made. ∞ (Sec. 3.3)



(a)



(b)

▲ **FIGURE 5.1 Refraction** A straw placed in a bowl of water appears bent (a) because the light from the part of the straw under the surface is refracted as it leaves the water and enters the air. (b) Consequently, the image formed in our eyes is displaced relative to the true position of the straw. (R. Megna/Fundamental Photographs, NYC)


## Refracting and Reflecting Telescopes

Optical telescopes fall into two basic categories: *refractors* and *reflectors*. **Refraction** is the bending of a beam of light as it passes from one transparent medium (e.g., air) into another (e.g., glass). Consider for example how a straw that is half immersed in a glass of water looks bent (Figure 5.1). The straw is straight, of course, but the light by which we see it is bent—refracted—as that light leaves the water and enters the air. When the light then enters our eyes, we perceive the straw as being bent.

A **refracting telescope** uses a *lens* to gather and concentrate a beam of light. Figure 5.2(a) shows how refraction at two faces of a prism can be used to change the direction of a beam of light. As illustrated in Figure 5.2(b), we can think of a lens as a series of prisms combined in such a way that all light rays arriving parallel to its axis (the imaginary line through the center of the lens), regardless of their distance from that axis, are refracted to pass through a single point, called the *focus*. The distance between the primary mirror and the focus is the *focal length*.

Figure 5.3 shows how a **reflecting telescope** uses a curved *mirror* instead of a lens to focus the incoming light. As shown in Figure 5.3(a), light striking a polished surface is reflected back, leaving the mirror at the same angle at which it arrived. The mirror in a reflecting telescope is constructed so that all light rays arriving parallel to its axis are reflected to pass through the focus (Figure 5.3b). In astronomical contexts, the mirror that collects the incoming light is usually called the *primary mirror*, because telescopes often contain more than one mirror. The focus of the primary mirror is referred to as the **prime focus**.

Astronomical telescopes are often used to make **images** of their field of view (simply, the portion of the sky that the



**▲ FIGURE 5.2 Refracting Lens** (a) Refraction by a prism changes the direction of a light ray by an amount that depends on the angle between the prism's faces. When the angle between the faces is large, the deflection is large; when the angle is small, so is the deflection. (b) A lens can be thought of as a series of prisms.

telescope “sees”). Figure 5.4 illustrates how that is accomplished, in this case by the mirror in a reflecting telescope. Light from a distant object (here, a comet) reaches us as parallel, or very nearly parallel, rays. Any ray of light entering the instrument parallel to the telescope's axis strikes the mirror and is reflected through the prime focus. Light coming from a slightly different direction—inclined slightly to the axis—is focused to a slightly different point. In this way, an image is formed near the prime focus. Each point on the image corresponds to a different point in the field of view.


The prime-focus images produced by large telescopes are actually quite small—the image of the entire field of view may be as little as 1 cm across. Often, the image is magnified with a lens known as an *eyepiece* before being observed by eye or, more likely, recorded as a photograph or digital image. The angular diameter of the magnified image is much greater than the telescope's field of view, allowing much more detail to be discerned. Figure 5.5(a) shows the basic design of a simple

refracting telescope, illustrating how a small eyepiece is used to view the image focused by the lens. Figure 5.5(b) shows how a reflecting telescope accomplishes the same function.


## Comparing Refractors and Reflectors

The two telescope designs shown in Figure 5.5 achieve the same result: Light from a distant object is captured and focused to form an image. On the face of it, then, it might appear that there is little to choose between the two in deciding which type to buy or build. However, as the sizes of telescopes have increased steadily over the years (for reasons to be discussed in Section 5.3), a number of important factors have tended to favor reflecting instruments over refractors:

1. The fact that light must pass through the lens is a major disadvantage of refracting telescopes. Just as a prism disperses white light into its component colors, the lens



**◀ FIGURE 5.3 Reflecting Mirror** (a) Reflection of light from a flat mirror occurs when light is deflected, depending on its angle of incidence. (b) Curved mirrors focus to a single point all rays of light arriving parallel to the mirror axis. The arrows indicate the directions of the incoming and reflected rays.




**▲ FIGURE 5.4 Image Formation** An image is formed by a mirror as rays of light coming from different points on a distant object focus to slightly different locations. Notice that the image is inverted (i.e., upside down).

in a refracting telescope tends to focus red and blue light differently. This deficiency is known as *chromatic aberration*. Careful design and choice of materials can largely correct this deficiency, but it is very difficult to eliminate entirely. Obviously, this problem does not occur with mirrors.


- As light passes through the lens, some of it is absorbed by the glass. This absorption is a relatively minor problem for visible radiation, but it can be severe for infrared and ultraviolet observations because glass blocks most of the radiation in those regions of the electromagnetic spectrum. Again, this problem does not affect mirrors.

- A large lens can be quite heavy. Because it can be supported only around its edge (so as not to block the incoming radiation), the lens tends to deform under its own weight. A mirror does not have this drawback because it can be supported over its entire back surface.
- A lens has two surfaces that must be accurately machined and polished—a task that can be very difficult indeed—but a mirror has only one.



For these reasons, *all* large modern telescopes use mirrors as their primary light gatherers. The largest refractor ever built, installed in 1897 at the Yerkes Observatory in Wisconsin and still in use today, has a lens diameter of just over 1 m (40 inches). By contrast,

**◀ FIGURE 5.5 Refractors and Reflectors** Comparison of (a) refracting and (b) reflecting telescopes. Both types are used to gather and focus electromagnetic radiation—to be observed by human eyes or recorded on photographs or in computers. In both cases, the image formed at the focus is viewed with a small magnifying lens called an eyepiece.



**FIGURE 5.6 Reflecting Telescopes** Four reflecting telescope designs: (a) prime focus, (b) Newtonian focus, (c) Cassegrain focus, and (d) Nasmyth/coudé focus. Each design uses a primary mirror at the bottom of the telescope to capture radiation, which is then directed along different paths for analysis. Notice that the secondary mirrors shown in (c) and (d) are actually slightly diverging, so that they move the focus outside the telescope.

many recently constructed reflecting telescopes have mirror diameters in the 10-m range, and still larger instruments are on the way.

## Types of Reflecting Telescope

Figure 5.6 shows some basic reflecting telescope designs. Radiation from a star enters the instrument, passes down the main tube, strikes the primary mirror, and is reflected back toward the prime focus, near the top of the tube. Sometimes astronomers place their recording instruments at the prime focus; however, it can be inconvenient, or even impossible, to suspend bulky pieces of equipment there. More often, the light is intercepted on its path to the focus by a *secondary mirror* and redirected to a more convenient location, as in Figure 5.6(b) through 5.6(d).

In a **Newtonian telescope** (named after Sir Isaac Newton, who invented this particular design), the light is intercepted before it reaches the prime focus and then is deflected by  $90^\circ$ , usually to an eyepiece at the side of the instrument. This is a popular design for smaller reflecting telescopes, such as those used by amateur astronomers, but it is relatively uncommon in large instruments. On a large telescope, the Newtonian focus may be many meters above the ground, making it an inconvenient place to attach equipment (or place an observer).


Alternatively, astronomers may choose to work on a rear platform where they can use equipment, such as a spectroscopy, that is too heavy to hoist to the prime focus. In this case, light reflected by the primary mirror toward the prime focus is intercepted by a smaller secondary mirror, which reflects it back down through a small hole at the center of the

primary mirror. This arrangement is known as a **Cassegrain telescope** (after Guillaume Cassegrain, a French lensmaker). The point behind the primary mirror where the light from the star finally converges is called the *Cassegrain focus*.

A more complex observational configuration requires starlight to be reflected by several mirrors. As in the Cassegrain design, light is first reflected by the primary mirror toward the prime focus and is then reflected back down the tube by a secondary mirror. Next, a third, much smaller, mirror reflects the light out of the telescope, where (depending on the details of the telescope's construction) the beam may be analyzed by a detector mounted alongside, at the *Nasmyth focus*, or it may be directed via a further series of mirrors into an environmentally controlled laboratory known as the *coudé room* (from the French word for “bent”). This laboratory is separate from the telescope itself, enabling astronomers to use very heavy and finely tuned equipment that cannot be placed at any of the other foci (all of which necessarily move with the telescope). The arrangement of mirrors is such that the light path to the coudé room does not change as the telescope tracks objects across the sky.

To illustrate some of these points, Figure 5.7(a) shows the twin 10-m-diameter optical/infrared telescopes of the Keck Observatory on Mauna Kea in Hawaii, operated jointly by the California Institute of Technology and the University of California. The diagram in part (b) illustrates the light paths and some of the foci. Observations may be made at the Cassegrain, Nasmyth, or coudé focus, depending on the needs of the user. As the size of the person in part (c) indicates, this is indeed a very large telescope—in fact, the two mirrors are among the largest on Earth. We will see numerous examples throughout this text of Keck's many important discoveries.






**▲ FIGURE 5.7 Keck Telescope** (a) The two 10-m telescopes of the Keck Observatory. (b) Artist's illustration of the telescope, the path taken by an incoming beam of starlight, and some of the locations where instruments may be placed. (c) One of the 10-m mirrors. (The odd shape is explained in Section 5.3.) Note the technician in orange coveralls at center. (*W. M. Keck Observatory*)

Perhaps the best-known telescope on (or near) Earth is the *Hubble Space Telescope (HST)*, named for one of America's most notable astronomers, Edwin Hubble. Placed in Earth orbit by NASA's space shuttle *Discovery* in 1990 and (as of 2013) still in operation, *HST* is a Cassegrain telescope in which all the instruments are located directly behind the 2.4-m primary mirror, as sketched in Figure 5.8(a). The telescope's detectors are capable of making measurements in the optical, infrared, and ultraviolet parts of the spectrum, from about 100 nm (UV) to 2200 nm (IR).

Soon after launch, astronomers discovered that the telescope's primary mirror had been polished to the wrong shape and was unable to focus light as accurately as expected. In 1993, in the first and most important of

five servicing missions carried out during the instrument's lifetime, astronauts aboard the space shuttle *Endeavour* visited *HST* and corrected the defect by installing an intricate set of small mirrors (each about the size of a coin) in the light path behind the primary mirror to compensate for its faulty construction. *Hubble's* sensitivity and resolution are now close to the original design specifications.

During two decades of operation, *Hubble* has revolutionized our view of the sky and helped rewrite more than one theory of the universe along the way. Figure 5.8(b) illustrates the telescope's improved image quality by comparing ground-based and *Hubble* images of the spiral galaxy M101. Many additional spectacular examples of the telescope's remarkable capabilities appear throughout this book.



(a)

**▲ FIGURE 5.8 Hubble Space Telescope** (a) This “see-through” diagram displays some of *HST*’s hardware surrounding its main mirror (in light blue). (b) These two images compare the majestic spiral galaxy M101 as observed with the large Mayall telescope on Kitt Peak Mountain (bottom) and with the *Hubble* telescope in orbit (top). (D. Berry; AURA/NASA)



(b)

### CONCEPT Check

- ✓ Why do all modern telescopes use mirrors to gather and focus light?

## 5.2 Telescope Size

Modern astronomical telescopes have come a long way from Galileo’s simple apparatus (see the Part 1 opening text). Their development over the years has seen a steady increase in *size*, for two main reasons. The first has to do with the amount of light a telescope can collect—its *light-gathering power*. The second is related to the amount of detail that can be seen—the telescope’s *resolving power*. Simply put, large telescopes can gather and focus more radiation than can their smaller counterparts, allowing astronomers to study fainter objects and to obtain more detailed information about bright ones. This fact has played a central role in determining the design of contemporary instruments.

### Light-Gathering Power

One important reason for using a larger telescope is simply that it has a greater **collecting area**, which is the total area capable of gathering radiation. The larger the telescope’s reflecting mirror (or refracting lens), the more light it collects, and the easier it is to measure and study an object’s radiative properties. Astronomers spend much of their time observing

very distant—and hence very *faint*—cosmic sources. In order to make detailed observations of such objects, very large telescopes are essential. Figure 5.9 illustrates the effect of increasing the size of a telescope by comparing images of the Andromeda Galaxy taken with two different instruments. A large collecting area is particularly important for spectroscopic work, as the radiation received in that case must be split into its component wavelengths for further analysis.

The observed brightness of an astronomical object is directly proportional to the area of our telescope’s mirror and therefore to the *square* of the mirror diameter. Thus, a 5-m telescope will produce an image 25 times as bright as a 1-m instrument, because a 5-m mirror has  $5^2 = 25$  times the collecting area of a 1-m mirror. We can also think of this relationship in terms of the length of *time* required for a telescope to collect enough energy to create a recognizable image on a photographic plate. Our 5-m telescope will produce an image 25 times faster than the 1-m device because it gathers energy at a rate 25 times greater. Put another way, a 1-hour exposure with a 1-m telescope is roughly equivalent to a 2.4-minute exposure with a 5-m instrument.

Until the 1980s, the conventional wisdom was that telescopes with mirrors larger than 5 or 6 m in diameter were simply too expensive and impractical to build. The problems involved in casting, cooling, and polishing a huge block of quartz or glass to very high precision (typically less than the width of a human hair) were just too great. However, new, high-tech manufacturing techniques, coupled with radically



(a)



(b)

▲ **FIGURE 5.9 Sensitivity** Telescope size affects the image of a cosmic source, in this case the Andromeda Galaxy. Both photographs had the same exposure time, but image (b) was taken with a telescope twice the size of that used to make image (a). Fainter detail can be seen as the diameter of the telescope mirror increases because larger telescopes are able to collect more photons per unit time, greatly extending our view of the universe. (Adapted from AURA)

new mirror designs, make the construction of telescopes in the 8- to 12-m range almost a routine matter. Experts can now make large mirrors much lighter for their size than was once thought feasible and can combine many smaller mirrors into the equivalent of a much larger single-mirror telescope.

The Keck telescopes, shown in detail in Figure 5.7 and in a larger view in Figure 5.10, are a case in point. Each telescope combines 36 hexagonal 1.8-m mirrors into the equivalent collecting area of a single 10-m reflector. The first Keck

telescope became fully operational in 1992; the second was completed in 1996. The large size of these devices and the high altitude at which they operate make them particularly well suited to detailed spectroscopic studies of very faint objects, in both the optical and infrared parts of the spectrum. Mauna Kea's 4.2-km (13,800 feet) altitude minimizes atmospheric absorption of infrared radiation, making this site one of the finest locations on Earth for infrared astronomy.


Numerous other large telescopes can be seen in the figure. Some are designed exclusively for infrared work; others, like Keck, operate in both the optical and the infrared. To the right of the Keck domes is the 8.3-m Subaru (the Japanese name for the Pleiades) telescope, part of the National Astronomical Observatory of Japan. Its mirror, shown in Figure 5.10(b), is the largest single mirror (as opposed to the segmented design used in Keck) yet built. Subaru saw “first light” in 1999. In the distance is another large single-mirror instrument: the 8.1-m Gemini North telescope, completed in 1999 by a consortium of seven nations, including the United States. Its twin, Gemini South, in the Chilean Andes, went into service in 2002.

In terms of total available collecting area, the largest telescope currently in operation is the European Southern Observatory's optical-infrared Very Large Telescope (VLT), located at Cerro Paranal, in Chile (Figure 5.11). The VLT consists of four separate 8.2-m mirrors that can function as a single instrument. The last of its four mirrors was completed in 2001.

## Resolving Power

A second advantage of large telescopes is their finer **angular resolution**. In general, *resolution* refers to the ability of any device, such as a camera or telescope, to form distinct, separate images of objects lying close together in the field of view. The finer the resolution, the better we can distinguish the objects and the more detail we can see. In astronomy, where we are always concerned with angular measurement, “close together” means “separated by a small angle in the sky,” so angular resolution is the factor that determines our ability to see fine structure. Figure 5.12 illustrates how the appearance of two objects—stars, say—might change as the angular resolution of our telescope varies. Figure 5.13 shows the result of increasing resolving power with views of the Andromeda Galaxy at several different resolutions.

What limits a telescope's resolution? One important factor is *diffraction*, the tendency of light—and all other waves, for that matter—to bend around corners. ∞ (Discovery 3-1) Because of diffraction, when a parallel beam of light enters a telescope, the rays spread out slightly, making it impossible to focus the beam to a sharp point, even with a perfectly constructed mirror. Diffraction introduces a certain “fuzziness,” or loss of resolution, into any optical system. The degree of fuzziness—the minimum angular separation that can be distinguished—determines



**▲ FIGURE 5.10 Mauna Kea Observatory** (a) The world's highest ground-based observatory, at Mauna Kea, Hawaii, is perched atop a dormant volcano more than 4 km (nearly 14,000 feet) above sea level. Among the domes visible in the picture are those housing the Canada–France–Hawaii 3.6-m telescope, the 8.1-m Gemini North instrument, the 2.2-m telescope of the University of Hawaii, Britain's 3.8-m infrared facility, and the twin 10-m Keck telescopes. To the right of the twin Kecks is the Japanese 8.3-m Subaru telescope. The thin air at this high-altitude site guarantees less atmospheric absorption of incoming radiation and hence a clearer view than at sea level, but the air is so thin that astronomers must occasionally wear oxygen masks while working. (b) The mirror in the Subaru telescope. (R. Wainscoat; NAOJ)




**▲ FIGURE 5.11 VLT Observatory** Located at the Paranal Observatory in Atacama, Chile, the European Southern Observatory's Very Large Telescope (VLT) is currently the world's largest optical telescope. Four 8.2-m reflecting telescopes are used in tandem to create the effective area of a single 16-m mirror. (ESO)

the angular resolution of the telescope. The amount of diffraction is proportional to the wavelength of the radiation and inversely proportional to the diameter of the telescope mirror. For a circular mirror and otherwise perfect optics, we can write (in convenient units):

$$\text{angular resolution (arcsec)} = 0.25 \frac{\text{wavelength } (\mu\text{m})}{\text{diameter (m)}}$$

where  $1 \mu\text{m}$  (1 micron) =  $10^{-6}\text{m}$  (see Appendix 2).

For a given telescope size, the amount of diffraction increases in proportion to the wavelength used, and observations in the infrared or radio range are often limited by its effects. For example, the best possible angular resolution of blue light (with a wavelength of 400 nm) that can be obtained using a 1-m telescope is about  $0.1''$ . This quantity is called the **diffraction-limited resolution** of the telescope. But if we were to use our 1-m telescope to make observations in the near infrared, at a wavelength of  $10 \mu\text{m}$  (10,000 nm), the best resolution we could obtain would be only  $2.5''$ . A 1-m radio telescope operating at a wavelength of 1 cm would have an angular resolution of just under  $1^\circ$ .




**Interactive** **FIGURE 5.12 Resolving Power** Two comparably bright light sources become progressively clearer when viewed at finer and finer angular resolution. When the angular resolution is much poorer than the separation of the objects, as in (a), the objects appear as a single fuzzy “blob.” As the resolution improves, through (b) and (c), the two sources become discernible as separate objects.

For light of any given wavelength, large telescopes produce less diffraction than small ones. A 5-m telescope observing in blue light would have a diffraction-limited resolution five times finer than the 1-m telescope just discussed—about  $0.02''$ . A 0.1-m (10-cm) telescope would have a diffraction limit of  $1''$ , and so on. For comparison, the angular resolution of the human eye in the middle of the visual range is about  $0.5'$ .


### PROCESS OF SCIENCE Check

- ✓ Give two reasons why astronomers need to build very large telescopes.

► **FIGURE 5.13 Resolution** Detail becomes clearer in the Andromeda Galaxy as the angular resolution improves some 600 times, from (a)  $10'$ , to (b)  $1'$ , (c)  $5''$ , and (d)  $1''$ . The resolution of the human eye is approximately that of part (b)—if only our eyes were sensitive enough to see this view. (Adapted from AURA)




(a)




(b)



(c)



(d)






## Image Processing

Computers are also widely used to reduce *background noise* in astronomical images. Noise is anything that corrupts the integrity of a message, such as static on an AM radio or “snow” on a television screen. The noise corrupting telescopic images has many causes. In part, it results from faint, unresolved sources in the telescope’s field of view and from light scattered into the line of sight by Earth’s atmosphere. It can also be caused by imperfections within the detector itself, which may result in an electronic “hiss” similar to the faint background hiss you might hear when you listen to a particularly quiet piece of music on your stereo.

Even though astronomers often cannot determine the origin of the noise in their observations, they can at least measure its characteristics. For example, if we observe a part of the sky where there are no known sources of radiation, then whatever signal we do receive is (almost by definition) noise. Once the properties of the signal have been measured, the effects of noise can be partially removed with the aid of high-speed computers, allowing astronomers to see features in their data that would otherwise remain hidden.

Using computer processing, astronomers can also compensate for known instrumental defects. In addition, the computer can often carry out many of the relatively simple, but tedious and time-consuming, chores that must be performed before an image (or spectrum) reaches its final “clean” form. Figure 5.15 illustrates how computerized image-processing techniques were used to correct for known instrumental problems in *HST*, allowing much of the planned resolution of the telescope to be recovered even before its repair in 1993 (see Section 5.1).




**▲ FIGURE 5.15 Image Processing** (a) Ground-based view of the star cluster R136, a group of stars in the Large Magellanic Cloud (a nearby galaxy). (b) The “raw” image of this same region as seen by the *Hubble Space Telescope* in 1990, before its first repair mission. (c) The same image after computer processing that partly compensated for imperfections in the mirror. (d) The same region as seen by the repaired *HST* in 1994, here observed at a somewhat shorter (bluer) wavelength. (AURA/NASA)

## Photometry

When a CCD is placed at the focus of a telescope to record an image of the instrument’s field of view, the telescope is acting, in effect, as a high-powered camera. However, astronomers often want to carry out more specific measurements of the radiation received from space.

One very fundamental property of a star (or any other astronomical object) is its *brightness*—the amount of light energy from the star striking our detector every second. The measurement of brightness is called **photometry** (literally, “light measurement”). In principle, determining a star’s brightness is just a matter of adding up the values in all the CCD pixels corresponding to the star (see Figure 5.14c). However, in practice, the process is more complicated, as stellar images may overlap, and computer assistance is generally needed to disentangle them.

Astronomers often combine photometric measurements with the use of colored *filters* in order to limit the wavelengths they measure. (A filter simply blocks out all incoming radiation, except in some specific range of wavelengths; see Section 17.3 for a more detailed discussion.) Many standard filters exist, covering various “slices” of the spectrum, from near-infrared through visible to near-ultraviolet wavelengths. By confining their attention to these relatively narrow ranges, astronomers can often estimate the shape of an object’s black-body curve and hence determine, at least approximately, the object’s temperature.  (Sec. 3.4) Filters are also used with CCD images in order to simulate natural color. For example, most of the visible-light *HST* images in this text are actually composites of *three* raw images, taken through red, green, and blue filters, respectively, and combined afterwards to reconstruct a single color frame.

Astronomical objects are generally faint, and most astronomical images entail long exposures—minutes to hours—in

order to see fine detail (Section 5.2). Thus, the brightness we measure from an image is really an average over the entire exposure. Short-term fluctuations (if any) cannot be seen. When highly accurate and rapid measurements of light intensity are required, a specialized device known as a **photometer** is used. The photometer measures the total amount of light received in all or part of the field of view. When only a portion of the field is of interest, that region is selected simply by masking (blocking) out the rest of the field of view. Using a photometer often means “throwing away” spatial detail—usually no image is produced—but in return, more information is obtained about the intensity and time variability of a source, such as a pulsating star or a supernova explosion.

## Spectroscopy

Often, astronomers want to study the *spectrum* of the incoming light. Large **spectrometers** work in tandem with optical telescopes. Light collected by the primary mirror may be redirected to the coudé room, defined by a narrow slit, dispersed (split into its component colors) by means of a prism or a diffraction grating, and then sent on to a detector—a process not so different in concept from the operation of the simple spectroscopy described in Chapter 4. ∞ (Sec. 4.1) The spectrum can be studied in real time (i.e., as it is being received at the telescope) or recorded using a CCD (or, less commonly nowadays, a photographic plate) for later analysis. Astronomers can then apply the analysis techniques discussed in Chapter 4 to extract detailed information from the spectral lines they record. ∞ (Sec. 4.5)

### CONCEPT Check

✓ Why aren't astronomers satisfied with just taking photographs of the sky?

## 5.4 High-Resolution Astronomy

Even large telescopes have limitations. For example, according to the discussion in the preceding section, the 10-m Keck telescope should have an angular resolution of around  $0.01''$  in blue light. In practice, however, without further technological advances, it could not do better than about  $1''$ . In fact, apart from instruments using special techniques developed to examine some particularly bright stars, *no* ground-based optical telescope built before 1990 can resolve astronomical objects to much better than  $1''$ . The reason is *turbulence* in Earth's atmosphere—small-scale eddies of swirling air all along the line of sight, which blur the image of a star even before the light reaches our instruments.

### Atmospheric Blurring

As we observe a star, atmospheric turbulence produces continual small changes in the optical properties of the air

between the star and our telescope (or eye). As a result, the light from the star is refracted slightly as it travels toward us, so the stellar image dances around on our detector (or on our retina). This continual deflection is the cause of the well-known “twinkling” of stars. It occurs for the same reason that objects appear to shimmer when viewed across a hot roadway on a summer day: The constantly shifting rays of light reaching our eyes produce the illusion of motion.

On a good night at the best observing sites, the maximum amount of deflection produced by the atmosphere is slightly less than  $1''$ . Consider taking a photograph of a star. After a few minutes of exposure (long enough for the intervening atmosphere to have undergone many small random changes), the image of the star has been smeared out over a roughly circular region an arc second or so in diameter. Astronomers use the term **seeing** to describe the effects of atmospheric turbulence. The circle over which a star's light (or the light from any other astronomical source) is spread is called the **seeing disk**. Figure 5.16 illustrates the formation of the seeing disk for a small telescope.\*

Atmospheric turbulence has less effect on light of longer wavelengths—ground-based astronomers generally “see” better in the infrared. However, offsetting this improvement in image quality is the fact that the atmosphere is wholly or partially opaque over much of the infrared range. ∞ (Sec. 3.3) For these reasons, to achieve the best possible observing conditions, telescopes are sited on mountaintops (to get above as much of the atmosphere as possible) in regions of the world where the atmosphere is known to be fairly stable and relatively free of dust and moisture. Another reason for the choice of remote locations is the growing problem of **light pollution** in populated areas—unwanted upward-directed light from streets, parking lots, homes, and businesses, that scatters back from dust in the air into our telescopes, literally drowning out the faint signals from distant stars and galaxies that astronomers want to observe.


In the continental United States, these sites tend to be in the desert Southwest. The U.S. National Observatory for optical astronomy in the Northern Hemisphere, completed in 1973, is located high on Kitt Peak near Tucson, Arizona. The site was chosen because of its many dry, clear nights. Seeing less than  $1''$  from such a location is regarded as good, and seeing a few arc seconds is tolerable for many purposes. Even better conditions are found on Mauna Kea, Hawaii (Figure 5.10), and at numerous sites in the Andes Mountains of Chile (Figures 5.11 and 5.17), which is why many large telescopes have recently been constructed at these exceptionally clear locations.

\*In fact, for a large instrument—more than about 1 m in diameter—the situation is more complicated, because rays striking different parts of the mirror have actually passed through different turbulent regions of the atmosphere. The end result is still a seeing disk, however.



► **FIGURE 5.16 Atmospheric**

**Turbulence** Light rays from a distant star strike a telescope detector at slightly different locations because of turbulence in Earth's atmosphere. Over time, the light covers a roughly circular region on the detector, and even the pointlike image of a star is recorded as a small disk, called the *seeing disk*.



An optical telescope placed in orbit about Earth or on the Moon could obviously overcome the limitations imposed by the atmosphere on ground-based instruments. Without atmospheric blurring, extremely fine resolution—close to the diffraction limit—can be achieved, subject only to the engineering restrictions associated with building or placing large structures in space. The 2.4-m mirror in the *Hubble Space Telescope* has a (blue-light) diffraction limit of only 0.05", giving astronomers a view of the universe as much as 20 times sharper than that normally available from even much larger ground-based instruments.

## Active Optics

Current techniques for producing ultrasharp images take the ideas of computer control and image processing (see Section 5.2) several stages further. By analyzing the image formed by a telescope *while the light is still being collected*, it is possible to adjust the telescope from moment to moment to avoid or compensate for the effects of mirror distortion, temperature changes in the dome, and even atmospheric turbulence.

Even under conditions of perfect seeing, most telescopes would not achieve diffraction-limited resolution. The temperature of the mirror or in the dome may fluctuate slightly during the many minutes or even hours required for the image to be exposed, and the precise shape of the mirror may change slightly as the telescope tracks a source across the sky. The effect of these changes is that the mirror's focus may shift from minute to minute, blurring the eventual image in much the same way as atmospheric turbulence creates a seeing disk (Figure 5.16). At the best observing sites, the seeing is often so good that these tiny effects may be the main cause of image blurring. The collection of techniques aimed at controlling such environmental and mechanical fluctuations is known as **active optics**.

The first telescope designed to incorporate active optics was the New Technology Telescope (NTT), constructed

in 1989 at the European Southern Observatory in Chile and upgraded in 1997. (NTT is the most prominent instrument visible in Figure 5.17.) This 3.5-m instrument, employing the latest in real-time telescope controls, can achieve a resolution as sharp as 0.2" by making minute modifications to the tilt of its mirror as its temperature and orientation change, thus maintaining the best possible focus at all times. Figure 5.18 shows how active optics can dramatically improve the resolution of an image. Active-optics techniques now include improved dome design to control airflow, precise control of the mirror temperature, and the use of pistons behind the mirror to maintain its precise shape. All of the large telescopes described earlier include active optics systems, improving their resolution to a few tenths of an arc second.

## Real-Time Control

With active optics systems in place, Earth's atmosphere once again becomes the main agent limiting a telescope's resolution. Remarkably, even this problem can now be addressed, using an approach known as **adaptive optics**, a technique that actually deforms the shape of a mirror's surface under computer control, while the image is being exposed, in order to undo the effects of atmospheric turbulence. The mirror in question is generally not the large primary mirror of the telescope. Rather, for both economic and technical reasons, a much smaller (typically 20- to 50-cm-diameter) mirror is inserted into the light path and manipulated to achieve the desired effect.

Adaptive optics presents formidable theoretical and practical problems, but the rewards are so great that it has been the subject of intense research since the 1970s. The effort received an enormous boost in the 1990s from declassified military technology from the Strategic




▲ **FIGURE 5.17 European Southern Observatory** Located in the Andes Mountains of Chile, the European Southern Observatory at La Silla is run by a consortium of European nations. Numerous domes house optical telescopes of different sizes, each with varied support equipment, making this one of the most versatile observatories south of the equator. The largest telescope at La Silla—the square building to the right of center—is the New Technology Telescope, a 3.5-m active optics device. (ESO)

Defensive Initiative, a Reagan-era missile defense program (dubbed “Star Wars” by its detractors) intended to target and shoot down incoming ballistic missiles. In the system shown in Figure 5.19, a laser probes the atmosphere above the telescope, creating an “artificial star” that allows astronomers to gauge atmospheric conditions and pass that information to a computer that modifies the telescope mirror thousands of times per second to compensate for poor seeing.

Adaptive corrections are somewhat easier to apply in the infrared than in the optical, because atmospheric distortions are smaller (stars “twinkle” less in the infrared) and because the longer infrared wavelengths impose less stringent requirements on the precise shape of the mirror. Infrared adaptive optics systems already exist in many large telescopes. For example, Gemini and Subaru have reported adaptive optics resolutions of around  $0.06''$  in the near

infrared—not quite at the diffraction limit ( $0.03''$  for an 8-m telescope at  $1\ \mu\text{m}$ , according to the earlier equation), but already better than the resolution of *HST* at the same wavelengths (see Figure 5.20a). Both Keck and the VLT incorporate adaptive optics instrumentation capable of producing diffraction-limited images at near-infrared wavelengths.


Visible-light adaptive optics has been demonstrated experimentally, and some astronomical telescopes are beginning to incorporate the technology. Figure 5.20(b) compares a pair of visible-light observations of a nearby double star called Castor. The observations were made with a relatively modest 1.5-m telescope. The adaptive optics system clearly distinguishes the two stars. Remarkably, adaptive optics techniques are giving astronomers the “best of both worlds,” achieving with large ground-based optical telescopes the kind of resolution once attainable only from space.



◀ **FIGURE 5.18 Active Optics** These false-color infrared photographs of part of the star cluster R136—the same object shown in Figure 5.15—contrast the resolution obtained (a) without and (b) with an active optics system. Both images were taken with the New Technology Telescope shown in Figure 5.17. (ESO)



◀ **FIGURE 5.19 Adaptive Optics System** In this daytime photo, a test is being conducted at the Lick Observatory 3-m Shane telescope in California. A laser is used to create an “artificial star” (light reflected from the atmosphere back into the telescope) to improve guiding. The laser beam probes the atmosphere above the telescope, allowing tiny computer-controlled changes to be made in the shape of the mirror surface thousands of times each second. (*Lick Observatory*)



▲ **FIGURE 5.20 Adaptive Optics in Action** (a) The star cluster NGC 6934 is resolved to a little less than 1" in this uncorrected visible-light image (left) from the 8-m Gemini North telescope in Hawaii. With adaptive optics applied (right), the resolution in the infrared is improved by nearly a factor of 10, allowing more stars to be seen much more clearly. (b) These visible-light images of the double star Castor were acquired at a military observatory on Mount Haleakala in Maui, Hawaii. The uncorrected image (left) is blurred over several arc seconds, giving only a hint of its binary nature. With adaptive optics (right), the resolution is improved to 0.1", and the two stars are clearly separated. (*NOAO; MIT Lincoln Laboratory*)

### CONCEPT Check

- ✓ What steps do optical astronomers take to overcome the obscuring and blurring effects of Earth's atmosphere?

## 5.5 Radio Astronomy

In addition to the visible radiation that penetrates Earth's atmosphere on a clear day, radio radiation also reaches the ground. Indeed, as indicated in Figure 3.8, the radio window in the electromagnetic spectrum is much wider than the optical window. ∞ (Sec. 3.3) Because the atmosphere is no hindrance to long-wavelength radiation, radio astronomers have built many ground-based **radio telescopes** capable of detecting radio waves reaching us from space. These devices have all been constructed since the 1950s—radio astronomy is a much younger subject than optical astronomy.

### Early Observations

The field of radio astronomy originated with the work of Karl Jansky at Bell Labs in 1931. Jansky was studying the causes of shortwave-radio interference when he discovered a faint static “hiss” that had no apparent terrestrial (Earthly) source. He noticed that the strength of the hiss varied in time and that its peak occurred about 4 minutes earlier each day. He soon realized that the peaks were coming exactly one *sidereal day* apart, and he concluded that the hiss was indeed not of terrestrial origin, but came from a definite direction in space. ∞ (Sec. 1.4) That direction is now known to correspond to the center of our Galaxy.

Some astronomers were intrigued by Jansky's discovery, but with the limited technology of the day—and even

more limited Depression-era budgets—progress was slow. Jansky himself was moved to another project at Bell Labs and never returned to astronomical studies. However, by 1940, the first systematic surveys of the radio sky were under way. After a series of technological breakthroughs made during World War II, these studies rapidly grew into a distinct branch of astronomy.

During the 1930s, astronomers became aware that the space between the stars in our Galaxy is not empty, but instead is filled with extremely diffuse (low-density) gas (see Chapter 18). The growing realization in the 1940s that this otherwise completely invisible part of the Galaxy could be observed and mapped in detail at radio wavelengths established the true importance of Jansky's pioneering work. Today he is regarded as the father of radio astronomy.

## Essentials of Radio Telescopes

Figure 5.21(a) shows the world's largest steerable radio telescope: the large 105-m-diameter (340-foot-diameter) telescope located at the National Radio Astronomy Observatory in West Virginia. Although much larger than reflecting optical telescopes, most radio telescopes are built in basically the same way. They have a large, horse-shoe-shaped mount supporting a huge curved metal dish


that serves as the collecting area. As illustrated in Figure 5.21(b), the dish captures incoming radio waves and reflects them to the focus, where a receiver detects the signals and channels them to a computer for storage and analysis.

Conceptually, the operation of a radio telescope is similar to the operation of an optical reflector with the detecting instruments placed at the prime focus (Figure 5.6a). However, unlike optical instruments, which can detect all visible wavelengths simultaneously, radio detectors normally register only a narrow band of wavelengths at any one time. To observe radiation with a different frequency, we must retune the equipment, much as we tune a radio or television set to a different channel.

Radio telescopes must be built large partly because cosmic radio sources are extremely faint. In fact, the total amount of radio energy received by Earth's entire surface is less than a trillionth of a watt. Compare this with the roughly 10 million watts our planet's surface receives in the form of infrared and visible light from any of the bright stars seen in the night sky. To capture enough radio energy to allow detailed measurements to be made, a large collecting area is essential. Figure 5.22 shows an even larger, but unmovable, radio telescope strung among the hills of Arecibo, Puerto Rico. Constructed in 1963 in a natural depression in the hillside, the Arecibo telescope is




(a)



(b)

▲ **FIGURE 5.21 Radio Telescope** (a) The 105-m-diameter device at the National Radio Astronomy Observatory in Green Bank, West Virginia, is 150 m tall—taller than the Statue of Liberty and nearly as tall as the Washington Monument. (b) Schematic diagram of the telescope shows the path taken by an incoming beam of radio radiation (colored blue). (NRAO)



▲ **FIGURE 5.22 Arecibo Observatory** An aerial photograph of the 300-m-diameter dish at the National Astronomy and Ionospheric Center near Arecibo, Puerto Rico. The left inset shows a close-up of the radio receivers hanging high above the dish. The right inset shows technicians adjusting the dish surface to make it smoother. (D. Parker/T. Acevedo/NAIC; Cornell)

approximately 300 m (1000 feet) in diameter. Its reflecting surface spans nearly 20 acres.

The angular resolution of radio telescopes is generally quite poor compared with that of their optical counterparts because of the effects of diffraction. Typical wavelengths of radio waves are about a million times longer than those of visible light, and these longer wavelengths impose a corresponding crudeness in angular resolution. (Recall from Section 5.2 that the longer the wavelength, the greater the amount of diffraction.) Even the enormous sizes of radio dishes only partly offset this effect. The radio telescope shown in Figure 5.21 can achieve a resolution of about 1' when receiving radio waves having wavelengths of around 3 cm. However, it was designed to operate most efficiently (i.e., it is most sensitive to radio signals) at wavelengths closer to 1 cm, where the resolution is approximately 20". The best angular resolution obtainable with a single radio telescope is about 10" (for the largest instruments operating at millimeter wavelengths)—at

least 100 times coarser than the capabilities of some large optical systems.

Radio telescopes can be built so much larger than their optical counterparts because their reflecting surfaces need not be as smooth as is necessary for light waves of shorter wavelength. Provided that surface irregularities (dents, bumps, and the like) are much smaller than the wavelength of the waves to be detected, the surface will reflect them without distortion. Because the wavelength of visible radiation is short (less than  $10^{-6}$  m), extremely smooth mirrors are needed to reflect the waves properly, and it is difficult to construct very large mirrors to such exacting tolerances. However, even rough metal surfaces can focus 1-cm waves accurately, and radio waves of wavelength a meter or more can be reflected and focused perfectly well by surfaces having irregularities even as large as your fist. The Arecibo instrument was originally surfaced with chicken wire, which was lightweight and cheap. Although fairly rough, the chicken wire was adequate for proper reflection because

the openings between adjacent strands of wire were much smaller than the long-wavelength radio waves that were to be detected.


The entire Arecibo dish was resurfaced with thin metal panels in 1974 and was further upgraded in 1997 so that it can now be used to study radio radiation of shorter wavelength. Since the 1997 upgrade, the panels can be adjusted to maintain a precise spherical shape to an accuracy of about 3 mm over the entire surface. At a frequency of 5 GHz (corresponding to a wavelength of 6 cm—the shortest wavelength that can be studied, given the properties of the dish surface), the telescope’s angular resolution is about  $1'$ . The huge size of the dish creates one distinct disadvantage, however: The Arecibo telescope cannot be pointed very well to follow cosmic objects across the sky. The detectors can move roughly  $10^\circ$  on either side of the focus, restricting the telescope’s observations to those objects that happen to pass within about  $20^\circ$  of overhead as Earth rotates.

Arecibo is an example of a rough-surfaced telescope capable of detecting long-wavelength radio radiation. At the other extreme, Figure 5.23 shows the 36-m-diameter Haystack dish in northeastern Massachusetts. Constructed of polished aluminum, this telescope maintains a parabolic curve to an accuracy of about a millimeter all the way across its solid surface. It can reflect and accurately focus radio radiation with wavelengths as short as a few millimeters. The telescope is contained within a protective shell, or radome, that protects the surface from the harsh New England weather. The radome acts much like the protective dome of an optical telescope, except that there is no slit through which the telescope “sees.” Incoming cosmic radio signals pass virtually unimpeded through the radome’s fiberglass construction.

## The Value of Radio Astronomy


Despite the inherent disadvantage of relatively poor angular resolution, radio astronomy enjoys many advantages. Radio telescopes can observe 24 hours a day. Darkness is not needed for receiving radio signals because the Sun is a relatively weak source of radio energy, so its emission does not swamp radio signals arriving at Earth from elsewhere in the sky. In addition, radio observations can often be made through cloudy skies, and radio telescopes can detect the longest-wavelength radio waves even during rain or snowstorms. Poor weather causes few problems because the wavelength of most radio waves is much larger than the typical size of atmospheric raindrops or snowflakes. Optical astronomy cannot be performed under these conditions because the wavelength of visible light is smaller than a raindrop, a snowflake, or even a minute water droplet in a cloud.

However, perhaps the greatest value of radio astronomy (and, in fact, of all astronomies concerned with non-visible regions of the electromagnetic spectrum) is that it



▲ **FIGURE 5.23 Haystack Observatory** Photograph of the Haystack dish, inside its protective radome. For scale, note the engineer standing at the bottom. Note also the dull shine on the telescope surface, indicating its smooth construction. Haystack is a poor optical mirror, but a superb radio telescope. It can be used to reflect and accurately focus radiation having short radio wavelengths, even as small as a fraction of a centimeter. (MIT)

opens up a whole new window on the universe. There are three main reasons for this. First, just as objects that are bright in the visible part of the spectrum (e.g., the Sun) are not necessarily strong radio emitters, many of the strongest radio sources in the universe emit little or no visible light. Second, visible light may be strongly absorbed by interstellar dust along the line of sight to a source. Radio waves, by contrast, are generally unaffected by intervening matter. Third, as mentioned earlier, many parts of the universe cannot be seen at all by optical means, but are easily detectable at longer wavelengths. The center of the Milky Way Galaxy is a prime example of such a totally invisible region—our knowledge of the Galactic center is based



▲ **FIGURE 5.24 Orion Nebula in Radio and Visible** The Orion Nebula is a star-forming region about 1500 light-years from Earth. (The nebula is located in the constellation Orion and can be seen as a small smudge in Figure 1.8.) The bright regions in this photograph are stars and clouds of glowing gas. The dark regions are not empty, but their visible emission is obscured by interstellar matter. Superimposed on the optical image is a radio contour map (blue lines) of the same region. Each curve of the contour map represents a different intensity of radio emission. Note how the radio contours in some places enclose regions that are dark in visible light, allowing us to “see through” the obscuring material. The resolution of the optical image is about  $1''$ ; that of the radio map is  $1'$ . (Background photo: AURA)

almost entirely on radio and infrared observations. Thus, these observations not only afford us the opportunity to study the same objects at different wavelengths, but also allow us to see whole new classes of objects that would otherwise be completely unknown.

Figure 5.24 shows an optical photograph of the Orion Nebula (a huge cloud of interstellar gas) taken with the 4-m telescope on Kitt Peak. Superimposed on the optical image is a radio map of the same region, obtained by scanning the Haystack radio telescope (Figure 5.23) back and forth across the nebula and taking many measurements of radio intensity. The map is drawn as a series of contour lines connecting locations of equal radio brightness, similar to pressure contours drawn by meteorologists on weather maps or height contours drawn by cartographers on topographic maps. The inner contours represent stronger radio signals, the outside contours weaker signals. Note, however, how the radio map is much less detailed than its optical counterpart; that’s because the acquired radio radiation has such a long wavelength compared to light.

The radio map in Figure 5.24 has many similarities to the visible-light image of the nebula. For instance, the radio emission is strongest near the center of the optical image and declines toward the nebular edge. But there are also subtle differences between the radio and optical images. The two differ chiefly toward the upper left of the main cloud, where visible light seems to be absent, despite the existence of radio waves. How can radio waves be detected from locations not showing any emission of light? The answer is that this particular nebular region is known to be especially dusty in its top left quadrant. The dust obscures the short-wavelength visible light, but not the long-wavelength radio radiation. Thus, we have a trade-off typical of many in astronomy: Although the long-wave radio signals provide a less resolved map of the region, those same radio signals can pass relatively unhindered through dusty regions, in this case allowing us to see the true extent of the Orion Nebula.

### CONCEPT Check

- ✓ In what ways does radio astronomy complement optical observations?

## 5.6 Interferometry

The main disadvantage of radio astronomy compared with optical work is its relatively poor angular resolution. However, in some circumstances, radio astronomers can overcome this limitation with a technique known as **interferometry**. This technique makes it possible to produce radio images of angular resolution higher than can be achieved with even the best optical telescopes, on Earth or in space.

In interferometry, two or more radio telescopes are used in tandem to observe the *same* object at the *same* wavelength and at the *same* time. The combined instruments together make up an **interferometer**. Figure 5.25 shows a large interferometer—many separate radio telescopes working together as a team. By means of electronic cables or radio links, the signals received by each antenna in the array making up the interferometer are sent to a central computer that combines and stores the data.

### A Radio Interferometer

Interferometry works by analyzing how the signals interfere with each other when added together. ∞ (Discovery 3-1) Consider an incoming wave striking two detectors (Figure 5.26). Because the detectors lie at different distances from the source, the signals they record will, in general, be out of step with one another. In that case, when the signals are combined, they will interfere destructively, partly canceling each other out. Only if the detected radio waves happen to be exactly in step will



◀ **FIGURE 5.25 VLA Interferometer**

(a) This large interferometer, located on the Plain of San Augustin in New Mexico, comprises 27 separate dishes spread along a Y-shaped pattern about 30 km across. The most sensitive radio device in the world, it is called the Very Large Array, or VLA for short. (b) A close-up view from ground level shows how some of the VLA dishes are mounted on railroad tracks so that they can be repositioned easily. (NRAO)

(a)



(b)


the signals combine constructively to produce a strong signal. Notice that the amount of interference depends on the *direction* in which the wave is traveling relative to the line joining the detectors. Thus—in principle, at least—careful analysis of the strength of the combined signal can provide an accurate measurement of the source's position in the sky.





As Earth rotates and the antennae track their target, the interferometer's orientation relative to the source changes, and a pattern

of peaks and troughs emerges. In practice, extracting positional information from the data is a complex task, as multiple antennae and several sources are usually involved. Suffice it to say that, after extensive computer processing, the interference pattern translates into a high-resolution image of the target object.

An interferometer is, in essence, a substitute for a single huge antenna. As far as resolving power is concerned, the effective diameter of an interferometer is the distance between its outermost dishes. In other words, two small dishes can act as opposite ends of an imaginary, but huge, single radio telescope, dramatically improving angular resolution. For example, a resolution of a few arc seconds can be achieved at typical radio wavelengths (such as 10 cm), either by using a single radio telescope 5 km in diameter (which is impossible to build) or by using two or more

▶ **FIGURE 5.26 Interferometry** Two detectors, A and B, record different signals from the same incoming wave because of the time it takes the radiation to traverse the distance between them. When the signals are combined, the amount of interference depends on the wave's direction of motion, providing a means of measuring the position of the source in the sky. Here, the dark-blue waves come from a source high in the sky and interfere destructively when captured by antennas A and B. But when the same source has moved because of Earth's rotation (light-blue waves), the interference can be constructive.



A  Waves arrive out of step  
B  = destructive interference  
A  Waves arrive in step  
B  = constructive interference



much smaller dishes separated by the same 5 km, but connected electronically. The larger the distance separating the telescopes—that is, the longer the *baseline* of the interferometer—the better is the resolution attainable.

Large interferometers like the instrument shown in Figure 5.25 now routinely attain radio resolution comparable to that of optical images. Figure 5.27 shows an interferometric radio map of a pair of colliding galaxies about 62 million light-years away, and a photograph of those same galaxies taken with a large optical telescope. The radio clarity is much better than in the contour map of Figure 5.24—in fact, the radio resolution in part (a) is comparable to that of the optical image in part (b). Note that the optical image in (b) is in true color, but the radio image in (a) is represented in *false color*, a commonly used technique for displaying data acquired in non-visible light. The radio “colors” do not represent actual wavelengths of the radiation emitted, but instead some other property of the source, in this case radio intensity, increasing from red to yellow.


See *Discovery 5-1* for a discussion of the latest powerful interferometer, just now coming online high in the Chilean Andes.

Astronomers have created radio interferometers spanning great distances, first across North America and later between continents. A typical very-long-baseline interferometry experiment (usually known by its abbreviation, VLBI) might use radio telescopes in North America, Europe, Australia, and Russia to achieve an angular resolution on the order of 0.001". It seems that even Earth's diameter is no limit: Radio astronomers have successfully used an antenna in orbit, together with several antennae on the ground, to construct an even longer baseline and achieve still better resolution. Proposals exist to place interferometers entirely in Earth orbit and even on the Moon.

## Interferometry at Other Wavelengths


Although the technique was originally developed by radio astronomers, interferometry is no longer restricted to the radio domain. Radio interferometry became feasible when electronic equipment and computers achieved speeds great enough to combine and analyze radio signals from separate radio detectors without loss of data. As the technology has improved, it has become possible to apply the same methods to radiation of higher frequency. Millimeter-wavelength interferometry has become an established and important observational technique, and both the Keck telescopes and the VLT are routinely used for near-infrared interferometry.

Perhaps the highest resolution interferometric instrument currently in operation is the six-telescope optical array operated by the Center for High Angular Resolution Astronomy (CHARA) on Mount Wilson in California



▲ **FIGURE 5.27 Radio–Optical Comparison** (a) ALMA radio “image” (or radiograph) of the colliding Antennae Galaxies observed at radio frequencies with an angular resolution of a few arc seconds. (b) Visible-light photograph of that same galaxy, made with the *Hubble* telescope and displayed on the same scale as (a). (ESO/NAOJ/NRAO; STScI)

(Figure 5.28). Although each telescope is only 1 m in diameter, the placement of the array over the mountain results in a combined light beam having resolution equivalent to a single telescope 300 m across. CHARA is not designed to produce images of the stars it studies. However, it can resolve details as small as 0.0002" across, allowing the positions, orbits, and even radii of some stars to be measured with exquisite accuracy.



**▲ FIGURE 5.28 Optical Interferometry** This aerial photo shows the facilities of the CHARA array intermingled with the existing equipment of the historic Mount Wilson Observatory, in California. The small 1-m telescopes in the array are numbered. (E. Simison/Sea West Enterprises)

### CONCEPT Check

- ✓ **What is the main reason for the poor angular resolution of radio telescopes? How do radio astronomers overcome this problem?**

## 5.7 Space-Based Astronomy

Optical and radio astronomy are the oldest branches of astronomy, but since the 1970s there has been a virtual explosion of observational techniques spanning the rest of the electromagnetic spectrum. Today, all portions of the spectrum are studied, from radio waves to gamma rays, to maximize the amount of information available about astronomical objects.

As noted earlier, the types of astronomical objects that can be observed differ quite markedly from one wavelength range to another. Full-spectrum coverage is essential not only to see things more clearly, but even to see some things at all. Because of the transmission characteristics of Earth's atmosphere, astronomers must study practically all regions of the electromagnetic spectrum, from gamma rays through X-rays to visible light, and on down to infrared and radio waves, from space. The rise of these "other astronomies" has therefore been closely tied to the development of the space program.

### Infrared Astronomy

Infrared studies are a vital component of modern observational astronomy. Infrared astronomy spans a broad range


of phenomena in the universe, from planets and their parent stars to the vast regions of interstellar space where new stars are forming, to explosive events occurring in faraway galaxies. Generally, **infrared telescopes** resemble optical telescopes, but their detectors are designed to be sensitive to radiation of longer wavelengths. Indeed, as we have seen, many ground-based "optical" telescopes are also used for infrared work, and some of the most useful infrared observing is done from the ground (e.g., from Mauna Kea—see Figure 5.10), even though the radiation is somewhat diminished in intensity by our atmosphere.

As with radio observations, the longer wavelength of infrared radiation often enables us to perceive objects that are partially hidden from optical view. As a terrestrial example of the penetrating properties of infrared radiation, Figure 5.29(a) shows a dusty and hazy

region in California, hardly viewable optically, but easily seen in the infrared (Figure 5.29b). Figures 5.29(c) and (d) show a similar comparison for an astronomical object—the dusty regions of the Orion Nebula, where visible light is blocked by interstellar clouds, but which is clearly distinguishable in the infrared.

Astronomers can make better infrared observations if they place their instruments above most or all of Earth's atmosphere using balloon-, aircraft-, rocket-, and satellite-based telescopes. As might be expected, orbiting infrared telescopes are generally somewhat smaller than the massive instruments found in ground-based observatories. In 2003, NASA launched the 0.85-m *Spitzer Space Telescope (SST)*, named in honor of Lyman Spitzer, Jr., a renowned astrophysicist and the first person to propose (in 1946) that a large telescope be located in space. *Spitzer's* detectors are designed to operate at wavelengths between 3.6 and 160  $\mu\text{m}$ , with resolution varying from 2.5" to 40" across this range. Unlike previous space-based observatories, *SST* does not orbit Earth, but instead follows our planet in its orbit around the Sun, trailing millions of kilometers behind in order to minimize Earth's heating effect on the detectors. It is currently drifting away from Earth at the rate of 0.1 AU per year. Figure 5.30 shows some spectacular (false-color) recent imagery from *SST*.

*Spitzer's* detectors were cooled to near absolute zero in order to observe infrared signals from space without interference from the telescope's own heat. Unfortunately, the liquid helium doing this could not be confined indefinitely, and (as expected) it slowly leaked away into space. In 2009, *SST* entered a new "warm" phase of operation as its temperature



▲ **FIGURE 5.29 Smog Revealed** An optical photograph (a) taken near San Jose, California, and an infrared photo (b) of the same area taken at the same time. Infrared radiation of long wavelength can penetrate smog much better than short-wavelength visible light. The same advantage pertains to astronomical observations: An optical view (c) of an especially dusty part of the central region of the Orion Nebula is more clearly revealed in this infrared image (d) showing a cluster of stars behind the obscuring dust. (*Lick Observatory; NASA*)

increased to roughly 30 K—still very cold by Earth standards, but warm enough for the telescope’s own thermal emission to overwhelm the long-wavelength detectors on board. (Recall from Wien’s law that the thermal emission of a 30-K object peaks at a wavelength of roughly  $100\ \mu\text{m}$ .) However, the shorter-wavelength detectors (at  $3.6$  and  $4.5\ \mu\text{m}$ ) should remain operational at least through 2014, and the telescope will continue to be a key astronomical resource during that time.


The latest—and largest—orbiting infrared telescope is the 3.5-m European *Herschel Space Observatory*, named for the British astronomer Sir William Herschel, who first demonstrated (in 1800) the existence of infrared radiation. Launched in 2009, *Herschel* is designed to operate in the far-infrared part of the spectrum at wavelengths between about  $50$  and  $700\ \mu\text{m}$ . The spacecraft is stationed at the  $L_2$  Lagrangian point of Earth’s orbit, a stable location roughly 1.5 million km from Earth, outward along the Sun–Earth line. Figure 5.31(a) is a *Herschel* view of a

nearby star-forming region called the Eagle Nebula. This false-color image combines data taken at three different infrared wavelengths ( $70\ \mu\text{m}$ ,  $160\ \mu\text{m}$ , and  $250\ \mu\text{m}$ ), represented as blue, green, and red, respectively. Figure 5.31(b) shows a visible-light image of the same field. Note how the infrared view shows huge clouds of warm dust and gas, critical components of the star-formation process that are completely dark at visible wavelengths. *Herschel* is expected to cease operations in late 2013, when its coolant runs out.

The present instrument package aboard the *Hubble Space Telescope* (see Section 5.1) also includes a high-resolution ( $0.1''$ ) near-infrared camera and spectroscope. NASA plans to launch *Hubble*’s successor, the *James Webb Space Telescope*, in 2018. With a 6.5-m segmented mirror and detectors optimized for use at near- and mid-infrared wavelengths, *JWST* is expected to become the premier instrument for infrared astronomy.



(a)



(b)


▲ **FIGURE 5.30 Spitzer Images** These images from the *Spitzer Space Telescope*, now in orbit around the Sun, clearly show its camera's capabilities. (a) This unnamed star-forming region displays much dust (red-orange) amid myriad stars (blue-white). (b) The much larger spiral galaxy M100 also radiates heat from its embedded dust. (JPL)

## Ultraviolet Astronomy

On the short-wavelength side of the visible spectrum lies the ultraviolet domain. Extending in wavelength from 400 nm (blue light) down to a few nanometers (“soft” X-rays), this



(a)



(b)



▲ **FIGURE 5.31 Infrared–Optical Comparison** (a) This infrared image of the Eagle Nebula was acquired with the *Herschel Space Observatory*. In this false-colored image, colors denote dust temperatures, descending from blue to red. (b) The same nebula in visible light shows clearly just how much of the region is obscured by the dust within it. (ESA; ESO)

spectral range has only recently begun to be explored. Because Earth’s atmosphere is partially opaque to radiation below 400 nm and is totally opaque below about 300 nm (due in part to the ozone layer), astronomers cannot conduct any useful ultraviolet observations from the ground, even from the highest mountaintop. Rockets, balloons, or satellites are therefore essential to any **ultraviolet telescope**—a device designed to capture and analyze that high-frequency radiation.

## DISCOVERY 5-1


## The ALMA Array

High in the South American mountains, a new and powerful telescope has been built by an international consortium of astronomers and engineers from the United States, Canada, Europe, East Asia, and Chile. The Atacama Large Millimeter Array (ALMA) is the largest astronomical project on Earth today, residing about 5000 m altitude on the Chajnantor plateau in northern Chile. There, in the Atacama Desert, the telescope scans the universe from one of the driest places on our planet, without clouds, radio interference, or light pollution. It's a dream come true for astronomers wishing to sense the radio universe with clarity and resolution equal to that seen by optical telescopes.

ALMA is a telescope of revolutionary design—actually an array of 66 radio antennas that operate in synchrony as a single device. Each high-precision antenna detects radiation of wavelength between 0.3 and 10 mm—the so-called millimeter band that is midway between the traditional radio and infrared parts of the electromagnetic spectrum. To date, astronomers haven't been able to probe the universe very much in this remote spectral domain and so far have had only glimpses of new science to come. The antennas all aim toward the same cosmic object at any one time, yet from slightly different perspectives the many antennas examine it with superb resolution. Furthermore, the antennas are mobile, moving around on the desert floor and enabling researchers to effectively “zoom” in on target objects. The photo shows one of the first deployments of some of the antennas in 2011.

The earliest images from ALMA began arriving in 2012, giving astronomers an inkling of the array's potential; even better

quality images are expected as the antennas are finely tuned and as more antennas join the array. One the most intriguing ALMA images is shown in Figure 5.27—two galaxies in collision. Another is the image here—a narrow dust ring around a young star system. This is Fomalhaut, shown here as the bright emission at the center of the ring, a star about 25 light-years away and thought to perhaps have planets now forming within its dusty ring. That ring or disk is shown here in two ways: The bluish-colored radiation at bottom was observed by the *Hubble Space Telescope* in the optical domain, and the narrow yellow ring at the top by ALMA (in false color). *Hubble* astronomers thought they had earlier spied a huge, Jupiter-sized planet within the dust, but the new, high-resolution ALMA image shows no such planet. If there are planets present there, they must be much smaller, perhaps comparable to Earth. The hunt for Earth-sized planets is heating up, and ALMA is at the forefront of that major effort.



(ESO/NAOJ/NRAO; STScI)

ALMA is expected to be a telescopic workhorse for the next generation of astronomers. As more antennas are added, perhaps ultimately totaling several hundred, this single most powerful telescope ever built is expected to extensively probe one of astronomy's last frontiers. A whole new window on the universe has now been opened, capturing never-before-seen details about the earliest stars and galaxies, the enigmatic black holes at the hearts of most galaxies, and perhaps directly imaging the formation of planets beyond our solar system.


One of the most successful ultraviolet space missions was the *International Ultraviolet Explorer (IUE)*, placed in Earth orbit in 1978 and shut down for budgetary reasons in late 1996. Like all ultraviolet telescopes, its basic appearance and construction were quite similar to those of optical and infrared devices. Several hundred astronomers from all over the world used *IUE*'s near-ultraviolet spectroscopes to explore a variety of phenomena in planets, stars, and galaxies. In subsequent chapters, we will learn what this relatively new window on the universe has shown us about the activity and even the violence that seems to pervade the cosmos.

Figure 5.32 shows images captured by two more recent ultraviolet satellites. Figure 5.32(a) shows an image of a supernova remnant—the remains of a violent stellar


explosion that occurred some 12,000 years ago—obtained by the *Extreme Ultraviolet Explorer (EUVE)* satellite, launched in 1992. Since its launch, *EUVE* has mapped out our local cosmic neighborhood as it presents itself in the far ultraviolet and has radically changed astronomers' conception of interstellar space in the vicinity of the Sun. Figure 5.32(b) shows an image of two relatively nearby galaxies, called M81 and M82, captured by the *Galaxy Evolution Explorer (GALEX)* satellite, launched in 2003. *HST*, best known as an optical telescope, is also a superb imaging and spectroscopic ultraviolet instrument.

### PROCESS OF SCIENCE Check

- ✓ Why are observations made at many different electromagnetic wavelengths useful to astronomers?



(a)



(b)




## High-Energy Astronomy

*High-energy* astronomy studies the universe as it presents itself to us in X-rays and gamma rays—the types of radiation whose photons have the highest frequencies and hence the greatest energies. How do we detect radiation of such short wavelengths? First, it must be captured high above Earth's atmosphere, because none of it reaches the ground. Second, its detection requires the use of equipment fundamentally different in design from that used to capture the relatively low-energy radiation discussed up to this point.

The difference in the design of **high-energy telescopes** comes about because X-rays and gamma rays cannot be reflected easily by any kind of surface. Rather, these rays tend to pass straight through, or be absorbed by, any material they strike. When X-rays barely graze a surface, however, they can be reflected from it in a way that yields an image, although the mirror design is fairly complex. As illustrated in Figure 5.33, to ensure that all incoming rays are reflected at grazing angles, the telescope is constructed as a series of nested cylindrical mirrors, carefully shaped to bring the X-rays to a sharp focus. For gamma rays, no such method of producing an image has yet been devised; present-day gamma-ray telescopes simply point in a specified direction and count the photons they collect.

◀ **FIGURE 5.32 Ultraviolet Images** (a) A camera on board the *Extreme Ultraviolet Explorer* satellite captured this image of the Cygnus loop supernova remnant, the result of a massive star that blew itself virtually to smithereens. The release of energy was enormous, and the afterglow has lingered for centuries. The glowing field of debris shown here within the telescope's circular field of view lies some 1500 light-years from Earth. (b) This false-color image of the spiral galaxy M81 and its companion M82, made by the *Galaxy Evolution Explorer* satellite, reveals stars forming in the blue arms well away from the galaxy's center. (NASA; GALEX)




▲ **FIGURE 5.33 X-Ray Telescope** (a) The arrangement of nested mirrors in an X-ray telescope allows the rays to be reflected at grazing angles and focused to form an image. (b) A cutaway 3-D rendition of the mirrors, showing their shape more clearly.


In addition, detection methods using photographic plates or CCD devices do not work well for hard (high-frequency) X-rays and gamma rays. Instead, individual photons are counted by electronic detectors on board an orbiting device, and the results are then transmitted to the ground for further processing and analysis. Furthermore, the number of photons in the universe seems to be inversely related to their frequency. Trillions of visible (starlight) photons reach the detector of an optical telescope on Earth each second, but hours or even days are sometimes needed for a single gamma-ray photon to be recorded. Not only are these photons hard to focus and measure, but they are also very scarce.

The *Einstein Observatory*, launched by NASA in 1978, was the first X-ray telescope capable of forming an image of its field of view. During its 2-year lifetime, this spacecraft drove major advances in our understanding of high-energy phenomena throughout the universe. The German *ROSAT* (short for “Röntgen Satellite,” after Wilhelm Röntgen, the discoverer of X-rays) was launched in 1991. During its 7-year lifetime, this instrument generated a wealth of high-quality observational data. It was turned off in 1999, a few months after its electronics were irreversibly damaged when the telescope was accidentally pointed too close to the Sun.

In July 1999, NASA launched the *Chandra X-Ray Observatory* (named in honor of the Indian astrophysicist Subramanyan Chandrasekhar and shown in Figure 5.34). With greater sensitivity, a wider field of view, and better resolution than either *Einstein* or *ROSAT*, *Chandra* is providing high-energy astronomers with new levels of observational detail. Figure 5.35 shows a typical image returned by *Chandra*: a supernova remnant in the constellation Cassiopeia. Known as Cas A, the ejected gas is all that now remains of a star that was observed to explode about



▲ **FIGURE 5.34 Chandra Observatory** The *Chandra* X-ray telescope is shown here during the final stages of its construction in 1998. The left end of the mirror arrangement, depicted in Figure 5.33, is at the bottom of the satellite as oriented here. *Chandra*'s effective angular resolution is 1", allowing this spacecraft to produce images of quality comparable to that of optical photographs. *Chandra* now occupies an elliptical orbit high above Earth; its farthest point from our planet, 140,000 km out, reaches almost one-third of the way to the Moon. (NASA)



◀ **FIGURE 5.35 X-Ray Image** This is a false-color *Chandra* X-ray image of the supernova remnant Cassiopeia A, a debris field of scattered, hot gases that were once part of a massive star. Roughly 10,000 light-years from Earth and barely visible in the optical part of the spectrum, Cas A is awash in brilliantly glowing X-rays spread across some 10 light-years. (CXC/SAO)


Gamma-ray astronomy is the youngest entrant into the observational arena. As just mentioned, imaging gamma-ray telescopes do not exist, so only fairly coarse ( $1^\circ$  resolution) observations can be made. Nevertheless, even at this resolution, there is much to be learned. Cosmic gamma rays were originally detected in the 1960s by the U.S. *Vela* series of satellites, whose primary mission was to monitor illegal nuclear detonations on Earth. Since then, several X-ray telescopes have also been equipped with gamma-ray detectors.

NASA's *Compton Gamma-Ray Observatory (CGRO)* was placed in orbit by the space shuttle in 1991. It scanned the sky and studied individual objects in much greater detail than had previously been attempted. Many examples of *CGRO*'s imagery appear throughout this book. The mission ended on June 4, 2000, when, following a failure of one of the satellite's three gyroscopes, NASA opted for a controlled reentry and dropped *CGRO* into the Pacific Ocean. In 2008, NASA launched the *Fermi Gamma-Ray Space Telescope* (Figure 5.36a). With greater sensitivity to a broader range of gamma-ray energies than *CGRO*, *Fermi*'s capabilities are greatly expanding astronomers' view of

320 years ago. The false-color image shows 50-million-kelvin gas in wisps of ejected stellar material; the bright white point at the very center of the debris may be a black hole. The *XMM-Newton* satellite is more sensitive to S-rays than is *Chandra* (that is, it can detect fainter X-ray sources), but it has significantly poorer angular resolution ( $5''$ , compared to  $0.5''$  for *Chandra*), making the two missions complementary to one another.



(a)



(b)

▲ **FIGURE 5.36 Gamma-Ray Astronomy** (a) The *Fermi Gamma-Ray Space Telescope*, shown here in an artist's conception, was named after Enrico Fermi, an Italian-American scientist who did pioneering work in high-energy physics. The wide arrays are solar panels to power the spacecraft; the box amidship contains layers of tungsten that detect the gamma rays. (b) A typical false-color gamma-ray image—this one showing the remains of a violent event (a supernova) in a region named W44. The gamma rays are shown mainly in magenta. (NASA)



the high-energy universe. Figure 5.36(b) shows a false-color gamma-ray image of a violent stellar explosion in a distant galaxy. An early all-sky image obtained by *Fermi* appears in Figure 5.37(e).

### CONCEPT Check


- ✓ List some scientific benefits of placing telescopes in space. What are the drawbacks of space-based astronomy?

## 5.8 Full-Spectrum Coverage

Many astronomical objects are now routinely observed at many different electromagnetic wavelengths. As we proceed through the text, we will discuss more fully the wealth of information that high-precision astronomical instruments can provide us.

It is reasonable to suppose that the future holds many further improvements in both the quality and the availability of astronomical data and that many new discoveries will be made. The current and proposed pace of technological progress presents us with the following exciting prospect: Within the next decade, if all goes according to plan, it will be possible, for the first time ever, to make *simultaneous* high-quality measurements of any astronomical object at *all* wavelengths, from radio ray to gamma ray. The consequences of this development for our understanding of the workings of the universe may be little short of revolutionary.

As a preview of the comparison that full-spectrum coverage allows, Figure 5.37 shows a series of images of our own Milky Way Galaxy. The images were made by several different instruments, at wavelengths ranging from radio to gamma rays, over a period of about 5 years. By comparing the features visible in each, we immediately see how multi-wavelength observations can complement one another, greatly extending our perception of the dynamic universe around us.




**Narrated FIGURE 5.37 Multiple Wavelengths** The Milky Way Galaxy as it appears at (a) radio, (b) infrared, (c) visible, (d) X-ray, and (e) gamma-ray wavelengths. Each frame is a panoramic view covering the entire sky. The center of our Galaxy, which lies in the direction of the constellation Sagittarius, is at the center of each map. (ESA; UMass/Caltech; A. Mellinger; MPI; NASA)

**The Big Question** Astronomy is a data-driven science. The most stunning discoveries are often made as new telescopes come online; some scopes are bigger, others operate in orbit, and almost all of them are better than any equipment available previously. The biggest advances have been made by instruments built to sense new domains of the electromagnetic spectrum. Will telescopes someday operate outside the electromagnetic spectrum, detecting new kinds of radiation or some still unknown particles? Whole new windows on the universe potentially beckon.

# Chapter Review

## SUMMARY


**1** A **telescope** (p. 100) is a device designed to collect as much light as possible from some distant source and deliver it to a detector for detailed study. A **refracting telescope** (p. 100) uses a lens to concentrate and focus the light; **reflecting telescopes** (p. 100) use mirrors. The **Newtonian** (p. 103) and **Cassegrain** (p. 103) reflecting telescope designs employ secondary mirrors to avoid placing detectors at the prime focus. More complex light paths are used to allow the use of large or heavy equipment that cannot be placed near the telescope. All large astronomical telescopes are reflectors, because large mirrors are lighter and much easier to construct than large lenses, and also suffer from fewer optical defects.




**2** The light-gathering power of a telescope depends on its **collecting area** (p. 105), which is proportional to the square of the mirror diameter. To study the faintest sources of radiation, astronomers must use large telescopes. Large telescopes also suffer least from the effects of diffraction and hence can achieve better **angular resolution** (p. 106) once the blurring effects of Earth's atmosphere are overcome. The amount of diffraction is proportional to the wavelength of the radiation under study and inversely proportional to the size of the mirror.



**3** Most modern telescopes use **charge-coupled devices**, or **CCDs** (p. 109), instead of photographic plates to collect their data. The field of view is divided into an array of millions of **pixels** (p. 109) that accumulate an electric charge when light strikes them. CCDs are many times more sensitive than photographic plates, and the resultant data are easily saved directly in digital form for later processing. The light collected by a telescope may be processed in a number of ways. It can be made to form an **image** (p. 100). **Photometry** (p. 111) may be performed either on a stored image or during the observation itself, using a specialized detector, or a **spectrometer** (p. 111) may be used to analyze the spectrum of the radiation received.




**4** The resolution of most ground-based optical telescopes is limited by **seeing** (p. 111)—the blurring effect of Earth's turbulent atmosphere, which smears the pointlike images of stars out into **seeing disks** (p. 111) a few arc seconds in diameter. Radio and space-based telescopes do not suffer from this effect, so their resolution is determined mainly by the effects of diffraction. Astronomers can greatly improve a telescope's resolution by using **active optics** (p. 112), in which a telescope's environment and focus are carefully monitored and controlled, and **adaptive optics** (p. 112), in which the blurring effects of atmospheric turbulence are corrected for in real time.




**5** **Radio telescopes** (p. 114) are conceptually similar in construction to optical reflectors, although they are generally much larger than optical instruments, in part because so little radio radiation reaches Earth from space, so a large collecting area is essential. Their main disadvantage is that diffraction of long-wavelength radio waves limits their resolution, even for very large radio telescopes. Their principal advantage is that they allow astronomers to explore a whole new part of the electromagnetic spectrum and of the universe—many radio emitters are completely undetectable in visible light. In addition, radio observations are largely unaffected by Earth's atmosphere and weather or by the position of the Sun.




**6** In order to increase the effective area of a telescope, and hence improve its resolution, several separate instruments may be combined into a device called an **interferometer** (p. 118), in which the interference pattern of radiation received by two or more detectors is used to reconstruct a high-precision map of the source. Using **interferometry** (p. 118), radio telescopes can produce images sharper than those from the best optical telescopes. Infrared and optical interferometric systems are now in use at many observatories.



**7** **Infrared** (p. 121) and **ultraviolet telescopes** (p. 123) are generally similar in design to optical systems. Observations in some parts of the infrared range can be carried out from the ground, but ultraviolet astronomy *must* be done from space. **High-energy telescopes** (p. 125) study the X-ray and gamma-ray regions of the electromagnetic spectrum. X-ray telescopes can form images of their field of view, although the mirror design is more complex than that of optical instruments. Gamma-ray telescopes simply point in a certain direction and count the photons they collect. Because the atmosphere is opaque at these short wavelengths, both types of telescopes must be placed in space.



**8** Different physical processes can produce very different types of electromagnetic radiation, and the appearance of an object at one wavelength may bear little resemblance to its appearance at another. Observations at wavelengths spanning the spectrum are essential to a complete understanding of astronomical events.



**MasteringAstronomy**<sup>®</sup> For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- LO1** List three advantages of reflecting telescopes over reflectors.
- LO2 POS** Cite two reasons that astronomers are continually building larger and larger telescopes.
- LO3** What are the advantages of a CCD over a photograph?
- LO4** How does Earth's atmosphere affect what is seen through an optical telescope?
- What advantages does the *Hubble Space Telescope* (*HST*) have over ground-based telescopes? List some disadvantages.
- What determines the resolution of a ground-based telescope?
- How do astronomers use adaptive optics to improve the resolution of telescopes?
- Why do radio telescopes have to be very large?
- LO5** Which astronomical objects are best studied with radio techniques?
- LO6** What is interferometry, and what problem in radio astronomy does it address? Is it limited to radio astronomy?
- Why do infrared satellites have to be cooled?
- LO7** Are there any ground-based ultraviolet observatories?
- In what ways do the mirrors in X-ray telescopes differ from those found in optical instruments?
- LO8 POS** What are the main advantages of studying objects at many different wavelengths of radiation?
- POS** Our eyes can see light with an angular resolution of  $1'$ . Suppose our eyes detected only infrared radiation, with  $1^\circ$  angular resolution. Would we be able to make our way around on Earth's surface? To read? To sculpt? To create technology?



## Conceptual Self-Test: Multiple Choice

- VIS** According to Figure 5.2 ("Refracting Lens") the thickest lenses deflect and bend light (a) the fastest; (b) the slowest; (c) the most; (d) the least.
- The main reason that most professional research telescopes are reflectors is that (a) mirrors produce sharper images than lenses do; (b) their images are inverted; (c) they do not suffer from the effects of seeing; (d) large mirrors are easier to build than large lenses.
- If telescope mirrors could be made of odd sizes, the one with the *most* light-gathering power would be (a) a triangle with 1-m sides; (b) a square with 1-m sides; (c) a circle 1 m in diameter; (d) a rectangle with two 1-m sides and two 2-m sides.
- VIS** The image shown in Figure 5.12 ("Resolution") is sharpest when the ratio of wavelength to telescope size is (a) large; (b) small; (c) close to unity; (d) none of these.
- The primary reason professional observatories are built on the highest mountaintops is to (a) get away from city lights; (b) be above the rain clouds; (c) reduce atmospheric blurring; (d) improve chromatic aberration.

6. Compared with radio telescopes, optical telescopes can (a) see through clouds; (b) be used during the daytime; (c) resolve finer detail; (d) penetrate interstellar dust.
7. When multiple radio telescopes are used for interferometry, resolving power is most improved by increasing (a) the distance between telescopes; (b) the number of telescopes in a given area; (c) the diameter of each telescope; (d) the electrical power supplied to each telescope.
8. The *Spitzer Space Telescope (SST)* is stationed far from Earth because (a) this increases the telescope's field of view; (b) the telescope is sensitive to electromagnetic interference from terrestrial radio stations; (c) doing so avoids the obscuring effects of Earth's atmosphere; (d) Earth is a heat source and the telescope must be kept very cool.
9. The best way to study warm (1000 K) young stars forming behind an interstellar dust cloud would be to use (a) X-rays; (b) infrared light; (c) ultraviolet light; (d) blue light.
10. The best frequency range in which to study the hot (million-kelvin) gas found among the galaxies in the Virgo galaxy cluster would be in the following region of the electromagnetic spectrum: (a) radio; (b) infrared; (c) X-ray; (d) gamma-ray.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

1. ● A certain telescope has a  $10' \times 10'$  field of view that is recorded using a CCD chip having  $2048 \times 2048$  pixels. What angle on the sky corresponds to 1 pixel? What would be the diameter, in pixels, of a typical seeing disk (1" radius)?
2. ● The *SST*'s initial operating temperature was 5.5 K. At what wavelength (in micrometers) does the telescope's own black-body emission peak? How does this wavelength compare with the wavelength range in which the telescope is designed to operate?  (*More Precisely 3-2*)
3. ● A 2-m telescope can collect a given amount of light in 1 hour. Under the same observing conditions, how much time would be required for a 6-m telescope to perform the same task? A 12-m telescope?
4. ● A space-based telescope can achieve a diffraction-limited angular resolution of  $0.05''$  for red light (wavelength 700 nm). What would the resolution of the instrument be (a) in the infrared, at  $3.5 \mu\text{m}$ , and (b) in the ultraviolet, at 140 nm?
5. ●● Two identical stars are moving in a circular orbit around one another, with an orbital separation of 2 AU.  (*Sec. 2.6*) The system lies 200 light-years from Earth. If we happen to view the orbit head-on, how large a telescope would we need to resolve the stars, assuming diffraction-limited optics at a wavelength of  $2 \mu\text{m}$ ?
6. ● What is the equivalent single-mirror diameter of a telescope constructed from two separate 10-m mirrors? Four separate 8-m mirrors?
7. ● The Moon lies about 380,000 km away. To what distances do the angular resolutions of *SST* ( $3''$ ), *HST* ( $0.05''$ ), and a radio interferometer ( $0.001''$ ) correspond at that distance?
8. ● Estimate the angular resolutions of (a) a radio interferometer with a 5000-km baseline, operating at a frequency of 5 GHz, and (b) an infrared interferometer with a baseline of 50 m, operating at a wavelength of  $1 \mu\text{m}$ .

## Activities

### Collaborative



1. Your group has been assigned to observe the region of the sky around Orion to look for hot, bright young stars hidden in molecular clouds. Explain which of the telescopes described in the text would be your best choice, and estimate the level of detail you might expect to see.
2. Determine the maximum size interferometer your group could build if you placed 2-m radio telescopes at each of your homes. What would be its resolution at a wavelength of 1 cm?

### Individual

1. Take some photographs of the night sky. You will need a location with a clear, dark sky; a good digital camera that

lets you control the exposure time; a tripod and cable release; and a watch with a seconds display that is visible in the dark. Set your camera to the "manual" setting for the exposure and attach the cable release so you can control it. Set the focus on infinity. Point the camera at the desired constellation, seen through the viewfinder, and take a 20- to 30-second exposure. Don't touch any part of the camera or hold on to the cable release during the exposure to minimize all vibration. Keep a log of your shots.


2. Which image of the Milky Way Galaxy in Figure 5.36 (Multiple Wavelengths) do you think provides the most interesting information? Explain your reasoning.



## PART TWO

Early photo and sketch of Mars (*Lowell Observatory*)

# Our Planetary System



Percival Lowell (*Lowell Observatory*)


**The year 1877** was an important one in the study of the planet Mars. The Red Planet came unusually close to Earth, affording astronomers an especially good view. Of particular note was the discovery, by U.S. Naval Observatory astronomer Asaph Hall, of the two moons circling Mars. But most exciting was the report of the Italian astronomer Giovanni Schiaparelli on his observations of a network of linear markings that he termed *canali*. In Italian, this word usually means “grooves” or “channels,” but it can also mean “canals.” Schiaparelli probably did not intend to imply that the *canali* were anything other than natural, but the word was translated into English as “canals,” suggesting that the grooves had been constructed by intelligent beings.

As often happens—then as now—the world’s press (especially in the United States) sensationalized Schiaparelli’s observations, and some astronomers began drawing elaborate maps of Mars, showing oases and lakes where canals met in desert areas. His work fueled the widespread idea more than a century ago that intelligent life exists on Mars.


Percival Lowell (1855–1916), a successful Boston businessman (and brother of the poet Amy Lowell and Harvard president Abbott Lawrence Lowell), became so fascinated by these reports that he abandoned his business and purchased a clear-sky site at Flagstaff, Arizona, where he built a private observatory. He devoted his fortune and energies until the day he died to achieving a better understanding of the Martian “canals.” In doing so, he championed the idea that intelligent inhabitants of a drying (and dying) Mars had constructed a planetwide system of canals to transport water from the polar ice caps to the arid equatorial deserts.

Lowell was no slouch. He had earlier served as a distinguished diplomat and wrote extensively about the Far East. Later in life, he made an elaborate mathematical study of the orbit of Uranus, predicting the presence of an unseen body beyond Neptune—which was eventually found and named Pluto by Lowell Observatory’s Clyde Tombaugh in 1930. And Lowell offered support to young astronomers, including Vesto Slipher, whose pioneering research on the recession of the galaxies helped found modern cosmology. Even so, Lowell is best remembered for his passionate belief in advanced Martian civilizations.





Mars today, via *Hubble* (STScI)



Northern Martian plains, via *Phoenix* lander (JPL)

**Today**, we know that Mars is cool, dry, and probably lifeless; it certainly houses no intelligent beings and almost certainly never has. The Martian valleys and channels photographed by robot spacecraft in the 1970s are far too small to be the *canali* that Schiaparelli, Lowell, and others thought they saw on Mars. And the most recent armada of spacecraft—including the European *Mars Express* orbiter and the U.S. *Mars Reconnaissance Orbiter*, as well as the *Spirit/Opportunity* and *Curiosity* landers—that arrived at Mars in the early years of the 21st century reconfirm there is no liquid water there now.

The entire episode of Lowell's "canals" represents a classic case in which well-intentioned observers, perhaps obsessed with the notion of life on other worlds, let their personal opinions and prejudices seriously affect their analysis of reasonable data. The pair of globes of Mars (above left) shows how surface features, which were probably genuinely observed by astronomers a century ago, might have been imagined to be connected. The figure on the left of the pair is a photograph of how Mars actually looked in the best telescopes at the end of the 19th century. The sketch to its right is an interpretation, done at the height of the canal hoopla, of that same view. The human eye, under physiological stress, tends to connect dimly observed, yet distinctly separated, features, causing old maps of Mars to have been as much a work of art as of science. Humans saw patterns and canals where none in fact existed—as noted by today's higher quality image of Mars (above right).

The chronicle of the Martian canals illustrates how the scientific method demands that we acquire new data to sort out sense from nonsense, fact from fiction. Rather than simply believing the claims about the Martian canals, scientists sought further observations to test Lowell's hypothesis. Eventually, improved observations, climaxing in several robotic missions to the Red Planet more than a century after all the fuss began, totally disproved the existence of artificial canals. It often takes time, but the scientific method does lead to progress in understanding reality.

Panoramic view of Mars, via *Curiosity* Rover (JPL)





# 6

# The Solar System

## COMPARATIVE PLANETOLOGY AND FORMATION MODELS

In less than a single generation, we have learned more about the solar system—the Sun and everything that orbits it—than in all the centuries that went before. By studying the eight major planets, their moons, and the countless fragments of material that orbit in interplanetary space, astronomers have gained a richer outlook on our own home in space.

Space missions have visited all the planets of the solar system, extending astronomers' reach from the Earth-like inner planets to the giant gaseous worlds orbiting far from the Sun. Instruments aboard unmanned robots have taken close-up photographs of the planets and their moons and in some cases have made on-site measurements. Astronomers have come to realize that all solar system objects, large and small, have vital roles to play in furthering our understanding of our cosmic neighborhood.

**The Big Picture** Planets are by no means the only residents of our solar system. Modern telescopes and robot space probes have revealed much about comets, asteroids, and many of the planets' moons. Ironically, it is the pristine trash in our cosmic neighborhood that is telling us most about the origin and development of our solar system. Like the seafarers of old who discovered new worlds beyond Europe, today's spacefarers are now grandly exploring alien worlds beyond Earth.

**LEFT:** Asteroids sometimes collide with Earth, so it is very much in our own interest to keep an eye on them! This image shows a close-up of the asteroid Itokawa, which is only 0.5 km long—about five soccer fields across. It was photographed as the Japanese spacecraft, *Hayabusa*, having launched from Earth in 2003, slowly approached the asteroid in 2005. The craft then soft-landed, scooped up some rocky debris, and took off for Earth, landing back home in 2010. A remarkable engineering achievement, this mission also scientifically proved that asteroids like this one are the source of most meteorites—the oldest matter in the solar system. (JAXA)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Explain the importance of comparative planetology to solar system studies.
- 2 Describe the overall scale and structure of the solar system.
- 3 Summarize the basic differences between the terrestrial and the jovian planets.
- 4 Identify and describe the major nonplanetary components of the solar system.
- 5 Outline the theory of solar system formation that accounts for the overall properties of our planetary system.
- 6 Explain how the terrestrial planets formed.
- 7 Contrast the leading theories for the formation of the jovian worlds.
- 8 Describe how comets and asteroids formed, and explain their role in determining planetary properties.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.



## 6.1 An Inventory of the Solar System

The Greeks and other astronomers of old were aware of the Moon, the stars, and five planets—Mercury, Venus, Mars, Jupiter, and Saturn—in the night sky. ∞ (Sec. 2.2) They also knew of two other types of heavenly objects that were clearly neither stars nor planets. *Comets* appear as long, wispy strands of light in the night sky that remain visible for periods of up to several weeks and then slowly fade from view. *Meteors*, or “shooting stars,” are sudden bright streaks of light that flash across the sky, usually vanishing less than a second after they first appear. These transient phenomena must have been familiar to ancient astronomers, but their role in the “big picture” of the solar system was not understood until much later.

### Discovering Our Planetary System

Human knowledge of the basic content of the solar system remained largely unchanged from ancient times until the early 17th century, when the invention of the telescope made more detailed observations possible. Galileo Galilei was the first to capitalize on this new technology. (His simple telescope is shown in Figure 6.1.) Galileo’s discovery of the phases of Venus and of four moons orbiting Jupiter early in the 17th century helped change forever humankind’s vision of the universe. ∞ (Sec. 2.4)

As technological advances continued, knowledge of the solar system improved rapidly. Astronomers began discovering objects invisible to the unaided human eye. By the end

of the 19th century, astronomers had found Saturn’s rings (1659), the planets Uranus (1781) and Neptune (1846), many planetary moons, and the first *asteroids*—“minor planets” orbiting the Sun, mostly in a broad band (called the *asteroid belt*) lying between Mars and Jupiter. Ceres, the largest asteroid and the first to be sighted, was discovered in 1801. A large telescope of mid-19th-century vintage is shown in Figure 6.2.


The 20th century brought continued improvements in optical telescopes. Thousands more asteroids were discovered, along with three more planetary ring systems, dozens of moons, and the first *Kuiper belt objects*, orbiting beyond Neptune. The century also saw the rise of both nonoptical—especially radio and infrared—astronomy and spacecraft exploration, each of which has made vitally important contributions to the field of planetary science.

The latter part of the 20th century also saw an entirely new avenue for planetary exploration—space flight. Astronauts have carried out experiments on the Moon (see Figure 6.3), and numerous unmanned probes have left Earth and traveled to all of the other planets. Figure 6.4 shows a view from the *Spirit* robot prospecting on the Martian surface in 2005, its solar panels most evident in this panoramic view from inside a shallow crater.

As currently explored, our **solar system** is known to contain one star (the Sun), eight planets, 169 moons (at last count) orbiting those planets, eight *asteroids* and more than 100 *Kuiper belt objects* larger than 300 km (200 miles) in diameter, tens of thousands of smaller (but well-studied) asteroids and Kuiper belt objects, myriad *comets* a few



▲ **FIGURE 6.1 Early Telescope** The refracting telescope with which Galileo made his first observations was simple, but its influence on astronomy was immeasurable. (*Museo della Scienza; Scala/Art Resource, NY*)



▲ **FIGURE 6.2 Nineteenth-Century Telescope** By the mid-19th century, telescopes had improved enormously in both size and quality. Shown here is the Newtonian reflector built and used by Irish nobleman and amateur astronomer the Earl of Rosse. For 75 years, this 72-inch-diameter instrument was the largest telescope on Earth. (*Birr Scientific & Heritage Foundation*)



▲ **FIGURE 6.3 Lunar Exploration** An *Apollo* astronaut does some lunar geology—prospecting near a huge boulder in the Mare Serenitatis during the final manned mission to the Moon in 1972. (NASA)

kilometers in diameter, and countless *meteoroids* less than 100 m across. The list will undoubtedly grow as we continue to explore our cosmic neighborhood.

## Comparative Planetology

As we proceed through the solar system in the next few chapters, we will seek to understand how each planet compares with our own and what each contributes to our knowledge of the solar system as a whole. We will use the powerful perspective of **comparative planetology**—comparing and contrasting the properties of the diverse worlds we encounter—to understand better the conditions under which planets form and evolve. Comparative planetology will be our indispensable guide as we proceed through the coming chapters.

Our goal is to develop a comprehensive theory of the origin and evolution of our planetary system—a theory that explains all, or at least most, of the solar system’s observed properties. We will seek to answer basic questions such as Why did planet X evolve in one way, while planet Y turned out completely different? and Why are the planets’ orbits so orderly when their individual properties are not? In addressing these issues, we will find many similarities and common features among planets. However, each planet will also present new questions and afford unique insights into the ways planets work.

As we unravel the origin of our solar system, we may hope to learn something about planetary systems beyond our own. Since the mid-1990s, astronomers have detected more than 800 *extrasolar planets*—planets orbiting stars other than our own Sun. Many new planets are discovered each year (see Chapter 15), and our observations of them provide critical tests of modern theories of planet formation. Before the discovery of extrasolar planets, those theories had necessarily been based on observations of only our own solar system. Now astronomers have a whole new set of “proving grounds” in which to compare theory with reality.

Curiously, current data suggest that many of the newly discovered systems have properties rather different from those of our own, adding fuel to the long-standing debate among astronomers on the prevalence of planets like Earth and the possible existence of life as we know it elsewhere in the universe. It will be some time before astronomers can make definitive statements about the existence (or nonexistence) of planetary systems like our own.

### PROCESS OF SCIENCE Check

- ✓ In what ways might observations of extrasolar planets help us understand our own solar system?



▲ **FIGURE 6.4 Spirit on Mars** The Mars rover *Spirit* took hundreds of images to create this true-color, 360° panorama of the Martian horizon from within Gusev crater. The robot, whose tracks into the shallow basin can be seen at right center, also measured the chemistry and mineralogy of soils and rocky outcrops. (JPL)

## 6.2 Measuring the Planets

Table 6.1 lists some basic orbital and physical properties of the eight planets, with a few other well-known solar system objects (the Sun, the Moon, Pluto, an asteroid, and a comet) included for comparison. Note that the Sun, with more than a thousand times the mass of the next most massive object (the planet Jupiter), is clearly the dominant member of the solar system. In fact, the Sun contains about 99.9 percent of all solar system material. The planets—including our own—are insignificant in comparison.

Most of the quantities listed in Table 6.1 can be determined using methods described in Chapters 1 and 2. Here we present a brief summary of the properties listed in Table 6.1 and the techniques used to measure them:

- The *distance* of each planet from the Sun is known from Kepler's laws once the scale of the solar system is set by radar ranging on Venus. [∞ \(Sec. 2.6\)](#)
- A planet's sidereal *orbital period* (that is, relative to the stars) can be measured from repeated observations of its location on the sky, so long as Earth's own motion around the Sun is properly taken into account. [∞ \(Sec. 2.5\)](#)
- A planet's *radius* is found by measuring the angular size of planet—the angle from one side to the other as

we view it on the sky—and then applying elementary geometry. [∞ \(More Precisely 1-2\)](#)

- The *masses* of planets with moons may be calculated by applying Newton's laws of motion and gravity, just by observing the moons' orbits around the planets. [∞ \(More Precisely 2-2\)](#) The *sizes* of those orbits, like the sizes of the planets themselves, are determined by geometry.
- The masses of Mercury and Venus (as well as those of our own Moon and the asteroid Ceres) are a little harder to determine accurately, because these bodies have no natural satellites of their own. Nevertheless, it is possible to measure their masses by careful observations of their gravitational influence on other planets or nearby bodies. Mercury and Venus produce small, but measurable, effects on each other's orbits, as well as on that of Earth. The Moon also causes small “wobbles” in Earth's motion as the two bodies orbit their common center of mass.
- These techniques for determining mass were available to astronomers well over a century ago. Today, the masses of most of the objects listed in Table 6.1 have been accurately measured through their gravitational interaction with artificial satellites and space probes launched from Earth. Only in the case of Ceres is the mass still poorly known, mainly because that asteroid's gravity is so weak.

**TABLE 6.1** Properties of Some Solar System Objects

Object	Orbital Semimajor Axis (AU)	Orbital Period (Earth Years)	Mass (Earth Masses)	Radius (Earth Radii)	Number of Known Satellites	Rotation Period * (days)	Average Density (kg/m <sup>3</sup> )	(g/cm <sup>3</sup> )
Mercury	0.39	0.24	0.055	0.38	0	59	5400	5.4
Venus	0.72	0.62	0.82	0.95	0	−243	5200	5.2
Earth	1.0	1.0	1.0	1.0	1	1.0	5500	5.5
Moon	—	—	0.012	0.27	—	27.3	3300	3.3
Mars	1.52	1.9	0.11	0.53	2	1.0	3900	3.9
Ceres (asteroid)	2.8	4.7	0.00015	0.073	0	0.38	2700	2.7
Jupiter	5.2	11.9	318	11.2	63	0.41	1300	1.3
Saturn	9.5	29.4	95	9.5	56	0.44	700	0.7
Uranus	19.2	84	15	4.0	27	−0.72	1300	1.3
Neptune	30.1	164	17	3.9	13	0.67	1600	1.6
Pluto (Kuiper belt object)	39.5	248	0.002	0.2	3	−6.4	2100	2.1
Hale-Bopp (comet)	180	2400	$1.0 \times 10^{-9}$	0.004	—	0.47	100	0.1
Sun	—	—	332,000	109	—	25.8	1400	1.4

\*A negative rotation period indicates retrograde (backward) rotation relative to the sense in which all planets orbit the Sun.

- A planet's *rotation period* may, in principle, be determined simply by watching surface features alternately appear and disappear as the planet rotates. However, with most planets this is difficult to do, as their surfaces are hard to see or may even be nonexistent. Mercury's surface features are hard to distinguish; the surface of Venus is completely obscured by clouds; and Jupiter, Saturn, Uranus, and Neptune have no solid surfaces at all—their atmospheres simply thicken and eventually become liquid as we descend deeper and deeper below the visible clouds. We will describe the methods used to measure these planets' rotation periods in later chapters.
- The final two columns in Table 6.1 list the *average density* of each object. **Density** is a measure of the “compactness” of matter. Average density is computed by dividing an object's mass (in kilograms, say) by its volume (in cubic meters, for instance). For example, we can easily compute Earth's average density. Earth's mass, as determined from observations of the Moon's orbit, is approximately  $6.0 \times 10^{24}$  kg. [↪ \(More Precisely 2-2\)](#) Earth's radius  $R$  is roughly 6400 km, so its volume is  $\frac{4}{3}\pi R^3 \approx 1.1 \times 10^{12}$  km<sup>3</sup>, or  $1.1 \times 10^{21}$  m<sup>3</sup>. [↪ \(Sec. 1.6\)](#) Dividing Earth's mass by its volume, we obtain an average density of approximately 5500 kg/m<sup>3</sup>.


On average, then, there are about 5500 kilograms of Earth matter in every cubic meter of Earth volume. For comparison, the density of ordinary water is 1000 kg/m<sup>3</sup>, rocks on Earth's surface have densities in the range 2000–3000 kg/m<sup>3</sup>, and iron has a density of some 8000 kg/m<sup>3</sup>. Earth's atmosphere (at sea level) has a density of only a few kilograms per cubic meter. Because many working astronomers are more familiar with the CGS (centimeter-gram-second) unit of density (grams per cubic centimeter, abbreviated g/cm<sup>3</sup>, where 1 kg/m<sup>3</sup> = 1000 g/cm<sup>3</sup>), Table 6.1 lists density in both SI and CGS units.

### CONCEPT Check


- ✓ How do astronomers go about determining the bulk properties (i.e., masses, radii, and densities) of distant planets?

## 6.3 The Overall Layout of the Solar System

By earthly standards, the solar system is immense. The distance from the Sun to the Kuiper belt outside Neptune's orbit is about 50 AU, more than a million times Earth's radius and roughly 20,000 times the distance from Earth to the Moon (Figure 6.5). Yet, despite the solar system's vast extent, the planets all lie very close to the Sun, astronomically speaking. Even the diameter of the Kuiper belt is just



▲ **FIGURE 6.5 Solar System** Major bodies of the solar system include the Sun, planets, and asteroids. Except for Mercury, the orbits of the planets are almost circular (a) and lie nearly in the same plane (b). The distance between adjacent orbits increases farther from the Sun. The entire solar system spans nearly 100 AU—roughly the diameter of the Kuiper belt—and is very flat.



▲ **FIGURE 6.6 Planetary Alignment** This image shows six planets—Mercury, Venus, Mars, Jupiter, Saturn, and Earth—during a planetary alignment in April 2002. The Sun and Moon are just below the horizon. As usual, the popular press contained many sensationalized predictions of catastrophes that would occur during this rare astronomical event. Also as usual, none came true. (*J. Lodriguss*)

1/1000 of a light-year, whereas the next nearest star is several light-years distant.

The planet closest to the Sun is Mercury. Moving outward, we encounter, in turn, Venus, Earth, Mars, Jupiter, Saturn, Uranus, and Neptune. In Chapter 2, we saw the basic properties of the planets' orbits. Their paths are all ellipses, with the Sun at (or very near) one focus. ∞ (Sec. 2.5) With the exception of the innermost world, Mercury, the planets move on low-eccentricity orbits. Accordingly, we can reasonably think of most planets' orbits as circles centered on the Sun. Note that the planetary orbits are not evenly spaced, becoming farther and farther apart as we move outward from the Sun.

All the planets orbit the Sun counterclockwise as seen from above Earth's North Pole and in nearly the same plane as Earth (the plane of the ecliptic). Mercury deviates somewhat from the latter statement: Its orbital planes lie at  $7^\circ$  to the ecliptic. Still, as illustrated in the figure, we can think of the solar system as being quite flat—its “thickness” perpendicular to the plane of the ecliptic is a

tiny fraction of its diameter. If we were to view the planets' orbits from a vantage point in the plane of the ecliptic about 50 AU from the Sun, none of the planets' orbits would be noticeably tilted. Figure 6.6 is a photograph of the planets Mercury, Venus, Mars, Jupiter, and Saturn taken during a chance planetary alignment in April 2002. These five planets can (occasionally) be found in the same region of the sky, in large part because their orbits lie nearly in the same plane in space.

### CONCEPT Check

✓ In what sense is the solar system “flat”?

## 6.4 Terrestrial and Jovian Planets


On large scales, the solar system presents us with a sense of orderly motion. The planets move nearly in a plane, on almost concentric (and nearly circular) elliptical paths, in the same direction around the Sun, at steadily increasing orbital intervals. Although the individual details of the planets are much less regular, their overall properties allow a natural division into two broad classes.

### Planetary Properties

Figure 6.7 compares the planets with one another and with the Sun. A clear distinction can be drawn between the inner and the outer members of our planetary system based on densities and other physical properties. The inner planets—Mercury, Venus, Earth, and Mars—are small, dense, and *solid*. The outer worlds—Jupiter, Saturn, Uranus, and Neptune—are large, of low density, and *gaseous*.

Because the physical and chemical properties of Mercury, Venus, and Mars are somewhat similar to Earth's, the four innermost planets are called the **terrestrial planets**. (The word *terrestrial* derives from the Latin word *terra*, meaning “land” or “earth.”) The four terrestrial planets all lie within about 1.5 AU of the Sun. All are small and of relatively low mass—Earth is the largest and most massive of the four—and all have a generally *rocky* composition and solid surfaces.

Jupiter, Saturn, Uranus, and Neptune are all similar to one another chemically and physically, and very different from the terrestrial worlds. They are labeled the **jovian planets**, after Jupiter, the largest member of the group. (The word *jovian* comes from *Jove*, another name for the Roman god Jupiter.) The jovian worlds all orbit far from the Sun. They are all much larger than the inner planets and quite different from them in both composition and structure. They have no solid surfaces, and their outer layers are composed predominantly of the light gases *hydrogen* and *helium*.



◀ **FIGURE 6.7 Sun and Planets**

Relative sizes of the planets and our Sun, drawn to scale. Notice how much larger the jovian planets are than Earth and the other terrestrial planets, and how much larger still is the Sun. Explaining this planetary dichotomy is an important goal of comparative planetology, although by no means the only one.

## Differences Among the Terrestrial Planets

Despite their similarities in orbits and composition, the terrestrial planets have many important differences, too. We list a few here, as prelude to the more detailed accounts presented in Chapters 7–10:

- All four terrestrial planets have *atmospheres*, but the atmospheres are about as dissimilar as we could imagine, ranging from a near vacuum on Mercury to a hot, dense inferno on Venus.
- Earth alone has *oxygen* in its atmosphere and *liquid water* on its surface.
- *Surface conditions* on the four planets are quite distinct from one another, ranging from barren, heavily cratered terrain on Mercury to widespread volcanic activity on Venus.
- Earth and Mars *spin* at roughly the same rate—one rotation every 24 (Earth) hours—but Mercury and Venus both take months to rotate just once, and Venus rotates in the opposite sense from the others.
- Earth and Mars have *moons*, but Mercury and Venus do not.
- Earth and Mercury have measurable *magnetic fields*, of very different strengths, whereas Venus and Mars have none.

Comparing the average densities of the terrestrial planets allows us to say something more about their overall *compositions*. However, before making the comparison, we must take into account how the weight of overlying layers compresses the interiors of the planets to different extents. When we do this, we find that the *uncompressed densities* of the terrestrial worlds—the densities they would have in the absence of any compression due to their own

gravity—decrease as we move outward from the Sun: 5300, 4400, 4400, and 3800 kg/m<sup>3</sup> for Mercury, Venus, Earth, and Mars, respectively. The amount of compression is greatest for the most massive planets, Earth and Venus, and much less for Mercury and Mars. Partly on the basis of these figures, planetary scientists conclude that Earth and Venus are quite similar in overall composition. Mercury's higher density implies that it contains a higher proportion of some dense material—most likely nickel or iron. The lower density of Mars probably means that it is deficient in that same material.

Finding the common threads in the evolution of these four diverse worlds is no simple task! The goal of comparative planetology is to understand how four planets with broadly similar overall physical properties came to differ so much in detail.

## Terrestrial–Jovian Comparison

For all their differences, the terrestrial worlds still seem similar compared with the jovian planets. Perhaps the simplest way to express the major differences between the terrestrial and jovian worlds is to say that the jovian planets are everything the terrestrial planets are not. We will discuss the jovian planets in more detail in Chapters 11–13. We highlight here some of the key differences between the terrestrial and jovian worlds:

- The terrestrial worlds lie close together, near the Sun; the jovian worlds are widely spaced through the outer solar system.
- The terrestrial worlds are small, dense, and rocky; the jovian worlds are large and gaseous, containing huge amounts of hydrogen and helium (the lightest elements), which are rare on the inner planets.

## DISCOVERY 6-1


## Gravitational “Slingshots”

*Celestial mechanics*—the study of the motions of gravitationally interacting objects—is an essential tool for scientists and engineers who wish to navigate manned and unmanned spacecraft throughout the solar system. Robot probes can now be sent on stunningly accurate trajectories, expressed in the trade with such slang phrases as “sinking a corner shot on a billion-kilometer pool table.” Near-flawless rocket launches, aided by occasional midcourse changes in flight paths, now enable interplanetary navigators to steer remotely controlled spacecraft through an imaginary “window” of space just a few kilometers wide and a billion kilometers away.

However, sending a spacecraft to another planet requires a lot of energy—often more than can be conveniently provided by a rocket launched from Earth or safely transported in a shuttle for launch from orbit. Faced with these limitations, mission scientists often use their knowledge of celestial mechanics to carry out “slingshot” maneuvers, which can boost an interplanetary probe into a more energetic orbit and also aid navigation toward the target, all at no additional cost!


The first figure illustrates a gravitational slingshot, or *gravity assist*, in action. A spacecraft approaches a planet, passes close by, and then escapes along a new trajectory. Obviously, the spacecraft’s *direction* of motion is changed by the encounter. Less obviously, the spacecraft’s *speed* is also altered as the planet’s gravity propels the spacecraft in the direction of the planet’s motion. By a careful choice of incoming trajectory, the craft can speed up (by passing “behind” the planet, as shown) or slow down (by passing in front), by as much as twice the planet’s orbital speed. Of course, there is no free lunch—the spacecraft gains energy from, or loses it to, the planet’s motion, causing the planet’s own orbit to change ever so slightly. However, since planets are so much more massive than spacecraft, the effect on the planet is tiny.

Such a slingshot maneuver has been used many times in missions to both the inner and the outer planets, as illustrated in the second figure, which shows the trajectories of the *Voyager* spacecraft through the outer solar system (see Chapters 11–13). The gravitational pulls of these giant worlds whipped the craft around at each visitation, enabling flight controllers to get considerable extra “mileage” out of the probes. *Voyager 1* is now high above the plane of the solar system, having been



deflected up and out following its encounter with Saturn. *Voyager 2* continued on for a “Grand Tour” of the four jovian planets. It is now outside the orbit of Pluto, in the Kuiper belt. More recently, the *Cassini* mission to Saturn (Chapter 12) which was launched in 1997 and arrived at its target in 2004, received *four* gravitational assists en route—two from Venus, one from Earth, and one from Saturn. Once in the Saturn system, *Cassini* used the gravity of Saturn and its moons to propel it through a complex series of maneuvers designed to bring it close to all the major moons, as well as to the planet itself. NASA’s current *New Horizons* mission used a gravity assist from Jupiter to propel the spacecraft out to Pluto’s orbit.

Every encounter with every planet or moon had a slingshot effect—sometimes accelerating and sometimes slowing the probe, but each time moving it into a different orbit—and every one of these effects was carefully calculated long before the spacecraft ever left Earth.



- The terrestrial worlds have solid surfaces; the jovian worlds have none (their dense atmospheres thicken with depth, eventually merging with their liquid interiors).
- The terrestrial worlds have weak magnetic fields, if any; the jovian worlds all have strong magnetic fields.
- All four jovian worlds are thought to contain large, dense “terrestrial” cores some 10 to 15 times the mass of Earth. These cores account for an increasing fraction of each planet’s total mass as we move outward from the Sun.
- The terrestrial worlds have only three *moons* among them; the jovian worlds have many moons each, no two of them alike and none of them like our own.
- Furthermore, all the jovian planets have *rings*, a feature unknown on the terrestrial planets.

The job of the planetary scientist is complicated by the fact that the same theory that accounts for the jovian planets must also explain the terrestrial ones, as well as the similarities and differences within each class. Sections 6.6 and 6.7 present an outline of the theory that does just that. Later, in Chapter 15, we will return to this theory in more depth and see how its predictions fare when faced with detailed observations of our planetary system and others. ∞ (Sec. 1.2)

### CONCEPT Check

- ✓ Why do astronomers draw such a clear distinction between the inner and the outer planets?

## 6.5 Interplanetary Matter

In the vast space among the eight known major planets move countless chunks of rock and ice, mostly small, some quite large. All orbit the Sun, many on highly eccentric paths. This final component of the solar system is the collection of **interplanetary matter**—cosmic “debris” ranging in size from the relatively large *asteroids* and members of the *Kuiper belt*, through the smaller *comets* and even smaller *meteoroids*, down to the smallest grains of *interplanetary dust* that litter our cosmic environment (see Chapter 14).

The dust arises when larger bodies collide and break apart into smaller pieces that, in turn, collide again and are slowly ground into microscopic fragments, which eventually settle into the Sun or are swept away by the *solar wind*, a stream of energetic charged particles that continually flows outward from the Sun and pervades the entire solar system. The dust is quite difficult to detect in visible light, but infrared studies reveal that interplanetary space contains surprisingly large amounts of it. Our solar system is an extremely good vacuum by terrestrial standards, but positively dirty by the standards of interstellar or intergalactic space.

Asteroids (Figure 6.8a) and meteoroids are generally rocky in composition, somewhat like the outer layers of the terrestrial planets. The distinction between the two is simply a matter of size: Anything larger than 100 m in diameter (corresponding to a mass of about 10,000 tons) is conventionally termed an asteroid; anything smaller is a meteoroid. Their total mass is much less than that of Earth’s Moon, so these objects play no important role in the present-day workings of the planets or their moons. Yet they are of crucial importance to our studies, for they provide the keys to answering some very fundamental questions about our planetary environment and what the solar system was like soon after its birth. Many of these bodies are made of material that has hardly evolved since the early days of the solar system. In addition, they often conveniently deliver themselves right to our doorstep, in the form of meteorites (the name we give them if they happen to survive the plunge through Earth’s atmosphere and find their way to the ground—see Section 14.4), allowing us to study them in detail without having to fetch them from space.

Comets (Figure 6.8b) are quite distinct from the other small bodies in the solar system. They are generally icy rather than rocky in composition (although they do contain some rocky material) and typically have diameters in the 1- to 10-km range. They are quite similar in chemical makeup to some of the icy moons of the outer planets. Even more so than the asteroids and meteoroids, comets represent truly ancient material—the vast majority probably have not changed in any significant way since their formation long ago along with the rest of the solar system (see Chapter 15). Comets striking Earth’s atmosphere do not reach the surface intact, so, until relatively recently (see Section 14.2), astronomers had no actual samples of cometary material. However, some comets do vaporize and emit radiation as their highly elongated orbits take them near the Sun (see Figure 6.8b), so scientists have long been able to determine their makeup by spectroscopic study of the radiation they give off as they are destroyed. ∞ (Sec. 4.2)


Finally, beyond the outermost jovian planet, Neptune, lies the Kuiper belt—an “outer asteroid belt” consisting of icy, cometlike bodies ranging in size from a fraction of a kilometer to more than 1000 km in diameter. Because of their small sizes and great distances, most known Kuiper belt objects have been discovered only recently, mainly since the mid-1990s. However, the best known member of this class—Pluto (Figure 6.8c)—has been known for much longer and has been the subject of recent controversy among astronomers. It was originally classified as a planet following its discovery in 1930. However, it simply doesn’t fit into the classification just described—in both mass and composition, it has much more in common with the icy jovian moons than with any terrestrial or jovian world. As a result, many astronomers questioned its “planetary” designation,




► **FIGURE 6.8 Asteroid and Comet** (a) Asteroids, like meteoroids, are generally composed of rocky material. This asteroid, Vesta, is nearly 500 km across and orbits between Mars and Jupiter. It was photographed by the *Dawn* spacecraft in 2011. (b) Most comets are composed largely of ice and so tend to be relatively fragile. This is comet McNaught, seen over the Pacific Ocean in 2007, with its vaporized tail extending away from the Sun for nearly a quarter of the way across the sky. (c) Pluto—seen here at center with three of its moons—is one of the largest members of the Kuiper belt. Formerly classified as a major planet, it was demoted during a heated debate among astronomers in 2006. (JPL; ESO; NASA)



(b)




(c)



an opinion that became stronger as the number of known Kuiper belt objects steadily increased.

The controversy came to a head with the detection of several Kuiper belt objects comparable in size to Pluto, and at least one *larger* body orbiting even farther from the Sun. As discussed further in Section 14.3, in 2006 astronomers decided that Pluto should no longer be classified as a major



(a)

planet, but rather as a *dwarf planet*. It is the largest known object in the Kuiper belt, but is no longer regarded as a planet on a par with the terrestrial or jovian worlds.

### CONCEPT Check

✓ Why are astronomers so interested in interplanetary matter?

## 6.6 How Did the Solar System Form?

During the past five decades, interplanetary probes have vastly increased our knowledge of the solar system (see *Discovery 6-2*), and their data form the foundation for much of the discussion in the next eight chapters. However, we can understand the overall organization of our planetary system—the basic properties presented in the previous three sections—without dwelling on these details. Indeed, some key elements of the modern theory of planetary formation predate the Space Age by many years. We present here the “standard” view of how the solar system came into being. This picture will underlie much of our upcoming discussion of the planets, their moons, and the contents of the vast spaces between them.

### Solar System Properties

Any theory of the origin and architecture of our planetary system must adhere to the known facts. We know of eight outstanding properties of our solar system as a whole:

1. *Each planet is relatively isolated in space.* The planets orbit at progressively larger distances from the central Sun; they are not bunched together. Each planet is roughly twice as far from the Sun as its next inward neighbor.


2. *The orbits of the planets are nearly circular.* In fact, apart from Mercury, which we will argue later is a special case, each planetary orbit is close to a perfect circle.
3. *The orbits of the planets all lie in nearly the same plane.* The planes swept out by the planets' orbits are accurately aligned to within a few degrees. Again, Mercury is a slight exception.
4. *The direction in which the planets orbit the Sun (counterclockwise as viewed from above Earth's North Pole) is the same as the direction in which the Sun rotates on its axis.* Virtually all the large-scale motions in the solar system (other than comets' orbits) are in the same plane and in the same sense. The plane is that of the Sun's equator; the sense is that of the Sun's rotation.
5. *Our planetary system is highly differentiated.* The inner, terrestrial planets are characterized by high densities and moderate atmospheres. By contrast, the jovian planets, farther from the Sun, have low densities and thick atmospheres.
6. *The asteroids are very old and exhibit a range of properties not characteristic of either the inner or the outer planets or their moons.* The asteroid belt shares, in rough terms, the bulk orbital properties of the planets. However, it appears to be made of ancient material, and the meteorites that strike Earth are the oldest rocks known.
7. *The Kuiper belt is a collection of asteroid-sized icy bodies orbiting beyond Neptune.* Pluto is the largest known member of this class.
8. *The Oort cloud comets are primitive, icy fragments that do not orbit in the plane of the ecliptic and reside primarily at large distances from the Sun.* While similar to the Kuiper belt in composition, the Oort cloud is a completely distinct part of the outer solar system.

All these observed facts, taken together, strongly suggest a high degree of order within our solar system, at least on large scales. The whole system is not a random assortment of objects spinning or orbiting this way or that. Rather, the overall organization points toward a single origin—an ancient but one-time event that occurred long ago.

## Nebular Contraction

One of the earliest heliocentric models of solar system formation may be traced back to the 17th-century French philosopher René Descartes. Imagine a large cloud of interstellar dust and gas (called a *nebula*) a light-year or so across. Now suppose that, due to some external influence, such as a collision with another interstellar cloud or perhaps the explosion of a nearby star, the nebula starts to contract under the influence of its own gravity. As it contracts, it becomes denser and hotter, eventually forming a star—the Sun—at its center (see Chapter 19). Descartes suggested that, while the Sun was forming in the cloud's hot core, the planets and their moons formed in the cloud's cooler outer regions. In other words, planets are by-products of the process of star formation.

In 1796, the French mathematician–astronomer Pierre Simon de Laplace developed Descartes' ideas in a more quantitative way. He showed mathematically that conservation of angular momentum (see *More Precisely 6-1*) demands that our hypothetical nebula spin faster as it contracts. A decrease in the size of a rotating mass must be balanced by an increase in its rotational speed. The latter, in turn, causes the nebula's *shape* to change as it collapses. Centrifugal forces (due to rotation) tend to oppose the contraction in directions perpendicular to the rotation axis, with the result that the nebula collapses most rapidly along that axis. As shown in Figure 6.9, the fragment eventually flattens into a pancake-shaped primitive solar system. This



**Interactive FIGURE 6.9 Nebular Contraction** (a) Conservation of angular momentum demands that a contracting, rotating cloud must spin faster as its size decreases. (b) Eventually, a small part of it destined to become the solar system came to resemble a gigantic pancake. The large blob at the center ultimately became the Sun. (c) The planets that formed from the nebula inherited its rotation and flattened shape.


# DISCOVERY 6-2

## Spacecraft Exploration of the Solar System

Since the 1960s, dozens of uncrewed space missions have traveled throughout the solar system. All of the planets have been probed at close range, and robot spacecraft have also visited numerous comets and asteroids. The impact of these missions on our understanding of our planetary system has been nothing short of revolutionary. The 10 chapters of Part 2 of this text display many examples of images radioed back to Earth by the robots that have explored our nearby environment in space.

The time line stretching across these two pages shows all major missions since the dawn of the space age in the early 1960s. Together, they have redefined our view of our cosmic back yard. Here we highlight a few of the most important.


Mercury has been visited by just two spacecraft to date. In the mid-1970s, *Mariner 10* executed a series of flybys of the planet, snapping thousands of images




Mercury from *Messenger* (2012)

revealing it to be almost as heavily cratered as Earth's Moon. More than 30 years later, NASA's *Messenger* probe went into orbit around Mercury in 2011, where it is now actively mapping the surface of this peculiar place. The figure at left is a mosaic of *Messenger* images.


More spacecraft have explored Venus than any other planet. The Soviet Union took the lead in the 1960s when nearly a dozen *Venera* probes orbited (and some landed on) Venus—the photograph below shows Russian engineers building one of the heavily armored craft. Since then, the United States (*Pioneer* and *Magellan*) and Europe (*Venus Express*) have sent several more craft to spy on this hellish world using radar. The new data have taught us much about Venus's surface and atmosphere, and the results have helped us better understand weather here on planet Earth.



Russian engineers with *Venera 11* (1978)  
(Sovfoto/Eastfoto)




*Magellan* launched by Space Shuttle *Atlantis* (1989)



All photos NASA except where noted.

Mars has been the target of very active robotic exploration. The United States has sent more than a dozen probes to orbit the “red planet” and often to land on its surface; Russia and Europe have also aimed craft at Mars, but most have missed the planet or crash landed. Several U.S. *Mariner* craft paved the way in the 1960s, showing the planet



Artist conception of *Viking* on Mars (1976)

to be surprisingly inhospitable yet geologically intriguing. The *Viking* program in the 1970s was one of NASA's finest missions, not only safely landing two craft (see photo at left) but also imaging the surface for the first time and searching for life (which it did not find).



*Spirit on Mars* (2005)

Since the late 1990s, a series of robots has orbited Mars and invaded its surface, sampling the air and dirt, drilling into stones, and digging for ice. The above true-color panoramic view was taken in 2005 from the rover *Spirit*, whose mother ship is shown in the foreground. The latest generation of rovers has mainly been searching for water, or evidence of it in the past. “Follow the water” has become a popular mantra in the search for life. So far, though, neither liquid water nor life of any kind has been found, although there seems to be ample evidence for water in Mars’s past.


1995 Jupiter  
*Galileo* (NASA)

1997 Mars  
*Mars Pathfinder* (NASA)

1997 Mars  
*Mars Global Surveyor* (NASA)

1990 Venus  
*Magellan* (NASA)

Two pairs of U.S. spacecraft launched in the 1970s—*Pioneer* and *Voyager*—rewrote our knowledge of the jovian planets. *Pioneer 10* and *11* took many photographs and made numerous scientific discoveries. The *Voyager* spacecraft (below) carried radio, visible-light, and infrared sensors as well as magnetometers to study planetary magnetic fields. *Pioneer 11*, and both *Voyager 1* and *2*, used Jupiter’s gravity to propel them onward to Saturn (see *Discovery 6-1*). *Voyager 1* was programmed to visit Titan, Saturn’s largest moon, and so did not come close enough to the planet to receive a gravity-assisted boost to Uranus. However,



Artist conception of *Voyager 1* (1979)

2011 Asteroid Belt  
*Dawn* (NASA)

2012 Mars  
*Mars Science Laboratory* (NASA)

2006 Venus  
*Venus Express* (ESA)


2008 Mars: *Phoenix* (NASA)  
2008 Mercury: *Messenger* (NASA)

2006 Mars  
*Mars Reconnaissance Orbiter* (NASA)

2004 Saturn  
*Cassini-Huygens* (NASA/ESA)

*Voyager 2* went on to visit both Uranus and Neptune in a spectacularly successful “Grand Tour” of the outer planets.


During the past 2 decades, exploration of the outer solar system has been the domain of the U.S. *Galileo* mission to Jupiter (a 7-year flight that arrived in 1995) and the U.S./European *Cassini* mission to Saturn (arriving in 2004, with multiple gravity assists; see *Discovery 6-1*), both to reach their target planets and subsequently to navigate among those planets’ moons. *Galileo*, shown at right during assembly, toured Jupiter’s moons, transforming our understanding of their structure and history. *Cassini* is still gliding among Saturn’s moons, providing spectacular images of the Saturn system, as illustrated in the backlit image below, as well as key new insights into the Saturn system.



*Galileo* during assembly (1988)



Saturn from *Cassini* (2006)



▲ **FIGURE 6.10 Beta Pictoris** (a) False color is used here to accentuate our view of a disk of warm matter surrounding the star Beta Pictoris. The bottom image is a close-up of the inner part of the disk, whose warp might be caused by the gravitational pull of unseen companions. In both images, the bright central star has been masked to reveal the much fainter disk surrounding it. (b) This artist's conception of the disk of clumped matter shows the warm disk with a young star at the center and several comet-sized or larger bodies already forming. Mottled dust pervades this scene since such protoplanetary regions are probably very “dirty.” (NASA; D. Berry)

swirling mass destined to become our solar system is usually referred to as the **solar nebula**.

If we now simply suppose that the planets formed out of this spinning material, with each planet accumulating much of the material in its vicinity, then we can already understand the origin of much of the large-scale architecture observed in our planetary system today, such as the isolation and circularity of the planets' orbits and the fact that they move in the same sense in nearly the same plane (Figure 6.9c). The planets inherited these properties (items 1 to 4 on our earlier list) from the rotating disk in which they were born. The idea that the planets formed from such a disk is called the **nebular theory**.

Astronomers are confident that the solar nebula formed such a disk because we see similar disks around other stars. Figure 6.10(a) shows a visible-light image of the region around the star Beta Pictoris, some 50 light-years from the Sun. When the light from the star itself is removed and the resulting image enhanced by a computer, a faint disk of matter (viewed almost edge-on here) can be seen. It is roughly 1000 AU across—about 10 times the diameter of the Kuiper belt. Astronomers think that we are witnessing Beta Pictoris pass through a formative stage similar to that experienced by our own Sun 4.6 billion years ago. Figure 6.10(b) shows an artist's conception of the disk. We assume that *all* planetary systems pass through this phase early in their evolution.

## The Condensation Theory

Scientific theories must continually be tested and refined as new data become available. ∞ (Sec. 1.2) Unfortunately for Laplace's nebular theory, although its description of the collapse and flattening of the solar nebula was basically correct, we now know that a disk of warm gas would *not* form

clumps of matter that would subsequently evolve into planets. In fact, modern computer calculations predict just the opposite: Clumps in the gas would tend to disperse, not contract further. However, the **condensation theory**—the model favored by most astronomers—rests squarely on the old nebular theory, combining its basic physical reasoning with new information about interstellar chemistry to avoid most of the original theory's problems.

The key new ingredient is the presence of *interstellar dust* in the solar nebula. Astronomers now recognize that the space between the stars is strewn with microscopic dust grains, an accumulation of the ejected matter of many long-dead stars (see Chapter 22). These dust particles probably formed in the cool atmospheres of old stars and then grew by accumulating more atoms and molecules from the interstellar gas within the Milky Way Galaxy. The end result is that our entire Galaxy is littered with miniature chunks of icy and rocky matter, having typical sizes of about  $10^{-5}$  m. Figure 6.11 shows one of many such dusty regions found in the vicinity of the Sun.

Dust grains play two important roles in the evolution of a gas cloud. First, dust helps to cool warm matter by efficiently radiating its heat away in the form of infrared radiation. ∞ (Sec. 3.4) As the cloud cools, its molecules move more slowly, reducing the internal pressure and allowing the nebula to collapse more easily under the influence of gravity. ∞ (More Precisely 3-1) Second, by acting as **condensation nuclei**—microscopic platforms to which other atoms can attach, forming larger and larger balls of matter—the dust grains greatly speed up the process of collecting enough atoms to form a planet. This is similar to the way raindrops form in Earth's atmosphere: Dust and soot in the air act as condensation nuclei around which water molecules cluster.

## MORE PRECISELY 6-1

### Angular Momentum

Most celestial objects rotate. Planets, moons, stars, and galaxies all have some **angular momentum**, which we can define as the tendency of a body to keep spinning or moving in a circle. Angular momentum is as important a property of an object as its mass or its energy.

Consider first a simpler motion—*linear momentum*, which is defined as the product of an object's mass and its velocity:

$$\text{linear momentum} = \text{mass} \times \text{velocity.}$$


Linear momentum is the tendency of an object to keep moving in a straight line in the absence of external forces. Picture a truck and a bicycle rolling equally fast down a street. Each has some linear momentum, but you would obviously find it easier to stop the less massive bicycle. Although the two vehicles have the same speed, the truck has more momentum. We see, then, that the linear momentum of an object depends on the mass of the object. It also depends on the speed of the object: If two bicycles were rolling down the street at different speeds, the slower one could be stopped more easily.


Angular momentum is an analogous property of objects that are rotating or revolving. It is a measure of the object's tendency to keep spinning, or, equivalently, of how much effort must be expended to stop the object from spinning. However, in addition to mass and (angular) speed, angular momentum also depends on the way in which an object's mass is distributed.

Intuitively, we know that the more massive an object, or the larger it is, or the faster it spins, the harder it is to stop. In fact, angular momentum depends on the object's *mass*, *rotation rate* (measured in, say, revolutions per second), and *radius*, in a very specific way:

$$\text{angular momentum} \propto \text{mass} \times \text{rotation rate} \times \text{radius}^2.$$

(Recall that the symbol  $\propto$  means “is proportional to”; the constant of proportionality depends on the details of how the object's mass is distributed.)

According to Newton's laws of motion, both types of momentum—linear and angular—must be *conserved* at all times.  (Sec. 2.8) In other words, both linear and angular momentum must remain *constant* before, during, and after a physical change in any object (so long as no external forces act on the object). For example, as illustrated in the first figure, if a spherical object having some spin begins to contract, the previous relationship demands that it spin faster so that the product  $\text{mass} \times \text{angular speed} \times \text{radius}^2$  remains constant. The sphere's mass



(Orban/Corbis/Sygma)

does not change during the contraction, yet the size of the object clearly decreases. Its rotation speed must therefore increase in order to keep the total angular momentum unchanged. This constancy is referred to as **conservation of angular momentum**.


Figure skaters use the principle of conservation of angular momentum, too. They spin faster by drawing in their arms (as shown in the second pair of figures) and slow down by extending them. Here, the mass of the human body remains the same, but its lateral size changes, causing the body's rotation speed to increase or decrease, as the case may be, to keep its angular momentum unchanged.

**EXAMPLE 1** Suppose the sphere has radius 1 m and starts off rotating at 1 revolution per minute. It then contracts to one-tenth its initial size. Conservation of angular momentum entails that the sphere's final angular speed  $A$  must satisfy the relationship

$$\text{mass} \times A \times (0.1 \text{ m})^2 = \text{mass} \times (1 \text{ rev/min}) \times (1 \text{ m})^2.$$

The mass is the same on either side of the equation and therefore cancels, so we find that  $A = (1 \text{ rev/min}) \times (1 \text{ m})^2 / (0.1 \text{ m})^2 = 100 \text{ rev/min}$ , or about 1.7 rev/s.

**EXAMPLE 2** Now suppose that the “sphere” is a large interstellar gas cloud that is about to collapse and form the solar nebula. Initially, let's imagine that it has a diameter of 1 light-year and that it rotates very slowly—once every 10 million years. Assuming that the cloud's mass stays constant, its rotation rate must *increase* to conserve angular momentum as the radius decreases. By the time it has collapsed to a diameter of 100 AU, the cloud has shrunk by a factor of  $(1 \text{ light-year}/100 \text{ AU}) \approx 630$ . Conservation of angular momentum then implies that the cloud's (average) spin rate has increased by a factor of  $630^2 \approx 400,000$ , to roughly 1 revolution every 25 years—about the orbital period of Saturn.



◀ **FIGURE 6.11 Dark Cloud** Interstellar gas and dark dust lanes mark this region of star formation. The dark cloud known as Barnard 86 (dark, empty space at left) flanks a cluster of young blue stars called NGC 6520 (right). Barnard 86 may be part of a larger interstellar cloud that gave rise to these stars. (*D. Malin/Anglo-Australian Telescope*)

The formation of condensation nuclei greatly hastened the critical process of forming the first small clumps of matter. Once these clumps formed, they grew rapidly by sticking to other clumps. (Imagine a snowball thrown through a fierce snowstorm, growing bigger as it encounters more snowflakes.) As the clumps grew larger, their surface areas increased and, consequently, the rate at which they swept up new material accelerated. Gradually, this process of **accretion**—the gradual growth of small objects by collision and sticking—produced objects of pebble size, baseball size, basketball size, and larger.

Simulations indicate that, in perhaps as little as 100,000 years, accretion resulted in objects a few hundred kilometers across the size of small moons. By that time, their gravitational pulls had become just strong enough to affect their neighbors. Astronomers call these objects **planetesimals**—the building blocks of the solar system. Figure 6.12 shows an infrared view of a relatively nearby star thought to be surrounded by a disk in which planetesimals are growing.

The planetesimals' gravity was now strong enough to sweep up material that would otherwise not have collided with them, and their rate of growth accelerated, allowing them to form still larger objects (Figures 6.13a and b). Because larger bodies have stronger gravity, eventually almost all the original material was swept up into a few large **protoplanets**—the accumulations of matter that would in time evolve into the planets we know today (Figure 6.13c). Notice how, as the number of bodies decreases, the orbits of the remainder become more widely spaced and more nearly circular. The fact that this particular simulation produced exactly four terrestrial planets is pure chance—the details of the accretion process are random. However, regardless of the precise number of planets formed, the computer models do generally reproduce both the planets' approximately circular orbits and their increasing orbital spacing as we move outward from the Sun.


As the protoplanets grew, a competing process became important. Their strong gravitational fields produced many

high-speed collisions between planetesimals and protoplanets. These collisions led to **fragmentation** as small objects broke into still smaller chunks that were then swept up by the larger protoplanets. Only a relatively small number of 10- to 100-km fragments escaped capture by a planet or a moon and became the asteroids and comets. After about 100 million years, the primitive solar system had evolved into eight protoplanets, dozens of protomoons, and a glowing **protosun** at the center.


Roughly a billion more years were required to sweep the system clear of interplanetary trash. This was a period of intense meteoritic bombardment whose effects on the Moon and elsewhere are still evident today.

## Differentiation of the Solar System

The condensation theory just described can account—in broad terms, at least—for the formation of the planets and the large-scale architecture of the solar system. What does



▲ **FIGURE 6.12 Newborn Solar System?** This infrared image, taken by the *Spitzer Space Telescope*, of the bright star Fomalhaut, some 25 light-years from Earth, shows a circumstellar disk in which the process of accretion is underway. The star itself is well inside the yellowish blob at center. The outer disk, which is falsely colored orange to match the cooler dust emission, is about three times the diameter of our solar system. (*NASA*)



◀ **FIGURE 6.13**  
**Accreting Planets**

Initially in the inner solar system, many moon-sized planetesimals orbited the Sun. Gradually, they collided and coalesced, forming a few large planets in roughly circular orbits.


it say about the differences between the terrestrial and the jovian planets? To understand why a planet's composition depends on its location in the solar system, we must consider the *temperature* of the solar nebula.

As the primitive solar system contracted under the influence of gravity, it heated up as it flattened into a disk. The density and temperature were greatest near the center and much lower in the outlying regions. Detailed calculations indicate that the gas temperature near the core of the contracting system was several thousand kelvins. At a distance of 10 AU, out where Saturn now resides, the temperature was only about 100 K.


The high temperatures in the warmer regions of the cloud caused dust grains to break apart into molecules, which in turn split into excited atoms. Because the extent to which the dust was destroyed depended on temperature, it therefore also depended on location in the solar nebula. Most of the original dust in the inner solar system disappeared at this stage, whereas the grains in the outermost parts probably remained largely intact.

As the dusty nebula radiated away its heat, its temperature decreased everywhere except in the very core, where the Sun was forming. As the gas cooled, new dust grains began to condense (or crystallize) out, much as raindrops, snowflakes, and hailstones condense from moist, cooling air here on Earth. It may seem strange that although there was plenty of interstellar dust early on, it was partly destroyed, only to form again later. However, a critical change had occurred. Initially, the nebular gas was uniformly peppered with dust grains of all compositions; when the dust re-formed later, the distribution of grains was very different.

Figure 6.14 plots the temperature in various parts of the primitive solar system just before accretion began. At any given location, the only materials to condense out were those able to survive the temperature there. As marked on



(a)



(b)

▲ **FIGURE 6.14** Temperature in the Early Solar Nebula

(a) Theoretically computed variation of temperature across the primitive solar nebula illustrated in (b), which shows half of an accreting disk (see Figure 6.15). Note the distinction between the inner rocky grains (colored here red and black) and the outer icy grains (blue).



the figure, in the innermost regions, around Mercury's present orbit, only metallic grains could form; it was simply too hot for anything else to exist. A little farther out, at about 1 AU, it was possible for rocky, silicate grains to form, too. Beyond about 3 or 4 AU, water ice could exist, and so on, with the condensation of more and more material possible at greater and greater distances from the Sun.

In the inner regions of the primitive solar system, condensation from gas to solid began when the average temperature was about 1000 K. The environment there was too hot for ice to survive. Many of the more abundant heavier elements, such as silicon, iron, magnesium, and aluminum, combined with oxygen to produce a variety of rocky materials. The dust grains in the inner solar system were therefore predominantly rocky or metallic in nature, as were the protoplanets and planets they eventually became.

In the middle and outer regions of the primitive planetary system, beyond about 5 AU from the center, the temperature was low enough for the condensation of several abundant gases—water (H<sub>2</sub>O), ammonia (NH<sub>3</sub>), and methane (CH<sub>4</sub>)—into solid form. After hydrogen (H) and helium (He), the elements carbon (C), nitrogen (N), and oxygen (O) are the most common materials in the universe. As a result, wherever icy grains could form, they greatly outnumbered the rocky and metallic particles that condensed out of the solar nebula at the same location. Consequently, the objects that formed at these distances were formed under cold conditions out of predominantly low-density, icy material. These ancestral fragments were destined to form the cores of the jovian planets.

Note that, in this scenario, the composition of the outer solar system is much more typical of the universe as a whole than are the rocky and metallic inner planets. The outer solar system is not deficient in heavy elements. Rather, because of the conditions under which it formed, the inner solar system is *underrepresented in light material*.

### CONCEPT Check


- ✓ How does the condensation theory account for the circular, planar orbits of the planets and the broad differences between terrestrial and the jovian worlds?

## 6.7 Jovian Planets and Planetary Debris



The condensation-accretion picture just described has become the accepted model for the formation of the terrestrial planets. However, although similar processes may also have occurred in the outer solar system, the origin of the giant jovian worlds is decidedly less clear. Two somewhat different views have emerged.

## Making the Jovian Worlds

The first, more conventional, scenario is illustrated in Figure 6.15. With raw material readily available in the form of abundant icy grains, protoplanets in the outer solar system grew quickly and soon became massive enough for their strong gravitational fields to capture large amounts of gas directly from the solar nebula. In this view, called the **core-accretion theory**, four large protoplanets became



▲ **FIGURE 6.15 Solar System Formation** The condensation theory of planet formation (not drawn to scale). (a) The solar nebula after it has contracted and flattened to form a spinning disk (Figure 6.9b). The large red blob in the center will become the Sun. Smaller blobs in the outer regions may become jovian planets. (b) Dust grains act as condensation nuclei, forming clumps of matter that collide, stick together, and grow into moon-size planetesimals. The composition of the grains and thus of the planetesimals depends on location within the nebula. (c) After a few million years, strong winds from the still-forming Sun begin expelling nebular gas, and some massive planetesimals in the outer solar system have already captured gas from the nebula. (d) With the gas ejected, planetesimals continue to collide and grow. The gas giant planets are already formed, and the Sun has become a genuine star. (e) Over the course of a hundred million years or so, planetesimals are accreted or ejected, leaving a few large planets that travel in roughly circular orbits.



◀ **FIGURE 6.16**  
**T Tauri Star**

(a) Strong stellar winds from the newborn Sun sweep away the gas disk of the solar nebula, (b) leaving only giant planets and planetesimals behind. This stage of stellar evolution occurs only a few million years after the formation of the nebula.

the cores of the jovian worlds; the captured gas became their thick atmospheres. The smaller, inner protoplanets never reached this stage—another reason why their masses remained relatively low.


Recently, some astronomers have highlighted a potentially serious snag in this picture: There may not have been enough *time* for these events to have taken place. Most young stars apparently go through a highly active evolutionary stage known as the *T Tauri* phase (see Chapter 19), in which their radiation and stellar winds become very intense. During this period, much of the nebular gas between the planets was blown away into interstellar space (Figure 6.16).

The problem is that the nebular disk was at most a few million years old when all this occurred, leaving very little time for the large jovian cores to grow and capture gas from the nebula before it was destroyed. Furthermore, some researchers argue that, in the relatively dense stellar environments in which most stars are born (see Chapter 19), close encounters between still-forming stars may destroy many disks even sooner than that, giving giant planets perhaps as little as a few hundred thousand years in which to form.

The second formation scenario suggests that the giant planets formed through *instabilities* in the cool outer regions of the solar nebula, where portions of the cloud began to collapse under their own gravity—a picture not so far removed from Laplace's original idea—mimicking, on small scales, the collapse of the initial interstellar cloud. In this alternative **gravitational instability theory**, illustrated in Figure 6.17, the jovian protoplanets formed directly from the nebular gas, skipping the initial condensation-and-accretion stage and perhaps taking no more than a thousand years to acquire much of their mass. Right from the start, these first protoplanets had gravitational fields strong enough to scoop up more gas and dust from the solar

nebula, allowing them to grow into the giants we see today before the gas supply dispersed.

If both these theories eventually lead to gas-rich jovian planets, how can we distinguish between them? One possible way involves the composition of their cores. Because the planets formed so quickly in the instability



**Interactive FIGURE 6.17 Jovian Condensation** As an alternative to the growth of massive protoplanetary cores followed by the accretion of nebular gas, some or all of the giant planets might have formed directly through instabilities in the cool gas of the outer solar nebula. Part (a) shows the same instant as Figure 6.15(a). (b) Only a few thousand years later, four gas giants have already formed (red blobs), circumventing the accretion process sketched in Figure 6.15(b) and (c). With the nebula gone (c), the giant planets have taken their place in the outer solar system. (See Figure 6.15d.)

theory, computer models suggest that their cores should contain no more than about six Earth masses of rocky material. The core-accretion theory, by contrast, predicts much larger core masses—up to 20 times that of Earth. Detailed measurements of the jovian interiors by future space missions could settle the argument.


Many of the moons of the jovian planets presumably also formed through accretion, but on a smaller scale, in the gravitational fields of their parent planets. Once the nebular gas began to accrete onto the large jovian protoplanets, conditions probably resembled a miniature solar nebula, with condensation and accretion continuing to occur. The large moons of the outer planets almost certainly formed in this way. The smaller moons are more likely captured planetesimals.

## Giant-Planet Migration

Many aspects of the formation of the giant planets remain unresolved. Interactions among the growing planets, and between the planets and their environment, probably played a critical role in determining just how and where the planets formed. One particularly intriguing scenario is the possibility that Jupiter—and maybe all four giant planets—formed considerably farther from the Sun than its present orbit and subsequently “migrated” inward. This supposed migration is indicated schematically in Figures 6.15 and 6.17 by the changing locations of the jovian protoplanets.

The idea of planetary migration has been around since the mid-1980s, when theorists realized that friction between massive planets and the nebula in which they moved would have caused just such an inward drift. Observational support came in 1999, when *Galileo* scientists (see *Discovery* 6-2) announced much higher than expected concentrations of the gases nitrogen, argon, krypton, and xenon in Jupiter’s atmosphere. These gases, which are thought to have been carried to the planet by captured planetesimals, could not have been retained in the planetesimal ice at temperatures typical of Jupiter’s current orbit. Instead, they imply that the planetesimals—and, presumably, Jupiter too—formed at much lower temperatures. Either the nebula was cooler than previously thought, or Jupiter formed out in what is now the Kuiper belt!

The events just described did not take long, astronomically speaking. The giant planets formed within a few million years of the appearance of the flattened solar nebula—the blink of an eye compared with the 4.6-billion-year age of the solar system. At that point, intense radiation and strong winds from violent activity on the surface of the newborn Sun ejected the nebular gas, halting further growth. Accretion in the inner solar system proceeded more slowly, taking perhaps 100 million years to form the planets we know today (Figure 6.15e). The rocky asteroids and icy comets are all that remain of the matter that originally condensed out of the solar nebula—the last surviving witnesses to the birth of our planetary system. To place all these formative processes in perspective, Figure 6.18 presents a simplified time line of the first billion years after the formation of the solar nebula.



◀ **FIGURE 6.18** Solar System Formation

Schematic time line of some key events occurring during the first billion years of our solar system. The various tracks show the evolution of the Sun and the solar nebula, as well as that of the inner and outer solar system. Note that the tracks are intended to illustrate approximate relationships between events, not the precise times at which they occurred. Planetary scientists think that all of the processes represented here should have occurred in other planetary systems, too.

**CONCEPT Check**

✓ Why is the rate at which the Sun formed important in a theory of the formation of the jovian planets?

**Asteroids and Comets**


In the inner solar system, planetesimal fragments that escaped capture by one of the terrestrial planets received repeated “gravity assists” from those bodies and were eventually boosted beyond the orbit of Mars (see *Discovery 6-1*). Roughly a billion years were required to sweep the inner solar system clear of interplanetary “trash.” The myriad rocks of the asteroid belt between Mars and Jupiter failed to accumulate into a planet. Probably, nearby Jupiter’s huge gravitational field caused them to collide too destructively to coalesce, or otherwise hindered the development of a protoplanet. The result is a band of rocky planetesimals, still colliding and occasionally fragmenting, but never coalescing into a larger body.

In the outer solar system, with the formation of the four giant jovian planets, the remaining planetesimals were subject to those planets’ strong gravitational fields. Over a period of hundreds of millions of years, interactions with the giant planets, especially Uranus and Neptune, flung many of the outer region’s interplanetary fragments into

orbits taking them far from the Sun (Figure 6.19). Astronomers think that those icy bodies now make up the Oort cloud, whose members occasionally visit the inner solar system as comets.

A key prediction of this model is that some of the original planetesimals remained behind, forming the broad band known as the Kuiper belt, lying beyond the orbit of Neptune, some 30 to 40 AU from the Sun. More than 1200 Kuiper belt objects, having diameters ranging from 50 km to more than 2000 km, are now known. Their existence lends strong support to the condensation theory of planetary formation.

Computer simulations reveal that the ejection of the planetesimals involved a remarkably complex interplay among the jovian planets, whereby Uranus and Neptune kicked some bodies out into the Kuiper belt but deflected others *inward* toward Jupiter and Saturn, whose strong gravitational fields then propelled the planetesimals out into the distant Oort cloud. As shown in Figure 6.19, the orbits of all four giant planets were significantly modified by these interactions. By the time the outer solar system had been cleared of comets, Jupiter had moved slightly closer to the Sun, its orbital semimajor axis decreasing by a few tenths of an AU. The other giant planets moved outward—Saturn by about 1 AU, Uranus by 3 or 4 AU, and Neptune by some 7–10 AU. Note that these orbital changes occurred long *after*



**▲ FIGURE 6.19 Planetesimal Ejection** The ejection of icy planetesimals help to form the Oort cloud and Kuiper belt. (a) Initially, once the giant planets had formed, leftover planetesimals were found throughout the solar system. Interactions with Jupiter and Saturn apparently “kicked” planetesimals out to very large radii (the Oort cloud). Interactions with Uranus and especially Neptune tended to keep the Kuiper belt populated, but also deflected many planetesimals inward to interact with Jupiter and Saturn. (b) After hundreds of millions of years and as a result of the inward and outward “traffic,” the orbits of all four giant planets were significantly modified by the time the planetesimals inside Neptune’s orbit had been ejected. As depicted here, Neptune was affected most and may have moved outward by as much as 10 AU.

the supposed inward migrations mentioned earlier. Life as a jovian planet is far from simple!

During this period, many icy planetesimals were also deflected into the inner solar system, where they played an important role in the evolution of the inner planets. A long-standing puzzle in the condensation theory's account of the formation of the inner planets has been where the water and other volatile gases on Earth and elsewhere originated. At the inner planets' formation, their surface temperatures were far too high, and their gravity too low, to capture or retain those gases. The most likely explanation seems to be that the water and other light gases found on Earth and

elsewhere in the inner solar system arrived there in the form of comets from the outer solar system. Kicked into eccentric orbits as the gravitational fields of the jovian planets cleared the outer solar system of leftover planetesimals, these icy fragments bombarded the newborn terrestrial worlds, supplying them with water *after* their formation.

### CONCEPT Check

✓ Would you expect to find comets and asteroids orbiting other stars?

**The Big Question** Hardly a decade ago, astronomers regarded our solar system as ordinary—typical of any other planetary system that might exist beyond Earth. The Copernican principle is a powerful idea, and we thought our solar system was nothing special. However, now that alien planets are being discovered around other stars, as we will see in Chapter 15, we are no longer sure how “normal” our solar system actually is. A basic question lingers: How common or different is our system of planets compared to all those planets orbiting other stars in the universe?


## Chapter Review

### SUMMARY


**1** The **solar system** (p. 136) consists of the Sun and everything that orbits it, including the eight major planets, the moons that orbit them, and the many small bodies found in interplanetary space. **Comparative planetology** (p. 137) contrasts the properties of the diverse bodies found in the solar system and elsewhere to understand the conditions under which planets form and develop.



**2** The major planets orbit the Sun in the same sense—counterclockwise as viewed from above Earth's North Pole—on roughly circular orbits that lie close to the plane of the ecliptic. The orbit of the innermost planet, Mercury, is the most eccentric and has the greatest orbital inclination. The spacing between planetary orbits increases as we move outward from the Sun. The diameter of Neptune's orbit is roughly 60 AU.



**3** The average **density** (p. 139) of a planet is obtained by dividing the planet's total mass by its volume. The innermost four planets in the solar system—Mercury, Venus, Earth, and Mars—have average densities comparable




to Earth's and are generally rocky in composition. They are called the **terrestrial planets** (p. 140). The outer **jovian planets** (p. 140)—Jupiter, Saturn, Uranus, and Neptune—have much lower densities than the terrestrial worlds and are made up mostly of gaseous or liquid hydrogen and helium. Compared with the terrestrial worlds, the jovian planets are larger and more massive, rotate more rapidly, and have stronger magnetic fields. In addition, the jovian planets all have ring systems and many moons orbiting them.

**4** The asteroids, or “minor planets,” are small bodies, none of them larger than Earth's Moon and most of which orbit in a broad band called the asteroid belt between the orbits of Mars and Jupiter. Comets are chunks of ice found chiefly in the outer solar system. Their importance to planetary astronomy lies in the fact that they are thought to be “leftover” material from the formation of the solar system and therefore contain clues to the very earliest stages of its development. The Kuiper belt is a band of icy bodies orbiting beyond the orbit of Neptune.




**5** According to the **nebular theory** (p. 148), a large cloud of dust and gas began to collapse under its own gravity. As it did so, it began to spin faster, to conserve angular




momentum, eventually forming a rotating disk—the **solar nebula** (p. 148)—out of which the planets arose. The **condensation theory** (p. 148) builds on the nebular theory by including the effects of dust and solar heating on planet formation. The temperature at any given location in the solar nebula determined which materials could condense out there, and hence also the composition of the **planetesimals** (p. 150) that are forming. The terrestrial planets are rocky because they formed in the hot inner regions of the solar nebula, near the Sun, where only rocky and metallic materials condensed out. Farther out, the nebula was cooler, and ices could also form.

**6** Rocky and metallic dust grains condensed out in the inner solar system. The terrestrial planets formed by accretion and merger of these planetesimals during the first 100 million years of the solar system. A few large bodies came to dominate, eventually becoming the four terrestrial planets we see today. The final stages of this process were likely marked by catastrophic collisions between planet-sized bodies.




**7** In the outer solar system, the nebula was cooler, and ices of water and ammonia could also form. According to the **gravitational instability theory** (p. 153), the jovian planets formed

directly and very rapidly through instabilities in the nebular disk. In the more standard **core-accretion theory** (p. 152), icy protoplanet cores became so large that they could capture hydrogen and helium gas from the nebula. Before the nebula was ejected by strong winds from the still-forming Sun, a few million years after the nebula formed, interactions between the giant planets and the gas probably caused the former to migrate inward from their initial orbits.



**8** The asteroid belt is a collection of rocky planetesimals that never coalesced into a planet because of Jupiter's strong gravity. Many leftover planetesimals in the outer solar system were ejected into the Kuiper belt and the Oort cloud by the gravitational fields of the outer planets. Some occasionally still visit the inner solar system as comets. The expulsion of the icy planetesimals may have significantly changed the giant planets' orbits; Uranus and Neptune probably migrated outward during this period. Much of Earth's water was carried to our world by planetesimals deflected from the outer solar system.



**MasteringAstronomy**® For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- LO1 POS** What is comparative planetology? Why is it useful? What is its ultimate goal?
- LO2** Name and describe all the different types of objects found in the solar system. Give one distinguishing characteristic of each. Include a mention of interplanetary space.
- Why is it necessary to know the distance to a planet in order to determine the planet's mass?
- LO3** Name three important differences between the terrestrial planets and the jovian planets.
- LO4 POS** Why are asteroids and meteoroids important to planetary scientists?
- Comets generally vaporize upon striking Earth's atmosphere. How, then, do we know their composition?
- LO5 POS** What is the key ingredient in the modern condensation theory of the solar system's origin that was missing or unknown in the nebular theory?
- Give three examples of how the condensation theory explains the observed features of the present-day solar system.
- LO6** Describe how the terrestrial planets formed.
- Why are the jovian planets so much more massive than the terrestrial planets?
- How did the temperature structure of the solar nebula determine planetary composition?
- LO7** Describe two possible ways in which the jovian planets may have formed. What role did the Sun play in the process?
- What happened to the outer planets as the solar system was cleared of icy planetesimals?
- LO8** How did the Kuiper belt and the Oort cloud form?
- Describe a possible history of a single comet now visible from Earth, starting with its birth in the solar nebula somewhere near the orbit of Jupiter.

## Conceptual Self-Test: Multiple Choice

1. A planet's mass can most easily be determined by measuring the planet's (a) moon's orbits; (b) angular diameter; (c) position in the sky; (d) orbital speed around the Sun.
2. If we were to construct an accurate scale model of the solar system on a football field with the Sun at one end and Neptune at the other, the planet closest to the center of the field would be (a) Earth; (b) Jupiter; (c) Saturn; (d) Uranus.
3. The inner planets tend to have (a) fewer moons; (b) faster rotation rates; (c) stronger magnetic fields; (d) higher gravity than the outer planets have.
4. A solar system object of rocky composition and comparable in size to a small city is most likely (a) a meteoroid; (b) a comet; (c) an asteroid; (d) a planet.
5. The asteroids are mostly (a) found between Mars and Jupiter; (b) just like other planets, only younger; (c) just like other planets, only smaller; (d) found at the very edge of our solar system.
6. In the leading theory of solar system formation, the planets (a) were ejected from the Sun following a close encounter with another star; (b) formed from the same flattened, swirling gas cloud that formed the Sun; (c) are much younger than the Sun; (d) are much older than the Sun.
7. The inner planets formed (a) when the Sun's heat destroyed all the smaller bodies in the inner solar system; (b) in the outer solar system and then were deflected inward by interactions with Jupiter and Saturn; (c) by collisions and mergers of planetesimals; (d) when a larger planet broke into pieces.
8. The solar system is differentiated because (a) all the heavy elements in the outer solar system have sunk to the center; (b) all the light elements in the inner solar system became part of the Sun; (c) all the light elements in the inner solar system were carried off in the form of comets; (d) only rocky and metallic particles could form close to the Sun.
9. **VIS** According to Figure 6.14, the temperature in the solar nebula at the location now at the center of the asteroid belt was (a) 2000 K; (b) 900 K; (c) 400 K; (d) 100 K.
10. **VIS** According to Figure 6.18 ("Solar System Formation"), the jovian planets formed (a) at the same time as the terrestrial planets; (b) after the terrestrial planets; (c) within a few million years of the formation of the Sun; (d) at the same time as the Oort cloud.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

1. ● Only planets Mercury and Mars have orbits that deviate significantly from circles. Calculate the perihelion and aphelion distances of these planets from the Sun. ∞ (More Precisely 2-1)
2. ● Use Newton's law of gravity to compute your weight (a) on Earth, (b) on Mars, (c) on the asteroid Ceres, and (d) on Jupiter (neglecting temporarily the absence of a solid surface on this planet!). ∞ (Sec. 2.7)
3. ● Suppose the average mass of each of the 7000 asteroids in the solar system is about  $10^{17}$  kg. Compare the total mass of all asteroids with the mass of Earth.
4. ●● A *short-period* comet is conventionally defined as a comet having an orbital period of less than 200 years. What is the maximum possible aphelion distance for a short-period comet with a perihelion of 0.5 AU? Where does this place the comet relative to the outer planets?
5. ●● An interstellar cloud fragment 0.2 light-year in diameter is rotating at a rate of one revolution per million years. It now begins to collapse. Assuming that the mass remains constant, estimate the cloud's rotation period when it has shrunk to (a) the size of the solar nebula, 100 AU across, and (b) the size of Earth's orbit, 2 AU across.
6. ●● Consider a planet growing as it accretes material from the solar nebula. As the planet grows, its density remains roughly constant. Does the force of gravity at the surface of the planet increase, decrease, or stay the same? What would happen to the surface gravity and escape speed as the radius of the planet doubled? Give reasons for your answer.
7. ● How many 100-km-diameter rocky ( $3000 \text{ kg/m}^3$ ) planetesimals would have been needed to form Earth?
8. ● Use the data given in the text to calculate Neptune's orbital period before interactions with planetesimals expanded the orbit to its present size.

## Activities

---

### Collaborative

1. As a group, decide which of the space missions described in *Discovery 6-2* produced the most interesting results about the solar system. Explain your reasoning. Go online and research one of the other missions mentioned in that box, and present its major objectives and findings to the class.
2. What should be the U.S. government's policy on the mining of minerals from asteroids? Justify the policy.

### Individual

1. You can begin to visualize the ecliptic—the plane of the planets' orbits—just by noticing the path of the Sun throughout the day and of the full Moon in the course of a single night. It helps if you watch from one spot, such as

your backyard or a rooftop. It's also good to have a general notion of direction. (West is where the Sun sets!) The movements of the Sun, Moon, and planets are confined to a narrow pathway across our sky. This pathway reflects the plane of the solar system, the ecliptic.

2. The only way to tell an asteroid from a star is to watch it over several nights. The magazines *Sky & Telescope* and *Astronomy* often publish charts for especially prominent asteroids. Look for Ceres, Pallas, or Vesta, the brightest asteroids. Use the chart to locate the appropriate star field and aim binoculars at that location in the sky. You may be able to pick out the asteroid from the chart. If you can't, make a rough drawing of the entire field. Come back a night or two later and look again. The "star" that has moved is the asteroid.





# Earth

## OUR HOME IN SPACE

Earth is the best-studied terrestrial planet. From the matter of our world sprang life, intelligence, culture, and all the technology we now use to explore the cosmos. We ourselves are “Earthstuff” as much as are rocks, trees, and air. Now, as humanity begins to explore the solar system, we can draw on our knowledge of Earth to aid our understanding of the other planets.

By cataloging Earth’s properties and attempting to explain them, we set the stage for our comparative study of the solar system. Local and regional events such as volcanoes, earthquakes, and weather, as well as global trends such as climate change and continental drift, help us decipher both our own world and other nearby planets. Every piece of information we glean about the structure and history of our planet plays a vital role in helping us understand the planetary system in which we live.

**The Big Picture** If we are to appreciate the full grandeur of the universe and its many varied contents, we must first come to know our own planet. Many textbooks on astronomy skip Earth, claiming it to be the purview of geology. Yet, Earth is the platform from which we observe the much bigger universe beyond; it is the only cosmic perspective we have to explore our place in the cosmic scheme of things, and it is a planetary body of importance in and of itself. Our study of astronomy begins at home.

**LEFT:** Photographs like this, showing Earth hanging in space like a “blue marble,” help us appreciate life on our planet—and our place in the universe. Centered on the Americas, this image reveals the air, water, land, and life on our planet as a complex, ever-changing interactive system. The more we learn about Earth, the better we can compare and contrast our home world with other planets and moons. This image is a mosaic of many photographs taken in 2012 by the first of a new generation of weather satellites orbiting Earth. It is the finest picture of our home in space ever made. (NASA/NOAA)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Summarize the physical properties of planet Earth.
- 2 Explain how Earth’s atmosphere helps to heat us, as well as protect us.
- 3 Outline our current model of Earth’s interior, and describe some of the experimental techniques used to establish the model.
- 4 Summarize the evidence for the phenomenon of “continental drift,” and discuss the physical processes that drive it.
- 5 Describe the nature and origin of Earth’s magnetosphere.
- 6 Explain how both the Moon and the Sun influence Earth’s surface and affect our planet’s spin.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.

## 7.1 Overall Structure of Planet Earth

Earth's physical and orbital properties are determined using techniques that are conceptually similar to those presented in Chapter 6: simple geometry to determine Earth's radius, the orbit of the Moon to measure our planet's mass, and so on. [∞ \(Sec. 6.2\)](#) Throughout the body of this text, we will use rounded-off numbers whenever possible, taking our planet's mass and radius to be  $6.0 \times 10^{24}$  kg and 6400 km, respectively.


Dividing mass by volume, we find that Earth's average density is around  $5500 \text{ kg/m}^3$ . This simple calculation allows us to make a very important deduction about the interior of our planet. From direct measurements, we know that the water that makes up much of Earth's surface has a density of  $1000 \text{ kg/m}^3$ , and the rock beneath us on the continents, as well as on the seafloor, has a density in the range from 2000 to  $4000 \text{ kg/m}^3$ . We can immediately conclude that, because the surface layers have densities much less than the average, much denser material must lie deeper, under the surface. Hence, we should expect that much of Earth's interior is made up of very dense matter, far more compact than the densest continental rocks on the surface.

On the basis of measurements made in many different ways—using aircraft in the atmosphere, satellites in orbit, gauges on the land, submarines in the ocean, and drilling gear below the rocky crust—scientists have built up

the following overall picture of our planet: As indicated in Figure 7.1, Earth may be divided into six main regions. In Earth's interior, a thick **mantle** surrounds a smaller, two-part **core**. At the surface, we have (1) a relatively thin **crust**, comprising the solid continents and the seafloor, and (2) the **hydrosphere**, which contains the liquid oceans and accounts for some 70 percent of our planet's total surface area. An **atmosphere** of air lies just above the surface. At much greater altitudes, a zone of charged particles trapped by Earth's magnetic field forms the **magnetosphere**. Virtually all our planet's mass is contained within the surface and the interior. The gaseous atmosphere and the magnetosphere contribute hardly anything—less than 0.1 percent—to the total.


## 7.2 Earth's Atmosphere

From a human perspective, probably the most important aspect of Earth's atmosphere is that we can breathe it. Air is a mixture of gases, the most common of which are nitrogen ( $\text{N}_2$ , 78 percent by volume), oxygen ( $\text{O}_2$ , 21 percent), argon (Ar, 0.9 percent), and carbon dioxide ( $\text{CO}_2$ , 0.03 percent). The amount of water vapor ( $\text{H}_2\text{O}$ ) varies from 0.1 to 3 percent, depending on location and climate. The presence of a large amount of oxygen makes our atmosphere unique in the solar system, and the presence of even trace amounts of water and carbon dioxide play vital roles in the workings of our planet.



◀ **FIGURE 7.1** The Main Regions of Planet Earth

Earth's inner core of radius 1300 km is surrounded by a 2200-km thick liquid outer core. Most of the rest of Earth's 13,000-km interior is taken up by the mantle, topped by a thin crust only a few tens of kilometers thick. The liquid portions of Earth's surface make up the hydrosphere. Above the hydrosphere and solid crust lies the atmosphere, most of it within 50 km of the surface. Earth's outermost region is the magnetosphere, extending thousands of kilometers into space.



◀ **FIGURE 7.2 Earth's Atmosphere** Diagram of Earth's atmosphere, showing the changes in temperature (blue curve, bottom axis) and pressure (right-hand axis) from the planet's surface to the bottom of the ionosphere. Pressure decreases steadily with increasing altitude, but the temperature may fall or rise, depending on height above the ground.


the surface is the **mesosphere**. Above about 80 km, in the **ionosphere**, the atmosphere is kept partly ionized by solar ultraviolet radiation. Note how the temperature gradient (decreasing or increasing with altitude) changes from one atmospheric region to the next.

Atmospheric density decreases steadily with increasing altitude, and as the right-hand vertical axis in Figure 7.2 shows, so does pressure. Climbing even a modest mountain—4 or 5 km high, say—clearly demonstrates the thinning of the air in the troposphere. Climbers must wear oxygen masks when scaling the tallest peaks on Earth.

The troposphere is the region of Earth's (or any other planet's) atmosphere where *convection* occurs, driven by the heat of Earth's warm surface. **Convection** is the constant upwelling of warm air and the concurrent downward flow of cooler air to take its place, a process that physically transfers heat from a lower (hotter) to a higher (cooler) level. In Figure 7.3, part of Earth's surface is heated by the Sun. The air immediately above the warmed surface is heated, expands a little, and becomes less dense. As a result, the hot air becomes buoyant and starts to rise. At higher

## Atmospheric Structure

Figure 7.2 shows a cross-section of our planet's atmosphere. Compared with Earth's overall dimensions, the extent of the atmosphere is not great. Half of it lies within 5 km of the surface, and all but 1 percent is found below 30 km. The portion of the atmosphere below about 12 km is called the **troposphere**. Above it, extending up to an altitude of 40 to 50 km, lies the **stratosphere**. Between 50 and 80 km from



◀ **FIGURE 7.3 Convection** Circulation currents resulting from convection are familiar to us as the winds in Earth's atmosphere, caused by the solar-heated ground. Over and over, hot air rises, cools, and falls back to Earth. Eventually, steady circulation patterns are established and maintained, provided that the source of heat (the Sun, in the case of Earth) remains intact.

altitudes, the opposite effect occurs: The air gradually cools, grows denser, and sinks back to the ground. Cool air at the surface rushes in to replace the hot buoyant air. In this way, a circulation pattern is established. These *convection cells* of rising and falling air not only contribute to atmospheric heating, but also are responsible for surface winds. The constant churning motion in convection cells is responsible for all the weather we experience.

Atmospheric convection can also create clear-air turbulence—the bumpiness we sometimes experience on aircraft flights. Ascending and descending parcels of air, especially below fluffy clouds (themselves the result of convective processes, when water vapor condenses out at the cool tops of convection cells), can cause a choppy ride. For this reason, passenger aircraft tend to fly above most of the turbulence, at the top of the troposphere or in the lower stratosphere, where the atmosphere is stable and the air is calm.

Above about 100 km, in the ionosphere, the atmosphere is significantly ionized by the high-energy portion of the Sun’s radiation spectrum, which breaks down molecules into atoms and atoms into ions. The degree of ionization increases with altitude. The presence of many free electrons makes this region of the upper atmosphere a good conductor of electricity, and the conductivity renders the ionosphere highly reflective to certain radio wavelengths. ∞ (Sec. 3.3) The reason that AM radio stations can be heard well beyond the horizon is that their signals bounce off the ionosphere before reaching a receiver. FM signals cannot be received from stations over the horizon, however, because the ionosphere is transparent to the somewhat shorter wavelengths of radio waves in the FM band.

## Atmospheric Ozone


Within the stratosphere is the **ozone layer**, where, at an altitude of around 25 km, incoming solar ultraviolet radiation is absorbed by atmospheric ozone and nitrogen. (Ozone [O<sub>3</sub>] consists of three oxygen atoms combined into a single molecule. Ultraviolet radiation breaks ozone down, forming molecular oxygen [O<sub>2</sub>] again.)

The ozone layer is one of the insulating spheres that serve to shield life on Earth from the harsh realities of outer space. Not so long ago, scientists judged space to be hostile to advanced life-forms because of what is missing out there: breathable air and a warm environment. Now most scientists regard outer space as harsh because of what is *present* out there: fierce radiation and energetic particles, both of which are injurious to human health. Without the protection of the ozone layer, advanced life (at least on Earth’s surface) would be at best unlikely and at worst impossible.

Human technology has reached the point where it has begun to produce measurable—and possibly permanent—changes to our planet. One particularly undesirable by-product of our ingenuity is a group of chemicals

known as chlorofluorocarbons (CFCs), relatively simple compounds once widely used for a variety of purposes—propellant in aerosol cans, solvents in dry-cleaning products, and coolant in air conditioners and refrigerators. In the 1970s, it was discovered that instead of quickly breaking down after use, as had previously been thought, CFCs accumulate in the atmosphere and are carried high into the stratosphere by convection. There they are broken down by sunlight, releasing chlorine, which quickly reacts with ozone, turning it into oxygen. In chemical terms, the chlorine is said to act as a *catalyst*—it is not consumed in the reaction, so it survives to react with many more ozone molecules. A single chlorine atom can destroy up to 100,000 ozone molecules before being removed by other, less frequent chemical reactions.

Thus, even a small amount of CFCs is extraordinarily efficient at destroying atmospheric ozone, and the net result of CFC emission is a substantial increase in ultraviolet radiation levels at Earth’s surface, with detrimental effects to most living organisms. Figure 7.4 shows a vast ozone “hole” over the Antarctic. The hole is a region where atmospheric circulation and low temperatures conspire each Antarctic spring to create a vast circumpolar cloud of ice crystals that act to promote the ozone-destroying reactions, resulting in ozone levels about 50 percent below normal for the region. Ozone depletion is not confined to the Antarctic, although



▲ **FIGURE 7.4 Antarctic Ozone Hole** This composite image constructed from satellite observations shows (in pink) a huge “hole” in the ozone layer over the Antarctic continent. The hole is a region where climatic conditions and human-made chemicals combine to rob our atmosphere of its protective ozone blanket. The depth and area of the hole have grown significantly since the hole was discovered in the 1980s. Its maximum size is now larger than North America.

## MORE PRECISELY 7-1

### Why Is the Sky Blue?

Is the sky blue because it reflects the color of the ocean, or is the ocean blue because it reflects the color of the surrounding sky? The answer is the latter, and the reason has to do with the way that light is *scattered* by air molecules and minute dust particles. By *scattering*, we mean the process by which radiation is absorbed and then reradiated by the material through which it passes.

As sunlight passes through our atmosphere, it is scattered by gas molecules in the air. The British physicist Lord Rayleigh first investigated this phenomenon about a century ago, and today it bears his name—it is known as *Rayleigh scattering*. The process turns out to be highly sensitive to the wavelength of the light involved.

Rayleigh found that blue light is much more easily scattered than red light, basically because the wavelength of blue light (400 nm) is closer to the size of air molecules than the wavelength of red light (700 nm). He went on to prove mathematically, on the basis of the laws of electromagnetism, that the amount of scattering is inversely proportional to the *fourth* power of the wavelength:

$$\text{scattering by molecules} \propto \frac{1}{\text{wavelength}^4}.$$

Rayleigh's formula applies to scattering by particles (such as molecules) that are smaller than the wavelength of the light involved. Larger particles, such as dust, also preferentially scatter blue light, but by an amount that depends only inversely on the wavelength:



$$\text{scattering by dust} \propto \frac{1}{\text{wavelength}}.$$

**EXAMPLE** Let's compare the relative scattering of blue (400 nm) and red (700 nm) light by atmospheric molecules and dust. For Rayleigh scattering, blue light is scattered  $(700/400)^4 \approx 9.4$  times more efficiently than red light. That is, blue photons are almost 10 times more likely to be scattered out of a beam of sunlight (taken out of the forward beam and redirected to the side) than are red photons. For scattering by dust, the corresponding factor is  $(700/400) = 1.75$ —not as big a differential, but still enough to have a large effect when the air happens to be particularly dirty.

When the Sun is high in the sky, the blue component of incoming sunlight is scattered much more than any other color component. Thus, some blue light is removed from the line of sight between us and the Sun and may scatter many times in the atmosphere before eventually entering our eyes, as shown in the first figure. Red or yellow light is scattered relatively little and arrives at our eyes predominantly along the line of sight to the Sun. The net effect is that the Sun is “reddened” slightly because of the removal of blue light, whereas the sky away from the Sun appears blue. In outer space, where there is no atmosphere, there is no Rayleigh scattering of sunlight, and the sky is black (although, as we will see in Chapter 18, light from distant stars is reddened in precisely the same way as it passes through clouds of interstellar gas and dust).

At dawn or dusk, with the Sun near the horizon, sunlight must pass through much more atmosphere before reaching our eyes—so much so, in fact, that the blue component of the Sun's light is almost entirely scattered out of the line of sight, and even the red component is diminished in intensity. Accordingly, the Sun itself appears orange—a combination of its normal yellow color and a reddishness caused by the subtraction of virtually all of the blue end of the spectrum—and dimmer than at noon.

At the end of a particularly dusty day (second figure), when weather conditions or human activities during the daytime hours have raised excess particles into the air, short-wavelength Rayleigh scattering can be so heavy that the Sun appears brilliantly red. Reddening is often especially evident when we look at the westerly “sinking” summer Sun over the ocean, where seawater molecules have evaporated into the air, or during the weeks and months after an active volcano has released huge quantities of gas and dust particles into the air—as was the case in North America when the Philippine volcano Mount Pinatubo exploded in 1991 in the most violent volcanic eruption in the past 100 years.



(NCAR/Science Source)

the effect is greatest there. Smaller holes have been observed in the Arctic, and occasional ozone depletions of up to 20 percent have been reported at lower northern latitudes.

In the late 1980s, when the effects of CFCs on the atmosphere were realized, the world moved rapidly to curtail their production and use, with the goal of phasing them out entirely by 2030. Substantial cuts have already been made, and the agreement to do so has become a model of international cooperation. Still, scientists think that, even if all remaining CFC emissions were to stop today, it would nonetheless take several decades for CFCs to leave the atmosphere completely.


## Surface Heating

Much of the Sun's radiation manages to penetrate Earth's atmosphere, eventually reaching the ground. (See *More Precisely 7-1* for more on how the atmosphere affects incoming sunlight before it reaches the surface.) Most of this energy takes the form of visible and near-infrared radiation—ordinary sunlight.  $\infty$  (Sec. 3.3) Essentially all of the solar radiation that is not absorbed by or reflected from clouds in the upper atmosphere is absorbed by Earth's surface. The result is that our planet's surface and most objects on it heat up considerably during the day. Earth cannot absorb this solar energy indefinitely, however. If it did, the surface would soon become hot enough to melt, and life on our planet would not exist.

As it heats up, Earth's surface reradiates much of the absorbed energy. This reemitted radiation follows the blackbody curve discussed in Chapter 3.  $\infty$  (Sec. 3.4) As the surface temperature rises, the amount of energy radiated increases rapidly, in accordance with Stefan's law. Eventually, Earth radiates as much energy back into space as it receives from the Sun, and a stable balance is struck. In the absence of any complicating effects, this balance would be achieved at an average surface temperature of about 250 K ( $-23^{\circ}\text{C}$ ). Wien's law tells us that, at that temperature, most of the reemitted energy is in the form of infrared (heat) radiation.

But there *are* complications. Infrared radiation is partially blocked by Earth's atmosphere, primarily because of the presence of molecules of water vapor and carbon dioxide, which absorb very efficiently in the infrared portion of the spectrum. Even though these two gases are only trace constituents of our atmosphere, they manage to absorb a large fraction of all the infrared radiation emitted from the surface. Consequently, only some of that radiation escapes back into space. The remainder is trapped within our atmosphere, causing the temperature to increase.

This partial trapping of solar radiation is known as the **greenhouse effect**. The name comes from the fact that a similar process operates in a greenhouse, where sunlight passes relatively unhindered through glass panes, but much of the infrared radiation reemitted by the plants is blocked by the glass and cannot get out. Consequently, the interior of



**Interactive FIGURE 7.5 Greenhouse Effect** Sunlight that is not reflected by clouds reaches Earth's surface, warming it up. Infrared radiation reradiated from the surface is partially absorbed by carbon dioxide (and also water vapor, not shown here) in the atmosphere, causing the overall surface temperature to rise.

the greenhouse heats up, and flowers, fruits, and vegetables can grow even on cold wintry days.\* The radiative processes that determine the temperature of Earth's atmosphere are illustrated in Figure 7.5. Earth's greenhouse effect makes our planet almost 40 K ( $40^{\circ}\text{C}$ ) hotter than would otherwise be the case.

The magnitude of the greenhouse effect is highly sensitive to the concentration of so-called **greenhouse gases** (that is, gases that absorb infrared radiation efficiently) in the atmosphere. Carbon dioxide and water vapor are the most important of these, although other atmospheric gases (such as methane) also contribute. The amount of carbon dioxide in Earth's atmosphere is increasing, largely as a result of the burning of fossil fuels (principally oil and coal) in the industrialized and developing worlds. Carbon dioxide levels have increased by over 20 percent in the last century, and they are continuing to rise at a present rate of 4 percent per decade. *Discovery 7-1* discusses the causes and

\*Note that although this process does contribute to warming the interior of a greenhouse, it is not the most important effect. A greenhouse works mainly because its glass panes prevent convection from carrying heat up and away from the interior. Nevertheless, the name "greenhouse effect" due to Earth's atmosphere has stuck.

## DISCOVERY 7-1

## The Greenhouse Effect and Global Warming


We saw in the text how greenhouse gases in Earth's atmosphere—notably, water vapor and carbon dioxide (CO<sub>2</sub>)—tend to trap heat leaving the surface, raising our planet's temperature by several tens of degrees Celsius. In and of itself, this *greenhouse effect* is not a bad thing—in fact, it is the reason that water exists in the liquid state on Earth's surface, and thus it is crucial to the existence and survival of life on our planet (see Chapter 28). However, if atmospheric greenhouse gas levels rise unchecked, the consequences could be catastrophic.

Since the Industrial Revolution in the 18th century, and particularly over the past few decades, human activities on Earth have steadily raised the level of carbon dioxide in our atmosphere. Fossil fuels (coal, oil, and gas), still the dominant energy source of modern industry, all release CO<sub>2</sub> when burned. At the same time, the extensive forests that once covered much of our planet are being systematically destroyed to make room for human expansion. Forests play an important role in this situation because vegetation absorbs carbon dioxide, thus providing a natural control mechanism for atmospheric CO<sub>2</sub>. Deforestation therefore also tends to increase the amount of greenhouse gases in Earth's atmosphere. The first figure shows atmospheric CO<sub>2</sub> levels over the past thousand years. Note the dramatic increase during the past two centuries.

*Global warming* is the slow rise in Earth's surface temperature caused by the increased greenhouse effect resulting from higher levels of atmospheric carbon dioxide. As shown in the second figure, average global temperatures have risen by about 0.5°C during the past century. This may not seem like much, but climate models predict that, if CO<sub>2</sub> levels continue to rise, a further increase of as much as 5°C is possible by the end of the 21st century. Such a rise would be enough to cause serious climate change on a global scale. Among the possible (some would say likely) consequences of such a temperature increase are the following phenomena.

- Melting of glaciers and the polar ice caps, leading to a rise in sea level of up to a meter by the year 2100, with the potential for widespread coastal flooding.
- Longer and more extreme periods of severe weather—heat waves, droughts and wildfires—yet with more precipitation (rain and snow) between them.
- Crop failures as Earth's temperate zones move toward the poles.
- Expansion of deserts in heavily populated equatorial regions.
- Increased numbers of mosquitoes and other pests spreading tropical diseases into unprotected populations.

The fossil record shows that major climatic changes have occurred many times before in Earth's history, but never at the rate predicted by these dire warnings. The unprecedented speed of the forecasted events may well be too rapid for many species (and some human societies) to survive.



In Section 7.2, we described the danger to Earth's ozone layer posed by CFCs—another product of modern technology with unexpected global consequences—and saw how, once their environmental impact was identified, rapid steps were taken to curb their use. A concerted international response to global warming has been much slower in coming. Most scientists see the human-enhanced greenhouse effect as a real threat to Earth's climate, and they urge prompt and deep reductions in CO<sub>2</sub> emissions, along with steps to slow and ultimately reverse deforestation.

Some, however—particularly those connected with the industries most responsible for the production of greenhouse gases—argue that Earth's long-term response to increased greenhouse emissions is too complex for simple conclusions to be drawn and that immediate action is unnecessary. They suggest that the current temperature trend may be part of some much longer cycle or that natural environmental factors may in time stabilize, or even reduce, the level of CO<sub>2</sub> in the atmosphere without human intervention. In part because of these objections, international agreements to limit carbon emissions have failed to win approval in the United States (currently the largest producer of greenhouse gases) and thus have had only limited success.

Given the stakes, it is perhaps not surprising that these debates have become far more political than scientific in tone—not at all like the deliberative scientific method presented elsewhere in this text! The basic observations and much of the basic science are generally not seriously questioned, but the interpretation, long-term consequences, and proper response are all hotly debated. Separating the two sometimes is not easy, but the outcome may be of vital importance to life on Earth.



some possible consequences of rising carbon dioxide levels in Earth's atmosphere.

In Chapter 9, we will see how a runaway increase in carbon dioxide levels in the atmosphere of the planet Venus radically altered conditions on its surface, causing its temperature to rise to over 700 K. Although no one is predicting that Earth's temperature will ever reach that of Venus, many scientists now think that our planet's increase in carbon dioxide levels, if left unchecked, may result in global temperature increases of several kelvins over the next half century—enough to cause dramatic, and possibly catastrophic, changes in Earth's climate, ranging from rising sea levels to the accelerated spread of disease.

## Origin of Earth's Atmosphere

Why is our atmosphere made up of its present constituents? Why is it not composed entirely of nitrogen, say, or of carbon dioxide, like the atmospheres of Venus and Mars? The origin and development of Earth's atmosphere was a fairly complex and lengthy process.

When Earth first formed, any *primary atmosphere* it might have had would have consisted of the gases most common in the early solar system: hydrogen, helium, methane, ammonia, and water vapor—a far cry from the atmosphere we enjoy today. Almost all this low-density material, and especially any hydrogen or helium, escaped into space during the first half-billion or so years after Earth was formed. (For more information on how planets retain or lose their atmospheres, see *More Precisely 8-1*.)

Subsequently, Earth developed a *secondary atmosphere*, which was *outgassed* (expelled) from the planet's interior as a result of volcanic activity. Volcanic gases are rich in water vapor, methane, carbon dioxide, sulfur dioxide, and compounds containing nitrogen (such as nitrogen gas, ammonia, and nitric oxide). Solar ultraviolet radiation split the lighter, hydrogen-rich gases into their component atoms, allowing the hydrogen to escape and liberating much of the nitrogen from its bonds with other elements. As Earth's surface temperature fell, the water vapor condensed and oceans formed. Much of the carbon dioxide and sulfur dioxide became dissolved in the oceans or combined with surface rocks. Oxygen is such a reactive gas that any free oxygen that appeared at early times was removed as quickly as it formed. An atmosphere consisting largely of nitrogen slowly appeared.

The final major development in the story of our planet's atmosphere is known so far to have occurred only on Earth. *Life* appeared in the oceans more than 3.5 billion years ago, and organisms eventually began to produce atmospheric oxygen. The ozone layer formed, shielding the surface from the Sun's harmful radiation. Eventually, life spread to the land and flourished. The fact that oxygen is a major constituent of the present-day atmosphere is a direct consequence of the evolution of life on Earth.

## CONCEPT Check

- ✓ Why is the greenhouse effect important for life on Earth?

## 7.3 Earth's Interior


Although we reside on Earth, we cannot easily probe our planet's interior. Drilling gear can penetrate rock only so far before breaking. No substance used for drilling—even diamond, the hardest known material—can withstand the pressure below a depth of about 10 km. That's rather shallow compared with Earth's 6400-km radius. Fortunately, geologists have developed other techniques that indirectly probe the deep recesses of our planet.

### Seismic Waves

A sudden dislocation of rocky material near Earth's surface—an *earthquake*—causes the entire planet to vibrate a little. Earth rings like a giant bell. These vibrations are not random, however. They are systematic waves, called **seismic waves** (after the Greek word for “earthquake”), that move outward from the site of the quake. Like all waves, they carry information. This information can be detected and recorded with sensitive equipment—a *seismograph*—designed to monitor Earth tremors.

Decades of earthquake research have demonstrated the existence of many kinds of seismic waves. Two are of particular importance to the study of Earth's internal structure. First to arrive at a monitoring site after a distant earthquake are the primary waves, or *P-waves*. These are *pressure* waves, a little like ordinary sound waves in air, that alternately expand and compress the medium (the core or mantle) through which they move. Seismic P-waves usually travel at speeds ranging from 5 to 6 km/s and can travel through both liquids and solids. Some time later (the actual delay depends on the distance from the earthquake site), secondary waves, or *S-waves*, arrive. These are *shear* waves. Unlike P-waves, which vibrate the material through which they pass back and forth along the direction of travel of the wave, S-waves cause side-to-side motion, more like waves in a guitar string. The two types of waves are illustrated in Figure 7.6. S-waves normally travel through Earth's interior at 3 to 4 km/s; however, they cannot travel through liquid, which absorbs them.

The speeds of both P- and S-waves depend on the density of the matter through which the waves are traveling. Consequently, if we can measure the time taken for the waves to move from the site of an earthquake to one or more monitoring stations on Earth's surface, we can determine the density of matter in the interior. Figure 7.7 illustrates some P- and S-wave paths away from the site of an earthquake. Seismographs located around the world measure the




▲ **FIGURE 7.6 P and S Waves** (a) Pressure (P) waves traveling through Earth's interior cause material to vibrate in a direction parallel to the direction of motion of the wave. Material is alternately compressed and expanded. (b) Shear (S) waves produce motion perpendicular to the direction in which the wave travels, pushing material from side to side.

times of arrival, as well as the strengths, of the seismic waves. Both observations contain much useful information—about the earthquake itself and about Earth's interior through which the waves pass. Notice that the waves do not travel in straight lines through the planet. Because the wave velocity increases with depth, waves that travel deeper tend to overtake those on slightly shallower paths, and the waves bend as they move through the interior.

A particularly important result emerged after numerous quakes were monitored several decades ago: Seismic stations on the side of Earth opposite a quake never detect S-waves—these waves are blocked by material within Earth's interior. Furthermore, although P-waves always arrive at stations diametrically opposite the quake, parts of Earth's surface receive almost none (see Figure 7.7). Most geologists think that S-waves are absorbed by a liquid core at Earth's center and that P-waves are refracted at the core boundary, much as light is refracted by a lens. The result is the S- and P-wave “shadow zones” we observe. The fact that every earthquake exhibits these shadow zones is the best evidence that the core of our planet is hot enough to be liquid.

The sizes of the shadow zones depend on the radius of the core, and careful analysis of the seismic data yield a




▲ **FIGURE 7.7 Seismic Waves** Earthquakes generate pressure (P, or primary) and shear (S, or secondary) waves that can be detected at seismographic stations around the world. The waves bend while moving through Earth's interior because of the variation in density and temperature within our planet. S-waves (colored red) are not observed by stations “shadowed” by the liquid core of Earth. P-waves (colored green) do reach the side of Earth opposite the earthquake, but their interaction with Earth's core produces another shadow zone, where almost no P-waves are detected.

core radius of about 3500 km. In fact, very faint P-waves *are* observed in the P-wave shadow zone indicated in Figure 7.7. These are thought to be reflected off the surface of a solid **inner core**, of radius 1300 km, lying at the center of the liquid **outer core**.

## Modeling Earth's Interior

Because earthquakes occur often and at widespread places across the globe, geologists have accumulated a large amount of data about shadow zones and seismic-wave properties. They have used these data, along with direct knowledge of surface rocks, to build mathematical models of Earth's interior. Our knowledge of the deepest recesses of our planet is based almost entirely on modeling and indirect observation. We will find many more examples of this powerful combination throughout the text.

Figure 7.8 presents a model that most scientists accept. According to this model, Earth's outer core is surrounded by a thick mantle and topped with a thin crust. The mantle is about 3000 km thick and accounts for the bulk (80 percent) of our planet's volume. The crust has an average thickness of only 15 km—a little less (around 8 km) under the oceans and somewhat more (20–50 km) under the



▲ **FIGURE 7.8 Earth's Interior** Computer models of Earth's interior imply that the density and temperature vary considerably through the mantle and the core. Seismic modeling is key to understanding the Earth in bulk.

continents. The average density of crust material is around  $3000 \text{ kg/m}^3$ . Density and temperature both increase with depth. Specifically, from Earth's surface to its very center, the density increases from roughly  $3000 \text{ kg/m}^3$  to a little more than  $12,000 \text{ kg/m}^3$ , and the temperature rises from just under  $300 \text{ K}$  to well over  $5000 \text{ K}$ . Much of the mantle has a density midway between the densities of the core and crust: about  $5000 \text{ kg/m}^3$ .

The high central density tells geologists that the inner parts of Earth must be rich in nickel and iron. Under the heavy pressure of the overlying layers, these metals (whose densities under surface conditions are around  $8000 \text{ kg/m}^3$ ) can be compressed to the high densities predicted by the model. The sharp increase in density at the mantle-core boundary results from the difference in composition between the two regions. The mantle is composed of dense, but *rocky*, material—compounds of silicon and oxygen. The

core consists primarily of even denser *metallic* elements. There is no similar jump in density or temperature at the inner core boundary—the material there simply changes from the liquid to the solid state.

The model suggests that the core must be a mixture of nickel, iron, and some other lighter element, possibly sulfur. Without direct observations, it is difficult to be absolutely certain of the light component's identity. All geologists agree that much of the core must be liquid. The existence of the shadow zone demands this (and, as we will see, our current explanation of Earth's magnetic field relies on it). However, despite the high temperature, the pressure near the center—about 4 million times the atmospheric pressure at Earth's surface—is high enough to force the material there into the solid state.


Because geologists have been unable to drill deeper than about  $10 \text{ km}$ , no experiment has yet recovered a sample of Earth's mantle. However, we are not entirely ignorant of the mantle's properties. In a *volcano*, hot lava upwells from below the crust, bringing a little of the mantle to us and providing some inkling of Earth's interior. Observations of the chemical and physical properties of newly emerged lava are generally consistent with the model sketched in Figure 7.8.

The composition of the upper mantle is probably quite similar to the iron-magnesium-silicate mixtures known as *basalt*. You may have seen some dark gray basaltic rocks scattered across Earth's surface, especially near volcanoes. Basalt is formed as material from the mantle upwells from Earth's interior as lava, cools, and then solidifies. With a density between  $3000 \text{ kg/m}^3$  and  $3300 \text{ kg/m}^3$ , basalt contrasts with the lighter *granite* (density  $2700\text{--}3000 \text{ kg/m}^3$ ) that constitutes much of the rest of Earth's crust. Granite is richer than basalt in the light elements silicon and aluminum, which explains why the surface continents do not sink into the interior. Their low-density composition lets the crust "float" atop the denser matter of the mantle and core below.

## Differentiation

Earth, then, is not a homogeneous ball of rock. Instead, it has a layered structure, with a low-density crust at the surface, an intermediate-density material in the mantle, and a high-density core. Such variation in density and composition is known as **differentiation**.

Why isn't our planet just one big, rocky ball of uniform density? The answer appears to be that much of Earth was *molten* at some time in the past. As a result, the higher density matter sank to the core and the lower density material was displaced toward the surface. A remnant of this ancient heating exists today: Earth's central temperature is nearly equal to the surface temperature of the Sun. What processes were responsible for heating the entire planet to that extent? To answer this question, we must try to visualize the past, as sketched in Figure 7.9.



According to current models of solar system formation, when Earth formed 4.6 billion years ago, it did so by capturing material from its surroundings, growing in mass as it swept up “preplanetary” chunks of matter in its vicinity (Sec. 6.6). As the young planet grew, its gravitational field strengthened and the speed with which newly captured matter struck its surface increased. This process generated a lot of heat—so much, in fact, that Earth may already have been partially or wholly molten by the time it reached its present size. As Earth began to differentiate and heavy material sank to the center, even more gravitational energy was released, and the interior temperature must have increased still further.

Later, Earth continued to be bombarded with debris left over from the formation process. At its peak about 3.8 billion years ago, this secondary bombardment was probably intense enough to keep the surface molten, but only down to a depth of a few tens of kilometers. Erosion by wind and water has long since removed all trace of this early period from the surface of Earth, but the Moon still bears visible scars of the onslaught.


A second important process for heating Earth soon after its formation was **radioactivity**—the release of energy by certain rare heavy elements, such as uranium, thorium, and plutonium (see *More Precisely 7-2*). These elements release energy and heat their surroundings as their complex heavy nuclei decay (break up) into simpler lighter ones. Although the energy produced by the decay of a single radioactive atom is tiny, Earth contained a lot of radioactive atoms, and a lot of time was available. Rock is such a poor conductor of heat that the energy would have taken a very long time to reach the surface and leak away into space, so the heat built up in the interior, adding to the energy left there by Earth’s formation.

Provided that enough radioactive elements were originally spread throughout the primitive Earth, rather like raisins in a cake, the entire planet—from crust to core—could have melted and remained molten for about a billion years. That’s a long time by human standards, but not so long in the cosmic scheme of things. Measurements of the ages of some surface rocks indicate that Earth’s crust finally began to solidify roughly 700 million years after it originally formed. Radioactive heating did not stop at that point, of course; it

**◀ FIGURE 7.9 Earth's Differentiation** Earth’s interior changed greatly throughout its early history. (a) At its origin, 4.6 billion years ago, the Earth was probably already partly molten owing to debris bombardment and continued gravitational infall in its formative stage. (b) A second period of heavy bombardment, at about 3.8 billion years ago, likely caused its cooling surface layers to again become completely molten to a depth of tens of kilometers. (c) Early on especially, yet continuing to lesser extent to the present, radioactive heating from within has caused much of Earth’s interior to liquefy, allowing its heavy metals to sink to the core while its lighter-weight rocks floated to the surface.

## MORE PRECISELY 7-2

### Radioactive Dating


In Chapter 4, we saw that atoms are made up of electrons and nuclei and that nuclei are composed of protons and neutrons.  (Sec. 4.2) The number of protons in a nucleus determines which element it represents. However, the number of neutrons can vary. In fact, most elements can exist in several *isotopic* forms, all containing the same number of protons, but different numbers of neutrons in their nuclei. The particular nuclei we have encountered so far—the most common forms of hydrogen, helium, carbon, and iron—are all *stable*. For example, left alone, a carbon-12 nucleus, consisting of six protons and six neutrons, will remain unchanged forever. It will not break up into smaller pieces, nor will it turn into anything else.

Not all nuclei are stable, however. Many nuclei—such as carbon-14 (containing 6 protons and 8 neutrons), thorium-232 (90 protons, 142 neutrons), uranium-235 (92 protons, 143 neutrons), uranium-238 (92 protons, 146 neutrons), and plutonium-241 (94 protons, 147 neutrons)—are inherently *unstable*. Left alone, they will eventually break up into lighter “daughter” nuclei, emitting some elementary particles and releasing some energy in the process. The change happens spontaneously, without any external influence. This instability is known as **radioactivity**. The energy released by the disintegration of the radioactive elements just listed is the basis for nuclear fission reactors (and atomic bombs).

Unstable heavy nuclei achieve greater stability by disintegrating into lighter nuclei, but they do not do so immediately. Each type of “parent” nucleus takes a characteristic amount of time to decay. The **half-life** is the name given to the time required for half of a sample of parent nuclei to disintegrate. Notice that this is really a statement of probability. We cannot say *which* nuclei of a given element will decay in any given half-life interval; we can say only that half of them are expected to do so. If a given sample of material has half-life  $T$ , then we can write down a simple expression for the amount of material remaining after time  $t$ :

$$\text{fraction of material remaining} = (1/2)^{t/T}.$$

Thus, if we start with a billion radioactive nuclei embedded in a sample of rock, a half-billion nuclei will remain after one half-life, a quarter-billion after two half-lives, and so on. The first figure illustrates the decline in the number of parent nuclei as a function of time.



Every radioactive isotope has its own half-life, and most of their half-lives are now well known from studies conducted since the 1950s. For example, the half-life of uranium-235 is 713 million years, and that of uranium-238 is 4.5 billion years. Some radioactive elements decay much more rapidly, others much more slowly, but these two types of uranium are particularly important to geologists because their half-lives are comparable to the age of the solar system. The second figure illustrates the half-lives and decay reactions for four unstable heavy nuclei.

The decay of unstable radioactive nuclei into more stable *daughter* nuclei provides us with a useful tool for measuring the ages of any rocks we can get our hands on. The first step is to measure the amount of stable nuclei of a given kind (e.g., lead-206, which results from the decay of uranium-238). This amount is then compared with the amount of remaining unstable parent nuclei (in this case, uranium-238) from which the daughter nuclei descended. Knowing the rate (or half-life) at which the disintegration occurs, the age of the rock then follows directly. If half of the parent nuclei of some element have decayed, so that the number of daughter nuclei equals the number of parents, the age of the rock must be equal to the half-life of the radioactive nucleus studied. Similarly, if only a quarter of the parent nuclei remain (three times as many daughters as parents), the rock’s age is twice the half-life of that element, and so on.

In practice, ages can be determined by these means to within an accuracy of a few percent. The most ancient rocks on Earth are dated at 3.9 billion years old. These rare specimens have been found in Greenland and Labrador.

continued even after Earth’s surface cooled and solidified. But radioactive decay works in only one direction, always producing lighter elements from heavier ones. Once gone, the heavy and rare radioactive elements cannot be replenished.

So the early source of heat diminished with time, allowing the planet to cool over the past 4 billion years. In this process, Earth has cooled from the outside in, much like a hot potato, since regions closest to the surface can most easily unload their excess heat into space. In that way,

the surface developed a solid crust, and the differentiated interior attained the layered structure now implied by seismic studies.

### PROCESS OF SCIENCE Check

- ✓ Would scientists be able to model Earth’s interior if our planet were geologically inactive, with no volcanos or earthquakes?


**EXAMPLE** Suppose that careful chemical analysis of a sample of rock reveals that, for every nucleus of uranium-238 remaining in the sample, there is 0.41 of a lead-206 nucleus. If we assume that there was no lead-206 initially present, and hence that every lead-206 nucleus is the decay product of a nucleus of uranium-238, we can easily calculate the fraction of uranium-238 nuclei remaining. The answer is

$$\begin{aligned} \text{fraction of uranium-238} &= \frac{1}{1 + 0.41} \\ &= 0.71 \approx \sqrt{\frac{1}{2}} = \left(\frac{1}{2}\right)^{0.5}. \end{aligned}$$

From this equation, it follows that the elapsed time must be 0.5 times the half-life of uranium-238, or 2.25 billion years. If we were to repeat the analysis with uranium-235 and lead-207,

we would expect to find consistent results within the measurement errors: 2.25 billion years is  $2250/713 = 3.2$  uranium-235 half-lives, so only  $(1/2)^{3.2} \approx 11$  percent of any uranium-235 should remain—daughter lead-207 nuclei should outnumber parent uranium-235 nuclei by more than eight to one.

The radioactive-dating technique rests on the assumption that the rock has remained *solid* while the radioactive decays have been going on. If the rock melts, there is no particular reason to expect the daughter nuclei to remain in the same locations their parents had occupied, and the whole method fails. Thus, radioactive dating indicates the time that has elapsed since the last time the rock in question solidified. Hence, the 3.9-billion-year value represents only a portion—a lower limit—of the true age of our planet. It does not measure the duration of Earth's molten existence.



## 7.4 Surface Activity

Earth is geologically alive today. Its interior seethes and its surface constantly changes. Figure 7.10 shows some examples of two kinds of surface geological activity: a volcano, where molten rock and hot ash upwell through fissures or cracks in the surface, and (the aftermath of) an earthquake, which occurs when the crust suddenly dislodges under great pressure. Catastrophic volcanoes and earthquakes are relatively rare

events these days, but geological studies imply that surface activity must have been more frequent, and probably more violent, long ago.

### Continental Drift

Many traces of past geological events are scattered across our globe. Erosion by wind and water has wiped away much of the evidence for ancient activity, but modern



▲ **FIGURE 7.10 Geological Activity** (a) An active volcano on Kilauea in Hawaii. Kilauea seems to be a virtually continuous eruption. (b) Other, more sudden eruptions, such as that of Mount St. Helens in Washington State on May 18, 1980, are rare catastrophic events that can release more energy than the detonation of a thousand nuclear bombs. (c) The aftermath of an earthquake that claimed more than 5000 lives and caused billions of dollars' worth of damage in Kobe, Japan, in January 1995. (D.Peebles/Alamy; AP Photo/J. Smith; H. Yamaguchi/Syigma)

exploration has documented most of the recent sites. Figure 7.11 is a map of the currently active areas of our planet. The red dots represent volcanoes and earthquakes. Nearly all these sites have experienced surface activity within the last century, often resulting in much damage and loss of life.

The intriguing aspect of Figure 7.11 is that the active sites are not spread evenly across our planet. Instead, they trace well-defined lines of activity, where crustal rocks dislodge (as in earthquakes) or mantle material upwells (as in volcanoes). In the mid-1960s, scientists realized that these lines are really the outlines of gigantic “plates,” or slabs of Earth’s surface.\* Most startling of all, the plates are slowly moving—literally drifting around the surface of our planet. These plate motions have created the surface mountains, oceanic trenches, and other large-scale features across the face of planet Earth and have shaped the continents themselves. The process is popularly known as “continental drift” or, more technically, as **plate tectonics**. The major plates of the world are marked on Figure 7.11.


Taken together, the plates make up Earth’s **lithosphere**, which contains both the crust and a small part of the upper mantle. The lithosphere is the portion of Earth that undergoes tectonic activity. The semisolid part of the mantle over

\*Not all volcanoes are found near plate boundaries. The Hawaiian Islands, near the center of the Pacific plate, are associated with a “hot spot” in Earth’s upper mantle that melts the crust above it. Over millions of years, the motion of the Pacific plate across the hot spot has created a chain of volcanic islands.

which the lithosphere slides is known as the **asthenosphere**. The relationships between these regions of Earth are shown in Figure 7.12.

The idea of continental drift was first suggested in 1912 by a German meteorologist named Alfred Wegener, who pointed out the remarkable geographic fit between the continents on either side of the Atlantic ocean. Note in Figure 7.11 how the Brazilian coast (on the easternmost part of South America) meshes nicely with the indented Ivory Coast along west Africa. In fact, most of the continental landmasses in the Southern Hemisphere fit together remarkably well. Following the arrows in the figure backwards, we can see that the fits are roughly consistent with the present locations of the plates involved. The fit appears not to be as good in the Northern Hemisphere, but it improves markedly if we consider the continental shelves (the continental borders, which are under water) instead of just the portions that happen to stick up above sea level.

Few took Wegener’s ideas seriously at the time, in part because there was no known mechanism that could drive the plates’ motions. Nearly all scientists thought it preposterous that large segments of rocky crust could be drifting across the surface of our planet. Those skeptical views persisted for more than half a century, when the accumulation of data in support of continental drift became overwhelming. Similar-looking fossils are found on opposite sides of the Atlantic Ocean at just the locations where the continents “fit together,” and studies of the seafloor near the center of the Atlantic (discussed in more detail below) indicate the



▲ **FIGURE 7.11 Global Plates** Red dots represent active sites where major volcanoes or earthquakes have occurred in the 20th century. Taken together, the sites outline vast “plates,” indicated in dark blue, that drift around on the surface of our planet. The white arrows show the general directions and speeds of the plate motions.

formation of new crust as plates separate. Today, backed by evidence, Wegener’s “crazy” theory forms the foundation for all geological studies of our planet’s outer layers. [\(Sec. 1.2\)](#)

The plates are not simply slowing to a stop after some ancient initial movements. Rather, they are still drifting today, although at an extremely slow rate. Typically, the speeds of the plates amount to only a few centimeters per year—about the same rate as your fingernails grow. Still, this is well within the measuring capabilities of modern equipment. Curiously, one of the best ways of monitoring plate motion on a global scale is by making accurate observations of very distant astronomical objects. Quasars (see Chapter 25), lying many hundreds of millions of light-years from Earth, never show any measurable apparent motion on the sky stemming from their own motion in space. Thus, any apparent change in their position (after correction for Earth’s motion) can be interpreted as arising from the motion of the telescope—that is, of the continental plate on which it is located!

On smaller scales, laser-ranging and other techniques now routinely track the relative motion of plates in many areas, such as California, where advance warning of earthquake activity is at a premium. During the course of Earth’s history, each plate has had plenty of time to move large distances, even at its sluggish pace. For example, a drift rate of


only 2 cm per year can cause two continents (e.g., Europe and North America) to separate by some 4000 km over the course of 200 million years. That may be a long time by human standards, but it represents only about 5 percent of the age of Earth.

A common misconception is that the plates are the continents themselves. Some plates are indeed made mostly of continental landmasses, but other plates are made of a continent plus a large part of an ocean. For example, the Indian plate includes all of India, much of the Indian Ocean, and all of Australia and its surrounding south seas (see Figure 7.11). Still other plates are mostly ocean. The seafloor itself is a slowly drifting plate, and the oceanic water merely fills in the depressions between continents. The southeastern portion of the Pacific Ocean, called the Nazca plate, contains no landmass at all. For the most part, the continents are just passengers riding on much larger plates.

## Effects of Plate Motion

As the plates drift around, we might expect collisions to be routine. Indeed, plates do collide, but unlike two automobiles that collide and then stop, the surface plates are driven by enormous forces. They do not stop easily. Instead, they just keep crunching into one another, reshaping the landscape and causing violent seismic activity.





► **FIGURE 7.12 Earth's Upper Mantle**

The outer layers of Earth's interior include the rocky lithosphere, which comprises all of the crust and part of the upper mantle. It is typically between 50 and 100 km thick. Below it lies the asthenosphere, a relatively soft part of the mantle over which the lithosphere slips.

Figure 7.13(a) shows a collision currently occurring between two continental landmasses: The subcontinent of India, on the prow of the northward-moving Indian plate, is crashing into the landmass of Asia, located on the Eurasian plate (see Figure 7.11). The resulting folds of rocky crust create mountains—in this case, the snow-covered Himalayan mountain range at the upper right. A peak like Mount Everest (Figure 7.13b) represents a portion of Earth's crust that has been lifted over 8800 m by the slow, but inexorable, force produced when one plate plows into another.

Not all colliding plates produce mountain ranges. At other locations, called **subduction zones**, one plate slides under the other, ultimately to be destroyed as it sinks into the mantle. Subduction zones are responsible for most of the deep trenches in the world's oceans.


Nor do all plates experience head-on collisions. As noted by the arrows of Figure 7.11, many plates slide or shear past one another. A good example is the most famous active region in North America: the San Andreas Fault in California (Figure 7.14). The site of much earthquake activity, this fault marks the boundary where the Pacific and North American plates are rubbing past each other. The motion of these two plates, like that of moving parts in a poorly oiled machine, is neither steady nor smooth. The sudden jerks that occur when they do move against each other are often strong enough to cause major earthquakes.

At still other locations, the plates are moving apart. As they recede, new material from the mantle wells up


between them, forming **midocean ridges**. Notice in Figure 7.11 the major boundary separating the North and South American plates from the Eurasian and African plates, marked by the thin strip down the middle of the Atlantic Ocean. Discovered after World War II by oceanographic ships studying the geography of the seafloor, this giant fault is called the *Mid-Atlantic Ridge*. It extends, like a seam on a giant baseball, all the way from Scandinavia in the North Atlantic to the latitude of Cape Horn at the southern tip of South America. The entire ridge is a region of seismic and volcanic activity, but the only major part of it that rises above sea level is the island of Iceland.

Robot submarines have retrieved samples of the ocean floor at a variety of locations on either side of the Mid-Atlantic Ridge, and the ages of the samples have been measured by means of radioactive dating techniques. As depicted in Figure 7.15, the ocean floor closest to the ridge is relatively young, whereas material farther away, on either side, is older—exactly as we would expect if hot molten matter is upwelling and solidifying as the plates on either side drift apart. The Atlantic Ocean has apparently been growing in this way for the past 200 million years, the age of the oldest rocks found on any part of the Atlantic seafloor.


Other studies of the Mid-Atlantic Ridge yield important information about Earth's magnetic field. As hot material (carrying traces of iron) from the mantle emerges from cracks in the oceanic ridges and solidifies,





(a)



(b)




◀ **FIGURE 7.13 Himalayas** (a) The subcontinent of India, imaged here by sensing infrared radiation from orbit, lies at the northernmost tip of the Indian plate. As this plate drifts northward, the Indian landmass collides with Asia, on the Eurasian plate. The impact causes Earth's crust to buckle and fold, thrusting up the Himalayan mountain range (covered with snow at the upper right). (b) The results of the ongoing process depicted in (a) can be seen in this view of the area near Mount Everest. (© Michael Klesius/National Geographic Image Collection)



it becomes slightly magnetized, retaining an imprint of Earth's magnetic field *at the time it cooled*. Thus, the ocean floor has preserved within it a record of Earth's magnetism during past times, rather like a tape recording. Samples of ocean floor close to the ridge are magnetized in the same sense as Earth's field today. This material is "young" basalt that upwelled and cooled fairly recently. However, samples retrieved farther from the ridge, corresponding to older material that upwelled long ago, are often magnetized with the *opposite* orientation. As we move away from the ridge, the imprinted magnetic field flips back and forth, more or less regularly and symmetrically on either side of the ridge.

The leading explanation of these different magnetic orientations is that they were caused by reversals in Earth's magnetic field that occurred as the plates drifted away from the central ridge. Taken in conjunction with the data on the age of the seafloor, these measurements allow us to time our planet's magnetic reversals. On average, Earth's magnetic field reverses itself roughly

◀ **FIGURE 7.14 Californian Fault** The San Andreas Fault is the result of two plates sliding past one another. The Pacific plate, which includes a large slice of the California coast, is drifting to the northwest relative to the North American plate. (D. Parker/Science Source)



◀ **FIGURE 7.15 Seafloor Spreading**

Samples of ocean floor retrieved by oceanographic vessels are youngest close to the Mid-Atlantic Ridge and progressively older farther away.

every half-million years. Current theory suggests that such reversals are part of the way in which all planetary magnetic fields are generated. As we will see in Chapter 16, similar phenomenon (with a reversal time of approximately 11 years) is also observed on the Sun.


## What Drives the Plates?

What process is responsible for the enormous forces that drag plates apart in some locations and ram them together in others? The answer is probably convection—the same physical process we encountered earlier in our study of the atmosphere. Figure 7.16 is a cross-sectional diagram of the top few hundred kilometers of our planet's interior. It depicts roughly the region in and around a midocean ridge. There, the ocean floor is covered with a layer of sediment—dirt, sand, and dead sea organisms that have fallen through the seawater for millions of years. Below the sediment lies about 10 km of granite, the low-density rock that makes up the crust. Deeper still lies the upper mantle, whose temperature increases with depth. Below

the base of the lithosphere, at a depth of perhaps 50 km, the temperature is sufficiently high that the mantle is soft enough to flow very slowly, although it is not molten. This region is the asthenosphere.

The setting is a perfect one for convection—warm matter underlying cool matter. The warm mantle rock rises, just as hot air rises in our atmosphere. Sometimes, the rock squeezes up through cracks in the granite crust. Every so often, such a fissure may open in the midst of a continental landmass, producing a volcano such as Mount St. Helens (see Figure 7.10b) or possibly a geyser like those at Yellowstone National Park. However, most such cracks are on the ocean floor. The Mid-Atlantic Ridge is a prime example.

Not all the rising warm rock in the upper mantle can squeeze through cracks and fissures. Some warm rock cools and falls back down to lower levels. In this way, large circulation patterns become established within the upper mantle, as depicted in Figure 7.16. Riding atop these convection patterns are the plates. The circulation is extraordinarily sluggish. Semisolid rock takes millions of years to complete one



**Interactive FIGURE 7.16 Plate Drift** The motion of Earth's tectonic plates is caused by convection—giant circulation patterns in the upper mantle that drag the plates across the surface.


convection cycle. Although the details are far from certain and remain controversial, many researchers suspect that the large-scale circulation patterns near plate boundaries drive the motions of the plates.

This constant recycling of plate material provides a natural explanation for the **rock cycle**—the process by which surface rock on our planet is continuously redistributed and transformed from one type into another. Deep below the surface, in the asthenosphere, temperatures are high enough that mantle rock exists in the form of molten *magma*. When this material cools and hardens, it forms *igneous rocks*. (Granite and basalt are familiar examples.) Igneous rocks are associated with volcanic activity (in which the magma is called *lava*) and spreading regions such as the Mid-Atlantic Ridge, where magma emerges as two plates separate. The weathering and erosion of surface rocks produce sandy grains that are deposited as sediments and may eventually become compacted into *sedimentary rocks* such as sandstone and shale. Subsequently, at high temperatures or pressures, igneous or sedimentary rocks may be physically or chemically transformed into *metamorphic rocks* (e.g., marble and slate). Such conditions occur as plates collide and form mountain ranges or as a plate dives deep into a subduction zone.

## Past Continental Drift

Figure 7.17 illustrates how all the continents nearly fit together like pieces of a puzzle. Geologists think that sometime in the past a single gargantuan landmass dominated our planet. This ancestral supercontinent, known as *Pangaea* (meaning “all lands”), is shown in Figure 7.17(a). The rest of the planet was presumably covered with water. The present locations of the continents, along with measurements of their current drift rates, suggest that *Pangaea* was the major land feature on Earth approximately 200 million years ago. Dinosaurs, which were then the dominant form of life, could have sauntered from Russia to Texas via Boston without getting their feet wet. *Pangaea* explains the geographical and fossil evidence (cited earlier) that first led scientists to the idea of continental drift. The other frames in Figure 7.17 show how *Pangaea* split apart, its separate pieces drifting across Earth’s surface, eventually becoming the familiar continents we know today.

There is nothing particularly special about a time 200 million years in the past. We don’t suppose that *Pangaea* remained intact for 4 billion years after the crust first formed, only to break up so suddenly and so recently. It is much more plausible that *Pangaea* formed after an earlier period during which other plates, carrying widely separated continental masses, were driven together by tectonic forces, merging their landmasses into a single supercontinent. There has probably been a long series of “*Pangaeas*” stretching back in time over much of Earth’s history, as tectonic forces have continually formed, destroyed, and re-formed our planet’s landmasses. There will likely be many more.



▲ **FIGURE 7.17 Pangaea** Given their current estimated drift rates and directions, the plate movements can be traced back into the past. About 200 million years ago, they would have been at the approximate positions shown in (a). The continents’ current positions are shown in (d).

**CONCEPT Check**

- ✓ Describe the causes, and some consequences, of plate tectonics on Earth.

## 7.5 Earth's Magnetosphere

Simply put, the *magnetosphere* is the region around a planet that is influenced by that planet's magnetic field. Discovered by artificial satellites launched in the late 1950s and sketched in Figure 7.18, Earth's magnetosphere extends far above the atmosphere, completely surrounding our planet. Close to Earth, the magnetic field is similar in overall structure to the field of a gigantic bar magnet (Figure 7.19). The *magnetic field lines*, which indicate the strength and direction of the field at any point in space, run from south to north, as indicated by the blue arrowheads in these figures.

The north and south *magnetic poles*, where the magnetic field lines intersect Earth's surface vertically, are roughly aligned with Earth's spin axis. Neither pole is fixed relative to our planet, however—both drift at a rate of some 10 km per year—nor are the poles symmetrically placed. At present, Earth's magnetic north pole lies in northern Canada, at a latitude of about 80° N, almost due north of the center of North America; the magnetic south pole lies at a latitude of about 60° S, just off the coast of Antarctica south of Adelaide, Australia.


Earth's magnetosphere contains two doughnut-shaped zones of high-energy charged particles, one located about 3000 km, and the other 20,000 km, above Earth's surface. These zones are named the **Van Allen belts**, after the American physicist whose instruments on board one of the first artificial satellites initially detected them. We call them “belts” because they are most pronounced near

Earth's equator and because they completely surround the planet. Figure 7.19 shows how these invisible regions envelop Earth, except near the North and South Poles.


The particles that make up the Van Allen belts originate in the solar wind—the steady stream of charged particles flowing from the Sun. ∞ (Sec. 6.5) Traveling through space, neutral particles and electromagnetic radiation are unaffected by Earth's magnetism, but electrically charged particles are strongly influenced. As illustrated in the inset to Figure 7.19, a magnetic field exerts a force on a moving charged particle, causing the particle to spiral around the magnetic field lines. In this way, charged particles—mainly electrons and protons—from the solar wind can become trapped by Earth's magnetism, and herded into the Van Allen belts. The outer belt contains mostly electrons; the much heavier protons accumulate in the inner belt.

We could never survive unprotected in the Van Allen belts. Unlike the lower atmosphere, on which humans and other life-forms rely for warmth and protection, much of the magnetosphere is subject to intense bombardment by large numbers of high-velocity, and potentially very harmful, charged particles. Colliding violently with an unprotected human body, these particles would deposit large amounts of energy wherever they made contact, causing severe damage to living organisms. Without sufficient shielding on the *Apollo* spacecraft, for example, the astronauts might not have survived the passage through the magnetosphere on their journey to the Moon.

Particles from the Van Allen belts often escape from the magnetosphere near Earth's north and south magnetic poles, where the field lines intersect the atmosphere. Their collisions with air molecules create a spectacular light show called an **aurora** (plural *aurorae*; Figure 7.20). This colorful display



◀ **FIGURE 7.18 Earth's Magnetosphere** The magnetosphere is the region surrounding a planet wherein particles from the solar wind are trapped by the planet's magnetic field. Far from Earth, the magnetosphere is greatly distorted by the solar wind, with a long “tail” extending from the nighttime side of Earth (here, at right) far into space. The magnetopause is the boundary of the magnetosphere in the sunward direction.



▲ **FIGURE 7.19 Van Allen Belts** Earth's magnetic field resembles the field of an enormous bar magnet buried deep inside our planet (but offset slightly from Earth's axis of rotation). High above Earth's atmosphere, the magnetosphere (light blue-green area) contains two doughnut-shaped regions (grayish areas) of magnetically trapped charged particles. These are the Van Allen belts. The convergence of the field lines near Earth's magnetic poles causes the particles to be reflected back toward the other pole.

results when atmospheric molecules, excited upon collision with the charged particles, fall back to their ground states and emit visible light. Many different colors are produced because each type of atom or molecule can take one of several possible paths as it returns to its ground state. ∞ (Sec. 4.3) Aurorae are most brilliant at high latitudes, especially inside the Arctic and Antarctic circles. In the north, the spectacle is called the *aurora borealis*, or *northern lights*. In the south, it is called the *aurora australis*, or *southern lights*.

Occasionally, particularly after a storm on the Sun (see Chapter 16), the Van Allen belts can become distorted by the solar wind and overloaded with many more particles than normal, allowing some particles to escape prematurely and at lower latitudes. For example, in North America, the *aurora borealis* is normally seen with any regularity only in

northern Canada and Alaska. However, at times of greatest solar activity, the display has occasionally been seen as far south as the southern United States.

As is evident from Figure 7.18, Earth's magnetosphere is not symmetrical. Satellite mapping reveals that it is quite distorted, forming a teardrop-shaped cavity. On the sunlit (daytime) side of Earth, the magnetosphere is compressed by the flow of high-energy particles in the solar wind. The boundary between the magnetosphere and this flow, known as the *magnetopause*, is found at about 10 Earth radii from our planet. On the side opposite the Sun, the field lines are extended away from Earth, with a long tail often reaching beyond the orbit of the Moon.


What is the origin of the magnetosphere and the Van Allen belts within it? Despite the artistic license in



(a)



(b)



▲ **FIGURE 7.20 Aurorae** (a) A colorful aurora flashes rapidly across the sky, resembling huge windblown curtains glowing in the dark. Aurorae result from the emission of light radiation after magnetospheric particles collide with atmospheric molecules. The colors are produced as excited ions, atoms, and molecules recombine and cascade back to their ground states. (b) An aurora high above Earth, as photographed from a space shuttle (visible at left). (NCAR; NASA)

Figure 7.19, Earth’s magnetism is not really the result of a huge bar magnet lying within our planet. In fact, geophysicists think that Earth’s magnetic field is not a “permanent” part of our planet at all. Instead, it is thought to be continuously generated within the outer core and to exist only because Earth is rotating. As in the dynamos that run industrial machines, Earth’s magnetism is produced by the spinning, electrically conducting, liquid metal core deep within our planet. The theory that explains planetary (and other) magnetic fields in terms of rotating, conducting material flowing in the planet’s interior is known as **dynamo theory**. Both rapid rotation *and* a conducting liquid core are needed for such a mechanism to work. This connection between internal structure and magnetism is very important for studies of the other planets in the solar system: We can tell a lot about a planet’s interior simply by measuring its magnetic field.

Earth’s magnetic field plays an important role in controlling many of the potentially destructive charged particles that venture near our planet. Without the magnetosphere, Earth’s atmosphere—and perhaps the surface, too—would be bombarded by harmful particles, possibly damaging many forms of life on our planet. Some researchers have even suggested that, had the magnetosphere not existed, life might never have arisen on our planet.

### PROCESS OF SCIENCE Check

- ✓ What does the existence of a planetary magnetic field tell us about a planet’s interior?

## 7.6 The Tides

Earth is unique among the planets in that it has large quantities of liquid water on its surface. Approximately three-quarters of Earth’s surface is covered by water, to an average depth of about 3.6 km. Only 2 percent of the water is contained within lakes, rivers, clouds, and glaciers. The remaining 98 percent is in the oceans, forming the *hydrosphere*.

Most people are familiar with the daily fluctuation in ocean level known as the **tides**. At most coastal locations on Earth, there are two low tides and two high tides each day. The “height” of the tides—the magnitude of the variation in sea level—can range from a few centimeters to many meters, depending on the location on Earth and the time of year. The height of a typical tide on the open ocean is about a meter, but if this tide is funneled into a narrow opening such as the mouth of a river, it can become much higher. For example, at the Bay of Fundy, on the U.S.–Canada border between Maine and New Brunswick, the high tide can reach nearly 20 m (approximately 60 feet, or the height of a six-story building) above the low-tide level. An enormous amount of energy is contained in the daily motion of the oceans. This energy is constantly eroding and reshaping our planet’s coastlines. In some locations, it has been harnessed as a source of electrical power for human activities.

### Gravitational Deformation

What causes the tides? A clue comes from the observation that they exhibit daily, monthly, and yearly cycles. In fact, the tides are a direct result of the gravitational influence of the Moon

and the Sun on Earth. We have already seen how gravity keeps Earth and the Moon in orbit about each other, and both in orbit around the Sun.  $\infty$  (Sec. 2.7) For simplicity, let's first consider just the interaction between Earth and the Moon.

Recall that the strength of the gravitational force depends on the distance separating any two objects. Thus, the Moon's gravitational attraction is greater on the side of Earth that faces the Moon than on the opposite side, some 12,800 km (Earth's diameter) farther away. This difference in the gravitational force is small—only about 3 percent—but it produces a noticeable effect—a **tidal bulge**. As illustrated in Figure 7.21, Earth becomes slightly elongated, with the long axis of the distortion pointing toward the Moon.

Earth's oceans undergo the greatest deformation, because liquid can most easily move around on our planet's surface. (A bulge *is* actually raised in the solid material of Earth, but it is about a hundred times smaller than the oceanic bulge.) Thus, the ocean becomes a little deeper in some places (along the line joining Earth to the Moon) and shallower in others (perpendicular to this line). The daily tides we experience result as Earth rotates beneath this deformation.

The variation in the Moon's gravity across Earth is an example of a *differential force*, or **tidal force**. The average gravitational force between two bodies determines their orbit around one another. However, the *tidal force*, superimposed on that average, tends to deform the bodies. The tidal influence of one body on another diminishes very rapidly with increasing distance—in fact, as the inverse *cube* of the separation. For example, if the distance from Earth to the Moon were to double, the tides resulting from the Moon's gravity would decrease by a factor of eight. This rapid decline with increasing distance means that one object has to be very close or very massive in order to have a significant tidal effect on another.

We will see many situations in this book where tidal forces are critically important in understanding astronomical phenomena. We still use the word *tidal* in these other contexts, even though we are not discussing oceanic tides and, possibly, not even planets at all. In general astronomical use, the term refers to the deforming effect that the gravity of one body has on another.


Notice in Figure 7.21 that the side of Earth opposite the Moon also exhibits a tidal bulge. The different gravitational pulls—greatest on that part of Earth closest to the Moon, weaker at Earth's center, and weakest of all on Earth's opposite side—cause average tides on opposite sides of our planet to be approximately equal in height. On the side nearer the Moon, the ocean water is pulled slightly toward the Moon. On the opposite side, the ocean water is left behind as Earth is pulled closer to the Moon. Thus, high tide occurs *twice*, not once, each day at any given location.

Both the Moon and the Sun exert tidal forces on our planet. Thus, instead of one tidal bulge, there are actually two—one pointing toward the Moon, the other toward the Sun. Even though the Sun is 375 times farther away from

Earth than is the Moon, the Sun's mass is so much greater (by a factor of 27 million) that its tidal influence is still significant—about half that of the Moon. The interaction between them accounts for the changes in the height of the tides over the course of a month or a year. When Earth, the Moon, and the Sun are roughly lined up (Figure 7.22a), the gravitational effects reinforce one another, so the highest tides are generally found at times of new and full moons. These tides are known as *spring tides*. When the Earth–Moon line is perpendicular to the Earth–Sun line (at the first and third quarters; Figure 7.22b), the daily tides are smallest. These are termed *neap tides*.


## Earth's Slowing Rotation

Earth rotates once on its axis (relative to the stars) in  $23^{\text{h}} 56^{\text{m}}$ —one sidereal day. However, we know from fossil measurements that Earth's rotation is gradually slowing down, causing the length of the day to increase by about 1.5 milliseconds (ms) every century—not much on the scale of a human lifetime, but over millions of years, this steady slowing of Earth's spin adds up. At this rate, half a billion years ago, the day was just over 22 hours long and the year contained 397 days.



▲ **FIGURE 7.21 Lunar Tides** This illustration, which is not to scale, shows how the Moon induces tides on both the near and far sides of Earth. (a) The lunar gravitational force is greatest on the side nearest the Moon and smallest on the opposite side. (b) The *differences* between the lunar forces experienced at the locations shown in part (a) and the force exerted by the Moon on Earth's center indeed make a difference. Closest to the Moon, the oceans tend to be pulled away from Earth; on the far side, Earth tends to be pulled away from the oceans. The result is tidal bulges that we on Earth experience each and every day.





**Interactive FIGURE 7.22 Solar and Lunar Tides** The combined effects of the Sun and the Moon produce variations in high and low tides. (a) When the Moon is either full or new, Earth, Moon, and Sun are approximately aligned, and the tidal bulges raised in Earth's oceans by the Moon and the Sun reinforce one another. (b) When the Moon is in its first or third quarter, the tidal effects of the Moon and the Sun partially cancel each other, and the tides are smallest.

shown in Figure 7.21. Instead, because of the effects of friction, both between the crust and the oceans and within Earth itself, Earth's rotation tends to drag the tidal bulge around with it, causing the bulge to be displaced by a small angle from the Earth–Moon line, in the same direction as Earth's spin (Figure 7.23). The net effect of the Moon's gravitational pull on this slightly offset bulge is to *reduce* our planet's rotation rate. At the same time, the Moon is spiraling slowly away from Earth, increasing its average distance from our planet by about 4 cm per year.

A number of natural biological clocks lead us to the conclusion that Earth's spin rate is decreasing. For example, each day a growth mark is deposited on a certain type of coral in the reefs off the Bahamas. These growth marks are similar to the annual rings found in tree trunks, except that in the case of coral, the marks are made daily, in response to the day–night cycle of solar illumination. However, they also show yearly variations as the coral's growth responds to Earth's seasonal changes, allowing us to perceive annual cycles. Coral growing today shows 365 marks per year, but ancient coral shows many more growth deposits per year. Fossilized reefs that are five hundred million years old contain coral with nearly 400 deposits per year of growth.


Why is Earth's spin slowing? The main reason is the tidal effect of the Moon. In reality, the tidal bulge raised in Earth by the Moon does *not* point directly at the Moon, as was

This process will continue until Earth rotates on its axis at exactly the same rate as the Moon orbits Earth. At that time, the Moon will always be above the same point on Earth and will no longer lag behind the bulge it raises. Earth's rotation period will be 47 of our present days, and the distance to the Moon will be 550,000 km (about 43 percent greater than at present). However, this will take a very long time—many billions of years—to occur.

### CONCEPT Check

✓ In what ways do tidal forces differ from the familiar inverse-square force of gravity?

► **FIGURE 7.23 Tidal Bulge** The tidal bulge raised in Earth by the Moon does not point directly at the Moon. Instead, because of the effects of friction, the bulge points slightly ahead of the Moon, in the direction of Earth's rotation. (The magnitude of the effect is greatly exaggerated in this diagram, which is not to scale.) Because the Moon's gravitational pull on the near-side part of the bulge is greater than the pull on the far side, the overall effect is to decrease Earth's rotation rate.




**The Big Question** Not long after Earth formed, debris bombardment from outside and radioactive heating from inside caused the whole planet to melt. Any water present early on would have evaporated and escaped. So, where did all the water now on Earth come from? Nearly three-quarters of Earth's surface is abundant in water, and to great ocean depths. In fact, Earth has so much water—just look at the chapter opening photo on page 160—that it might have been more properly called Aqua. One possibility is that comets, which are hardly more than dirty ice balls, delivered the water. Another is that water upwelled from inside our planet during early volcanism. No one knows for sure.


# Chapter Review

## SUMMARY

**1** The six main regions of Earth are (from inside to outside) a central metallic **core** (p. 162), which is surrounded by a thick rocky **mantle** (p. 162), topped with a thin **crust** (p. 162). The liquid oceans on our planet's surface make up the **hydrosphere** (p. 162). Above the surface is the **atmosphere** (p. 174), which is composed primarily of nitrogen and oxygen and thins rapidly with altitude. Surface winds and weather in the **troposphere** (p. 163), the lowest region of Earth's atmosphere, are caused by **convection** (p. 163), the process by which heat is moved from one place to another by the upwelling or downflow of a fluid, such as air or water. Higher above the atmosphere lies the **magnetosphere** (p. 162), where charged particles from the Sun are trapped by Earth's magnetic field.




**2** At high altitudes, in the **ionosphere** (p. 163), the atmosphere is kept ionized by the absorption of high-energy radiation and particles from the Sun. In the **stratosphere** (p. 163), just above the troposphere, lies the **ozone layer** (p. 164), where incoming solar ultraviolet radiation is absorbed. Both the ionosphere and the ozone layer help protect us from dangerous radiation from space. The **greenhouse effect** (p. 166) is the absorption and trapping of infrared radiation emitted by Earth's surface by atmospheric gases (primarily carbon dioxide and water vapor). It makes our planet's surface some 40 K warmer than would otherwise be the case. Earth's atmosphere was outgassed from our planet's interior by volcanoes and was then altered by solar radiation and, finally, by the emergence of life.



**3** We study Earth's interior by observing how **seismic waves** (p. 168), produced by earthquakes just below Earth's surface, travel through the mantle. We can also study the upper mantle by analyzing the material brought to the surface when a volcano erupts. Earth's center is dense and extremely hot. The planet's iron core consists of a solid **inner core** (p. 169) surrounded by a liquid **outer core** (p. 169). The process by


which heavy material sinks to the center of a planet and lighter material rises to the surface is called **differentiation** (p. 170). Earth's differentiation implies that our planet must have been at least partially molten in the past. One way in which this could have occurred is by the heat released during Earth's formation and subsequent bombardment by material from interplanetary space. Another possibility is the energy released by the decay of **radioactive** (p. 171) elements present in the material from which Earth formed.



**4** Earth's surface is made up of about a dozen enormous slabs, or plates. The slow movement of these plates across the surface is called continental drift or **plate tectonics** (p. 174). Earthquakes, volcanism, and mountain building are associated with plate boundaries, where plates may collide, move apart, or rub against one another. The motion of the plates is thought to be driven by convection in Earth's mantle. The rocky upper layer of Earth that makes up the plates is the **lithosphere** (p. 174). The semisolid region in the upper mantle over which the plates slide is called the **asthenosphere** (p. 174). The constant recycling and transformation of crust material as plates separate, collide, and sink into the mantle is called the **rock cycle** (p. 179). Evidence for past plate motion can be found in the geographical fit of continents, in the fossil record, and in the ages and magnetism of surface rocks.



**5** Earth's magnetic field extends far beyond the surface of our planet. Charged particles from the solar wind are trapped by Earth's magnetic field lines to form the **Van Allen belts** (p. 180) that surround our planet. When particles



from the Van Allen belts hit Earth's atmosphere, they heat and ionize the atoms there, causing them to glow in an **aurora** (p. 180). According to **dynamo theory** (p. 182), planetary magnetic fields are produced by the motion of rapidly rotating, electrically conducting fluid (such as molten iron) in the planet's core.

**6** The daily **tides** (p. 182) in Earth's oceans are caused by the gravitational effect of the Moon and the Sun, which

raise **tidal bulges** (p. 183) in the hydrosphere. The tidal effect of the Moon is almost twice that of the Sun. The size of the tides depends on the orientations of the Sun and the Moon relative to Earth. The tidal interaction between Earth and the Moon is causing Earth's spin to slow.



**MasteringAstronomy**<sup>®</sup> For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- LO1** By comparison with Earth's average density, what do the densities of the water and rocks in Earth's crust tell us about Earth's interior?
- POS** How do geologists use earthquakes to obtain information about Earth's interior?
- LO2** What is the greenhouse effect, and what effect does it have on Earth's surface temperature?
- LO3 POS** Give two reasons geologists think that part of Earth's core is liquid.
- What clue to our planet's history does Earth's differentiation provide?
- What is convection? What effect does it have on (a) Earth's atmosphere? (b) Earth's interior?
- How did radioactive decay heat Earth early in its history? When did this heating end?
- LO4** What process is responsible for surface mountains, oceanic trenches, and other large-scale features on Earth's surface?
- What conditions are needed to create a dynamo in Earth's interior? What effect does this dynamo have?
- LO5** Give a brief description of Earth's magnetosphere, and tell how it was discovered. How does the magnetosphere protect us from the harsh realities of interplanetary space?
- POS** How do we know that Earth's magnetic field has undergone reversals in the past? How do you think Earth's magnetic field reversals might have affected the evolution of life on our planet?
- LO6** Explain how the Moon produces tides in Earth's oceans.
- If the Moon had oceans like Earth's, what would the tidal effect be like there? How many high and low tides would there be during a "day"? How would the variations in height compare with those on Earth?
- POS** If Earth had no moon, do you think we would know anything about tidal forces?
- Is the greenhouse effect operating in Earth's atmosphere helpful or harmful? Give examples. What are the consequences of an enhanced greenhouse effect?

## Conceptual Self-Test: Multiple Choice

- If you were making a scale model of Earth representing our planet by a 12-inch basketball, the inner core would be about the size of (a) a  $\frac{1}{2}$ -inch ball bearing; (b) a 2-inch golf ball; (c) a 4-inch tangerine; (d) a 7-inch grapefruit.
- Earth's average density is about the same as that of (a) a glass of water; (b) a heavy iron meteorite; (c) an ice cube; (d) a chunk of black volcanic rock.
- VIS** According to Figure 7.2 ("Earth's Atmosphere"), commercial jet airplanes flying at 10 km are in (a) the troposphere; (b) the stratosphere; (c) the ozone; (d) the mesosphere.
- If there were significantly more greenhouse gases, such as  $\text{CO}_2$ , in Earth's atmosphere, then (a) the ozone hole would close; (b) the ozone hole would get larger; (c) Earth's average temperature would change; (d) plants would grow faster than animals could eat them.
- If seismometers registered P- and S-waves everywhere on the Moon, they would suggest that the Moon had (a) the same layered structure as Earth; (b) no molten core; (c) no moonquakes; (d) the same density throughout.
- The deepest that geologists have drilled into Earth is about the same as (a) the height of the Statue of Liberty; (b) the altitude most commercial jet airplanes fly; (c) the distance between New York and Los Angeles; (d) the distance between the United States and China.
- Due to plate tectonics, the width of the Atlantic Ocean is separating at a rate about the same as the growth of

- (a) grass; (b) human hair; (c) human fingernails; (d) dust in a typical home.
8. At Earth's geographic North Pole, a magnetic compass needle would point (approximately) (a) toward Alaska; (b) toward Kansas City; (c) toward Paris; (d) straight down.
9. If Earth had no Moon, then tides would (a) not occur; (b) occur more often and with more intensity; (c) still occur, but not really be measurable; (d) occur with the same frequency, but would not be as strong.
10. Which of the following statements is true? Because of the tides, (a) Earth's rotation rate is increasing; (b) the Moon is spiraling away from Earth; (c) Earth will eventually drift away from the Sun; (d) earthquake activity is increasing.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

- Approximating Earth's atmosphere as a layer of gas 7.5 km thick, with uniform density  $1.3 \text{ kg/m}^3$ , calculate the total mass of the atmosphere. Compare your result with Earth's mass.
- Most of Earth's ice is found in Antarctica, where permanent ice caps cover approximately 0.5 percent of Earth's total surface area and are 3 km thick, on average. Earth's oceans cover roughly 71 percent of our planet, to an average depth of 3.6 km. Assuming that water and ice have roughly the same density, estimate by how much sea level would rise if global warming were to cause the Antarctic ice caps to melt.
- On the basis of the data presented in the text, estimate the fractions of Earth's volume represented by (a) the inner core, (b) the outer core, (c) the mantle, and (d) the crust.
- Following an earthquake, how long would it take a P-wave, moving in a straight line with a speed of 5 km/s, to reach Earth's opposite side?
- At 3 cm/yr, how long would it take a typical plate to traverse the present width of the Atlantic Ocean, about 6000 km?
- A certain sample of rock is found to contain three times as many lead-207 nuclei as uranium-235 nuclei. On the basis of the data given in *More Precisely 7-2*, what ratio of uranium-238 to lead-206 nuclei would you expect?
- In a second sample of rock, it is found that 25 percent of uranium-238 nuclei have decayed into lead-206. On the basis of the data given in *More Precisely 7-2*, estimate the age of the rock sample.
- You are standing on Earth's surface, and the full Moon is directly overhead. By what fraction is your weight decreased due to the combination of the Sun's and the Moon's tidal gravitational forces?

## Activities

### Collaborative

- Go online and read about global warming. How much carbon dioxide is produced each year by human activities? How does this compare with the total amount of carbon dioxide in Earth's atmosphere? Do all—or most—scientists agree that global warming is an inevitable consequence of carbon dioxide production? What political initiatives are currently under way to address the problem? As a group, which, if any, do you think are likely to succeed?
- Using a ruler and self-stick or taped-on labels, create a scale model of Earth on the shortest member of your group. Use the group member's height in inches divided by Earth's diameter (12,800 km) as the scale factor along with

Figure 7.1. For example, if the selected group member is 65 inches tall then the 50 km maximum depth of Earth's crust is  $(65 \text{ inches}/12,800 \text{ km}) \times (50 \text{ km}) = 0.25 \text{ inches}$  from the top of the head and 0.25 inches from the bottom of the feet.

### Individual

- Go to a sporting goods store and get a tide table; many stores near the ocean provide them free. Choose a month, and plot the height of one high and one low tide versus the day of the month. Now mark the dates when the primary phases of the Moon occur. How well does the phase of the Moon predict the tides?



# 8

# The Moon and Mercury

## SCORCHED AND BATTERED WORLDS

The Moon is Earth's only natural satellite. Mercury, the smallest terrestrial world, is the planet closest to the Sun. These two small bodies have much in common—indeed, at first glance, you might even mistake one for the other. Both have heavily cratered, ancient surfaces, littered with boulders and pulverized dust. Both lack atmospheres to moderate day-to-night variations in solar heating and experience wild temperature swings as a result. Both are geologically dead.

Yet despite their superficial similarity, the Moon and Mercury differ significantly from each other in many important ways. And it is precisely those differences that make these desolate worlds so interesting to planetary scientists. Why is the Moon so unlike our own planet, despite its nearness to us, and why did Mercury turn out so different from both Earth and the Moon? In this chapter, we explore the properties of these two worlds as we begin our comparative study of the planets and moons that make up our solar system.

**The Big Picture** The Moon, Mercury, and Earth are all quite different places. Smaller, colder, hotter, drier, among other detrimental qualities, crater-scarred Moon and Mercury resemble nothing like the warm, lush, comfortable planet on which we live. Although there's no life on these alien worlds, it doesn't necessarily mean there never will be, especially on our neighboring Moon. Despite its magnificent desolation, the Moon might one day harbor a colony of human settlers, or be a way station (or rest area) in humankind's future voyages to more hospitable planets and moons.

**LEFT:** America's manned exploration of the Moon was arguably the greatest engineering feat of the 20th century, perhaps one of the greatest of all time. Nine crewed missions were launched to the Moon, a dozen astronauts were landed, and all returned safely to Earth. Here, an *Apollo 16* astronaut is prospecting near the rim of Plum Crater for rock samples that might help reveal the origin of the Moon. The "rover" that carried him several kilometers from his landing craft can be seen in the left background. Given the lack of wind and water on the Moon, the bootprints in the foreground are destined to survive for more than a million years. (NASA)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Specify the general characteristics of the Moon and Mercury, and compare them with those of Earth.
- 2 Describe the surface features of the Moon and Mercury, and recount how those two bodies were formed by events early in their history.
- 3 Explain how the Moon's rotation is influenced by its orbit around Earth and Mercury's by its orbit around the Sun.
- 4 Explain how observations of cratering can be used to estimate the age of a body's surface.
- 5 Describe the evidence for ancient volcanism on the Moon and Mercury.
- 6 Compare the Moon's interior structure with that of Mercury.
- 7 Summarize the leading theory of the formation of the Moon.
- 8 Outline how astronomers have pieced together the story of the Moon's evolution, and compare its evolutionary history with that of Mercury.


### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.

## 8.1 Orbital Properties

We begin our study of the Moon and Mercury by examining their orbits. This knowledge will, in turn, aid us in determining and explaining the other properties of these worlds.

### The Moon


Parallax methods, described in Chapter 1, can provide us with quite accurate measurements of the distance to the Moon, using Earth's diameter as a baseline.  (Sec. 1.6) Radar ranging yields more accurate distances. The Moon is much closer than any of the planets, and the radar echo bounced off the Moon's surface is strong. A radio telescope receives the echo after a round trip of 2.56 seconds. Dividing this time by 2 and multiplying it by the speed of light (300,000 km/s) gives us a distance of 384,000 km. (The actual distance at any particular time depends on the Moon's location in its slightly elliptical orbit around Earth.)

Current laser-ranging technology, using reflectors placed on the lunar surface by *Apollo* astronauts to reflect laser beams fired from Earth, allows astronomers to measure the round-trip time with submicrosecond accuracy. Repeated measurements have allowed astronomers to determine the Moon's orbit to within a few centimeters. This precision is necessary for programming unmanned spacecraft to land successfully on the lunar surface.


### Mercury

Viewed from Earth, Mercury never strays far from the Sun. As illustrated in Figure 8.1(a), the planet's 0.4-AU orbital semimajor axis means that its angular distance from the Sun never exceeds  $28^\circ$ . Consequently, the planet is visible to the naked eye only when the Sun's light is blotted out—just before dawn or just after sunset (or, much less frequently, during a total solar eclipse)—and it is not possible to follow Mercury through a full cycle of phases. In fact, although Mercury was well known to ancient astronomers, they originally believed that this companion to the Sun was two different objects, and the connection between the planet's morning and evening appearances took some time to establish. However, later Greek astronomers were certainly aware that the “two planets” were really different alignments of a single body. Figure 8.1(b), a photograph taken just after sunset, shows Mercury above the western horizon, along with three other planets and the Moon.


Because Earth rotates at a rate of  $15^\circ$  per hour, Mercury is visible for at most 2 hours on any given night, even under the most favorable circumstances. For most observers at most times of the year, Mercury is generally visible for a much



(a)




(b)



**▲ FIGURE 8.1 Evening Sky** (a) Mercury's orbit has a semimajor axis of just 0.4 AU, so the planet can never be farther than  $28^\circ$  from the Sun, as seen from Earth. Mercury's eccentric orbit means that this maximum separation is achieved only for the special configuration shown here, in which the Earth-Sun line is perpendicular to the long axis of Mercury's orbit and Mercury is near aphelion (its greatest distance from the Sun). (b) Four planets, together with the Moon, are visible in this photograph taken shortly after sunset. (*J. Sanford/Science Source*)

shorter period. Nowadays, large telescopes can filter out the Sun's glare and observe Mercury even during the daytime, when the planet is higher in the sky and atmospheric effects are reduced. (The amount of air that the light from the planet has to traverse before reaching our telescope decreases as



**FIGURE 8.2 Phases of Mercury**  
Mercury's appearance, as seen by us on Earth, changes at different points along its orbit. The best images of the planet (insets) are taken when it is at its maximum elongation (greatest apparent distance from the Sun) and show a "half Mercury" (cf. Figure 2.12a). (R. Beebe)

the height of the planet above the horizon increases.) In fact, some of the best views of Mercury have been obtained in this way. The naked-eye or amateur astronomer is generally limited to nighttime observations, however.

In all cases, it becomes progressively more difficult to view Mercury the closer (in the sky) its orbit takes it to the Sun. The best images of the planet therefore show a "half Mercury," close to its maximum angular separation from the Sun, or *maximum elongation*, as illustrated in Figure 8.2. (A planet's *elongation* is just its angular distance from the Sun, as seen from Earth.)

## 8.2 Physical Properties

From Earth, the Moon's angular diameter is about  $0.5^\circ$ . Knowing that and the distance to the Moon, we can easily calculate our satellite's true size, as discussed in Chapter 1. [∞ \(More Precisely 1-2\)](#) The Moon's radius is about 1700 km, roughly one-fourth that of Earth. More precise measurements yield a lunar radius of 1738 km. We can determine Mercury's radius by similar reasoning. At its closest approach to Earth, at a distance of about 0.52 AU, Mercury's angular diameter is measured to be  $13''$  (arc seconds), implying a radius of about 2450 km, or 0.38 of Earth's radius. More accurate measurements by unmanned space probes yield a result of 2440 km.

Even before the Space Age, the masses of both the Moon and Mercury were already quite well known from studies of their effects on Earth's orbit. [∞ \(Sec. 6.2\)](#) The mass of the Moon is  $7.3 \times 10^{22}$  kg, approximately one-eightieth (0.012) the mass of Earth. The mass of Mercury is  $3.3 \times 10^{23}$  kg—about 0.055 Earth mass.

The Moon's average density of  $3300 \text{ kg/m}^3$  contrasts with the average Earth value of about  $5500 \text{ kg/m}^3$ ,

suggesting that the Moon contains fewer heavy elements (such as iron) than Earth does. In contrast, despite its many other similarities to the Moon, Mercury's mean density is  $5400 \text{ kg/m}^3$ , only slightly less than that of Earth. Assuming that surface rocks on Mercury are of similar density to surface rocks on Earth and the Moon, we are led to the conclusion that the interior of Mercury must contain a lot of high-density material, most probably iron. In fact, since Mercury is considerably less massive than Earth, its interior is squeezed less by the weight of overlying material, so Mercury's iron core must actually contain a much larger fraction of the planet's mass than does our own planet's core. [∞ \(Sec. 6.2\)](#)

Because the Moon and Mercury are so much less massive than Earth, their gravitational fields are also weaker. The force of gravity on the lunar surface is only about one-sixth that on Earth; Mercury's surface gravity is a little stronger—about 0.4 times Earth's. Thus, an astronaut weighing 180 lb on Earth would weigh a mere 30 lb on the Moon and 72 lb on Mercury. Those bulky space suits used by the *Apollo* astronauts on the Moon were not nearly as heavy as they appeared!

Astronomers have never observed any appreciable atmosphere on the Moon or Mercury, either spectroscopically from Earth or during close approaches by spacecraft. This is a direct consequence of these bodies' weak gravitational fields, as discussed in *More Precisely 8-1* (p. 196). Simply put, a massive object has a better chance of retaining an atmosphere because the more massive an object is, the larger is the speed needed for atoms or molecules to escape from the object's gravitational pull. The Moon's escape speed is only 2.4 km/s, compared with 11.2 km/s for Earth; Mercury's escape speed is 4.2 km/s. Any early atmospheres those worlds might have had are gone forever. [∞ \(Sec. 7.2\)](#)



During its flybys of Mercury in 1974 and 1975, the U.S. space probe *Mariner 10* found traces of what was at first thought to be an atmosphere on the planet. ∞ (Discovery 6-2) However, this gas is now known to be temporarily trapped hydrogen and helium “stolen” from the solar wind by the planet’s gravity. Mercury captures this gas and holds it for just a few weeks before it leaks away again into space. More recently, NASA’s *Messenger* probe found that, while the gas is indeed composed largely of hydrogen and helium, it also contains more massive atoms of sodium, potassium, and magnesium. In fact, both the Moon and Mercury have extremely tenuous atmospheres (less than a trillionth the density of Earth’s atmosphere) of such relatively heavy atoms. Scientists think that these atoms have been kicked off the surface by interactions with the solar wind; they do not constitute a true atmosphere in any sense. Thus, neither the Moon nor Mercury has any protection against the harsh environment of interplanetary space. This fact is crucial in understanding their surface evolution and present-day appearance.

Lacking the moderating influence of an atmosphere, both the Moon and Mercury are characterized by wide variations in surface temperature. Noontime temperatures at the Moon’s equator can reach 400 K, well above the boiling point of water. Because of its proximity to the Sun, Mercury’s daytime temperature is even higher—radio observations of the planet’s thermal emissions indicate that it can reach 700 K. ∞ (Sec. 3.4) But at night or in the shade, temperatures on both worlds fall to about 100 K, well below water’s freezing point. Mercury’s 600-K temperature range is the largest of any planet or moon in the solar system.

### CONCEPT Check

✓ Why do the Moon and Mercury have no significant atmospheres, unlike Earth?

## 8.3 Surface Features on the Moon and Mercury

### Lunar Terrain

The first observers to point their telescopes at the Moon—most notable among them Galileo Galilei—saw large dark areas resembling (they thought) Earth’s oceans. They also saw light-colored areas resembling the continents. Both types of regions are clear in Figure 8.3, a *mosaic* (a composite image constructed from many individual photographs) of the full Moon. The light and dark surface features are also evident to the naked eye, creating the face of the familiar “man in the Moon.”

Today we know that the dark areas are not oceans, but extensive flat areas that resulted from lava flows during




▲ **FIGURE 8.3 Full Moon, Near Side** A photographic mosaic of the full Moon, north pole at the top. Because the Moon emits no visible radiation of its own, we can see it only by the reflected light of the Sun. Some prominent maria are labeled. (UC/Lick Observatory)

a much earlier period of the Moon’s evolution. Nevertheless, they are still called **maria**, a Latin word meaning “seas” (singular: *mare*). There are 14 maria, all roughly circular. The largest of them (Mare Imbrium) is about 1100 km in diameter. The lighter areas, originally dubbed *terrae*, from the Latin word for “land,” are now known to be elevated several kilometers above the maria. Accordingly, they are usually called the lunar **highlands**.

The smallest lunar features we can distinguish with the naked eye are about 200 km across. Telescopic observations further resolve the surface into numerous bowl-shaped depressions, or **craters** (after the Greek word for “bowl”). Most craters apparently formed eons ago, primarily as the result of meteoritic impact. In Figures 8.4(a) and (b), craters are particularly clear near the *terminator* (the line that separates day from night on the surface), where the Sun is low in the sky and casts long shadows that enable us to distinguish quite small surface details.


Due to the blurring effects of our atmosphere, the smallest lunar objects that telescopes on Earth’s surface can resolve are about 1 km across (see Figure 8.4c). Much more detailed photographs have been taken by orbiting spacecraft and, of course, by visiting astronauts (see *Discovery 8-1*, p. 198). Figure 8.5 is a view of some lunar craters taken from an orbiting spacecraft, showing features as small as 500 m across. Craters are found everywhere on the Moon’s surface, although they are much more prevalent in the highlands.



**▲ FIGURE 8.4 Moon, Close Up** (a) The Moon near third quarter makes visible surface features near the *terminator*, where sunlight strikes at a sharp angle and the lit lunar globe changes to dark. (b) This magnified view of a region near the terminator, as seen from Earth through a large telescope, shows the central dark area called Mare Imbrium, which is ringed at the bottom by the Apennine mountains. (c) Enlargement of a portion of (b). The smallest craters visible here have diameters of about 2 km, about twice the size of the Barringer crater on Earth shown in Figure 8.18. (UC/Lick Observatory; Palomar)

They come in all sizes—the largest are hundreds of kilometers in diameter; the smallest are microscopic.



Based on studies of lunar rock brought back to Earth by *Apollo* astronauts and unmanned Soviet landers, geologists have identified important differences in both *composition* and *age* between the highlands and the



maria. The highlands are made largely of rocks rich in aluminum, making them lighter in color and lower in density ( $2900 \text{ kg/m}^3$ ) than the material in the maria, which contains more iron, giving it a darker color and greater density ( $3300 \text{ kg/m}^3$ ). Loosely speaking, the highlands represent the Moon's crust, whereas the maria are made of mantle material. Maria rock is quite similar to terrestrial basalt, and geologists think that it arose on the Moon much as basalt did on Earth, from the upwelling of molten material through the crust. ∞ (Sec. 7.3) Radioactive dating indicates ages of 4 to 4.4 billion years for highland rocks and from 3.2 to 3.9 billion years for those from the maria. ∞ (More Precisely 7-2)

All of the Moon's significant surface features have names. The 14 maria bear fanciful Latin names—Mare Imbrium (“Sea of Showers”), Mare Nubium (“Sea of Clouds”), Mare Nectaris (“Sea of Nectar”), and so on. Most mountain ranges in the highlands bear the names of terrestrial mountain ranges—the Alps, the Carpathians, the Apennines, the Pyrenees, and so on. Most of the craters are named after great scientists or philosophers, such as Plato, Aristotle, Eratosthenes, and Copernicus.

**◀ FIGURE 8.5 Moon from Apollo** The Moon, as seen from the *Apollo 8* orbiter during the first human circumnavigation of our satellite in 1968. Craters ranging in size from 20 km to 500 m (also the width of the long fault lines) can be seen. (NASA)



▲ **FIGURE 8.6 Full Moon, Far Side** The far side of the Moon, as photographed by the *Apollo 16* manned mission. The large, dark region at center bottom outlines the South Pole–Aitken Basin, the largest and deepest impact basin known in the solar system. Only a few small maria exist on the far side. (NASA)

Because the Moon rotates once on its axis in exactly the same time it takes to complete one orbit around Earth, the Moon has a “near” side, which is always visible from Earth, and a “far” side, which never is (see Section 8.4). To the surprise of most astronomers, when the far side of the Moon was mapped, first by Soviet and later by U.S. spacecraft (see *Discovery 8-1*), no major maria were found there. The lunar far side (Figure 8.6) is composed almost entirely of highlands. This fact has great bearing on our theory of how the Moon’s surface terrain came into being, for it implies that the processes involved could *not* have been entirely internal in nature. Earth’s presence must somehow have played a role.

### CONCEPT Check

- ✓ Describe three important ways in which the lunar maria differ from the highlands.


## The Surface of Mercury

Mercury is difficult to observe from Earth because of Mercury’s closeness to the Sun. Even with a fairly large telescope, we see it only as a slightly pinkish disk. Figure 8.7 is one of the few photographs of Mercury taken from Earth that shows any evidence of surface markings.


Astronomers could only speculate about the faint, dark markings in the days before *Mariner 10*’s arrival. We now know that these markings are much like those seen by an observer gazing casually at Earth’s Moon. The largest ground-based telescopes can resolve surface features on Mercury about as well as we can perceive features on the Moon with our unaided eyes.

Figure 8.8 is a global *Messenger* view of the planet as we know it today; Figure 8.9 shows a close-up of the surface. There are no signs of clouds, rivers, dust storms, or other aspects of weather, recent or ancient. Indeed, much of Mercury’s cratered surface bears a strong resemblance to the Moon’s highlands. However, Mercury’s craters are less densely packed than their lunar counterparts, and extensive, gently rolling **intercrater plains** cover some 40 percent of the planet’s surface. These plains are the oldest visible parts of the planet’s surface—they formed just under 4 billion years ago—and appear to have covered many earlier craters. They are distributed roughly uniformly over the planet’s surface.


In addition, Mercury has numerous *smooth plains*, akin to the lunar maria, where lava has filled depressions produced by large meteoritic impacts. They are less obvious than the maria on the Moon simply because their color is comparable to the rest of Mercury’s surface. For the most part, the smooth plains formed hundreds of millions of years after the intercrater plains.



▲ **FIGURE 8.7 Mercury** This photograph of Mercury, taken from Earth with a large ground-based optical telescope, shows only a few faint surface features. (Palomar Observatory/Caltech)



▲ **FIGURE 8.8 Mercury, Up Close** Mercury is imaged here as a mosaic of photographs—a composite image constructed from many individual images—taken by the *Messenger* spacecraft in 2008 as it bypassed the planet. Notice the young, extensively rayed craters, here imaged with a resolution of about 5 km. (NASA)



▲ **FIGURE 8.9 Mercury, Very Close** Another photograph of Mercury by *Messenger*; this one taken at much higher resolution (about 300 m). The dark material around the crater at lower left is typical of many large craters on Mercury. The cause of the dark halos is not known. (NASA)

## 8.4 Rotation Rates


The spins of both the Moon and Mercury are strongly influenced by their proximity to their parent bodies—Earth and the Sun, respectively. By studying the processes responsible for the rotation rates observed today, astronomers learn about the role of tidal forces in shaping the details of the solar system.

### The Rotation of the Moon

As mentioned earlier, the Moon's rotation period is precisely equal to its period of revolution about Earth—27.3 days—so the Moon keeps the same side facing Earth at all times (see Figure 8.10). To an astronaut standing on the Moon's near-side surface, Earth would appear almost stationary in the sky (although our planet's daily rotation would be clearly evident). This condition, in which the spin of one body is precisely equal to (or *synchronized* with) its revolution around another body, is known as a **synchronous orbit**.

The fact that the Moon is in a synchronous orbit around Earth is no accident. It is an inevitable consequence of the gravitational interaction between those two bodies. Just as the Moon raises tides on Earth, Earth also produces a tidal bulge in the Moon. Indeed, because Earth is so much more massive, the tidal force on the Moon is about 20 times greater than that on Earth, and the Moon's tidal bulge is correspondingly larger.

In Chapter 7, we saw how lunar tidal forces are causing Earth's spin to slow and how, as a result, Earth will eventually rotate on its axis at the same rate as the Moon revolves around Earth. ∞ (Sec. 7.6) Earth's rotation will not become synchronous with the Earth–Moon orbital period for



▲ **Interactive FIGURE 8.10 The Moon's Synchronous Rotation** As the Moon orbits Earth, it keeps one face permanently pointed toward our planet. To the astronaut depicted here, Earth is always directly overhead. In fact, the Moon is slightly elongated in shape (highly exaggerated here) owing to Earth's tidal pull on it, with its long axis perpetually pointing toward Earth. It is often useful to think of the Earth and the Moon as a single system.

## MORE PRECISELY 8-1

### Why Air Sticks Around

Why do some planets and moons have atmospheres, while others do not, and what determines the composition of the atmosphere if one exists? Why does a layer of air, made up mostly of nitrogen and oxygen, lie just above Earth's surface? After all, experience shows that most gas naturally expands to fill all the volume available. Perfume in a room, fumes from a poorly running engine, and steam from a teakettle all disperse rapidly until we can hardly sense them. Why doesn't our planet's atmosphere similarly disperse by floating away into space?

The answer is that *gravity* holds it down. Earth's gravitational field exerts a pull on all the atoms and molecules in our atmosphere, preventing them from escaping. However, gravity is not the only influence acting, for if it were, all of Earth's air would have fallen to the surface long ago. *Heat*—the rapid random motion of the molecules in a gas—competes with gravity to keep the atmosphere buoyant. Let's explore this competition between gravity and heat in a little more detail.

All gas molecules are in constant random motion. The temperature of any gas is a direct measure of this motion: The hotter the gas, the faster the molecules are moving. [∞ \(More Precisely 3-1\)](#) The Sun continuously supplies heat to our planet's atmosphere, and the resulting rapid movement of heated molecules produces *pressure*, which tends to oppose the force of gravity, preventing our atmosphere from collapsing under its own weight.

An important measure of the strength of a body's gravity is the body's *escape speed*—the speed needed for any object to escape forever from its surface. [∞ \(Sec. 2.8\)](#) This speed increases with increased mass or decreased radius of the parent body (often a moon or a planet). In convenient (Earth) units, it can be expressed as

$$\begin{aligned} \text{escape speed (in km/s)} \\ = 11.2 \sqrt{\frac{\text{mass of body (in Earth masses)}}{\text{radius of body (in Earth radii)}}}. \end{aligned}$$

Thus, Earth's escape speed is  $11.2\sqrt{1/1} = 11.2$  km/s. If the *mass* of the parent body is quadrupled, the escape speed doubles. If the

parent body's *radius* quadruples, then the escape speed is halved. In other words, you need high speed to escape the gravitational attraction of a very massive or very small body, but you can escape from a less massive or larger body at lower speeds.

To determine whether a planet will retain an atmosphere, we must compare the planet's escape speed with the *molecular speed*, which is the average speed of the gas particles making up the planet's atmosphere. This speed actually depends not only on the temperature of the gas, but also on the mass of the individual molecules—the hotter the gas or the smaller the molecular mass, the higher is the average speed of the molecules:

$$\begin{aligned} \text{average molecular speed (in km/s)} \\ = 0.157 \sqrt{\frac{\text{gas temperature (K)}}{\text{molecular mass (hydrogen atom masses)}}}. \end{aligned}$$

Thus, increasing the absolute temperature of a sample of gas by a factor of four—for example, from 100 K to 400 K—doubles the average speed of its constituent molecules, and, at a given temperature, molecules of hydrogen ( $\text{H}_2$ : molecular mass = 2) in air move, on average, four times faster than molecules of oxygen ( $\text{O}_2$ : molecular mass = 32), which are 16 times heavier.

**EXAMPLE 1** For nitrogen ( $\text{N}_2$ : molecular mass = 28) and oxygen ( $\text{O}_2$ : molecular mass = 32) in Earth's atmosphere, where the temperature near the surface is nearly 300 K, the preceding formula yields the following average molecular speeds:

$$\begin{aligned} \text{nitrogen: } & 0.157 \text{ km/s} \times \sqrt{\frac{300}{28}} = 0.51 \text{ km/s;} \\ \text{oxygen: } & 0.157 \text{ km/s} \times \sqrt{\frac{300}{32}} = 0.48 \text{ km/s.} \end{aligned}$$

These speeds are far smaller than the 11.2 km/s needed for a molecule to escape into space. As a result, Earth is able to retain its nitrogen–oxygen atmosphere. On the whole, *our planet's gravity simply has more influence than the heat of our atmosphere.*

hundreds of billions of years. In the case of the Moon, however, the process has already gone to completion. The Moon's much larger tidal deformation caused it to evolve into a synchronous orbit long ago, and the Moon is said to have become *tidally locked* to Earth. Most of the moons in the solar system are similarly locked by the tidal fields of their parent planets.

Actually, the size of the lunar bulge is too great to be produced by Earth's present-day tidal influence. The explanation seems to be that, long ago, the distance from Earth to the Moon may have been as little as two-thirds of its current value, or about 250,000 km. Earth's tidal force on the Moon would then have been more than three times greater than it is today

and could have accounted for the Moon's elongated shape. The resulting distortion could have “set” when the Moon solidified, thus surviving to the present day, and at the same time accelerating the synchronization of the Moon's orbit.

### Measurement of Mercury's Spin


In principle, the ability to discern surface features on Mercury should allow us to measure its rotation rate simply by watching the motion of a particular region around the planet. In the mid-19th century, an Italian astronomer named Giovanni Schiaparelli did just that. He concluded

In reality, the situation is a little more complicated than a simple comparison of speeds. Atmospheric molecules can gain or lose speed by bumping into one another or by colliding with objects near the ground. Thus, although we can characterize a gas by its average molecular speed, the molecules do not *all* move at the same speed, as illustrated in the accompanying figure. A tiny fraction of the molecules in any gas have speeds much greater than average—one molecule in two million has a speed more than three times the average, and one in  $10^{16}$  exceeds the average by more than a factor of five. This means that at any instant, *some* molecules are moving fast enough to escape, even when the average molecular speed is much less than the escape speed. The result is that all planetary atmospheres slowly leak away into space.

Don't be alarmed—the leakage is usually very gradual! As a rule of thumb, if the escape speed from a planet exceeds the average speed of a given type of molecule by a factor of


six or more, then molecules of that type will not have escaped from the planet's atmosphere in significant quantities in the 4.6 billion years since the solar system formed. Conversely, if the escape speed is less than six times the average speed of molecules of a given type, then most of them will have escaped by now, and we should not expect to find them in the atmosphere.

For air on Earth, the mean molecular speeds of oxygen and nitrogen that we just computed are comfortably below one-sixth of the escape speed. However, if the Moon originally had an Earth-like atmosphere, that lunar atmosphere would have been heated by the Sun to much the same temperature as Earth's air today, so the average molecular speed would have been about 0.5 km/s. Because the Moon's escape speed is only  $11.2\sqrt{0.012/0.27} = 2.4$  km/s—less than six times the average molecular speed—any original lunar atmosphere long ago dispersed into interplanetary space. Mercury's escape speed is  $11.2\sqrt{0.055/0.38} = 4.2$  km/s. However, its peak surface temperature is around 700 K, corresponding to an average molecular speed for nitrogen or oxygen of about 0.8 km/s, more than one-sixth of the escape speed, so there has been ample time for those gases to escape.



**EXAMPLE 2** We can use the foregoing arguments to understand some aspects of atmospheric *composition*. Hydrogen molecules ( $H_2$ : molecular mass = 2) move, on average, at about 1.9 km/s in Earth's atmosphere at sea level, so they have had time to escape since our planet formed ( $6 \times 1.9$  km/s = 11.4 km/s, which is greater than Earth's 11.2-km/s escape speed). Consequently, we find very little hydrogen in Earth's atmosphere today. However, on the planet Jupiter, with a lower temperature (about 100 K), the speed of hydrogen molecules is correspondingly slower—about 1.1 km/s. At the same time, Jupiter's escape speed is 60 km/s, over five times higher than Earth's. For those reasons, Jupiter has retained its hydrogen—in fact, hydrogen is the dominant ingredient of Jupiter's atmosphere.

that Mercury always keeps one side facing the Sun, much as our Moon perpetually presents only one face to Earth. The explanation suggested for this supposed synchronous rotation was the same as that for the Moon: The tidal bulge raised in Mercury by the Sun had modified the planet's rotation rate until the bulge always pointed directly at the Sun. Although the surface features could not be seen clearly, the combination of Schiaparelli's observations and a plausible physical explanation was enough to convince most astronomers, and the belief that Mercury rotates synchronously with its revolution about the Sun (i.e., once every 88 Earth days) persisted for almost half a century.

In 1965, astronomers making observations of Mercury from the Arecibo radio telescope in Puerto Rico (see Figure 5.21) discovered that this long-held view was in error. They used the Arecibo instrument as a giant radar gun, sending out pulses of radio waves toward the planet and waiting for the echoes to return. (See Figure 2.18 for a similar measurement of the planet Venus.)  (Sec. 2.6) The returning pulses were much weaker than the original outgoing beam, but the huge size of the Arecibo dish allowed the researchers to detect the reflected signal and then analyze it to determine Mercury's rotation rate.

## DISCOVERY 8-1

### Lunar Exploration

The Space Age began in earnest on October 4, 1957, with the launch of the Soviet satellite *Sputnik 1*. Thirteen months later, on January 4, 1959, the Soviet *Luna 1*, the first human-made craft to escape Earth's gravity, passed the Moon. *Luna 2* crash-landed on the surface in September of that year, and *Luna 3* returned the first pictures of the far side a month later. The long-running *Luna* series established a clear Soviet lead in the early “space race” and returned volumes of detailed information about the Moon's surface. Several of the *Luna* missions landed and returned surface material to Earth.

The U.S. lunar exploration program got off to a rocky start. The first six attempts in the *Ranger* series, between 1961 and 1964, failed to accomplish their objective of just hitting the Moon. The last three were successful, however. *Ranger 7* collided with the lunar surface (as intended) on June 28, 1964. Five U.S. *Lunar Orbiter* spacecraft, launched in 1966 and 1967, were successfully placed in orbit around the Moon, and they relayed high-resolution images of much of the lunar surface back to Earth. Between 1966 and 1968, seven *Surveyor* missions soft-landed on the Moon and performed detailed analyses of the surface.

Many of these unmanned U.S. missions were performed in support of the manned *Apollo* program. On May 25, 1961, at a time when the U.S. space program was in great disarray, President John F. Kennedy declared that the United States would “send a man to the Moon and return him safely to Earth” before the end of the decade, and the *Apollo* program was born. On July 20, 1969, less than 12 years after *Sputnik* and only 8 years after the statement of the program's goal, *Apollo 11* commander Neil Armstrong became the first human to set foot on the Moon, in Mare Tranquilitatis (the Sea of Tranquility). Three-and-a-half years later, on December 14, 1972, scientist-astro-naut Harrison Schmitt, of *Apollo 17*, was the last.

The astronauts who traveled in pairs to the lunar surface in each lunar lander (shown in the first photograph) performed numerous geological and other scientific studies on the surface. The later landers brought with them a “lunar rover”—a small golf cart-sized vehicle that greatly expanded the area the astronauts could cover. Probably the most important single aspect of the *Apollo* program was the collection of samples of surface



(NASA)

rock from various locations on the Moon. In all, some 382 kg of material was returned to Earth. Chemical analysis and radioactive dating of these samples revolutionized our understanding of the Moon's surface history. No amount of Earth-based observations could have achieved the same results.

Each *Apollo* lander left behind a nuclear-powered package of scientific instruments called the *Apollo Lunar Surface Experiments Package (ALSEP)* to monitor the solar wind, measure heat flow in the Moon's interior, and, perhaps most important, record lunar seismic activity. With several *ALSEPs* on the surface, scientists could determine the location of “moonquakes” by triangulation and map the Moon's inner structure, obtaining information critical to our understanding of the Moon's evolution.

By any standards, the *Apollo* program was a spectacular success. It represents a towering achievement of the human race. The project's goals were met on schedule and within budget, and

To illustrate the basic method, Figure 8.11 shows a radar pulse reflecting from the surface of a hypothetical planet. The reflected signal as a whole may be redshifted or blueshifted by the Doppler effect, depending on the overall radial velocity of the planet relative to Earth.  $\infty$  (Sec. 3.5) But in addition, if the planet is rotating, the radiation reflected from the side moving toward us returns at a slightly higher frequency than the radiation reflected from the receding side. (Think of the two hemispheres as being separate sources of radiation moving at slightly different velocities, one toward us and one away.) The effect is very similar to the rotational line broadening discussed in Chapter 4, except that in this case the radiation


is not emitted by the planet, but only reflected from its surface.  $\infty$  (Sec. 4.5) Thus, even if the original beam consists of radiation of a single frequency, the reflected signal contains a spread of frequencies on either side of the original. By measuring that spread we can determine the planet's rotational speed.

In this way, the Arecibo researchers found that the rotation period of Mercury is not 88 days, as had previously been thought, but 59 days, exactly two-thirds of the planet's orbital period. Because there are exactly three rotations for every two revolutions, we say that there is a 3:2 *spin-orbit resonance* in Mercury's motion. In this context, the term **resonance** just means that two characteristic times—here, Mercury's day

our knowledge of the Moon, Earth, and the solar system increased enormously. But the “Age of *Apollo*” was short lived. Public interest quickly waned. Over half a billion people breathlessly watched on television as Neil Armstrong set foot on the Moon, yet barely 3 years later, when the program was abruptly canceled for largely political (rather than scientific, technological, or economic) reasons, the landings had become so routine that they no longer excited the interest of the American public. Unmanned space science moved away from the Moon and toward the other planets, and the manned space program foundered. Perhaps one of the most amazing—and saddest—facts about the *Apollo* program is that today, four decades on, no nation on Earth currently has the technical capacity to replicate the feat.

In 1994, the suitcase-sized U.S. military satellite *Clementine* was placed into lunar orbit mainly to test new imaging devices, a by-product of which was a highly detailed survey of the lunar surface. In 1998, NASA returned to the Moon for the first time in a quarter century with the launch of *Lunar Prospector*, another small satellite on a 1-year mission to study the Moon’s structure and origins. Both missions were successful, demonstrating the wealth of information that can be obtained by low-budget spacecraft. In 2009, NASA’s *Lunar Reconnaissance Orbiter* entered polar orbit just 50 km above the Moon’s surface. Its mission was to map the lunar surface with even greater precision, particularly in the polar regions where *Clementine* and *Prospector* had previously reported possible evidence of water ice in permanently shadowed craters (see Section 8.5).

In recent years, the Moon has been visited by spacecraft from several countries, including Japan, India, and China, in addition to missions from NASA and the European Space Agency. The Chinese *Chang’e* program is an ambitious series of orbiters and landers intended to lead to a sample return mission (*Chang’e 5*) in 2017 and a possible manned landing sometime after 2020. India’s *Chandrayan-1* orbiter and impactor reached the Moon in 2008, while *Chang’e 2* went into orbit around the Moon in 2010. NASA’s most recent lunar mission was the *Gravity Recovery and Interior Laboratory (GRAIL)*, comprising twin spacecraft orbiting 200 km apart some 50 km above the lunar surface. By monitoring the deviations between the two orbits, project scientists measured the Moon’s



(NASA)

gravitational field and internal structure with unprecedented accuracy. The second figure shows a high-resolution *GRAIL* map of the Moon’s gravity (with red denoting mass excesses and blue mass deficiencies). In addition to showing many prominent surface features on the far side, such maps also contain vital information about the lunar crust and upper mantle. The *GRAIL* mission ended in December 2012, but researchers are still analyzing the data to try to probe the Moon’s deep interior.

It seems that, after years of decline, a broad-based era of lunar exploration is now under way. Ambitious plans exist to establish permanent human colonies on the Moon, both for commercial ventures, such as mining, and for scientific research, such as an astronomical observatory. A confirmed discovery of water on the lunar surface would alleviate at least one major logistical problem associated with such an undertaking. However, it remains to be seen whether the political will and economic resources needed to make this dream a reality actually exist anywhere on Earth.


and year—are related to each other in a simple way. An even simpler example of a spin–orbit resonance is the Moon’s orbit around Earth. In that case, the rotation is synchronous with the revolution, and the resonance is said to be 1:1.

Figure 8.12 illustrates some implications of Mercury’s curious rotation for a hypothetical inhabitant of the planet. Mercury’s solar day—the time from noon to noon, say—is 2 Mercury years long! The Sun stays “up” in the black Mercury sky for almost 3 Earth months at a time, after which follow nearly 3 Earth months of darkness. At any given point in its orbit, Mercury presents the same face to the Sun, not every time it revolves, but *every other* time.

## Explanation of Mercury’s Rotation

Mercury’s 3:2 spin–orbit resonance did not occur by chance. What mechanism establishes and maintains it? In the case of the Moon orbiting Earth, the 1:1 resonance is the result of tidal forces. In essence, the lunar rotation period, which probably started off much shorter than its present value, has lengthened so that the tidal bulge created by Earth is fixed relative to the body of the Moon. Tidal forces (this time due to the Sun) are also responsible for Mercury’s 3:2 resonance, but in a much more subtle way.





▲ **FIGURE 8.11 Planetary Radar** A radar beam (blue waves) reflected from a rotating planet yields information about both the planet's line-of-sight motion and its rotation rate.

Mercury cannot settle into a 1:1 resonance because its orbit around the Sun is quite eccentric. By Kepler's second law, Mercury's orbital speed is greatest at perihelion (closest approach to the Sun) and least at aphelion (greatest distance from the Sun). ∞ (*More Precisely 2-1*) A moment's thought shows that, because of these variations in the planet's orbital speed, there is no way that the planet (rotating at a constant rate) can remain in a synchronous orbit. If its rotation were synchronous near perihelion, it would be too rapid at aphelion, and synchronism at aphelion would produce too slow a rotation at perihelion.


Tidal forces always act so as to synchronize the rotation rate with the instantaneous orbital speed, but such

synchronization cannot be maintained over Mercury's entire orbit. What happens? The answer is found when we realize that tidal effects diminish very rapidly with increasing distance. The tidal forces acting on Mercury at perihelion are much greater than those at aphelion, so perihelion "won" the struggle to determine the rotation rate. In the 3:2 resonance, Mercury's orbital and rotational motion are almost exactly synchronous *at perihelion*, so that particular rotation rate was naturally "picked out" by the Sun's tidal influence on the planet. Notice that even though Mercury rotates through only  $180^\circ$  between one perihelion and the next (see Figure 8.12), the appearance of the tidal bulge is the *same* each time around.

Resonances such as these occur quite frequently in the solar system. Many additional examples can be found in the motion of the planets, their moons and rings, as well as in orbits of many asteroids and Kuiper belt objects. The rotation of Mercury is one of the simplest nonsynchronous resonances known. Many resonances are much more complex. These intricate interactions are responsible for much of the fine detail observed in the motion of our planetary system.

The Sun's tidal influence also causes Mercury's rotation axis to be exactly perpendicular to its orbital plane. As a result, and because of Mercury's eccentric orbit and the spin-orbit resonance, some points on the surface get much hotter than others. In particular, the two (diametrically opposite) points on the equator where the Sun is directly overhead at perihelion get hottest of all. They are called the *hot longitudes*. The peak temperature of 700 K mentioned earlier occurs at noon at those two locations. At the *warm longitudes*, where the Sun is directly overhead at aphelion, the peak temperature is about 150 K cooler—a mere 550 K.

By contrast, the Sun is always on the horizon as seen from the planet's poles, so temperatures there never reach the sizzling levels of the equatorial regions. Earth-based



▲ **Interactive FIGURE 8.12 Mercury's Rotation** Mercury's orbital and rotational motions combine to produce a solar day that is 2 Mercury years long.

radar studies carried out during the 1990s suggest that Mercury's polar temperatures may be as low as 125 K and that, despite the planet's scorched equator, the poles may be covered with extensive sheets of water ice. (See Section 8.5 for similar findings regarding the Moon.)

### CONCEPT Check

✓ How has gravity influenced the rotation rates of the Moon and Mercury?





## 8.5 Lunar Cratering and Surface Composition

On Earth, the combined actions of wind and water erode our planet's surface and reshape its appearance almost daily. Coupled with the never-ending motion of Earth's surface plates, the result is that most of the ancient history of our planet's surface is lost to us. The Moon, in contrast, has no air, no water, no plate tectonics, and no ongoing volcanic or seismic activity. Consequently, features dating back almost to its formation are still visible today.

### Meteoritic Impacts

The primary agent of change on the lunar surface is interplanetary debris, in the form of *meteoroids*. This material, much of it rocky or metallic in composition, is strewn throughout the solar system, orbiting the Sun in interplanetary space, perhaps for billions of years, until it happens to collide with some planet or moon. ∞ (Sec. 6.5) On Earth, most meteoroids burn up in the atmosphere, producing the streaks of light known as *meteors*, or “shooting stars.” But the Moon, without an atmosphere, has no protection against this onslaught. Large and small meteoroids zoom in and collide with the surface, sometimes producing huge craters. Over billions of years, these collisions have scarred, cratered, and sculpted the lunar landscape. Craters are still being formed today—even as you read this—all across the surface of the Moon.

Meteoroids generally strike the Moon at speeds of several kilometers per second. At these speeds, even a small piece of matter carries an enormous amount of energy. For example, a 1-kg object hitting the Moon's surface at 10 km/s releases as much energy as the detonation of 10 kg of TNT! As illustrated in Figure 8.13, the impact of a meteoroid with the surface causes sudden and tremendous




**MA** **Interactive FIGURE 8.13 Meteoroid Impact** Several stages in the formation of a crater by meteoritic impact. (a) A meteoroid strikes the surface, releasing a large amount of energy. (b, c) The resulting explosion ejects material from the impact site and sends shock waves through the underlying surface. (d) Eventually, a crater results, surrounded by a blanket of ejected material.

pressures to build up, heating the normally brittle rock and deforming the ground like heated plastic. The ensuing explosion pushes previously flat layers of rock up and out, forming a crater.


The diameter of the eventual crater is typically 10 times that of the incoming meteoroid; the depth of the crater is about twice the meteoroid's diameter. Thus, our 1-kg meteoroid, measuring perhaps 10 cm across, would produce a crater about 1 m in diameter and 20 cm deep. Shock waves from the impact pulverize the lunar surface to a depth many times that of the crater itself. Numerous rock samples brought back by the *Apollo* astronauts show patterns of repeated shattering and melting—direct evidence of the violent shock waves and high temperatures produced in meteoritic impacts. The material thrown out by the explosion surrounds the crater in a layer called an *ejecta blanket*. The ejected debris ranges in size from fine dust to large boulders. Figure 8.14(a) shows the result of one particularly large meteoritic impact on the Moon. As shown in Figure 8.14(b), the larger pieces of ejecta may themselves form secondary craters.

In addition to the bombardment by meteoroids with masses of a gram or more, a steady “rain” of *micrometeoroids* (debris with masses ranging from a few micrograms up to about 1 gram) also eats away at the structure of the lunar surface. Some examples can be seen in Figure 8.15, a photomicrograph (a photograph taken through a microscope) of some glassy “beads” brought back to Earth by *Apollo* astronauts. The beads themselves were formed during the explosion following the impact of a meteoroid, when surface rock was melted, ejected, and rapidly cooled. Note how several of the beads also display fresh miniature craters caused by micrometeoroids that struck the beads after they had cooled and solidified.

In fact, the *rate* of cratering decreases rapidly with the size of the crater—fresh large craters are scarce, but small craters are common. The reason for this is simple: There just aren't very many large chunks of debris in interplanetary space, so their collisions with the Moon are rare. At present average rates, one new 10-km (diameter) lunar crater is formed roughly every 10 million years, a



**▲ FIGURE 8.14 Large Lunar Craters** (a) The meteorite that produced this large lunar crater, called Orientale Basin, thrust up much surrounding matter, which can be seen as concentric rings of cliffs called the Cordillera Mountains. The outermost ring is nearly 1000 km in diameter. Notice the smaller, sharper, younger craters that have impacted this ancient basin in more recent times. (b) Two smaller craters called Reinhold and Eddington sit amid the secondary cratering resulting from the impact that created the 90-km-wide Copernicus crater (near the horizon) about a billion years ago. The ejecta blanket from crater Reinhold, 40 km across and in the foreground, can be seen clearly. This view was obtained by looking northeast from the lunar module during the *Apollo 12* mission. (NASA)



▲ **FIGURE 8.15 Miniature Craters** Craters of all sizes litter the lunar landscape. Some shown here, embedded in glassy beads retrieved by *Apollo* astronauts, measure only 0.01 mm across. (The scale at the top is in millimeters.) The beads themselves were formed during the explosion following a meteoroid impact, when surface rock was melted, ejected, and rapidly cooled. (NASA)

new meter-sized crater is created about once a month, and centimeter-sized craters are formed every few minutes.

## Cratering History of the Moon

Astronomers can use the known ages (from radioactive dating) of Moon rocks to estimate the rate of cratering in the past. One very important result of this work is the discovery that the Moon was subjected to an extended period of intense meteoritic bombardment roughly 4 billion years ago. Indeed, this is a key piece of evidence supporting the condensation theory of solar system formation. ∞ (Sec. 6.6)

As we have seen, the heavily cratered highlands are older than the less-cratered maria, but the difference in cratering is not simply a matter of exposure time. Astronomers now think that the Moon, and presumably the entire inner solar system, experienced a sudden drop in meteoritic bombardment about 3.9 billion years ago. The highlands solidified and received most of their craters before that time, whereas the maria solidified afterward. The rate of cratering has remained relatively low ever since.

The great basins that comprise the maria are thought to have been created during the final stages of the heavy bombardment, between about 4.1 and 3.9 billion years ago. Subsequent volcanic activity filled the craters with lava, ultimately creating the formations we see today as the lava turned into solid rock. In a sense, then, the maria *are* oceans—ancient seas of molten lava, now solidified.

Not all these great craters became flooded with lava, however. One of the youngest craters is the Orientale Basin (Figure 8.14a), which formed about 3.9 billion years ago. This crater did not undergo much subsequent volcanism, and we can recognize its structure as an impact crater rather than as another mare. Similar “unflooded” basins are seen on the lunar far side (Figure 8.6).


Apart from meteorites found on Earth, the Moon is the only solar system object for which we have accurate age measurements, from radioactive dating of samples returned to Earth. However, studies of lunar cratering provide astronomers with an important alternative means of estimating ages in the solar system. By counting craters on a planet, moon, or asteroid and using the Moon to calibrate the numbers, an approximate age for the surface can be obtained. In fact, this is how most of the ages presented in the next few chapters are determined. Note that, as with radioactive dating, the technique measures only the time since the surface in question last solidified—all cratering is erased and the clock is reset if the rock melts. ∞ (More Precisely 7-2)

## Lunar Dust


Meteoroid collisions with the Moon are the main cause of the layer of pulverized ejecta—also called lunar dust, or *regolith* (meaning “fine rocky layer”)—that covers the lunar landscape to an average depth of about 20 m. This microscopic dust has a typical particle size of about 0.01 mm. In consistency, it is rather like talcum powder or ready-mix dry mortar. Figure 8.16 shows an *Apollo* astronaut’s boot prints in the regolith, which is thinnest on the maria (10 m) and thickest on the highlands (over 100 m deep in places).

The constant barrage from space results in a slow, but steady, erosion of the lunar surface. The soft edges of the craters visible in the foreground of Figure 8.17 are the result of this process. In the absence of erosion, those features would still be as jagged and angular today as they were just after they formed. Instead, the steady buildup of dust due to innumerable impacts has smoothed their outlines and will probably erase them completely in about 100 million years.

From the known dependence of the cratering rate on the size of a crater, planetary scientists can calculate how many small craters they would expect to find, given the numbers of large craters actually observed. When they make this calculation, they find a shortage of craters less than about 20 m deep. These “missing” craters have been filled in by erosion over the lifetime of the Moon. This gives



▲ **FIGURE 8.16 Regolith** The lunar soil, or regolith, is a layer of powdery dust covering the lunar surface to a depth of roughly 20 m. Here, an *Apollo* astronaut is adjusting some instruments for testing the composition of soil near Mount Hadley. His boot prints show how the astronaut's weight has compacted the regolith to a depth of a few centimeters. (NASA)



▲ **FIGURE 8.17 Lunar Surface** Despite the complete lack of wind and water on the airless Moon, its surface is slowly eroded by the constant “rain” of meteoroids, especially micrometeoroids. Note the soft edges of the background hills of this image and the *Apollo* astronaut's boot prints surrounding his landing craft (left) and tire tracks from the “lunar buggy” (right). (NASA)

us a very rough estimate of the average erosion rate: about 5 m per billion years, or roughly 1/10,000 the rate on Earth.

The current lunar erosion rate is very low because meteoritic bombardment on the Moon is a much less effective erosive agent than are wind and water on Earth. For comparison, the Barringer Meteor Crater (Figure 8.18) in the Arizona desert, one of the largest meteoroid craters on Earth, is only 25,000 years old, but has already undergone noticeable erosion. It will probably disappear completely in just a few million years, quite a short time geologically. If a crater that size had formed on the Moon even 4 billion years ago, it would still be plainly visible today. Even the shallow boot prints shown in Figure 8.16 are likely to remain intact for several million years.

## Lunar Ice?

In contrast to Earth's soil, the lunar regolith contains no organic matter like that produced by biological organisms. No life whatsoever exists on the Moon. Nor were any fossils found in *Apollo* samples. Lunar rocks are barren of life and apparently always have been. NASA was so confident of

this fact that the astronauts were not even quarantined on their return from the last few *Apollo* landings. Furthermore, all the lunar samples returned by the U.S. and Soviet Moon programs were bone dry—they didn't even contain minerals having water molecules locked within their crystal structure. Terrestrial rocks, by contrast, are almost always 1 or 2 percent water. The main reasons for this lack of water are the Moon's lack of an atmosphere and the high (up to 400 K) daytime temperatures found over most of the lunar surface.

Some regions of the Moon *may* contain water, however—in the form of ice. As early as the 1960s, some scientists had considered the theoretical possibility that ice might be found near the lunar poles. Since the Sun never rises more than a few degrees above the horizon, as seen from the Moon's polar regions, temperatures on the permanently shaded floors of craters near the poles never exceed about 100 K. Consequently, those scientists theorized, any water ice there could have remained permanently frozen since the very early days of the solar system, never melting or vaporizing and hence never escaping into space.

In 1996, controllers of the *Clementine* mission (see *Discovery 8-1*) reported that radar echoes from an old, deep crater



▲ **FIGURE 8.18 Barringer Crater** The Barringer Meteor Crater, near Winslow, Arizona, is 1.2 km in diameter and 0.2 km deep. (Note the access road at right for scale.) Geologists think that a large meteoroid whacked Earth and formed this crater about 25,000 years ago. The meteoroid was probably about 50 m across and likely weighed around 200,000 tons. The inset shows a close-up of one of the interior walls of the crater. (*U.S. Geological Survey*)

near the lunar south pole implied deposits of low-density material, probably water ice, at a depth of a few meters. In 1998, NASA's *Lunar Prospector* reported large amounts of ice—possibly totaling trillions of tons—at both lunar poles. At first, it appeared that the ice was mainly in the form of tiny crystals mixed with the lunar regolith, spread over many tens of thousands of square kilometers of deeply shadowed crater floors. Later analysis suggested that much of the ice might instead be in the form of smaller, but more concentrated “lakes” of material lying just below the surface.

Given the potential importance of this finding, NASA wanted to gain more information about possible lunar ice. In 1999, as *Lunar Prospector* neared the end of its operational lifetime, mission controllers crashed it into Shoemaker crater, one of the deep craters near the lunar south pole in which ice was suspected to hide. The hope was that telescopes on Earth might detect spectroscopic signatures of water vapor released by the impact. However, no water vapor was seen.

In 2009, NASA tried again, on a larger scale, with the *Lunar CRater Observation and Sensing Satellite (LCROSS)* mission, launched along with the *Lunar Reconnaissance Orbiter* mission (*LRO*; see *Discovery 8-1*). The Centaur rocket that boosted both missions into lunar orbit was crashed into Cabeus, another deeply shadowed crater near the lunar south pole, while *LCROSS* watched from just a few thousand kilometers away, radioing its spectroscopic data back to Earth via *LRO* before it too impacted the Moon minutes later. A few weeks later, NASA scientists announced that detailed analysis of the *LCROSS* data had confirmed the presence of water molecules in the ejecta.


The amount of water was not great—only about 1 part in 100,000, less than in the desert sand on Earth—but enough to support the earlier reports.

At the leading edge of scientific discovery, however, nothing is clear-cut. ∞ (Sec. 1.2) Detailed *LRO* observations, released in 2012, of Shackleton, yet another south polar crater, were inconclusive on the subject of lunar ice, suggesting that subsurface ice was one possible explanation, but not the only, or even the most likely, way to account for the observations.

If lunar ice exists, where did it come from? Most likely, it was brought to the lunar surface by meteoroids and comets, just as we saw earlier with water on Earth. ∞ (Sec. 6.7) Any ice that survived the impact would have been scattered across the surface. Over most of the Moon, that ice would have rapidly vaporized and escaped, but in the deep basins near the poles, it survived and built up over time. Whatever its origin, polar ice may be a crucial component of any serious attempt at human colonization of the Moon: The anticipated cost of transporting a kilogram of water from Earth to the Moon is between \$2,000 and \$20,000.

## Lunar Volcanism

Only a few decades ago, debate raged in scientific circles about the origin of lunar craters, with most scientists of the opinion that the craters were the result of volcanic activity. We now know that almost all lunar craters are actually meteoritic in origin. However, a few apparently are not. Figure 8.19 shows an intriguing alignment of several craters in a *crater-chain* pattern so straight that it is highly unlikely




▲ **FIGURE 8.19 Crater Chain** This “chain” of well-ordered craters was photographed by an *Apollo 14* astronaut. The largest crater, called Davy, is located on the western edge of Mare Nubium. (NASA)

to have been produced by the random collision of meteoroids with the surface. Instead, the chain probably marks the location of a subsurface fault—where cracking or shearing of the surface allowed molten matter to well up from below. As the lava cooled, it formed a solid “dome” above each fissure. Subsequently, the underlying lava receded and the centers of the domes collapsed, forming the craters we see today. Similar features have been observed on Venus by the *Magellan* probe (see Chapter 9). ∞ (Discovery 6-2)

Many other examples of lunar volcanism are known, both in telescopic observations from Earth and in the close-up photographs taken during the *Apollo* missions. Figure 8.20 shows a volcanic **rille**, a ditch where molten lava once flowed. There is good evidence for surface volcanism early in the Moon’s history, and volcanism explains the presence of the lava that formed the maria. However, whatever volcanic activity once existed on the Moon ended long ago. The measured ages for rock samples returned from the Moon are all greater than 3 billion years. (Recall from *More Precisely 7-2* that the radioactivity clock starts “ticking” when the rock solidifies.) Apparently, the maria solidified over 3 billion years ago, and the Moon has been dormant ever since.

### CONCEPT Check

- ✓ How has meteoritic bombardment affected the surface of the Moon?




▲ **FIGURE 8.20 Lunar Volcanism** A volcanic rille, photographed from the *Apollo 15* spacecraft orbiting the Moon, can be seen clearly here (bottom and center) winding its way through one of the maria. Called Hadley Rille, this system of valleys runs along the base of the Apennine Mountains (lower right) at the edge of the Mare Imbrium (to the left). The shadow-sided, most prominent peak at lower right, Mount Hadley, rises almost 5 km high. (NASA)

## 8.6 The Surface of Mercury



Like craters on the Moon, almost all craters on Mercury are the result of meteoritic bombardment. However, Mercury’s craters are not as deep as those on the Moon, their walls are generally not as high, and the ejected material landed closer to the impact sites—exactly as would be expected given Mercury’s greater surface gravity (which is a little more than twice that of the Moon).

Following *Mariner 10*’s visit, the leading explanation for Mercury’s relative lack of craters was that the older craters were filled in by volcanic activity, in much the same way as the Moon’s maria filled in older craters as they formed. More detailed observations by *Messenger* support that conclusion, and many geologists think that much of Mercury’s crust may have formed through repeated volcanic eruptions. However, this *flood volcanism* was apparently not associated with impact basins, suggesting that Mercury’s volcanic past was significantly different from the Moon’s.


Mercury has at least two types of surface feature not found on the Moon. Figure 8.21(a) shows a **scarp**, or cliff, on the surface that does not appear to be the result of volcanic or any other familiar geological activity. It cuts across several craters, indicating that whatever produced it occurred *after*



(a)



(b)



**▲ FIGURE 8.21 Mercury's Surface** (a) Scarps, or ridges, on Mercury's surface were photographed by *Messenger*. This cliff seems to have formed when the planet's crust cooled and shrank early in its history, causing a crease in the surface. (b) *Messenger* also discovered peculiar "hollows"—land features not seen anywhere else in the solar system. They appear here as lighter-colored depressions on a crater's rim and floor. (NASA)


most of the meteoritic bombardment ended about 4 billion years ago. Mercury shows no evidence of crustal motions like plate tectonics on Earth—no fault lines, spreading sites, or indications of plate collisions are seen. ∞ (Sec. 7.4) The scarps, of which several are known from *Mariner* and *Messenger* images, probably formed when the planet's interior cooled and shrank long ago, much as wrinkles form on the skin of an old, shrunken apple.

The second unique aspect of Mercury's landscape, discovered by *Messenger*, is a set of features collectively known as *hollows*—small, shallow, irregularly shaped, rimless depressions that are often found in clusters near the centers of impact craters (Figure 8.21b). Surprisingly, many have a bright, fresh appearance, suggesting that they may be actively forming today. They are not impact craters, but they do appear to be a secondary result of crater formation. Scientists theorize that meteoritic impacts can excavate material that becomes


**► FIGURE 8.22 Caloris Basin** Mercury's most prominent geological feature—the Caloris Basin—spans 1400 km and is ringed by concentric mountain ranges that reach more than 3 km high in places. This huge circular basin, shown here in orange in this false-colored visible image from *Messenger*, is similar in size to the Moon's Mare Imbrium and spans more than half of Mercury's radius. (NASA)

unstable when exposed to the harsh environment on Mercury's surface. The intense heat and solar wind may vaporize some light minerals, weakening the remaining rock and causing it to subside. If so, then these features are a direct result of Mercury's proximity to the Sun.


Figure 8.22 shows what may have been the last great geological event in the history of Mercury: an immense bull's-eye crater called the Caloris Basin, formed eons ago by the



1000 km







▲ **FIGURE 8.23 Weird Terrain** The refocusing of seismic waves after the Caloris Basin impact may have created the weird terrain on the opposite side of the planet.

impact of a large asteroid. (The basin is so called because it lies in Mercury’s “hot longitudes”—see Section 8.3—close to the planet’s equator; *calor* is the Latin word for “heat.”) Compare this basin with the Orientale Basin on the Moon (Figure 8.14a). The impact crater structures are quite similar, but even here there is a mystery: The patterns visible on the Caloris floor are unlike any seen on the Moon. Their origin, like the composition of the floor itself, is unknown.

So large was the impact that created the Caloris Basin that it apparently sent strong seismic waves reverberating throughout the entire planet. On the opposite side of Mercury from

Caloris, there is a region of oddly rippled and wavy surface features, often referred to as *weird* (or *jumbled*) terrain. Scientists theorize that this terrain was produced when seismic waves from the Caloris impact traveled around the planet and converged on the diametrically opposite point, causing large-scale disruption of the surface there, as illustrated in Figure 8.23.

### CONCEPT Check

- ✓ How do scarps on Mercury differ from geological faults on Earth?

## 8.7 Interiors


In Chapter 7 we saw how geologists combine bulk measurements of Earth’s density, gravity, and magnetic field with seismic studies and mathematical models to build up a detailed model of the planet’s interior. ∞ (Secs. 7.1, 7.3, 7.5) Planetary scientists attempt to do much the same with the Moon and Mercury, but since less detailed data are available, the conclusions are correspondingly less precise.

### The Moon

The Moon’s average density, about  $3300 \text{ kg/m}^3$ , is similar to the measured density of lunar surface rock, virtually eliminating any chance that the Moon has a large, massive, and very dense nickel–iron core like that of Earth. In fact, the low density implies that the entire Moon is deficient in iron and other heavy metals compared with their abundance on our planet.

There is no evidence for any large-scale lunar magnetic field. *Lunar Prospector* detected some very weak surface magnetic fields—less than a thousandth of Earth’s field—apparently associated with some large impact basins, but these are not thought to be related to conditions in the lunar core. As we saw in Chapter 7, researchers think that planetary magnetism requires a rapidly rotating liquid metal core, like Earth’s. ∞ (Sec. 7.5) Thus, the absence of a lunar magnetic field could be a consequence of the Moon’s slow rotation, the absence of a liquid core, or both.

Based on a combination of seismic data, gravitational and magnetic measurements, and a good deal of mathematical modeling resting on assumptions about the Moon’s interior composition, Figure 8.24 presents a schematic diagram of the Moon’s interior structure. The models suggest a central core about 330 km in radius, surrounded by a roughly 400-km-thick inner mantle of semi-solid rock having properties similar to Earth’s asthenosphere. ∞ (Sec. 7.4) Above the inner



▲ **FIGURE 8.24 Lunar Interior** Cutaway diagram of the Moon. Unlike Earth’s rocky lithosphere, the Moon’s is very thick—about 900 km. Below the lithosphere is the inner mantle, or lunar asthenosphere, a semisolid layer similar to the upper regions of Earth’s mantle.

mantle lies a 900-km-thick outer mantle of solid rock, topped by a 30-km-thick crust, as measured by *GRAIL* (see *Discovery 8-1*)—only slightly thicker than that of Earth. Together, these outer layers constitute the Moon’s lithosphere.

Data from the gravity experiment aboard *Lunar Prospector*, combined with magnetic measurements made as the Moon passed through Earth’s magnetic “tail” (see Figure 7.18), imply that the lunar core is denser and more iron rich than the rest of the Moon. Theoretical models predict a central temperature as low as 1500 K, too cool to melt rock or iron, but recent (2011) reanalysis of seismic data collected by sensitive equipment left on the surface by *Apollo* astronauts (see *Discovery 8-1*) suggest that the inner parts of the core may be at least partially molten, implying a somewhat higher temperature. As best we can tell, the Moon has a solid, mainly iron, inner core roughly 240 km in radius. The rest of the core, as well as the innermost 150 km of the surrounding inner mantle, is liquid.

Outside the core, the mantle seems to be of almost uniform density, although it is chemically differentiated (i.e., its chemical properties change from the deep interior to near the surface). The crust material, which forms the lunar highlands, is lighter than the mantle, which is similar in composition to the lunar maria.

The *GRAIL* mission found that the Moon’s battered crust is fractured and shattered to a depth of several kilometers, giving it a somewhat porous structure and a lower density (about 2600 kg/m<sup>3</sup>) than had previously been thought. In addition, while the crust averages about 30 km thick, its thickness is quite variable, ranging from as much as 60 km in some locations on the far side to almost zero under the larger near-side basins (see Figure 8.24). On average, the crust on the lunar far side is some 10–15 km thicker than that on the side facing Earth. If we assume that lava takes the line of least resistance in getting to the surface, then we can readily understand why the far side of the Moon has no large maria: Volcanic activity did not occur on the far side simply because the crust was too thick to allow it to occur there.

Why is the far-side crust thicker? The answer is probably related to Earth’s gravity. Just as heavier material tends to sink to the center of Earth, the denser lunar mantle tended to sink below the lighter crust in Earth’s gravitational field. The effect of this tendency was that the crust and the mantle became slightly off center with respect to each other. The mantle was pulled a little closer to Earth, while the crust moved slightly away. Thus, the crust became thinner on the near side and thicker on the far side.


## Mercury

Mercury’s magnetic field, discovered by *Mariner 10*, is about a hundredth that of Earth. Actually, the discovery that Mercury has any magnetic field at all came as a surprise to planetary scientists. Having detected no

magnetic field in the Moon (and, in fact, none in Venus or Mars, either), they had expected Mercury to have no measurable magnetism. Mercury does not rotate rapidly (as was thought necessary for a planetary dynamo to work), yet a magnetic field undeniably surrounds it. [☞ \(Sec. 7.5\)](#) Although weak, the field is strong enough to deflect the solar wind and create a small magnetosphere around the planet.

Before *Messenger*’s arrival, scientists thought it most likely that Mercury’s magnetic field was a “fossil remnant” dating back to the distant past when the planet’s core solidified. However, detailed observations by *Messenger* now suggest that the field is generated by dynamo action in the planet’s molten outer core, as on Earth. [☞ \(Sec. 7.5\)](#) In fact, an early surprise for the *Messenger* team was the degree to which the planet’s magnetosphere changed between the first two flybys, in January and October 2008. How such a relatively strong and variable field can be produced by a slowly rotating planet is unknown. Also unexplained is the fact that the entire field pattern is shifted almost 500 km from the planet’s center, toward the north pole.

Mercury’s magnetic field and large average density together imply that the planet is differentiated. Based on *Messenger* measurements, the planet seems to have a solid inner core with a radius of perhaps 1600 km, surrounded by a liquid outer core of radius 2100 km. A less dense lunar-like mantle some 350 km thick lies above the core. About 60 percent of the volume of Mercury, or 80 percent of its mass, is contained in its huge iron core. The ratio of core volume to total planet volume is greater for Mercury than for any other object in the solar system. Figure 8.25 illustrates the relative sizes and internal structures of Earth, the Moon, and Mercury.



▲ **FIGURE 8.25 Terrestrial Interiors** The internal structures of Earth, the Moon, and Mercury, drawn to the same scale. Note how large a fraction of Mercury’s interior is core. Planetary interiors are key to the global subject of comparative planetology.

**PROCESS OF SCIENCE Check**

✓ Why would we not expect strong magnetic fields on the Moon or Mercury?

**8.8 Origin of the Moon**

Over the years, many theories have been advanced to account for the origin of the Moon. However, both the similarities *and* the differences between the Moon and Earth conspire to confound many promising attempts to explain the Moon's existence.

**Theories of Lunar Formation**

One theory (the *sister*, or *coformation*, theory) suggests that the Moon formed as a separate object near Earth in much the same way as our own planet formed—the “blob” of material that eventually coalesced into Earth gave rise to the Moon at about the same time. The two objects thus formed as a double-planet system, each revolving about a common center of mass. Although once favored by many astronomers, this idea suffers from a major flaw: The Moon differs in both density and composition from Earth, making it hard to understand how both could have originated from the same preplanetary material.

A second theory (the *capture* theory) maintains that the Moon formed far from Earth and was later captured by it. In this way, the density and composition of the two objects need not be similar, for the Moon presumably materialized in a quite different region of the early solar system. The objection to this theory is that the Moon's capture would be an extraordinarily difficult event; it might even be an impossible one. Why? Because the mass of our Moon is so large relative to that of Earth. It is not that our Moon is the largest natural satellite in the solar system, but it is unusually large compared with its parent planet. Mathematical modeling suggests that it is quite implausible that Earth and the Moon could have interacted in just the right way for the Moon to have been captured during a close encounter sometime in the past. Furthermore, although there are indeed significant differences in composition between our world and its companion, there are also many similarities—particularly between the mantles of the two bodies—that make it unlikely that they formed entirely independently of one another.

A third, older, theory (the *daughter*, or *fission*, theory) speculates that the Moon originated out of Earth itself. The Pacific Ocean basin has often been mentioned as the place from which protolunar matter may have been torn—the result, perhaps, of the rapid spin of a young, molten Earth. Indeed, there are some chemical similarities between the matter in the Moon's outer mantle and that in Earth's Pacific basin. However, this theory offers no solution to the fundamental mystery of how Earth could have been

spinning so fast that it ejected an object as large as our Moon. Also, computer simulations indicate that the ejection of the Moon into a stable orbit simply would not have occurred. As a result, the daughter theory, in this form at least, is no longer taken seriously.

**The Impact Theory**


Today, many astronomers favor a hybrid of the capture and daughter themes. This idea—often called the **impact theory**—postulates a collision by a large, Mars-sized object with a youthful and molten Earth. Such collisions may have been quite frequent in the early solar system. ∞ (Sec. 6.7) The collision presumed by the impact theory would have been more a glancing blow than a direct impact. The matter dislodged from our planet then reassembled to form the Moon.

Computer simulations of such a catastrophic event show that most of the bits and pieces of splattered Earth could have coalesced into a stable orbit. Figure 8.26 shows some of the stages of one such calculation. If Earth had already formed an iron core by the time the collision occurred, then the Moon would indeed have ended up with a composition similar to that of Earth's mantle. During the collision, any iron core in the colliding object itself would have been left behind in Earth, eventually to become part of Earth's core. Thus, both the Moon's overall similarity to that of Earth's mantle and its lack of a dense central core are naturally explained.

Over the past two decades, planetary scientists have come to realize that collisions like this probably played important roles in the formation of all the terrestrial planets. ∞ (Sec. 6.6) Because of the randomness inherent in such events, as well as the Moon's unique status as the only large satellite in the inner solar system, it seems that the Moon may not provide a particularly useful model for studies of the other moons in the solar system. Instead, as we will see, a moon's properties depend greatly on the characteristics of its parent planet.

Nevertheless, the quest to understand the origin of the Moon highlights the interplay between theory and observation that characterizes modern science. ∞ (Sec. 1.2) Detailed data from generations of unmanned and manned lunar missions have allowed astronomers to discriminate between competing theories of the formation of the Moon, discarding some and modifying others. At the same time, the condensation theory of solar system formation provides a natural context in which the currently favored impact theory can occur. ∞ (Sec. 6.6) Indeed, without the idea that planets formed by collisions of smaller bodies, such an impact might well have been viewed as so improbable that the theory would never have gained ground.

Finally, do not think that every last detail of the Moon's formation is understood or agreed upon by experts. That is far from the case. Some important aspects of the Moon's physical and chemical makeup are still inadequately explained. For example, the degree to which the Moon



melted during its formation remains uncertain, and current models have some difficulty accounting for the observed lunar composition—although *GRAIL*'s recent revision of the density of the lunar crust has significantly improved the agreement between theory and observation on this latter point. The impact theory may well not be the last word on the subject. Still, past experience of the scientific method gives us confidence that the many twists and turns still to come will in the end lead us to a more complete understanding of our nearest neighbor in space.

### PROCESS OF SCIENCE Check

- ✓ How does the currently favored theory of the Moon's origin account for the Moon's observed lack of heavy materials compared with Earth and for the similarity in composition between the lunar crust and that of Earth?


## 8.9 Evolutionary History of the Moon and Mercury

Given all the data, can we construct reasonably consistent histories of the Moon and Mercury? The answer seems to be yes. Many specifics are still debated, but a broad consensus exists. Planned future missions to both bodies will continue to test and refine the pictures presented below.

### The Moon

The Moon formed about 4.6 billion years ago. ∞ (Sec. 6.6) The approximate age of the oldest rocks discovered in the lunar highlands is 4.4 billion years, so we know that at least part of the crust must already have solidified by that time and survived to the present. At its formation, the Moon was already depleted in heavy metals compared with Earth. Examine Figure 8.27 while studying the details that follow.

During the earliest phases of the Moon's existence—roughly the first half billion years or so—meteoritic bombardment must have been frequent enough to heat and remelt most of the *surface* layers of the Moon, perhaps to a depth of 400 km in places. The early solar system was surely populated with lots of interplanetary matter, much of it in the form of boulder-sized fragments that were capable of generating large amounts of energy upon colliding with planets and their moons. But the intense heat derived from such collisions could not have penetrated very far into the lunar interior: Rock simply does not conduct heat well.

 **Narrated FIGURE 8.26 Moon Formation** This sequence shows a simulated collision between Earth and an object the size of Mars. The sequence proceeds from top to bottom and zooms out dramatically. Note how most of the impactor's metallic core becomes part of Earth, leaving the Moon composed mainly of rocky material. (*W. Benz*)

This situation resembles the surface melting we suspect occurred on Earth from meteoritic impacts during the first billion years or so. But the Moon is much less massive than Earth and did not contain enough radioactive elements to heat it much further. Radioactivity probably heated the Moon a little, but not sufficiently to transform it from a warm, semisolid object to a completely liquid one. The chemical differentiation now inferred in the Moon's interior must have occurred during this period. If the Moon has a small iron core, that core also formed at this time.

The *GRAIL* mission has added an intriguing new ingredient to studies of the Moon's early history by uncovering evidence of linear "dikes" in the lunar surface that apparently *predate* many of the large impact features. The dikes are thought to be sheets of cooled lava that welled up through fissures in the crust during a previously unknown period of interior expansion during the first few hundred million years of the Moon's existence.

About 3.9 billion years ago, around the time that Earth's crust solidified, the heaviest phase of the meteoritic bombardment ceased. The Moon was left with a solid crust, which would ultimately become the highlands, dented with numerous large basins, soon to flood with lava and become the maria (Figure 8.27a). Between 3.9 and 3.2 billion years ago, lunar volcanism filled the maria with the basaltic material we see today. The age of the youngest maria—3.2 billion years—indicates the time when the volcanic activity subsided. The maria are the sites of the last extensive lava flows on the Moon, over 3 billion years ago. Their smoothness, compared with the older, more rugged highlands, disguises their great age.

Small objects cool more rapidly than large ones because their interior is closer to the surface, on average.


Being so small, the Moon rapidly lost its internal heat to space. As a consequence, it cooled much faster than Earth. As the Moon cooled, the volcanic activity ended and the thickness of the solid surface layer increased. With the exception of a few meters of surface erosion from eons of meteoritic bombardment (Figure 8.27c), the lunar landscape has remained more or less structurally frozen for the past 3 billion years. The Moon is dead now, and it has been dead for a long time.

## Mercury

Like the Moon, Mercury seems to have been a geologically dead world for much of the past 4 billion years. On both the Moon and Mercury, the absence of ongoing geological activity is a consequence of a thick, solid mantle that prevents volcanism or tectonic motion. Because of the *Apollo* program, the Moon's early history is much better understood than Mercury's, which remains somewhat speculative. Indeed, what we do know about Mercury's history is gleaned mostly through comparison with the Moon.

Mercury's unexpectedly large iron core seems to be a consequence of the planet's location in the hot inner regions of the early solar system when it formed some 4.6 billion years ago—much as described in our discussion of the condensation theory in Chapter 6, although the sheer size of the core still poses some problems to theorists. [∞ \(Sec. 6.6\)](#)

Before *Messenger*, some researchers had suggested that extreme conditions in the early solar nebula might have vaporized most of Mercury's original outer rocky layers or that a violent collision could have stripped away much of the planet's light mantle. However, both these



**▲ FIGURE 8.27 Lunar Evolution** Paintings of the Moon (a) about 4 billion years ago, after much of the meteoritic bombardment had subsided and the surface had somewhat solidified; (b) about 3 billion years ago, after molten lava had made its way up through surface fissures to fill the low-lying impact basins and create the smooth maria; and (c) today, with much of the originally smooth maria now heavily pitted with craters formed at various times within the past 3 billion years. (*U.S. Geological Survey*)

scenarios predict very high temperatures and imply that the present crust should be deficient in certain volatile compounds containing potassium, sulfur, and other elements, which would have been preferentially lost. Instead, new *Messenger* data indicate that the composition of the crust is remarkably “normal”—similar to the makeup of the other terrestrial planets—effectively ruling out these alternative theories of Mercury’s core size.

### ∞ (Sec. 1.2)

During the next half-billion years, Mercury melted and differentiated, like the other terrestrial worlds. It suffered the same intense meteoritic bombardment as the Moon. Being more massive than the Moon, Mercury cooled more slowly, so its crust was thinner and flood

volcanic activity more common at early times. Many craters were erased, resulting in the intercrater plains. The smooth plains around several major impact sites were produced late in the bombardment, some 3.8 billion years ago.


As Mercury’s large iron core formed and then cooled, the planet began to shrink, compressing the crust. This compression produced the scarps seen on Mercury’s surface and may have prematurely terminated volcanic activity by squeezing shut the cracks and fissures on the surface. Thus, the extensive volcanic outflows that formed the lunar maria did not take place on Mercury. Despite its larger mass and greater internal temperature, Mercury has probably been geologically inactive even longer than the Moon.

**The Big Question** While Mercury seems hopelessly uninhabitable, we still often wonder, Will humans eventually colonize the Moon—perhaps even “terraforming,” or changing, it to suit our purposes? Fifty years ago, humankind appeared well on its way toward establishing permanent lunar habitats, but those early (and spectacular) exploratory programs stalled. Today, the political will or economic means to mount human missions to even our closest neighbor in space has faded. Should governments, even the UN, lead the return to the Moon, or will it be best accomplished with private entrepreneurial ventures like those that settled the Americas centuries ago?

# Chapter Review

## SUMMARY

**1** The Moon orbits Earth; Mercury is the closest planet to the Sun. Both the Moon and Mercury are airless, virtually unchanging worlds that exhibit extremes in temperature. Mercury has no permanent atmosphere, although it does have a thin envelope of gas temporarily trapped from the solar wind. Both bodies are smaller and less massive than Earth and have weaker gravities. The absence of atmospheric blankets results in hot dayside temperatures and cold nightside temperatures on the Moon and Mercury. Sunlight strikes the polar regions of both the Moon and Mercury at such an oblique angle that temperatures there are very low, with the result that both bodies may have significant amounts of water ice near their poles.



**2** The main surface features on the Moon are the dark **maria** (p. 192) and the lighter colored **highlands** (p. 192). Highland rocks are less dense than rocks from the maria and are thought to represent the Moon’s crust. Maria rocks are thought to have originated in the lunar mantle. The surfaces of both the Moon and Mercury are covered with **craters** (p. 192) of all sizes, caused by meteoroids striking from space.



Lunar dust, called regolith, is made mostly of pulverized lunar rock, mixed with a small amount of material from impacting meteorites.

**3** The tidal interaction between Earth and the Moon is responsible for the Moon’s **synchronous orbit** (p. 195), in which the same side of the Moon always faces our planet. The large lunar equatorial bulge probably indicates that the Moon once rotated more rapidly and orbited closer to Earth. Mercury’s rotation rate is strongly influenced by the tidal effect of the Sun. Because of Mercury’s eccentric orbit, the planet rotates not synchronously, but exactly three times for every two orbits around the Sun. The condition in which a body’s rotation rate is simply related to its orbital period around some other body is known as **spin-orbit resonance** (p. 198).




**4** Meteoritic impacts are the main source of erosion on the surfaces of both the Moon and Mercury. The lunar highlands are older than the maria and are much more heavily cratered. The rate at which craters are formed decreases rapidly with increasing crater size. By measuring the ages of lunar rocks returned to Earth by *Apollo* astronauts, astronomers have deduced the rate of cratering in the past. They then use the amount of cratering to deduce the ages of




regions on the Moon (and elsewhere) from which surface samples are unavailable.

**5** Evidence for past volcanic activity on the Moon is found in the form of crater chains and solidified lava channels called **rilles** (p. 206). Mercury's surface features bear a striking similarity to those of the Moon. The planet is heavily cratered, much like the lunar highlands. Among the differences between Mercury and the Moon are Mercury's lack of lunarlike maria, its extensive **intercrater plains** (p. 194), and the great cracks, or **scarps** (p. 206), in its crust. The plains were caused by extensive lava flows early in Mercury's history. The scarps were apparently formed when the planet's core cooled and shrank, causing the surface to crack. Mercury has a large impact crater called the Caloris Basin, whose diameter is comparable to the radius of the planet. The impact that formed the crater apparently sent violent shock waves around the entire planet, buckling the crust on the opposite side.




**6** The Moon's average density is not much greater than that of its surface rocks, probably because the Moon cooled more rapidly than the larger Earth and solidified sooner, so there was less time for differentiation to occur, although the Moon likely has a small iron-rich core. The lunar crust is too thick and the mantle




too cool for plate tectonics to occur. Mercury's average density is considerably greater—similar to that of Earth—implying that Mercury contains a large high-density core, probably composed primarily of iron. The Moon has no measurable large-scale magnetic field, a consequence of its slow rotation and lack of a molten metallic core. Mercury's weak magnetic field seems to have been “frozen in” long ago, when the planet's iron core solidified.

**7** The most likely explanation for the formation of the Moon is that the newly formed Earth was struck by a large (Mars-sized) object. Part of the colliding body remained behind as part of our planet. The rest ended up in orbit as the Moon.



**8** The absence of a lunar atmosphere and any present-day lunar volcanic activity are both consequences of the Moon's small size. Lunar gravity is too weak to retain any gases, and lunar volcanism was stifled by the Moon's cooling mantle shortly after extensive lava flows formed the maria more than 3 billion years ago. The crust on the far side of the Moon is substantially thicker than the crust on the near side. As a result, there are almost no maria on the lunar far side. Mercury's evolutionary path was similar to that of the Moon for half a billion years after they both formed. Mercury's volcanic period probably ended before that of the Moon.



**MasteringAstronomy**® For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- How is the distance to the Moon most accurately measured?
- LO1** Compare and contrast the bulk properties of Earth, the Moon, and Mercury.
- Employ the concept of escape speed to explain why the Moon and Mercury have no significant atmospheres.
- LO2** Why is the surface of Mercury often compared with that of the Moon? List two similarities and two differences between the surfaces of Mercury and the Moon.
- LO3** What does it mean to say that Mercury has a 3:2 spin-orbit resonance? Why didn't Mercury settle into a synchronous orbit around the Sun, as the Moon did around Earth?
- What is a scarp? How are scarps thought to have formed? Why do scientists think that the scarps on Mercury formed after most meteoritic bombardment ended?
- What is the primary source of erosion on the Moon? Why is the average rate of lunar erosion so much less than on Earth?
- What evidence do we have for ice on the Moon?
- LO4 POS** Name two pieces of evidence indicating that the lunar highlands are older than the maria.
- LO5** When were the periods of greatest volcanic activity on the Moon and Mercury?
- LO6** What evidence do we have that the Moon and Mercury have liquid cores?
- LO7 POS** Describe the theory of the Moon's origin favored by many astronomers.
- LO8** How is Mercury's evolutionary history like that of the Moon? How is it different?
- The best place to aim a telescope or binoculars on the Moon is along the terminator line—the line between the Moon's light and dark hemispheres. Why is this? If you were standing on the lunar terminator, where would the Sun be in your sky? What time of day would it be if you were standing on Earth's terminator line?
- How is the varying thickness of the lunar crust related to the presence or absence of maria on the Moon?

## Conceptual Self-Test: Multiple Choice

- Compared with the diameter of Earth's Moon, the diameter of Mercury is (a) larger; (b) smaller; (c) nearly the same.
- In relation to the density of Earth's Moon, Mercury's density suggests that the planet (a) has an interior structure similar to that of the Moon; (b) has a dense metal core; (c) has a stronger magnetic field than the Moon; (d) is younger than the Moon.
- Compared with the phases of Earth's Moon, Mercury goes from new phase to full phase (a) faster; (b) more slowly; (c) in about the same time.
- Compared with the surface of Mercury, the surface of Earth's Moon has significantly (a) bigger craters; (b) more atmosphere; (c) more maria; (d) deeper craters.
- VIS** According to Figure 8.10 ("The Moon's Synchronous Rotation"), every two times Earth's Moon rotates on its axis, it orbits Earth (a) less than twice; (b) exactly two times; (c) more than twice; (d) three times.
- Planets and moons showing the most craters have (a) the oldest surfaces; (b) been hit by meteors the most times; (c) the strongest gravity; (d) molten cores.
- Compared with the Moon, Mercury has (a) a much smaller core; (b) a much larger core; (c) a similar-sized core.
- The most likely theory of the formation of Earth's Moon is that it (a) was formed by the gravitational capture of a large asteroid; (b) formed simultaneously with Earth's formation; (c) was created from a collision scooping out the Pacific Ocean; (d) formed from a collision of Earth with a Mars-sized object.
- Mercury, being smaller than Mars, probably cooled and solidified (a) faster, because it is smaller; (b) slower, because it is closer to the Sun; (c) in about the same time, because space is generally cold.
- On the scale of the 5-billion-year age of the solar system, the Moon is (a) about the same age as Earth; (b) much younger than Earth; (c) much older than Earth.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

- How long does a radar signal take to travel from Earth to Mercury and back when Mercury is at its closest point to Earth?
- The Moon's mass is one-eightieth that of Earth, and the lunar radius is one-fourth Earth's radius. On the basis of these figures, calculate the total weight on the Moon of a 100-kg astronaut with a 50-kg space suit and backpack. What would be the same astronaut's weight on Mercury?
- What is the angular diameter of the Sun, as seen from Mercury, at perihelion? At aphelion?
- The *Hubble Space Telescope* has a resolution of about 0.05". What is the size of the smallest feature it can distinguish on the surface of the Moon (distance = 380,000 km)? On Mercury, at closest approach to Earth?
- What was the orbital period of the *Apollo 11* command module, orbiting 10 km above the lunar surface?
- Compare the gravitational tidal acceleration of the Sun on Mercury (at perihelion; solar mass =  $2 \times 10^{30}$  kg) with the tidal effect of Earth on the Moon (at perigee).  $\infty$  (Sec. 7.6)
- (a) Using the rate given in the text for the formation of 10-km craters on the Moon, estimate how long would be needed for the entire Moon to be covered with new craters of that size. How much higher must the cratering rate have been in the past to cover the entire lunar surface with such craters in the 4.6 billion years since the Moon formed? (b) Repeat part (a) for meter-size craters.
- Assume that a planet will have lost its initial atmosphere by the present time if the average molecular speed exceeds one-sixth of the escape speed (see *More Precisely 8-1*). What would Mercury's mass have to be in order for it to still have a nitrogen atmosphere? The molecular weight of nitrogen is 28.

## Activities


### Collaborative

- The estimated cost of transporting a gallon of water from Earth to the Moon is about \$100,000. By determining how much water each group member uses in a single day, estimate the cost of taking a single day's supply of water for your group to the Moon.
- Observe the Moon during an entire cycle of phases. Take turns, and on each observing night sketch the appearance of the Moon and keep a log of the date and time. When does the Moon rise, set, and appear highest in the sky at each major phase? What is the interval of time between each phase?

### Individual

- Try to spot Mercury in the morning or evening twilight—not an easy task! From the Northern Hemisphere, the best evening sightings of the planet take place in the spring; the best morning sightings take place in the fall.
- Watch the Moon over a period of hours on a night when you can see one or more bright stars near it. Estimate how many Moon diameters it moves per hour, relative to the stars. Knowing the Moon is about  $0.5^\circ$  in diameter, how many degrees per hour does it move? Based on this, what is your estimate of the Moon's orbital period?





# Venus

## EARTH'S SISTER PLANET

Venus seems almost a carbon copy of our own world. The two planets are similar in size, density, and chemical composition. They orbit at comparable distances from the Sun. At formation, they must have been almost indistinguishable from one another. Yet they are now about as different as two terrestrial planets can be. Whereas Earth is a vibrant world, teeming with life, Venus is an uninhabitable inferno, with a dense, hot atmosphere of carbon dioxide, lacking any trace of oxygen or water.

Somewhere along their respective evolutionary paths, Venus and Earth diverged, and diverged radically. How did this occur? What were the factors leading to Venus's present condition? Why are Venus's surface, atmosphere, and interior so different from Earth's? In answering these questions, we will discover that a planet's environment, as well as its composition, can play a critical role in determining its future.

**The Big Picture** Earth and Venus started out much the same, but ended up very different. The same physical processes that keep our planet warm, comfortable, and habitable turned our sister planet into an unimaginable inferno. Astronomy is not a very practical subject, but here it is very much in our own interest to know why Venus became so hellishly hot. By probing the histories of these two comparably sized worlds, astronomers seek to understand why Venus and Earth changed so differently—and whether our home in space might someday undergo a similar climate catastrophe.

**LEFT:** Often called Earth's sister planet because of their comparable sizes, Venus is actually nothing like our own world. Surface conditions on Venus have changed radically over time due to geological activity and environmental change, and today the planet's surface temperature is hot enough (730 K) to melt lead, while the atmosphere rains sulfuric acid. This global view of the surface of Venus was created when radar data from the *Magellan* spacecraft were mapped onto a computer-generated globe. The color here is probably close to reality. (JPL)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Summarize Venus's general orbital and physical properties.
- 2 Describe the characteristics of Venus's atmosphere and contrast it with that of Earth.
- 3 Compare the large-scale surface features and geology of Venus with those of Earth and the Moon.
- 4 Present the evidence for ongoing volcanic activity on Venus.
- 5 Explain why the greenhouse effect has produced conditions on Venus very different from those on Earth.
- 6 Describe Venus's magnetic field and internal structure.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.


## 9.1 Orbital Properties

Venus is the second planet from the Sun. Its orbit lies within Earth's, so Venus, like Mercury, is always found fairly close to the Sun in the sky—our sister planet is never seen more than  $47^\circ$  from the Sun. Given Earth's rotation rate of  $15^\circ$  per hour, this means that Venus is visible above the horizon for at most 3 hours before the Sun rises or after it sets. Because we can see Venus from Earth only just before sunrise or just after sunset, the planet is often called the “morning star” or the “evening star,” depending on where it happens to be in its orbit. Figure 9.1 shows Venus in the western sky just after sunset.

Venus is the third-brightest object in the entire sky (after the Sun and the Moon). It appears more than 10 times brighter than the brightest star, Sirius. You can see Venus even in the daytime if you know just where to look. On a moonless night away from city lights, Venus casts a faint shadow. The planet's brightness stems from the fact that Venus is highly reflective. Nearly 70 percent of the sunlight reaching Venus is reflected back into space. (Compare this percentage with roughly 10 percent in the case of Mercury and the Moon.) Most of the sunlight is reflected from clouds high in the planet's atmosphere.


We might expect Venus to appear brightest when it is “full”—that is, when we can see the entire sunlit side. However, because Venus orbits between Earth and the Sun, Venus is full when it is at its greatest distance from us—1.7 AU away on the other side of the Sun, as illustrated in Figure 9.2. Recall from Chapter 2 that this alignment is known as *superior conjunction*, where the term “conjunction” simply indicates that two objects are close together in the sky.  $\infty$  (Sec. 2.2)

When Venus is closest to us, the planet is in the new phase, lying between Earth and the Sun (at *inferior conjunction*), and we again can't see it, because now the sunlit side faces away from us; only a thin ring of sunlight, caused by refraction in Venus's atmosphere, surrounds the planet. As Venus moves away from inferior conjunction, more and more of it becomes visible, but its distance from us also





**▲ FIGURE 9.1 Venus at Sunset** The Moon and Venus in the western sky just after sunset. Venus clearly outshines even the brightest stars in the sky. (J. Schad/Science Source)

continues to increase. Venus's maximum brightness, as seen from Earth, actually occurs about 36 days before or after its closest approach to our planet. At that time, Venus is about  $39^\circ$  from the Sun and 0.47 AU from Earth, and we see it as a rather fat crescent.



**MA** **Narrated FIGURE 9.2 Venus's Brightness** Venus appears full when it is at its greatest distance from Earth, on the opposite side of the Sun from us (superior conjunction). As its distance decreases, less and less of its sunlit side becomes visible. When closest to Earth, it lies between us and the Sun (inferior conjunction), so we cannot see the sunlit side of the planet at all. Venus appears brightest when it is about  $39^\circ$  from the Sun. (Compare Figure 2.12.) (Insets: UC/Lick Observatory)



## 9.2 Physical Properties

### Radius, Mass, and Density

We can determine Venus's radius from simple geometry, just as we did for Mercury and the Moon. [∞ \(Sec. 8.2\)](#) At closest approach, when Venus is only 0.28 AU from us, its angular diameter is 64". From this observation, we can determine the planet's radius to be about 6000 km. More accurate measurements from spacecraft give a value of 6052 km, or 0.95 Earth radii.

Like Mercury, Venus has no moon. Before the Space Age, astronomers calculated its mass by indirect means—through studies of its small gravitational effect on the orbits of the other planets, especially Earth. Now that spacecraft have orbited the planet, we know Venus's mass very accurately from measurements of its gravitational pull: Venus has a mass of  $4.9 \times 10^{24}$  kg, or 0.82 the mass of Earth.

From its mass and radius, we find that Venus's average density is  $5200 \text{ kg/m}^3$ . As far as these bulk properties are concerned, then, Venus seems similar to Earth. If the planet's overall composition were similar to Earth's as well, we could then reasonably conclude that Venus's internal structure and evolution were basically Earth-like. We will review what evidence there is on this subject later in the chapter.

### Rotation Rate

The same clouds whose reflectivity makes Venus so easy to see in the night sky also make it impossible for us to discern any surface features on the planet, at least in visible light. As a result, until the advent of suitable radar techniques in the 1960s, astronomers did not know the rotation period of Venus. Even when viewed through a large optical telescope, the planet's cloud cover shows few features, and attempts to determine Venus's period of rotation by observing the cloud layer were frustrated by the rapidly changing nature of the


clouds themselves. Some astronomers argued for a 25-day period, while others favored a 24-hour cycle.

Controversy raged until, to the surprise of all, radar observers announced that the Doppler broadening of their returned echoes implied a sluggish 243-day rotation period! [∞ \(Sec. 8.4\)](#) Furthermore, Venus's spin was found to be *retrograde*—that is, in a sense opposite that of Earth and most other solar system objects and opposite that of Venus's orbital motion.

Planetary astronomers define “north” and “south” for each planet in the solar system by the convention that planets *always* rotate from west to east. With this definition, Venus's retrograde spin means that the planet's north pole lies *below* the plane of the ecliptic, unlike any of the other terrestrial worlds. Venus's axial tilt—the angle between its equatorial and orbital planes—is  $177.4^\circ$  (compared with  $23.5^\circ$  in the case of Earth). However, astronomical images of solar system objects conventionally place objects lying above the ecliptic at the top of the frame. Thus, with the preceding definition of north and south, all the images of Venus shown in this chapter have the *south* pole at the top.


Figure 9.3 illustrates Venus's retrograde rotation and compares it with the rotation of its neighbors Mercury, Earth, and Mars. Because of the planet's slow retrograde rotation, its solar day (from noon to noon) is quite different from its sidereal rotation period of 243 Earth days (the time for one “true” rotation relative to the stars). [∞ \(Sec. 1.4\)](#) In fact, as illustrated in Figure 9.4, one Venus day is a little more than half a Venus year (225 Earth days).

Why is Venus rotating “backward” and why so slowly? At present, the best explanation planetary scientists can offer is that early in Venus's evolution, the planet was struck by a large body, much like the one that may have hit Earth and formed the Moon, and that impact was sufficient to reduce the planet's spin almost to zero. [∞ \(Sec. 8.8\)](#) Whatever its cause, the planet's rotation poses practical problems for Earth-bound observers. As luck would have it, Venus rotates



◀ **FIGURE 9.3 Terrestrial Planets' Spin**

The inner planets of the solar system—Mercury, Venus, Earth, and Mars—display widely different rotational properties. All orbit the Sun in the same direction and in nearly the same plane, but Venus rotates clockwise as seen from above the plane of the ecliptic, whereas Mercury, Earth, and Mars all spin counterclockwise. This is a perspective view, roughly halfway between a flat edge-on view and a direct overhead view.



▲ **FIGURE 9.4 Venus's Solar Day** Venus's orbit and retrograde rotation combine to produce a solar day on Venus equal to 117 Earth days, or slightly more than half a Venus year. The red arrows denote a fixed location, or an observer standing, on the planet's surface. The numbers in the figure mark time in Earth days.

almost exactly five times between one closest approach to Earth and the next. As a result, *Venus always presents nearly the same face to Earth at closest approach*. This means that observations of the planet's surface cover one side—the one facing us at closest approach—much more thoroughly than the other side, which we can see only when the planet is close to its maximum distance from Earth.

### CONCEPT Check

- ✓ **What is peculiar about Venus's rotation, and why does Venus rotate that way?**

## 9.3 Long-Distance Observations of Venus

Because Venus, of all the other planets, most nearly matches Earth in size, mass, and density, and because its orbit is closest to us, it is often called Earth's sister planet. But unlike Earth, Venus has a dense atmosphere and thick clouds that are opaque to visible radiation, making its surface completely invisible from the outside at optical wavelengths. Figure 9.5 shows one of the best photographs of Venus taken with a large telescope on Earth. The planet presents an almost featureless white-yellow disk, although it shows occasional hints of cloud circulation.


Atmospheric patterns on Venus are much more evident when the planet is examined with equipment capable



▲ **FIGURE 9.5 Venus** This photograph, taken from Earth, shows Venus with its creamy yellow mask of clouds. No surface detail can be seen because the clouds completely obscure our view of what lies beneath them. (AURA)

of detecting ultraviolet radiation. Some of Venus's atmospheric constituents absorb this high-frequency radiation, greatly increasing the cloud contrast. Figure 9.6(a) shows an ultraviolet image taken in 1979 by the U.S. *Pioneer Venus* spacecraft at a distance of 200,000 km from the planet's surface; Figure 9.6(b) shows a 2006 mosaic of infrared images from the European *Venus Express* orbiter, whose cameras partially penetrate the planet's thick haze. The large, fast-moving cloud patterns resemble Earth's high-altitude jet stream more than the great whirls characteristic of Earth's low-altitude clouds. The upper deck of clouds on Venus move at almost 400 km/h, encircling the planet in just 4 days—much faster than the planet itself rotates!

Early spectroscopic studies of sunlight reflected from Venus's clouds revealed the presence of large amounts of carbon dioxide, but provided little evidence for any other atmospheric gases. Until the 1950s, astronomers generally believed that observational difficulties alone prevented them from seeing other atmospheric components. The hope lingered that Venus's clouds were actually predominantly water vapor, like those on Earth, and that below the cloud



▲ **FIGURE 9.6 Venus, Up Close** (a) This image of Venus was made when the *Pioneer* spacecraft captured solar ultraviolet radiation reflected from the planet's upper clouds, which are probably composed mostly of sulfuric acid droplets, much like the corrosive acid in a car battery. (b) Venus in the infrared, as seen by *Venus Express* on approach to the planet. The longer infrared wavelength allows us to "see" deeper into Venus's lower clouds. (NASA; ESA)

cover Venus might be a habitable planet similar to our own. Indeed, in the 1930s, scientists had measured the temperature of the atmosphere spectroscopically at about 240 K, not much different from the temperature of our own upper atmosphere. ∞ (Secs. 4.5, 7.2) Calculations of the planet's surface temperature—taking into account the cloud cover and Venus's proximity to the Sun, and assuming an atmosphere much like our own—suggested that Venus should have a surface temperature only 10 or 20 degrees higher than Earth's.

These hopes for an Earth-like Venus were dashed in 1956, when radio observations of the planet were used to measure its thermal energy emission. Unlike visible light, radio waves easily penetrate the cloud layer—and they gave the first indication of conditions on or near the surface: The radiation emitted by the planet has a blackbody spectrum characteristic of a temperature near 730 K! ∞ (Sec. 3.4) Almost overnight, the popular conception of Venus changed from that of a lush tropical jungle to an arid, uninhabitable desert.

Radar observations of the surface of Venus are routinely carried out from Earth with the Arecibo radio telescope. ∞ (Sec. 5.5) With careful signal processing, this instrument can achieve a resolution of a few kilometers, but it can adequately cover only a fraction (roughly 25 percent) of the planet. The telescope's view of Venus is limited by the peculiar spin-orbit coincidence described in the previous section (which means that only one side of the planet can be studied) and also because radar reflections from regions near the "edge" of the planet are hard to obtain. However, the Arecibo data can usefully be combined with information received from probes orbiting Venus to build up a detailed picture of the planet's surface. Only with the arrival of the *Magellan* probe were more accurate data obtained.


### PROCESS OF SCIENCE Check

- ✓ Why did early studies of Venus lead astronomers to such an inaccurate picture of the planet's surface conditions?

## 9.4 The Surface of Venus

Although the planet's clouds are thick and the terrain below them totally shrouded, we are by no means ignorant of Venus's surface. Detailed radar observations have been made both from Earth and from the *Venera*, *Pioneer Venus*, and *Magellan* spacecraft. ∞ (Sec. 6.6) Analysis of the radar echoes yields a map of the planet's surface. Except for the last two figures, all the views of Venus in this section are "radargraphs" (as opposed to photographs) created in this way.

As Figure 9.7(a) illustrates, the early maps of Venus suffered from poor resolution; however, more recent probes—especially *Magellan*—have provided much sharper views. As in all the *Magellan* images, the light areas in Figure 9.7(b) represent regions where the surface is rough and efficiently scatters *Magellan's* sideways-looking radar beam back to the detector. Smooth areas tend to reflect the beam off into space instead and so appear dark. The strength of the returned signal as *Magellan* passed by thus results in a map of the planet's surface.



**▲ FIGURE 9.7 Venus Mosaics** (a) This image of the surface of Venus was made by a radar transmitter and receiver on board the *Pioneer* spacecraft, which is still in orbit about the planet, but is now inoperative. The two continent-sized landmasses are named Ishtar Terra (upper left) and Aphrodite (lower right). Colors represent altitude: Blue is lowest, red highest. The spatial resolution is about 25 km. (b) A planetwide mosaic of *Magellan* images, colored in roughly the same way as part (a). The largest “continent” on Venus, Aphrodite Terra, is the yellow dragon-shaped area across the center of this image. See also the full-page, chapter-opening photo on page 216. (NASA)

## Large-Scale Topography


Figure 9.8(a) shows basically the same *Pioneer Venus* data of Venus as Figure 9.7, except that this figure has been flattened out into a more conventional map. The altitude of the surface relative to the average radius of the planet is indicated by the use of color, with white representing the highest elevations and blue the lowest. (Note that the blue has nothing to do with oceans, nor does white indicate snow-capped mountains!) Figure 9.8(b) shows a map of Earth to the same scale and at the same spatial resolution. Some of Venus’s main features are labeled in Figure 9.8(c).

The surface of Venus appears to be relatively smooth, resembling rolling plains with modest highlands and lowlands. Two continent-sized features, called Ishtar Terra and Aphrodite Terra (named after the Babylonian and Greek counterparts, respectively, of Venus, the Roman goddess of love), adorn the landscape and contain mountains comparable in height to those on Earth. The elevated “continents” occupy only 8 percent of Venus’s total surface area. For comparison, continents on Earth make up about 25 percent of the surface. The remainder of Venus’s surface is classified as lowlands (27 percent) or rolling plains (65 percent), although there is probably little geological difference between the two terrains. Note that, although Earth’s tectonic plate boundaries are evident in Figure 9.8(b), no similar features can be

seen in Figure 9.8(a). ∞ (Sec. 7.4) There simply appears to be no large-scale plate tectonics on Venus.

Ishtar Terra (“Land of Ishtar”) lies in the southern high latitudes (at the *tops* of Figures 9.7a and 9.8a—recall our earlier discussion of Venus’s retrograde rotation). The projection used in Figure 9.8 makes Ishtar Terra appear larger than it really is—it is actually about the same size as Australia. This landmass is dominated by a great plateau known as Lakshmi Planum (Figure 9.9), some 1500 km across at its widest point and ringed by mountain ranges, including the Maxwell Montes range, which contains the highest peak on the planet, rising some 14 km above the level of Venus’s deepest surface depressions. Again for comparison, the highest point on Earth (the summit of Mount Everest) lies about 20 km above the deepest section of Earth’s ocean floor (Challenger Deep, at the bottom of the Marianas Trench on the eastern edge of the Philippines plate).

Figure 9.9(a) shows a large-scale *Venera* image of Lakshmi Planum, at a resolution of about 2 km. The “wrinkles” are actually chains of mountains, hundreds of kilometers long and tens of kilometers apart. The red area immediately to the right of the plain is Maxwell Montes. On the western (right-hand) slope of the Maxwell range lies a great crater, called Cleopatra, about 100 km across. Figure 9.9(b) shows a *Magellan* image of Cleopatra, which was originally thought to be volcanic in origin. Close-up views of the crater’s structure, however, have led planetary




▲ **FIGURE 9.8 Venus Maps** (a) Radar map of the surface of Venus, based on *Pioneer Venus* data. Color represents elevation, with white the highest areas and blue the lowest. (b) A similar map of Earth, at the same spatial resolution. (c) Another version of (a), with major surface features labeled. Compare with Figure 9.7, and notice how the projection exaggerates the size of surface features near the poles. (NASA)

scientists to conclude that the crater is meteoritic in origin, although some volcanic activity was apparently associated with its formation when the colliding body temporarily breached the planet's crust. Notice the dark (smooth) lava flow emerging from within the inner ring and cutting across the outer rim at the upper right.

It is now conventional to name features on Venus after famous women—Aphrodite, Ishtar, Cleopatra, and so on. However, the early nonfemale names (e.g., Maxwell

Montes, named after the Scottish physicist James Clerk Maxwell) predating this convention have stuck, and they are unlikely to change. Venus's other continent-sized formation, Aphrodite Terra, is located on the planet's equator and is comparable in size to Africa. Before *Magellan's* arrival, some researchers had speculated that Aphrodite Terra might have been the site of something akin to seafloor spreading at the Mid-Atlantic ridge on Earth—a region where two lithospheric plates moved






**▲ FIGURE 9.9 Ishtar Terra** (a) A *Venera* orbiter image of a plateau known as Lakshmi Planum in Ishtar Terra. The Maxwell Montes mountain range (orange) lies on the western margin of the plain, near the right-hand edge of the image. A meteor crater named Cleopatra is visible on the western slope of the Maxwell range. Note the two larger craters in the center of the plain itself. (b) A *Magellan* image of Cleopatra showing a double-ringed structure that identifies the feature to geologists as an impact crater. (NASA)

apart and molten rock rose to the surface in the gap between them, forming an extended ridge. ∞ (Sec. 7.4) With the low-resolution data then available, the issue could not be settled at the time.

The *Magellan* images now seem to rule out even this small-scale tectonic activity, and the Aphrodite region gives no indication of spreading. Figure 9.10 shows a portion of Aphrodite Terra called Ovda Regio. The crust appears buckled and fractured, with ridges running in two distinct directions across the image, suggesting that large compressive forces are distorting the crust. There seem to have been repeated periods of extensive lava flows. The dark regions are probably solidified lava flows. Some narrow lava channels, akin to rilles on the Moon, also appear. ∞ (Sec. 8.5) Such lava channels appear to be quite common on Venus. Unlike lunar rilles, however, they can be extremely long—hundreds or even thousands of kilometers. These lava “rivers” often have lava “deltas” at their mouths, where they deposited their contents into the surrounding plains.

Figure 9.11 shows a series of angular cracks in the crust, thought to have formed when lava welled up from a deep fissure, flooded the surrounding area, and then retreated below the planet’s surface. As the molten lava withdrew, the thin, new crust of solidified material collapsed under its own weight, forming the cracks we now see. Even taking into account the differences in temperature and composition between Venus’s crust and Earth’s, this terrain is not at all what we would expect at a spreading site similar to the

Mid-Atlantic Ridge. ∞ (Sec. 7.4) Although there is no evidence for plate tectonics on Venus, it is likely that the stresses in the crust that led to the large mountain ranges were caused by convective motion within Venus’s mantle—the same basic process that drives Earth’s plates. Lakshmi Planum, for example, is probably the result of a “plume” of upwelling mantle material that raised and buckled the planet’s surface.



**▲ FIGURE 9.10 Aphrodite Terra** A *Magellan* image of Ovda Regio, part of Aphrodite Terra. The intersecting ridges indicate repeated compression and buckling of the surface. The dark areas represent regions that have been flooded by lava upwelling from cracks like those shown in Figure 9.11. (NASA)



◀ **FIGURE 9.11 Lava Flows** These cracks in Venus's surface, detected by *Magellan* in another part of Aphrodite Terra, have allowed lava to reach the surface and flood the surrounding terrain. The dark regions are smooth lava flows. The network of fissures visible here is about 50 km long. (NASA)

Shield volcanoes, such as the Hawaiian Islands on Earth, are not associated with plate boundaries. Instead, they form when lava wells up through a “hot spot” in the crust and are built up over


long periods of time by successive eruptions and lava flows. A characteristic of shield volcanoes is the formation of a *caldera*, or crater, at the summit when the underlying lava withdraws and the surface collapses. The distribution of volcanoes over the surface of Venus appears random—quite different from the distribution on Earth, where volcanic activity clearly traces out plate boundaries (see Figure 7.11)—consistent with the view that plate tectonics is absent on Venus.

More volcanic features are visible in Figure 9.13, which shows a series of seven pancake-shaped **lava domes**, each about 25 km across. They probably formed when lava oozed out of the surface, formed the dome, and then withdrew,


## Volcanism and Cratering

On Earth, the principal agent of long-term, planetwide surface change is plate tectonics, driven by convection in our planet's mantle. ∞ (Secs. 7.3, 7.4) Volcanic and seismic activity are predominantly (although not exclusively) associated with plate boundaries. On Venus, without global plate tectonics, large-scale recycling of the crust by plate motion is not a factor in changing the planet's surface. Nevertheless, many areas of Venus have extensive volcanic features.

Most volcanoes on the planet are of the type known as **shield volcanoes**. Two large shield volcanoes, called Sif Mons and Gula Mons, are shown in Figure 9.12.



▲ **FIGURE 9.12 Volcanism on Venus** (a) Two larger volcanoes, known as Sif Mons (left) and Gula Mons, appear in this *Magellan* image. Color indicates height above a nominal planetary radius of 6052 km as indicated by the scale at left. In Section 9.5, we will see how past volcanism has played a crucial role in determining current conditions in Venus's atmosphere and surface. (b) This computer-generated view of Gula Mons, as seen from ground level (with colors based on data returned from Soviet landers) has a greatly exaggerated vertical scale (by about a factor of 40), so the mountain looks much taller relative to its width than it really is; Venus is actually a remarkably flat place. (NASA)



▲ **FIGURE 9.13 Lava Dome** (a) These dome-shaped structures on Venus resulted when viscous molten rock bulged out of the ground and then retreated, leaving behind a thin, solid crust that later cracked and subsided. *Magellan* found features like these in several locations on Venus. (b) This three-dimensional computer representation of four of the domes shows the view looking toward the right from near the center of the image in part (a). Colors in (b) are approximately correct, based on data returned by Soviet *Venera* landers. (NASA)

leaving the crust to crack and subside. Lava domes such as these are found in numerous locations on Venus. The largest volcanic structures on the planet are huge, roughly circular regions known as **coronae** (singular: *corona*). A large corona, called Aine, can be seen in Figure 9.14, another large-scale mosaic of *Magellan* images. Coronae are unique to Venus. They appear to have been caused by upwelling mantle material, perhaps similar to the uplift that resulted in Lakshmi Planum but on a somewhat smaller scale. They generally have volcanoes both in and around them, and closer inspection of the rims usually shows evidence for extensive lava flows into the plains below.

There is overwhelming evidence for past surface activity on Venus. Has this activity now stopped, or is it still going on? Two pieces of indirect evidence suggest that volcanism continues today. First, the level of sulfur dioxide above Venus's clouds shows large and fairly frequent fluctuations. It is quite possible that these variations result from volcanic eruptions on the surface. If so, volcanism may be the primary cause of Venus's thick cloud cover. Second, both the *Pioneer Venus* and the *Venera* orbiters observed bursts of radio energy from Aphrodite and

other regions of the planet's surface. The bursts are similar to those produced by lightning discharges that often occur in the plumes of erupting volcanoes on Earth, again suggesting ongoing activity. However, while these pieces of evidence are quite persuasive, they are still only circumstantial. No “smoking gun” (or erupting volcano) has yet been seen, so the case for active volcanism is not yet complete.



► **FIGURE 9.14 Venus Corona** This corona, called Aine, lies in the plains south of Aphrodite Terra and is about 300 km across. Coronae probably result from upwelling mantle material, causing the surface to bulge outward. Note the pancake-shaped lava domes at top, the many fractures in the crust around the corona, and the large impact craters with their surrounding white (rough) ejecta blankets that stud the region. (NASA)


The *Venus Express* spacecraft has been orbiting Venus since 2006. Its mission is currently scheduled to end in 2014. The probe does not carry instruments capable of imaging the surface, but its infrared sensors are designed to make precise temperature measurements at various levels in the atmosphere. One important mission goal is to map the planet's surface temperature with sufficient precision to detect hot spots on the surface, associated with volcanism or tectonic plumes, that are not evident in the *Magellan* maps. In 2010, mission scientists announced that three volcanic regions first identified by *Magellan* are significantly hotter than their surroundings, consistent with volcanic activity sometime during the past few million years—very recent by geological standards, although still far from the ongoing eruption many would like to see.

Not all the craters on Venus are volcanic in origin: Some, like Cleopatra (Figure 9.9b), were formed by meteoritic impact. Large impact craters on Venus are generally circular, but those less than about 15 km in diameter can be quite asymmetric in appearance. Figure 9.15(a) shows a *Magellan* image of a relatively small meteoritic impact crater, about 10 km across, in Venus's southern hemisphere. Geologists think that the light-colored region is the ejecta blanket—material ejected from the crater following the impact. The odd shape may be the result of a large meteoroid's breaking up just before impact into pieces that hit the surface near one another. Making craters such as these seems to be a fairly common fate for medium-sized bodies


(1 km or so in diameter) that plow through Venus's dense atmosphere. Figure 9.15(b) shows the largest known impact feature on Venus: the 280-km-diameter crater called Mead. Its double-ringed structure is in many ways similar to the Moon's Mare Orientale (Figure 8.14a). Numerous impact craters (identifiable by their ejecta blankets) can also be discerned in Figure 9.14.

Venus's atmosphere is sufficiently thick that small meteoroids do not reach the ground, so there are no impact craters smaller than 2–3 km across, and atmospheric effects probably also account for the relative scarcity of impact craters less than 25 km in diameter. Contrast this with Earth, where even 10-m-sized craters are formed quite frequently (every few years) by small meteoroids striking the ground. (We don't see huge numbers of such craters on Earth because they are eroded away by wind and water quite rapidly—within a few tens of thousands of years.) On average, the number of large-diameter craters on Venus's surface per square kilometer is only about one-tenth that in the lunar maria. Applying similar crater-age estimates to Venus as we do to Earth and the Moon suggests that much of the surface of Venus is quite young—less than a billion years old, and perhaps as little as 200 or 300 million years in some places, such as the region shown in Figure 9.12. [∞ \(Sec. 8.5\)](#)

Overall, the long-term degree of volcanism on Venus seems to be comparable to, but not as great as, that on Earth. However, planetary scientists think that the two planets differ in both the frequency and the severity of volcanic eruptions.




(a)



(b)

**▲ FIGURE 9.15 Impact Craters on Venus** (a) A *Magellan* image of a multiple-impact crater in Venus's southern hemisphere shows irregularly shaped, light-colored ejecta debris that was probably caused by a meteoroid that fragmented just prior to impact. The dark regions in the crater may be pools of solidified lava. (b) Venus's largest crater, named after anthropologist Margaret Mead, clearly displays a double-ringed structure. (NASA)



On Earth, near-continuous volcanic activity at plate boundaries provides a natural “release valve,” allowing energy from the interior to escape steadily through the surface in many small-scale volcanic events. On Venus, on the other hand, with no plate tectonics there is no such release mechanism, and heat from the planet’s interior tends to build up in the upper mantle. Although small-scale volcanoes may still form and erupt from time to time, most of this pent-up energy seems to be released catastrophically in planetwide volcanic eruptions every few hundred million years. Although erosion by the planet’s atmosphere may play some part in obliterating surface features, the main erosive agent on Venus is volcanism, which appears to have “resurfaced” much of the planet roughly 500 million years ago.


### Data from the Soviet Landers

The 1975 soft landings of the Soviet *Venera 9* and *Venera 10* spacecraft directly established that Venus’s surface is dry and dusty. Figure 9.16(a) shows one of the first photographs of the surface of Venus radioed back to Earth. Each craft lasted only about an hour before overheating, their electronic circuitry literally melting in this planetary oven. Typical rocks in the photo measure about 50 cm by 20 cm across—a little like flagstones on Earth. Sharp-edged and slablike, these rocks show little evidence of erosion.


*About as much sunlight penetrates Venus’s clouds as on a heavily overcast day on Earth.*



(a)



(b)



Apparently, they are quite young rocks, again supporting the idea of ongoing surface activity of some kind on Venus.

Later *Venera* missions took more detailed photographs, as shown in Figure 9.16(b). The presence of small rocks and finer material indicates the effects of erosive processes. These later missions also performed simple chemical analyses of the surface of Venus. The samples studied by *Venera 13* and *Venera 14* were predominantly basaltic in nature, again implying a volcanic past. However, not all the rocks were found to be basaltic: The *Venera 17* and *Venera 18* landers also found surface material resembling terrestrial granite, probably (as on Earth) part of the planet’s ancient crust.


### CONCEPT Check

- ✓ Are volcanoes on Venus associated mainly with the movement of tectonic plates, as on Earth?

## 9.5 The Atmosphere of Venus

Measurements made by the *Venera* and *Pioneer Venus* spacecraft have allowed astronomers to paint a fairly detailed picture of Venus’s atmosphere. ∞ (Sec. 6.6) The planet’s hot, dense, carbon dioxide atmosphere contrasts sharply with that of Earth, even though, as we will see, the two may have had comparable beginnings. ∞ (Discovery 6-2)

◀ **FIGURE 9.16 Venus In Situ** (a) The first direct view of the surface of Venus, radioed back to Earth from the Soviet *Venera 9* spacecraft, which made a soft landing on the planet in 1975. (b) Another view of Venus, in true color, from *Venera 14*. Flat rocks like those visible in part (a) are seen among many smaller rocks and even fine soil on the surface. The peculiar filtering effects of whatever light does penetrate the clouds make Venus’s air and ground appear peach colored—in reality, they are most likely gray, like rocks on Earth. (Russian Space Agency)



▲ **FIGURE 9.17 Venus's Atmosphere** The structure of the atmosphere of Venus, as determined by U.S. and Soviet probes. (One bar is the atmospheric pressure at sea level on Earth.) The planet's hot, dense atmosphere contrasts greatly with that of Earth, giving us important insights into the evolution of both.


## Atmospheric Structure

Figure 9.17 shows the variation of temperature and pressure with height. (Compare this figure with Figure 7.2, which gives similar information for Earth). The atmosphere

of Venus is about 90 times more massive than Earth's, and it extends to a much greater height above the surface. On Earth, 90 percent of the atmosphere lies within about 10 km of sea level. On Venus, the 90 percent level is found at an altitude of 50 km instead. The surface temperature and pressure of Venus's atmosphere are much greater than Earth's. However, the temperature drops more rapidly with altitude, and the upper atmosphere of Venus is actually colder than our own (the coldest measured so far, by *Venus Express* in 2012, was 100 K at 125 km altitude).

Venus's troposphere extends up to an altitude of nearly 100 km. The reflective clouds that block our view of the surface lie between 50 and 70 km above the surface. Data from the *Pioneer Venus* multiprobe indicate that the clouds may actually be separated into three distinct layers within that altitude range. Below the clouds, extending down to an altitude of some 30 km, is a layer of haze. Below 30 km, the air is clear. Above the clouds, a high-speed "jet stream" blows from west to east at about 300–400 km/h, fastest at the equator and slowest at the poles. This high-altitude flow is responsible for the rapidly moving cloud patterns seen in ultraviolet light.

Figure 9.18 shows a sequence of three ultraviolet images of Venus in which the variations in the cloud patterns can be seen. Note the characteristic V-shaped appearance of the clouds—a consequence of the fact that, despite their slightly lower speeds, the winds near the poles have a shorter distance to travel in circling the planet and so are always forging ahead of winds at the equator. Near the surface, the dense atmosphere moves more sluggishly—indeed, the fluid flow bears more resemblance to that in Earth's oceans than to the flow in Earth's air. Surface wind speeds on Venus are typically less than 2 m/s (roughly 4 mph).





▲ **FIGURE 9.18 Atmospheric Circulation** Three ultraviolet views of Venus, taken by the *Pioneer Venus* orbiter at about 10-hour intervals, shows the changing cloud patterns in the planet's upper atmosphere. The wind flow is from right to left (or clockwise from above), in the direction opposite the sideways "V" in the clouds. Notice the motion of the dark region marked by the blue arrow. Venus's retrograde rotation means that north is at the bottom of these images and west to the right. (NASA)

One of *Venus Express*'s main missions is to study atmospheric circulation on Venus. In 2006 the orbiter returned a series of intriguing images of the planet's south pole, showing a **polar vortex** of swirling winds there (Figure 9.19). Polar vortices are well known to atmospheric scientists. Although they may look like giant hurricanes, they are not storms in the usual sense. They are relatively stable, long-lived flows circling the polar regions. They are expected in any body (planet or moon) with an atmosphere, although the details depend on the properties of the atmosphere and the body's rotation rate. Earth's south polar vortex, for example, plays an important role in confining and concentrating the gases responsible for our planet's antarctic ozone hole. ∞ (Sec. 7.2)



NASA's *Pioneer Venus* orbiter discovered the planet's north polar vortex in 1978, and when *Venus Express* reached Venus in 2006, the search for the southern vortex was one of its top priorities. The Venus vortices present a puzzle to scientists because of their peculiar “double-lobed” structure (see right side of Figure 9.19), which is unique to Venus. This structure is not well understood. By making repeated observations of the southern vortex and watching how it changes in time, scientists hope to understand the forces driving it and gain clues to the global circulation of Venus's atmosphere.

structure (see right side of Figure 9.19), which is unique to Venus. This structure is not well understood. By making repeated observations of the southern vortex and watching how it changes in time, scientists hope to understand the forces driving it and gain clues to the global circulation of Venus's atmosphere.



▲ **FIGURE 9.19**

**Venus Polar Vortex** The visible image at left shows Venus during the day when sunlight reflects from its cloud tops. By contrast, the false-color insets at right are nighttime views of infrared radiation arising from deeper layers near Venus's south pole, emphasizing the dynamic swirls and vortices of its lower-atmospheric cloud structures. These images of the southern vortex were taken a few hours apart by Europe's *Venus Express* spacecraft. (ESA)



## Atmospheric Composition

Carbon dioxide ( $\text{CO}_2$ ) is the dominant component of Venus's atmosphere, accounting for 96.5 percent of it by volume. Almost all of the remaining 3.5 percent is nitrogen ( $\text{N}_2$ ). Trace amounts of other gases, such as water vapor, carbon monoxide, sulfur dioxide, and argon, are also present. This composition is clearly radically different from that of Earth's atmosphere. The absence of oxygen is perhaps not surprising, given the absence of life. (Recall our discussion of Earth's atmosphere in Chapter 7.) ∞ (Sec. 7.2) However, there is no sign of the large amount of water vapor we would expect to find if a volume of water equivalent to Earth's oceans had evaporated and remained in the planet's atmosphere. If Venus started off with an Earth-like composition, then something has happened to its water—Venus is now a very dry planet.

For a long time, the chemical makeup of the reflective cloud layer surrounding Venus was unknown. At first, scientists assumed that the clouds were water vapor or ice, as on Earth, but the reflectivity of the clouds at different wavelengths didn't match that of water ice. Later infrared observations carried out in the 1970s showed that the clouds (or at least the top layer of clouds) are actually composed of *sulfuric acid*, created by reactions between water and sulfur dioxide. Sulfur dioxide is an excellent absorber of ultraviolet radiation and could be responsible for many of the cloud patterns seen in ultraviolet light. Spacecraft observations confirmed the presence of all three compounds in the atmosphere and also indicated that there may be particles of sulfur suspended in and near the cloud layers, which may account for Venus's characteristic yellowish hue.


## The Greenhouse Effect on Venus

Given the distance of Venus from the Sun, the planet was not expected to be such a pressure cooker. As mentioned earlier, calculations based on Venus's orbit and reflectivity indicated a temperature not much different from Earth's, and early measurements of the cloud temperatures seemed to concur. Certainly, scientists reasoned, Venus could be no hotter than the

sunward side of Mercury, and it should probably be much cooler. This reasoning was obviously seriously in error.

Why is Venus's atmosphere so hot? And if, as we think, Venus started off like Earth, why is it now so different? The answer to the first question is fairly easy: Given the present composition of its atmosphere, Venus is hot because of the

Arrow thickness is proportional to the amount of radiation moving in and out.



**▲ FIGURE 9.20 Greenhouse Effect on Earth and Venus** Because Venus's atmosphere is much deeper and denser than Earth's, a much smaller fraction of the infrared radiation leaving the planet's surface escapes into space. The result is a much stronger greenhouse effect than on Earth and a correspondingly hotter planet. The reasons for this critical difference between the two planets are of vital importance for life on Earth.

greenhouse effect. Recall from our discussion in Chapter 7 that “greenhouse gases” in Earth's atmosphere—particularly water vapor and carbon dioxide—serve to trap heat from the Sun. ∞ (Sec. 7.2) By inhibiting the escape of infrared radiation reradiated from Earth's surface, these gases increase the planet's equilibrium temperature, in much the same way as an extra blanket keeps you warm on a cold night. Continuing the analogy a little further, the more blankets you place on the bed, the warmer you will become. Similarly, the more greenhouse gases there are in the atmosphere, the hotter the surface will be.

The same effect occurs naturally on Venus, whose dense atmosphere is made up almost entirely of a primary greenhouse gas, carbon dioxide. As illustrated schematically in Figure 9.20, the thick carbon dioxide blanket absorbs nearly 99 percent of all the infrared radiation released from the surface of Venus and is the immediate cause of the planet's sweltering 730 K surface temperature. Furthermore, the temperature is nearly as high at the poles as at the equator, and there is not much difference between the temperatures on the day and night sides. The circulation of the atmosphere spreads energy efficiently around the planet, making it impossible to escape the blazing heat, even during the planet's 2-month-long night.

## The Runaway Greenhouse Effect

But *why* is Venus's atmosphere so different from Earth's? Let's assume that the two planets started off with basically similar compositions. Why, then, is there so much carbon dioxide in the atmosphere of Venus, and why is the planet's atmosphere so dense? To address these questions, we must consider the processes that created the atmospheres of the terrestrial planets and then determined their evolution. In fact, we can turn the question around and ask instead, “Why is there so *little* carbon dioxide in Earth's atmosphere compared with that of Venus?”

Earth's atmosphere has evolved greatly since it first appeared. Our planet's secondary atmosphere was outgassed from the interior by volcanic activity 4 billion years ago. ∞ (Sec. 7.2) Since then, it has been reprocessed, in part by living organisms, into its present form. On Venus, the initial stages probably took place in more or less the same way, so that at some time in the past, Venus might well have had an atmosphere similar to the primitive secondary atmosphere on Earth, containing water, carbon dioxide, sulfur dioxide, and nitrogen-rich compounds. What happened on Venus to cause such a major divergence from subsequent events on our own planet?




On Earth, sunlight split the nitrogen-rich compounds, releasing nitrogen into the air. Meanwhile, the water condensed into oceans, and much of the carbon dioxide and sulfur dioxide eventually became dissolved in them. Most of the remaining carbon dioxide combined with surface rocks. Thus, much of the secondary outgassed atmosphere quickly became part of the surface of the planet. If all the dissolved or chemically combined carbon dioxide were released back into Earth's present-day atmosphere, its new composition would be 98 percent carbon dioxide and 2 percent nitrogen, and it would have a pressure about 70 times its current value. In other words, apart from the presence of oxygen (which appeared on Earth only after the development of life) and water (the absence of which on Venus will be explained shortly), Earth's atmosphere would be a lot like that of Venus! The real difference between Earth and Venus, then, is that Venus's greenhouse gases never left the atmosphere the way they did on Earth.

When Venus's secondary atmosphere appeared, the temperature was higher than on Earth, simply because Venus is closer to the Sun. However, the Sun was probably somewhat dimmer then (see Chapter 22)—perhaps only half its present brightness—so there is some uncertainty as to exactly how much hotter than Earth Venus actually was. If the temperature was already so high that no oceans condensed, the outgassed water vapor and carbon dioxide would have remained in the atmosphere, and the full greenhouse effect would have gone into operation immediately. If (as now seems more likely) oceans did form and most of the greenhouse gases left the atmosphere, as they did on Earth,\* the temperature must still have been sufficiently high that a process known as the **runaway greenhouse effect** came into play.

To understand the runaway greenhouse effect, imagine that we took Earth from its present orbit and placed it in Venus's orbit, some 30 percent closer to the Sun. At that distance from the Sun, the amount of sunlight striking Earth's surface would be about twice its present level, so the planet would warm up. More water would evaporate from the oceans, leading to an increase in atmospheric water vapor. At the same time, the ability of both the oceans and surface rocks to hold carbon dioxide would diminish, allowing more carbon dioxide to enter the atmosphere. As a result, greenhouse heating would increase, and the planet would warm still further, leading to a further increase in atmospheric greenhouse gases, and so on. Once started, the process would “run away,” eventually leading to the complete evaporation of the oceans, restoring all the original greenhouse gases to the atmosphere. Although the details are quite complex, basically the same thing would have happened on Venus long ago, ultimately resulting in the planetary inferno we see today.

\*In fact, careful study of the Magellan images reveals no sign of ancient seashores or ocean basins, nor evidence of erosion by rivers on Venus. However, it is unclear whether such features would have survived the heavy volcanism known to have occurred in the planet's more recent past.

The presence of atmospheric water vapor meant that the greenhouse effect on Venus was even more extreme in the past. By intensifying the blanketing effect of the carbon dioxide, the water vapor helped the surface of Venus reach temperatures perhaps twice as hot as current temperatures. At the high temperatures of the past, the water vapor was able to rise high into the planet's upper atmosphere—so high that it was broken up by solar ultraviolet radiation into its components, hydrogen and oxygen. The light hydrogen rapidly escaped, the reactive oxygen quickly combined with other atmospheric gases, and virtually all of the water on Venus was lost forever. This is the reason that Venus lacks water today. The tail end of this process is still ongoing. The *Venus Express* orbiter has detected minuscule quantities of hydrogen and oxygen escaping into space, consistent with the trace amount of water remaining in the atmosphere today.


Although it is highly unlikely that global warming will ever send Earth down the path taken by Venus, this episode highlights the relative fragility of the planetary environment.  (*Discovery 7-1*) No one knows how close to the Sun Earth could have formed before a runaway greenhouse effect would have occurred. But in comparing our planet with Venus, we have come to understand that there is an orbital limit, presumably between 0.7 and 1.0 AU, inside of which Earth would have suffered a similar catastrophic runaway. We must consider this “greenhouse limit” when we assess the likelihood that planets harboring life formed elsewhere in our Galaxy (see Chapter 28).

### CONCEPT Check

✓ If Venus had formed at Earth's distance from the Sun, what might its climate be like today?

## 9.6 Venus's Magnetic Field and Internal Structure

In 1962, *Mariner 2* flew by Venus, carrying, among other instruments, magnetometers to measure the strength of the planet's magnetic field. None was detected, and subsequent Soviet and U.S. missions, carrying more sensitive detectors, have confirmed this finding. *Pioneer Venus* did detect a weak “induced” magnetic field produced by the interaction between the planet's upper atmosphere and the solar wind, but Venus apparently has no intrinsic magnetic field of its own.

Venus, with an average density similar to that of Earth, probably has a similar overall composition and a partially molten iron-rich core. The lack of any significant magnetic field on Venus, then, is almost surely the result of the planet's extremely slow rotation and consequent lack of dynamo action.  (*Sec. 7.5*) Having no magnetosphere, Venus has no protection from the solar wind. The planet's upper atmosphere is continually bombarded by high-energy particles from the Sun, keeping the topmost layers permanently

ionized. However, the great thickness of the atmosphere prevents any of these particles from reaching the surface.

None of the *Venera* landers carried seismic equipment, so no direct measurements of the planet's interior have been made, and theoretical models of the interior have little hard data to constrain them. However, to many geologists, the surface of Venus resembles that of the young Earth, at an age of perhaps a billion years. At that time, volcanic activity had begun, but the crust was still relatively thin and the convective processes in the mantle that drive plate tectonic motion were not yet fully established. Measurements of the planet's gravitational field suggest that Venus lacks an asthenosphere, the semisolid part of the upper mantle over which Earth's lithosphere slides. ∞ (Sec. 7.4)

Why has Venus remained in that immature state and not developed plate tectonics as Earth did? That question remains to be answered. Some planetary geologists have

speculated that the high surface temperature has inhibited Venus's evolution by slowing the planet's cooling. Possibly the high surface temperature has made the crust too thin, or the mantle too fluid, for Earth-style plate motion to develop. Or perhaps the high temperature and thin crust have led to more volcanism, tapping the energy that might otherwise go into convective motion. It may also be that the presence of water plays an important role in lubricating convection in the mantle and plate motion, so that arid Venus could never have evolved along the same path as Earth.

### CONCEPT Check

- ✓ If the interior of Venus is quite Earth-like, and Venus has a molten iron core, why doesn't the planet have a magnetic field as Earth does?

**The Big Question** The biggest question confronting humans living on a globally warming Earth is obvious: Could Earth someday heat up as much as Venus? The fact that Venus is just little closer to the Sun than is Earth probably made all the difference. Venus's warmer water escaped early in its history and thus could not help remove CO<sub>2</sub> from the atmosphere by trapping it in rocks and oceans, as occurs on Earth even today; Earth's cooler water remained and reduced this famous greenhouse gas to trace amounts. Even so, we might wonder: How much CO<sub>2</sub> dare we humans release into our own air? Are we placing our planet on a path to become like Venus?


## Chapter Review

### SUMMARY

1 The interior orbit of Venus with respect to Earth's means that Venus never strays far from the Sun in the sky. Because of its highly reflective cloud cover, Venus is brighter than any star in the sky, as seen from Earth. The planet's mass and radius are similar to those of Earth. The planet's rotation is slow and retrograde, most likely because of a collision between Venus and some other solar system body during the late stages of the planet's formation.




2 The extremely thick atmosphere of Venus is nearly opaque to visible radiation, making the planet's surface invisible at optical wavelengths from the outside. Venus's atmosphere is nearly 100 times denser than Earth's and consists mainly of carbon dioxide. The temperature of the upper atmosphere is much like that of Earth's upper atmosphere, but the surface




temperature of Venus is a searing 730 K. The planet's high-level winds circulate rapidly around the planet, and there are peculiarly shaped **polar vortices** (p. 230) at both poles.

3 Venus's surface has been thoroughly mapped by radar from Earth-based radio telescopes and orbiting satellites. The planet's surface is mostly smooth, resembling rolling plains with modest highlands and lowlands. Two elevated continent-sized regions are called Ishtar Terra and Aphrodite Terra. There is no evidence for plate tectonic activity as on Earth. Features called **coronae** (p. 226) are thought to have been caused by an upwelling of mantle material. For unknown reasons, the upwelling never developed into full convective motion. The surface of the planet appears to be relatively young, resurfaced by volcanism within the past few hundred million years. Many **lava domes** (p. 225) and **shield volcanoes** (p. 225) were found by the *Magellan* orbiter on Venus's surface, but none of the volcanoes has yet proved to be currently active.




**4** Some craters on Venus are due to meteoritic impact, but the majority appear to be volcanic in origin. The evidence for currently active volcanoes on Venus includes surface features resembling those produced in Earthly volcanism, fluctuating levels of sulfur dioxide in Venus's atmosphere, and bursts of radio energy similar to those produced by lightning discharges that often occur in the plumes of erupting volcanoes on Earth. However, no actual eruptions have been seen. Soviet spacecraft that landed on Venus photographed surface rocks with sharp edges and a slablike character. Some rocks appear predominantly basaltic in nature, implying a volcanic past, others resemble terrestrial granite and are probably part of the planet's ancient crust.



**5** Although Venus and Earth may have started off with fairly similar surface conditions, their atmospheres are now very different. The total mass of Venus's atmosphere is about 90 times greater

than Earth's. The greenhouse effect caused by the large amount of carbon dioxide in Venus's atmosphere is the cause of the planet's current high temperatures. Almost all the water vapor and carbon dioxide initially present in Earth's early atmosphere quickly became part of the oceans or surface rocks. Because Venus orbits closer to the Sun than does Earth, surface temperatures on Venus were initially higher, and the planet's greenhouse gases never left the atmosphere. The **runaway greenhouse effect** (p. 232) caused all the planet's greenhouse gases—carbon dioxide and water vapor—to end up in the atmosphere, leading to the extreme conditions we observe today.



**6** Venus has no detectable magnetic field, almost certainly because the planet's rotation is too slow for any appreciable dynamo effect to have developed. To some planetary geologists, Venus's interior structure suggests that of the young Earth, before convection became established in the mantle.

**MasteringAstronomy**® For instructor-assigned homework go to [MasteringAstronomy.com](https://www.masteringastronomy.com).

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- Why does Venus appear so bright to the eye? Upon what factors does the planet's brightness depend?
- LO1** Explain why Venus is always found in the same general part of the sky as the Sun.
- POS** What is our current best explanation of Venus's slow, retrograde spin?
- POS** How did radio observations of Venus made in the 1950s change our conception of the planet?
- What did ultraviolet images returned by *Pioneer Venus* show about the planet's high-level clouds?
- LO2** Name three ways in which the atmosphere of Venus differs from that of Earth.
- What are the main constituents of Venus's atmosphere? What are clouds in the upper atmosphere made of?
- LOS** What component of Venus's atmosphere causes the planet to be so hot? Explain why there is so much of this gas in the atmosphere of Venus, compared with its presence in Earth's atmosphere. What happened to all the water that Venus must have had when the planet formed?
- If Venus had formed at Earth's distance from the Sun, what do you imagine its climate would be like today? Why do you think so?
- LO3** How do the "continents" of Venus differ from Earth's continents?
- How are the impact craters of Venus different from those found on other bodies in the solar system?
- LO4 POS** What evidence exists that volcanism of various types has changed the surface of Venus?
- What is the evidence for active volcanoes on Venus?
- LO6** Given that Venus, like Earth, probably has a partially molten iron-rich core, why doesn't Venus also have a magnetic field?
- Do you think that Earth is in any danger of being subject to a runaway greenhouse effect like that on Venus?

## Conceptual Self-Test: Multiple Choice

- VIS** According to Figure 9.2 ("Venus's Brightness"), Venus is never seen at midnight because (a) it is closer to the Sun than is Earth; (b) it will be in its new phase then; (c) it is visible only at sunset; (d) it will be at superior conjunction.
- Venus's permanent retrograde rotation about its axis results in the planet (a) always rising in the western sky; (b) orbiting the Sun in the opposite direction from Earth; (c) having its north pole below the plane of the ecliptic; (d) being brighter than any other planet.

3. Compared with Earth, Venus is (a) much smaller; (b) much larger; (c) about the same size.
4. Venus's surface is permanently obscured by clouds. As a result, the surface has been studied primarily by (a) robotic landers (b) orbiting satellites using radar; (c) spectroscopy; (d) radar signals from Earth.
5. Compared with Earth, Venus has a level of plate tectonic activity that is (a) much more rapid; (b) virtually nonexistent; (c) about the same.
6. Venus's atmosphere (a) has almost the same chemical composition as Earth's; (b) shows very high levels of humidity; (c) is composed mostly of carbon dioxide; (d) is predominantly made of acid droplets.
7. Compared with Earth's atmosphere, most of Venus's atmosphere is (a) compressed much closer to the surface; (b) spread out much farther from the surface; (c) similar in extent and structure.
8. **VIS** According to Figure 9.17 ("Venus's Atmosphere"), Venus's atmospheric temperature is (a) about the same as Earth's; (b) cooler than temperatures on the planet Mercury; (c) hotter than temperatures on Mercury; (d) high due to the presence of sulfuric acid.
9. Carbon dioxide on Venus (a) is all in the atmosphere; (b) was absorbed in surface water and has evaporated into space; (c) has dissolved in the atmospheric acid; (d) is integrated into the surface rocks.
10. Venus lacks a planetary magnetic field because (a) it rotates very slowly; (b) it does not have a molten core; (c) there are no plate tectonics on the planet; (d) the core contains little or no iron.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

1. ●● Using the data given in the text, calculate Venus's angular diameter, as seen by an observer on Earth, when the planet is (a) at its brightest, (b) at greatest elongation, and (c) at the most distant point in its orbit.
2. ●● Seen from Earth, through how many degrees per night (relative to the stars) does Venus move around the time of inferior conjunction (closest approach to Earth)?
3. ● How long does a radar signal take to travel from Earth to Venus and back when Venus is brightest? Compare this time with the round-trip time when Venus is at its closest point to Earth.
4. ● What is the size of the smallest feature that could be distinguished on the surface of Venus (at closest approach) by the Arecibo radio telescope at an angular resolution of 1"?
5. ● Could an infrared telescope with an angular resolution of 0.1" distinguish impact craters on the surface of Venus?
6. ● Approximating Venus's atmosphere as a layer of gas 50 km thick, with uniform density  $21 \text{ kg/m}^3$ , calculate the total mass of the atmosphere. Compare your answer with the mass of Earth's atmosphere (Chapter 7, Problem 1) and with the mass of Venus.
7. ● *Pioneer Venus* observed high-level clouds moving around Venus's equator in 4 days. What was their speed in km/h? In mph?
8. ● According to Stefan's law (see Section 3.4), how much more radiation—per square meter, say—is emitted by Venus's surface at 730 K than is emitted by Earth's surface at 300 K?

## Activities

### Collaborative

1. In 1994, NASA elected to turn off the *Magellan Orbiter*, even though it was working perfectly and still sending back valuable data. Does your group agree or disagree with NASA's decision? Explain your reasoning.

### Individual

1. Consult an almanac (or look online) to determine the next time Venus will pass between Earth and Sun. How many days before and after this event can you glimpse the planet with the eye alone?
2. Find out when Venus will next pass between Earth and the Sun. How many days before and after this event can you glimpse the planet with the naked eye? Using binoculars or a small telescope, examine Venus as it goes through its phases. Note the phase and the relative size (you can compare its size to the field of view in a telescope; always use the same eyepiece). Observe it every few days or once a week. Make a table of its shape, size, and relative brightness. Can you see the correlations between these properties first recognized by Galileo?



# Mars

## A NEAR MISS FOR LIFE?

Named by the ancient Romans for their bloody god of war, Mars is for many people the most intriguing of all celestial objects. Over the years, it has inspired speculation that life—perhaps intelligent and possibly hostile—may exist there. With the dawn of the Space Age, those notions had to be abandoned. Visits by robot spacecraft have revealed no signs of life of any sort on Mars, even at the microbial level.

Still, the planet’s properties are close enough to those of Earth that Mars is even now widely regarded as the second most hospitable environment for the appearance of life in the solar system, after Earth itself. At about the same time as Earth’s “twin,” Venus, was evolving into a searing inferno, the Mars of long ago may have had running water and blue skies. If life ever arose there, however, it must be long extinct. The Mars of today appears to be a dry, dead world.

**The Big Picture** Astronomers are a unique breed, in many ways commissioned by the public to keep an eye on the universe. And they have kept their eyes on Mars for a long time. Mars has probably been reconnoitered by humans more than any other cosmic object beyond the Earth and Moon. This alien world intrigues us, frustrates us, and even invites us to visit. Although Mars today seems as dry as any desert on Earth, many scientists think it was much wetter billions of years ago, when the Martian atmosphere was thicker and the climate was warmer. It does seem likely that some day, humans will colonize this planet, even “terraform” it, thus creating a second “Earth” for our species.

**LEFT:** This true-color mosaic of many images was made in 2013 by the *Curiosity* spacecraft that soft-landed in Mars’s equatorial region. Amid scattered rocks, which hold “memories” of the ancient events that formed them, this view shows wind-swept areas inside Gale crater, which was once probably filled with water and where subsurface soil and ice were scooped up and tested in an onboard mini-chemistry lab. (NASA/JPL)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Summarize the general orbital and physical properties of Mars.
- 2 Describe the observational evidence for seasonal changes on Mars.
- 3 Compare the surface features and geology of Mars with those of the Moon and Earth, and account for these characteristics in terms of Martian history.
- 4 Present the evidence that Mars once had a much denser atmosphere and running water on its surface.
- 5 Explain where that ancient water on Mars may be found today.
- 6 Compare the atmosphere of Mars with those of Earth and Venus, and explain why the evolutionary histories of these three worlds diverged so sharply.
- 7 Outline what is known of the internal structure of Mars.
- 8 Describe the characteristics of the Martian moons, and explain their probable origin.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.


## 10.1 Orbital Properties

Mars is the fourth planet from the Sun and the outermost of the four terrestrial worlds in the solar system. It lies outside Earth's orbit, as illustrated in Figure 10.1(a), which shows the orbits of both planets drawn to scale. Because of its superior (exterior) orbit, Mars ranges in our sky from a position that appears close to the Sun (*conjunction*, with Earth and Mars at points A in the figure) to one on the opposite side of the sky from the Sun (*opposition*, at points B). ∞ (Sec. 2.2) Contrast this orbit with Mercury's and Venus's inferior orbits, which ensure that we never see those planets far from the Sun in the nighttime sky. ∞ (Secs. 8.1, 9.1) From our earthly viewpoint, Mars appears to traverse a great circle in the sky, keeping close to the ecliptic and occasionally executing retrograde loops. ∞ (Sec. 2.2)

Mars's orbital eccentricity is 0.093, much larger than that of most other planets—only the innermost planet, Mercury, has a more elongated orbit. Because of this relatively large eccentricity, Mars's perihelion distance from the Sun—1.38 AU (207 million km)—is substantially smaller than its aphelion distance—1.67 AU (249 million km)—resulting in a large variation in the amount of sunlight striking the planet over the course of its year. In fact, the intensity of sunlight on the Martian surface is almost 45 percent greater when the planet is at perihelion than when it is at aphelion. As we will see, this has a substantial effect on the Martian climate.

Mars is largest and brightest in the night sky at opposition, when Earth lies between Mars and the Sun (location B in Figure 10.1a). If this happens to occur near Martian perihelion, the two planets can come as close as 0.37 AU (56 million km). The separation is less than 0.38 AU because Earth's orbit is slightly eccentric and our planet actually lies about 1.01 AU from the Sun when such an opposition occurs. (As indicated in Figure 10.1, Earth reaches aphelion in early July, whereas opposition at Martian perihelion happens in late August.) The angular size of Mars under these most favorable circumstances is about  $25''$ . Ground-based observations of the planet at those times can distinguish surface features as small as 100 km across—about the same resolution as the unaided human eye can achieve when viewing the Moon. Notice that, at this resolution, the extensive “canal systems” that once fueled such rampant speculation about life on Mars (see the Part 2 Opener on p. 134) could not possibly actually have been observed. They were strictly a figment of the human imagination.

Figure 10.1(b), shows the dates and configurations of eight successive Martian oppositions between January 1993 and December 2007. Oppositions occur at roughly 780-day intervals with corrections for the fact that, in accordance with Kepler's second law, the planets do not move at constant speeds around their orbits. ∞ (Sec. 2.5, *More Precisely 9-1*) Oppositions near Martian perihelion are less frequent,



▲ **FIGURE 10.1 Mars Orbit** (a) The orbit of Mars compared with that of Earth. Note that Mars's orbit is noticeably off center, unlike Earth's, whose eccentricity is barely perceptible here. When the planets are on opposite sides of the Sun, as at the points marked A, Mars is said to be at conjunction. The planets are at their closest at opposition, when Earth and Mars are aligned and on the same side of the Sun, as at the points marked B. (b) Several oppositions of Mars, including the particularly favorable (close) configuration of August 2003 and the unfavorable oppositions of February 1995 and March 1997.

occurring roughly once every 15 years. The most recent such event, on August 28, 2003, was one of the closest ever, with Mars just 0.373 AU from Earth, affording unprecedented observing conditions for amateurs and professionals alike. On average, the two planets come within 0.38 AU of one another only about three times per century.

Although Mars is quite bright and easily seen at opposition, the planet is still considerably fainter than Venus. This faintness results from a combination of three factors. First, Mars is more than twice as far from the Sun as is Venus, so each square meter on the Martian surface receives less than one-quarter the amount of sunlight that strikes each square meter on Venus. Second, the surface area of Mars is only about 30 percent that of Venus, so there are fewer square meters to intercept the sunlight. Finally, Mars is much less reflective than Venus—only about 15 percent of the sunlight striking the planet is reflected back into space, compared with nearly 70 percent in the case of Venus. Still, at its brightest, Mars is brighter than any star. Its characteristic red color, visible even to the naked eye, makes the planet easily identifiable in the night sky.

### CONCEPT Check

✓ Why do the closest views of Mars from Earth occur roughly only once every 15 years?

## 10.2 Physical Properties

As with Mercury and Venus, we can determine the radius of Mars by means of simple geometry. From the data given earlier for the planet's size and distance, we obtain a radius of about 3400 km. More accurate measurements give a result of 3394 km, or 0.53 Earth radii.

Unlike Mercury and Venus, Mars has two small moons in orbit around it, both visible (through telescopes) from Earth. Named Phobos (Fear) and Deimos (Panic) for the sons of Ares (the Greek name for the war god known to the Romans as Mars) and Aphrodite (the Greek name for Venus, goddess of love), these moons are little more than large rocks trapped by the planet's gravity. We will return to their individual properties at the end of the chapter. The larger of the two, Phobos, orbits at a distance of just 9378 km from the center of the planet once every 459 minutes. Applying the modified version of Kepler's third law (which states that the square of a moon's orbital period is proportional to the cube of its orbital semimajor axis divided by the mass of the planet it orbits), we find that the mass of Mars is  $6.4 \times 10^{23}$  kg, or 0.11 times that of Earth. ∞ (Sec. 2.8) Naturally, the orbit of Deimos yields the same result.

From the planet's mass and radius, it follows that the average density of Mars is  $3900 \text{ kg/m}^3$ , only slightly greater

than that of the Moon. If we assume that Martian surface rocks are similar to those on the other terrestrial planets, this average density suggests the existence of a substantial higher density core within the planet. Planetary scientists suspect that this core is composed largely of iron sulfide (a compound about twice as dense as Martian surface rock) and has a diameter of about 2500 km.

Surface markings easily seen on Mars allow astronomers to track the planet's rotation. Mars rotates once on its axis every 24.6 hours. One Martian day is thus similar in length to one Earth day. The planet's equator is inclined to the orbital plane at an angle of  $24.0^\circ$ , again similar to Earth's inclination of  $23.5^\circ$ . Thus, as Mars orbits the Sun, we find both daily and seasonal cycles, just as on Earth. In the case of Mars, however, the seasons are complicated somewhat by variations in solar heating due to the planet's eccentric orbit—southern summer occurs around the time of Martian perihelion and so is significantly warmer than summer in the north.


## 10.3 Long-Distance Observations of Mars

At opposition, when Mars is closest to us and most easily observed, we see it as full, so the Sun's light strikes the surface almost vertically, casting few shadows and preventing us from seeing any topographic detail, such as craters or mountains. Even through a large telescope, Mars appears only as a reddish disk, with some light and dark patches and prominent polar caps. These surface features undergo slow seasonal changes over the course of a Martian year. We saw in Chapter 1 how the inclination of Earth's axis produces similar seasonal changes. ∞ (Sec. 1.4) Figure 10.2 shows some of the best images of Mars ever made from Earth or Earth orbit, along with a photograph taken by one of the U.S. *Viking* spacecraft en route to the planet.

When the planet is viewed from Earth, the most obvious Martian surface features are the bright polar caps (see Figure 10.2a), growing and diminishing according to the seasons and almost disappearing at the time of the Martian summer. The dark surface features on Mars also change from season to season, although their variability probably has little to do with the melting of the polar ice caps. To the more fanciful observers around the start of the 20th century, these changes suggested the seasonal growth of vegetation on the planet. It was but a small step from seeing polar ice caps and speculating about teeming vegetation to imagining a planet harboring intelligent life, perhaps not unlike us.

But those speculations and imaginings were not to be confirmed. Spectroscopic observations from Earth and from Earth orbit revealed that the changing caps are mostly





**▲ FIGURE 10.2 Mars** (a) A deep-red (800-nm) image of Mars, taken in 1991 at Pic-du-Midi, an exceptionally clear site in the French Alps. One of the planet's polar caps appears at the top and a few other surface markings are evident in this ground-based telescopic view. (b) A visible-light *Hubble* image of Mars, taken from orbit while the planet was near opposition in 2003. (c) A view of Mars taken from a *Viking* spacecraft while approaching the planet in 1976. (CNRS; NASA)

frozen carbon dioxide (i.e., dry ice), not water ice, as at Earth's North and South Poles. ∞ (Sec. 4.4) The polar caps do contain water, but it remains permanently frozen, and the dark markings seen in Figure 10.2, once thought (by some) to be part of a network of canals dug by Martians for irrigation purposes, are actually highly cratered and eroded areas around which surface dust occasionally blows. From a distance, the repeated covering and uncovering of these landmarks gives the impression of surface variability, but it's only the thin dust cover that changes.

The powdery Martian surface dust is borne aloft by strong winds that often reach hurricane proportions (hundreds of kilometers per hour). In fact, when the U.S. *Mariner 9* spacecraft went into orbit around Mars in 1971, a planetwide dust storm obscured the entire landscape. Had the craft been on a flyby mission (for a quick look) instead of an orbiting mission (for a longer view), its visit would have been a failure. Fortunately, the storm subsided, enabling the craft to radio home detailed information about the planet's surface.

### CONCEPT Check

✓ Does Mars have seasons like those on Earth?

## 10.4 The Martian Surface

Maps of the surface of Mars returned by orbiting spacecraft show a wide range of geological features. Mars has huge volcanoes, deep canyons, vast dune fields, and many other geological wonders. Orbiters have performed large-scale surveys of much of the planet's surface, and lander data

have complemented these planetwide studies with detailed information on (so far) six specific sites. ∞ (Sec. 6.6) The current focus of Martian exploration, both from space and on the ground, is the ongoing search for water on or below the planet's surface. ∞ (Discovery 6-2)

### Large-Scale Topography



Figure 10.3 shows a planetwide mosaic of thousands of images taken in the 1970s by the *Viking* orbiters. The images show some of the planet's topographic features in true color. More recently, *Mars Global Surveyor* has mapped out the Martian surface to an accuracy of a few meters, using an instrument called a *laser altimeter*, which analyzes pulses of laser light to measure the distance between the spacecraft and the planet's surface. Figure 10.4 shows a Martian map based on those measurements, with spacecraft landing sites and some prominent surface features marked. Color indicates altitude, with blue representing the lowest-lying regions, white the highest.

A striking feature of the terrain of Mars, easily seen in Figure 10.4, is the marked difference between the planet's northern and southern hemispheres. The northern hemisphere is made up largely of rolling volcanic plains not unlike the lunar maria—indeed, this similarity was key to their identification as lava-flow features. Much larger than their counterparts on Earth or the Moon, these extensive lava plains were formed by eruptions involving enormous volumes of material. They are strewn with blocks of volcanic rock, as well as with boulders blasted out of impact areas by infalling meteoroids. (The Martian atmosphere is too thin to offer much resistance to incoming debris.) The southern

► **FIGURE 10.3 Mars Globe** This highly detailed mosaic of Mars is based on images taken by a *Viking* spacecraft while orbiting the planet. Mars's Tharsis region, 5000 km across, bulges out from the equator, rising to a height of about 10 km. The two large volcanoes on the left mark the approximate peak of the Tharsis bulge. Dominating the center of the field of view is a vast "canyon" known as Valles Marineris—the Mariner Valley. (NASA)


hemisphere consists of heavily cratered highlands lying some 5 km above the level of the lowland north. Most of the dark regions visible from Earth are mountainous regions in the south. Figure 10.5 contrasts typical terrains in the two hemispheres.

The northern plains are much less cratered than the southern highlands. On the basis of the arguments presented in Chapter 8, this smoother surface suggests that the northern surface is younger. ∞ (Sec. 8.5) Its age is perhaps 3 billion years, compared with 4 billion in the south. In places, the boundary between the southern highlands and the northern plains is quite sharp—the surface level can drop by as much as 4 km in a horizontal distance of 100 km or so. Most scientists assume that the southern terrain is the original crust of the planet. How most of the northern



This scale depicts height, ranging from dark blue 8 km below the surface to white 8 km above.

▲ **Interactive FIGURE 10.4 Mars Map** The *Mars Global Surveyor* data displayed as a flat map. The Mariner Valley (see also Figure 10.6) can be seen at left, whereas the opposite side of the planet is dominated by the giant Hellas impact basin. Note the great difference in elevation between the northern and southern hemispheres. Some surface features are labeled, as are the *Viking*, *Pathfinder*, *Exploration Rover*, *Phoenix*, and *Curiosity* robot landing sites. (NASA)



**▲ FIGURE 10.5 Mars Up Close** (a) The northern hemisphere of Mars, like this one near Chryse Planitia, consists of rolling, volcanic plains. (b) The southern Martian highlands, like this one in Hellas Planitia, are heavily cratered. Both of these *Mars Express* photographs are in true color. (ESA)

hemisphere could have been lowered in elevation and subsequently flooded with lava remains a mystery.

The major geological feature on the planet is the Tharsis bulge (marked in Figure 10.3). Roughly the size of North America, Tharsis lies on the Martian equator and rises some 10 km higher than the rest of the Martian surface. To its east lies Chryse Planitia (the “Plains of Gold”), to the west a region called Isidis Planitia (the “Plains of Isis,” an Egyptian goddess). These features are wide depressions, hundreds of kilometers across and up to 3 km deep. If we wished to extend the idea of “continents” from Earth and Venus to Mars, we would conclude that Tharsis is the only continent on the Martian surface. However, as on Venus, there is no sign of plate tectonics on Mars—the absence of fault lines or other evidence of plate motion tells geologists that the “continent” of Tharsis is not drifting as its Earthly counterparts are. ∞ (Sec. 7.4) Tharsis appears to be even less heavily cratered than the northern plains, making it the youngest region on the planet, an estimated 2 to 3 billion years old.


Almost diametrically opposite Tharsis, in the southern highlands, lies the Hellas Basin, which, paradoxically, contains the *lowest* point on Mars. (Hellas is clearly visible and labeled in Figure 10.4.) Some 3000 km across, the floor of the basin lies nearly 9 km below the basin’s rim and over 6 km below the average level of the planet’s surface. Its shape and structure identify the Hellas Basin as an impact feature. The formation of the Hellas Basin must have caused a major redistribution of the young Martian crust—perhaps even enough to account for a substantial portion of the highlands around it, according to some researchers. The basin’s heavily cratered floor indicates that the impact occurred very early on in Martian history—some 4 billion years ago—during

the heavy bombardment that accompanied the formation of the terrestrial planets. ∞ (Secs. 6.6, 8.5)

The giant Borealis Basin around the Martian north pole—most of the blue region at the top of Figure 10.4 (see also Figure 10.11a)—may be the result of one of the largest known impacts in the solar system. Computer simulations of the collision, combined with detailed data from the *Mars Global Surveyor* and *Mars Reconnaissance Orbiter* spacecraft, suggest that the basin could have formed when a giant impactor some 2000 km across—twice the size of the largest asteroid, Ceres—struck the planet a grazing blow 3 billion years ago. ∞ (Sec. 6.6) These ideas are still debated by planetary scientists, but the resulting impact feature would be comparable in size to the observed basin, and the collision could also explain why the northern hemisphere of Mars is so much lower than and differs so radically from the south.

## The Martian “Grand Canyon”

A particularly prominent feature associated with the Tharsis bulge is a great “canyon” known as Valles Marineris (the Mariner Valley). Shown in its entirety in Figure 10.3 and in more detail in Figure 10.6, this feature is not really a canyon in the terrestrial sense because running water played no part in its formation. Planetary astronomers theorize that it was formed by the same crustal forces that caused the entire Tharsis region to bulge outward, making the surface split and crack. The resulting cracks, called *tectonic fractures*, are found all around the Tharsis bulge. Valles Marineris is the largest of them. Cratering studies suggest that the cracks are at least 2 billion years old; age estimates for Valles Marineris range up to 3.5 billion years. ∞ (Sec. 8.5) Similar (but much



**▲ FIGURE 10.6 Valles Marineris** (a) The Mariner Valley is a huge canyon, 120 km wide and 7 km deep. Its length is about 4000 km, or nearly the full breadth of the continental United States. (b) A close-up view shows the complexity of the valley walls and dry tributaries. (c) A comparison, to scale, with Earth's Grand Canyon, which is a mere 20 km wide and 2 km deep, suggests just how big the Mariner Valley is. (NASA)

smaller) cracks, originating from similar causes, have been found in the Aphrodite Terra region of Venus. ∞ (Sec. 9.4)

Valles Marineris runs for almost 4000 km along the Martian equator, about one-fifth of the way around the planet. At its widest, it is some 120 km across, and it is as deep as 7 km in places. Like many Martian surface features, it simply dwarfs Earthly competition. The Grand Canyon in Arizona would easily fit into one of its side “tributary” cracks. Valles Marineris is so large that it can even be seen from Earth—in fact, it was one of the few “canals” observed by 19th-century astronomers that actually corresponded to a real feature on the planet’s surface. (It was known as the Coprates canal.) We must reemphasize, however, that this Martian feature was not constructed by intelligent beings, nor was it carved by a river, nor is it a result of Martian plate tectonics. For some reason, the crustal forces that formed it never developed into full-fledged plate motion as exists on Earth.


## Volcanism on Mars

Mars contains the largest known volcanoes in the solar system. Three very large volcanoes are found on the Tharsis bulge, two of them visible on the left-hand side of Figure 10.3. The largest volcano of all is Olympus Mons (Figure 10.7),

northwest of Tharsis, lying just over the left (western) horizon of Figure 10.3. This volcano measures some 700 km in diameter at its base—only slightly smaller than the state of Texas—and rises to a height of 25 km above the surrounding plains. The caldera, or crater, at its summit, measures 80 km across. The other three large volcanoes are a little smaller—a “mere” 18 km high—and lie near the top of the bulge.

Like Maxwell Mons on Venus, none of these volcanoes is associated with plate motion on Mars—as just mentioned, there is none. Instead, they are shield volcanoes, sitting atop a hot spot in the underlying Martian mantle. ∞ (Sec. 9.4) All four show distinctive lava channels and other flow features similar to those found on shield volcanoes on Earth. *Viking* and *Mars Global Surveyor* images of the Martian surface reveal many hundreds of volcanoes. Most of the largest are associated with the Tharsis bulge, but many smaller volcanoes are found in the northern plains.

The great height of Martian volcanoes is a direct consequence of the planet’s low surface gravity. As lava flows and spreads to form a shield volcano, its eventual height depends on the new mountain’s ability to support its own weight. The lower the gravity, the less is the weight and the higher is the mountain. It is no accident that Maxwell Mons on Venus and the Hawaiian shield volcanoes on Earth rise to roughly



◀ **FIGURE 10.7 Olympus Mons** The largest volcano known on Mars or anywhere else in the solar system, Olympus Mons is nearly three times taller than Mount Everest on Earth, measuring about 700 km across its base and rising 25 km high at its peak. This Martian mountain seems currently inactive and may have been extinct for at least several hundred million years. By comparison, the largest volcano on Earth, Hawaii’s Mauna Loa, measures a mere 120 km across and peaks just 9 km above the Pacific Ocean floor. (NASA)

the same height (about 10 km) above their respective bases—Earth and Venus have similar surface gravity. Mars’s surface gravity is only 40 percent that of Earth, so volcanoes rise roughly 2.5 times as high.

Are the Martian shield volcanoes still active? Scientists have found no direct evidence for recent or ongoing eruptions. However, if these volcanoes have been around since the Tharsis uplift (as the formation of the Tharsis bulge is known) and were active as recently as 100 million years ago (an age estimate based on the extent of impact cratering on their slopes), some of them may still be at least intermittently active. Millions of years, though, may pass between eruptions.

## Impact Cratering

The *Mariner* spacecraft found that the surfaces of Mars and its two moons are pitted with impact craters formed by meteoroids falling in from space. On Mars, as on Venus, there is a lack of small impact craters, less than roughly 5 km in diameter. ∞ (Sec. 9.4) This time, though, the explanation is not that such craters do not form—small meteoroids have no trouble penetrating the thin Martian atmosphere. Instead, thin or not, the atmosphere is an efficient erosive agent, transporting dust from place to place and erasing small impact craters faster than they can form.

Overall, erosion on Mars is about 100 times slower than on Earth, but still far faster than on the Moon or Venus. For comparison, a 1-km diameter crater might survive for 100 million years on Mars. On Earth, it would be gone in a million years or so, but it would remain intact for tens of billions of years on the Moon before being obliterated by meteoritic

erosion. On Venus, the crater would most likely survive until the next large-scale volcanic resurfacing event. ∞ (Sec. 9.4)

As on the Moon and Venus, the extent of large impact cratering (craters too big to have been filled in by erosion since they formed) serves as an age indicator for the Martian surface. ∞ (Sec. 8.5) The ages quoted earlier, ranging from 4 billion years for the southern highlands to a few hundred million years in the youngest volcanic areas, were obtained in this way.


## PROCESS OF SCIENCE Check

- ✓ How do we know that the northern Martian lowlands are younger than the southern highlands?

## 10.5 Water on Mars

Mars today is apparently dry and desolate, yet astronomers have strong evidence that that was not always the case. Unlike Earth, where water is abundant, and Venus, which (as we saw in Chapter 9) has been devoid of water for billions of years, Mars offers intriguing hints that it may once have harbored liquid water on its surface. ∞ (Sec. 9.5) The relatively low rate of surface erosion on Mars means that many surface features formed billions of years ago are still detectable, providing astronomers with a unique opportunity—in principle, at least—to probe the presence of water on that planet over the entire span of Martian history.

Because water is such a vital ingredient to the development of life on Earth (see Chapter 28), its presence on Mars has important implications for life there too. Let’s take a closer look at conditions on the Martian surface since the planet formed.



◀ **FIGURE 10.8 Martian Channel** (a) This runoff channel on Mars is about 400 km long and up to 5 km wide in places. (b) The Red River on Earth runs from the Texas Panhandle to the Mississippi River. The two differ mainly in that there is currently no liquid water in this, or any other, Martian valley, but the networks of runoff channels strongly suggest the presence of running water on Mars in the distant past. (ESA; NASA)

## Evidence for Past Running Water

Although the great surface cracks in the Tharsis region are not really canyons and were not formed by running water, photographic evidence reveals that liquid water once existed in great quantity on the surface of Mars. Two types of flow feature are seen: the *runoff channels* and the *outflow channels*.

The **runoff channels** (one of which is shown in Figure 10.8a) are found in the southern highlands. They are extensive systems—sometimes hundreds of kilometers in total length—of interconnecting, twisting channels that seem to merge into larger, wider channels. They bear a strong resemblance to river systems on Earth (Figure 10.8b), and many geologists think that this is just what they are—the dried-up beds of long-gone rivers that once carried water on Mars from the mountains down into the valleys. The systems of runoff channels are often referred to as *valley networks*.


Were the runoff channels river systems like those on Earth, part of a planetwide water cycle, in which rain fell, forming rivers that drained into lakes or oceans, which in turn evaporated to form clouds and more rain? Or were they formed by something more modest, such as periodically melting underground ice, with no associated lakes or other extended bodies of water? Astronomers have debated this issue for years, but based on the accumulated data from the latest crop of Mars probes, the evidence now seems to favor the former view.

Many planetary scientists think that Mars may have enjoyed an extended early period during which rivers, lakes, and perhaps even oceans adorned its surface. The valley networks speak of a time 4 billion years ago (the age of the Martian highlands), when the atmosphere was thicker, the

surface warmer, and liquid water widespread. Figure 10.9 is a *Mars Global Surveyor* image showing what mission specialists think may be a delta—a fan-shaped network of channels and sediment deposits where a river once flowed into a larger body of water, in this case a lake filling a crater in the southern highlands.

Further evidence for this early warm period comes from many sources. Recent chemical studies of Martian meteorites like those discussed in *Discovery 10-1* strongly suggest warm (above freezing) temperatures, and in 2010 both *Mars Reconnaissance Observer* and *Mars Express* sensors have detected deposits of clay all across the southern highlands. Many researchers regard clay as strong evidence for liquid water on the surface, although here, too, experts disagree—some maintain that the clay can be produced by other means, and does not necessarily imply the presence of water when it formed. Finally, analysis of volcanic rocks studied by the *Spirit* rover (see below) in 2009 suggest that, when they were ejected 3.5 billion years ago, the Martian atmosphere was up to a factor of 20 denser than it is today.

The **outflow channels** (Figure 10.10a) are probably relics of slightly more recent catastrophic flooding on Mars. They appear only in equatorial regions and generally do not form the extensive interconnected networks that characterize the runoff channels. Instead, they are probably the paths taken by huge volumes of water draining from the southern highlands into the northern plains. The onrushing water arising from these flash floods probably also formed the odd teardrop-shaped “islands” (resembling the miniature versions seen in the wet sand of our beaches at low tide) that have been found on the plains close to the ends




▲ **FIGURE 10.9 Martian River Delta** Did this fan-shaped region of twisted streams form as a river flowed into a larger sea? If it did, the *Mars Global Surveyor* image supports the idea that Mars once had large bodies of liquid water on its surface. Not all scientists agree with this interpretation, however. (NASA)


of the outflow channels (Figure 10.10b). Judging from the width and depth of the channels, the flow rates must have been truly enormous—perhaps as much as a hundred times greater than the  $10^5$  tons per second carried by the Amazon River, the largest river system on Earth. Flooding shaped the outflow channels about 3 billion years ago, about the same time as the northern volcanic plains formed.

The discussion of water on ancient Mars splits naturally into two periods—the early period 4 billion years ago, associated with the runoff channels as just discussed, and a later period roughly 3 billion years ago, marked by the outflow channels. Many researchers think that the data provide evidence for large open expanses of water on the Martian surface during this later “wet” period. Figure 10.11(a) is a computer-generated view of the Martian north polar region, based on *Mars Global Surveyor* images, showing the extent of what may have been an ancient ocean covering much of the northern lowlands. The Hellas basin (Figure 10.4) is another possible candidate for an ancient Martian sea.


Both the existence and the longevity of these oceans remain controversial. Proponents point to features such as the terraced “beaches” shown in Figure 10.11(b), which might conceivably have been left behind as a lake or ocean evaporated and the shoreline receded. But detractors maintain that the terraces could also have been created by geological activity, perhaps related to the tectonic forces that depressed the northern hemisphere far below the level of the south, in which case they have nothing whatever to do with Martian water. Furthermore, *Mars Global Surveyor* data released in 2003, as well as later chemical analyses by the Martian landers, seem to indicate that the Martian surface contains too few *carbonate* rock layers—compounds



(a)



(b)



▲ **FIGURE 10.10 Martian Outflow** (a) An outflow channel near the Martian equator bears witness to a catastrophic flood that occurred about 3 billion years ago. (b) The onrushing water that carved out the outflow channels was responsible for forming these oddly shaped “islands” as the flow encountered obstacles—impact craters—in its path. Each “island” is about 40 km long. (NASA)


containing carbon and oxygen that should have been formed in abundance in ancient oceans. Their absence would support the picture of a cold, dry Mars that never experienced the extended mild period required to form lakes and oceans.

However, as more data have been obtained, the evidence in favor of ancient lakes or seas has strengthened. The *Mars Express* orbiter has detected “hydrated” chemical compounds in surface rocks over broad swaths of the planet, strongly suggesting that those regions were wet for extended periods of time. In 2012, mission scientists reported that radar studies indicated sedimentary material and ice in the

topmost 100 meters of regions (like Figure 10.11a) previously identified as oceans. In addition, as discussed below, direct chemical analyses made by the most recent NASA landers indicate that their landing sites also experienced long periods in the past during which liquid water existed on the surface.


## Subsurface Ice

As far as we can tell, there is no liquid water on the Martian surface today. However, the detailed appearance of Martian impact craters provides an important piece of information about conditions just below the planet’s surface. The ejecta blankets surrounding many Martian craters look quite different from their lunar counterparts. Figure 10.12 compares the Copernicus crater on the Moon with the (fairly typical) crater Yuty on Mars. The material surrounding the lunar crater is just what one would expect from an explosion ejecting a large volume of dust, soil, and boulders. However, the ejecta blanket on Mars gives the distinct impression of a liquid that has splashed or flowed out of the crater. Geologists think that this *fluidized ejecta* crater indicates that a layer of **permafrost**, or water ice, lies just a few meters under the surface. The explosive impact heated and liquefied the ice, resulting in the fluid appearance of the ejecta.





(a)


◀ **FIGURE 10.11 Ancient Ocean?** (a) A possible ancient Martian ocean might once have spanned the polar regions. The blue areas in this computer-generated map indicate depth below the average radius of the planet, thus approximating possible ancient ocean extent. (The color elevation scale is nearly the same as in Figure 10.4.) (b) This high-resolution image shows tentative evidence for erosion by standing water in the floor of Holden Crater, about 140 km across. Notice the layering in the true-color inset, suggestive of beach sand dunes. (NASA)



(b)







**▲ FIGURE 10.12 Moon–Mars Crater Comparison** (a) The large lunar impact crater Copernicus is typical of those found on Earth’s Moon. Its ejecta blanket appears to be composed of dry, powdery material. (b) The ejecta from the Martian crater Yuty evidently was once liquid. This type of crater is sometimes called a “splosh” crater, but most of the water remaining on Mars today probably exists in the form of underground ice. (NASA)

More direct evidence for subsurface ice was obtained in 2002, when a gamma-ray spectrometer aboard the *Mars Odyssey* orbiter detected extensive deposits of water ice crystals (actually, the hydrogen they contain) mixed with the Martian surface layers at high latitudes (more than  $50^\circ$  north and south of the equator). In some locations ice appears to compose as much as 50 percent by volume of the planet’s soil. The instrument was similar in design to the one carried by *Lunar Prospector*, which found ice crystals in the regolith near the lunar poles. [∞ \(Sec. 8.5\)](#) Radar aboard *Mars Express* has confirmed these results and also suggests deep deposits of ice extending hundreds of meters below the surface in many locations. In 2010, *Mars Reconnaissance Observer* reported what appear to be “glacial melts” less than a few hundred million years old. Why the ice should have melted so relatively recently is unknown.


### “Recent” Water on the Martian Surface?

Prior to the arrival of *Mars Global Surveyor* in 2000, astronomers thought that all the water below the Martian surface existed in the form of ice. However, since then, *Surveyor* mission scientists have reported the discovery of numerous small-scale “gullies” in Martian cliffs and crater walls that apparently were carved by running water in the relatively recent past. These features are too small to have been resolved by the *Viking* cameras. One such gully, found in the

inner rim of a Martian impact crater in the southern highlands, is shown in Figure 10.13(a). Its structure has many similarities to the channels carved by flash floods on Earth.

The ages of these intriguing Martian features are uncertain and might be as great as a million years in some cases. However, the *Surveyor* team speculates that some of them may still be active today and that liquid water might exist in some regions of Mars at depths of less than 500 m. Some scientists dispute this interpretation, arguing that the “fluid” responsible for the gullies could have been solid (granular) or even liquid carbon dioxide, expelled under great pressure from the Martian crust. Others point to features such as that shown in Figure 10.13(b), a very similar looking gully found in an impact crater in the Arctic—perhaps the closest we can come on Earth to replicating the harsh conditions on Mars—which formed when subsurface ice became exposed to sunlight and melted. Perhaps the same occurred on Mars, in which case, even if the gullies were created by flowing water, that does not necessarily imply liquid water below the surface.


Data from *Surveyor*’s cameras have deepened the mystery of the Martian surface flows. Figure 10.14 shows two images, taken 6 years apart, of an unnamed impact crater in the southern highlands. The white streak in the second image is thought (by some) to be a frozen mudslide, where liquid water briefly flowed down the inside of the crater wall, carrying rocky debris with it, then froze on the chilly Martian surface. In 2011, *Mars Reconnaissance Observer* observed



(a)



(b)




◀ **FIGURE 10.13 Running Water on Mars?** (a) This high-resolution *Mars Global Surveyor* view (left) of a crater wall (right) near the Mariner Valley shows evidence of “gullies” apparently formed by running water in the relatively recent past. (b) Similar gullies in the Haughton impact crater on the Arctic’s Devon Island formed when underground ice temporarily melted, causing streams of water to flow on the surface. (NASA; Gordon Osinski/Canadian Space Agency)

transient “finger-like” flows hundreds of meters long, that might conceivably contain salty subsurface water. The origin and composition of these features are uncertain, but one thing is clear—whatever they are, they formed *recently*, demonstrating that their production is an ongoing process.

### PROCESS OF SCIENCE Check

- ✓ Describe how the scientific debate over water on Mars has evolved since the 1960s as more data have become available.



▲ **FIGURE 10.14 Recent Martian Outflow** This comparison between two *Mars Global Surveyor* images taken 6 years apart of a Martian impact crater shows that something—the white streak (lower right), possibly water—flowed across the surface. If so, then the flows responsible for the gullies in Figure 10.13 are not just a thing of the past—the activity is ongoing. (NASA)

## The Martian Polar Caps

We have already noted that the Martian polar caps are composed predominantly of carbon dioxide frost—dry ice—and show seasonal variations. Each cap in fact consists of two distinct parts—the **seasonal cap**, which grows and shrinks each year, and the **residual cap**, which remains permanently frozen. At maximum size, in southern midwinter, the southern seasonal cap is some 4000 km across. Half a Martian year later, the northern cap is at its largest, reaching a diameter of roughly 3000 km. The two seasonal polar caps do not have the same maximum size because of the eccentricity of Mars’s orbit around the Sun. During southern winter, Mars is considerably farther from the Sun than half a year later, in northern winter. Thus the southern winter season is longer and colder than that of the north, and the polar cap grows correspondingly larger.

## DISCOVERY 10-1

### Life on Mars?

Even before the *Viking* missions reached Mars in 1976, most astronomers had abandoned hope of finding life there. Scientists knew that Mars had no large-scale canal systems, no surface water, almost no oxygen in its atmosphere, and no seasonal vegetation changes. The current absence of liquid water on Mars especially dims the chances for life there now. However, running water and, possibly, a dense atmosphere in the past may have created conditions suitable for the emergence of life long ago. (See Chapter 28 for a fuller discussion of what constitutes “life,” scientifically speaking, and why water plays such an important role.)

In the hope that some form of microbial life—perhaps bacteria or other microscopic organisms—might have survived to the present day, the *Viking* landers carried out experiments designed to detect biological activity. The accompanying pair of photographs shows the Martian surface before and after the robot arm of one of the landers dug a shallow trench to scoop up soil samples. The arm is visible in the first frame.

All three *Viking* biological experiments assumed some basic similarity between hypothetical Martian bacteria and those found on Earth. A *gas-exchange experiment* offered a nutrient broth to any residents of a sample of Martian soil and looked for gases that would signal metabolic activity. A *labeled-release experiment* added compounds containing radioactive carbon to the soil and then waited for results signaling that Martian organisms had either eaten or inhaled the carbon. Finally, a *pyrolytic-release experiment* added radioactively tagged carbon dioxide to a sample of Martian soil and atmosphere, waited a while, and then removed


the gas and tested the soil (by heating it) for signs that something had absorbed the tagged gas.

Initially, all three experiments appeared to be giving positive signals! However, subsequent careful studies showed that all the results could be explained by inorganic (i.e., nonliving) chemical reactions. Thus, at present, we have no irrefutable evidence for even microbial life on the Martian surface. Most scientists think that the *Viking* robots detected peculiar reactions that mimicked the basic chemistry of living organisms in some ways, but they did not detect life itself.

A criticism of the *Viking* experiments is that they searched only for life now living. Today, Mars seems locked in an ice age—the kind of numbing cold that would prohibit sustained life as we know it. If bacterial life did arise on an Earth-like early Mars, however, then we might be able to find its fossilized remains preserved on or near the Martian surface. Surprisingly, one place to look for life on Mars is right here on Earth. Scientists think that some meteorites found on Earth’s surface come from the Moon and from Mars. These meteorites were apparently blasted off these bodies long ago during an impact of some sort, thrown into space, and eventually trapped by Earth’s gravity, ultimately to fall to the ground. The most fascinating of the rocks are surely those from the Red Planet—for one of them may harbor fossil evidence for past life on Mars!

The next figure shows ALH84001, a blackened 2-kg meteorite about 17 cm across, found in 1984 in Antarctica. On the basis of estimates of the cosmic-ray exposure it received before reaching Earth, the rock is thought to have been blasted off Mars about 16 million years ago. Looking at this specimen through a microscope (inset), scientists can see rounded orange-brown “globules” of carbonate minerals on the rock’s shiny crust. Because carbonates form only in the presence of water, the presence of these globules suggests that carbon dioxide gas and liquid water existed near ground level at some point in Mars’s history, a conclusion that planetary scientists had drawn earlier from studies of *Viking*’s orbital images of valleys apparently carved by water when the Martian climate was wetter and warmer.

In a widely viewed press conference in Washington, DC, in 1996, a group of scientists argued, on the basis of all the data accumulated from studies of ALH84001, that they had discovered fossilized evidence for life on Mars. The key pieces of evidence they presented for primitive Martian life were as follows: (1) Bacteria on Earth can produce structures similar to the globules shown in the inset. (2) The meteorite contains traces of *polycyclic aromatic hydrocarbons*—a tongue-twisting name for a class of complex organic molecules (usually abbreviated PAHs) that, although not directly involved in known biological cycles on Earth, occur among the decay products of plants and other organisms. (3) High-powered electron microscopes show that ALH84001 contains tiny teardrop-shaped crystals of magnetite and iron sulfide embedded in places where the carbonate has dissolved. On Earth, bacteria are known to manufacture similar chemical crystals. (4) On very small scales, elongated and egg-shaped structures are seen within the carbonate globules. The researchers interpret these minute structures as fossils of primitive organisms.



(Before)

(After)

(NASA)



(NASA)

The photomicrograph in the second figure shows this fourth, and most controversial, piece of evidence—curved, rod-like structures that resemble bacteria on Earth. Scale is crucial here, however. The structures are only about  $0.5\ \mu\text{m}$  across, 30 times smaller than ancient bacterial cells found fossilized on Earth. Furthermore, several key tests have not yet been done, such as cutting through the suspected fossilized tubes to search for evidence of cell walls, semipermeable membranes, or any internal cavities where body fluids would have resided. Nor has anyone yet found in ALH84001 any amino acids, the basic building blocks of life as we know it (see Section 28.1).


These results remain highly controversial. Many experts do not agree that life has been found on Mars—not even fossilized life. Skeptics maintain that all the evidence could be the result of chemical reactions not requiring any kind of biology. Carbonate compounds are common in all areas of chemistry; PAHs are found in many lifeless places (glacial ice, asteroid-belt meteorites, interstellar clouds, and even the exhaust fumes of automobiles); bacteria are not needed to produce crystals; and it remains unclear whether the tiny tubular structures shown are animal, vegetable, or merely mineral. In addition, there is the huge problem of contamination—after all, ALH84001 was found on Earth and apparently sat in the Antarctic ice fields for 13,000 years before being picked up by meteorite hunters.

During 1999, the team released a new analysis of a second meteorite, named Nakhla (shown in the third figure), discovered in the Sahara Desert in 1911 and also thought to have come from Mars. Again, the scientists reported evidence for microbial life, in the form of clusters of minute spheres and ovals found within tiny filled cracks deep inside the meteorite, having similarities in size, shape, and arrangement to known bacteria on Earth. Since Nakhla is a volcanic basalt rock that solidified about 1.3 billion years ago (as opposed to 4 billion years ago for ALH84001), the new work suggests that life might have spanned the entire history of Mars. If


so, then life might still be present there today—but the opponents remain largely unconvinced.

As things now stand, it's a matter of interpretation—at the frontiers of science, issues are usually not as clear-cut as we would hope. The scientific method requires that we continually test and retest the competing theories to try to determine the truth. Only additional analysis and new data—perhaps in the form of samples returned directly from the Martian surface—will tell conclusively whether primitive Martian life existed long ago, although most workers in the field seem to have concluded that, taken as a whole, the results do not support the claim of ancient life on Mars. Still, even some skeptics concede that as much as 20 percent of the organic (carbon-based) molecules in ALH84001 could have originated on the Martian surface—although that is a far cry from proving the existence of life there.

Should the claim of life on Mars hold up against the weight of healthy skepticism in the scientific community, these findings may go down in history as one of the greatest scientific discoveries of all time. We are—or at least were—not alone in the universe! Maybe.



(NASA)



◀ **FIGURE 10.15 Martian Polar Caps** The southern (a) and northern (b) polar caps of Mars are shown to scale in these mosaics of *Mariner 9* images. These are the residual (permanent) caps, seen here during their respective summers half a Martian year apart. The southern cap is some 350 km across and is made up mostly of frozen carbon dioxide. The northern cap is about 1000 km across and is composed mostly of water ice. The inset shows greater detail in the southern cap. (ESA; NASA)

an opinion strengthened by spectroscopic observations that show an increase in the concentration of water vapor above the north pole in northern summer as some small fraction of its water ice evaporates in the Sun's heat. (Note that, in this terminology, Earth's polar caps are both residual, and are composed entirely of water ice.) The thickness of the Martian caps is uncertain, but it is likely that, as on Earth, they represent a significant storehouse for water on the planet.

Why is there such a temperature difference (at least 50 K) between the two residual polar caps, and why is the northern cap warmer, despite the

The seasonal caps are composed entirely of carbon dioxide. Their temperatures are never greater than about 150 K ( $-120^{\circ}\text{C}$ ) the point at which dry ice can form. During the Martian summer, when sunlight striking a cap is most intense, carbon dioxide evaporates into the atmosphere, and the cap shrinks. In the winter, atmospheric carbon dioxide refreezes, and the cap reforms. As the caps grow and shrink, they cause substantial variations (up to 30 percent) in the Martian atmospheric pressure—a large fraction of the planet's atmosphere freezes out and evaporates again each year. From studies of these atmospheric fluctuations, scientists can estimate the amount of carbon dioxide in the seasonal polar caps. The maximum thickness of the seasonal caps is thought to be about 1 m.

The residual caps (Figure 10.15) are smaller and brighter than the seasonal caps and show an even more marked north–south asymmetry. The southern residual cap is about 350 km across and, like the seasonal caps, is probably made mostly of carbon dioxide, although it also contains some water ice. Its temperature remains below 150 K at all times. Its composition, long suspected by theorists, was finally established only in 2004 by spectroscopic imaging observations made by the *Mars Express* orbiter.

The northern residual cap is much larger—about 1000 km across—and warmer, with a temperature that can exceed 200 K in northern summertime. Planetary scientists think that the northern residual cap is made mostly of water ice,

fact that the planet's northern hemisphere is generally cooler than the south (see Section 10.2)? The reason is not fully understood, but it seems to be related to the giant dust storms that envelop the planet during southern summer. These storms, which last for a quarter of a Martian year (about 6 Earth months), tend to blow the dust from the warmer south into the cooler northern hemisphere. The northern ice cap becomes dusty and less reflective. As a result, it absorbs more sunlight and warms up.

## Climate Change on Mars

Aside from the gullies and transient flows mentioned earlier, which are suggestive but by no means conclusive, astronomers have no direct evidence for liquid water anywhere on the surface of Mars today, and the amount of water vapor in the Martian atmosphere is tiny. Yet even setting aside the hints of ancient oceans, the extent of the outflow channels clearly implies that a huge total volume of liquid water must have existed on Mars in the distant past. Where did all that water go? Some of it may have entered the atmosphere and escaped into space, but, as best we can tell, most of it today is locked in the permafrost layer under the Martian surface, with a little more contained in the polar caps.

Planetary scientists trace the history of water on Mars along the following broad lines. Early on, conditions on the planet were much warmer—perhaps even Earth-like—and


liquid water was widespread, forming the runoff channels as rainfall drained into river valleys. Roughly 4 billion years ago, for reasons discussed below, climatic conditions changed and the water began to freeze, forming the permafrost and drying out the riverbeds. Mars remained frozen for about a billion years, until volcanic activity, or meteoritic impact, that formed the northern lowlands heated large regions of the surface, melting the subsurface ice and causing flash floods that created the outflow channels.

Subsequently, activity subsided, the water refroze, and Mars once again became a dry world. The lifetime of the oceans, if they existed, might have been quite short—as little as a few million years, according to some estimates. The present level of water vapor in the Martian atmosphere is the maximum possible, given the atmosphere's present density and temperature. Estimates of the total amount of water stored as permafrost and in the polar caps are uncertain, but it is thought that if all the water on Mars were to become liquid, it would cover the surface to a depth of roughly 10 meters.



## The View from the Martian Landers

Remote sensing—taking images and other measurements from orbit—has been vitally important to our understanding of Mars, but in some cases there is just no substitute for a close-up look. To date, seven U.S spacecraft have successfully landed on the Martian surface (Discovery 6-2). Their landing sites, marked on Figure 10.4, spanned a variety of Martian terrains. Their goals included detailed geological and chemical analysis of Martian surface rocks, the search for life, and the search for water.


*Viking 1* landed in Chryse Planitia, a broad depression to the east of Tharsis. The view that greeted its cameras (Figure 10.16) was a windswept, gently rolling, rather desolate plain, littered with rocks of all sizes, not unlike a high desert on Earth. The surface rocks visible in Figure 10.16 are probably part of the ejecta blanket of a nearby impact crater. *Viking 2* landed somewhat farther north, in a region of Mars called Utopia, chosen in part because mission planners anticipated greater seasonal climatic variations there. The plain on which *Viking 2* landed was flat and featureless (Figure 10.17). The views that the two landers recorded seem to be quite typical of the low-latitude northern plains.



▲ **FIGURE 10.16** *Viking 1* This is the view from the *Viking 1* spacecraft now parked on the surface of Mars. The fine-grained soil and the reddish rock-strewn terrain stretching toward the horizon contain substantial amounts of iron ore; the surface of Mars is literally rusting away. The sky is pale pink, the result of airborne dust. (NASA)



▲ **FIGURE 10.17** *Viking 2* Another view of the Martian surface, this one rock strewn and flat, as seen through the camera aboard the *Viking 2* robot that soft-landed on the northern Utopian plains. The discarded canister is about 20 cm long. The 0.5-m scars in the dirt were made by the robot's shovel. (NASA)



(a)



(b)

**▲ FIGURE 10.18 Mars Panorama** (a) A panoramic view of the terrain where NASA's *Opportunity* rover landed on Mars in 2004. This is Endurance crater, roughly 130 m across, about the size of a football field. (b) A close-up of a clump of rocks at the bottom of Endurance crater, where *Opportunity* found evidence for extensive liquid water in Mars's past. The blue is false color and is not indicative of water currently on the surface. (NASA)

The *Viking* landers performed numerous chemical analyses of the Martian regolith. One important finding of these studies was the high iron content of the planet's surface. Chemical reactions between the iron-rich surface soil and free oxygen in the atmosphere is responsible for the iron oxide ("rust") that gives Mars its characteristic color. Although the surface layers are rich in iron relative to Earth's surface, the overall abundance is similar to Earth's average iron content. On Earth, much of the iron has differentiated to the center. Chemical differentiation does not appear to have been nearly so complete on Mars.

The next successful mission to the Martian surface was *Mars Pathfinder*. During the unexpectedly long lifetime of its mission in 1997 (it lasted almost 3 months instead of the


anticipated 1), the lander performed measurements of the Martian atmosphere and atmospheric dust while its robot rover *Sojourner* carried out chemical analyses of the soil and rocks within about 50 m of the parent craft. ∞ (Sec. 6.6) ∞ (Discovery 6-2) In addition, more than 16,000 images of the region were returned to Earth. The *Sojourner* lander found that the soil at its landing site was similar to that found by the *Viking* landers. However, analyses of nearby rocks revealed a chemical makeup different from that of the Martian meteorites found on Earth (see *Discovery 10-1*).

The landing site for the *Pathfinder* mission had been carefully chosen to lie near the mouth of an outflow channel, and the size distribution and composition of the many rocks and boulders surrounding the lander were consistent with their having been deposited there by flood waters. In addition, the presence of numerous rounded pebbles strongly suggested the erosive action of running water at some time in the past.

The twin landers *Spirit* and *Opportunity* of the *Mars Exploration Rover* mission have made extensive chemical and geological studies of rocks within a few kilometers of their landing sites on opposite sides of the planet, with the primary goal of finding evidence for liquid water on the surface at some time in the past. ∞ (Sec. 6.6) ∞ (Discovery 6-2) Their findings provide the best evidence yet for standing water on the ancient Martian surface and have changed the minds of many skeptical scientists on the subject of water on Mars.

*Spirit's* landing site was rocky and similar in many ways to the terrains encountered by earlier landers, although closer study by the science team reveals that most of the rocks in the lander's vicinity do appear to have been extensively altered by water long ago. Halfway around the planet, however, *Opportunity* appears to have hit the jackpot in its quest, finding itself surrounded by rocks showing every chemical and geological indication of having been very wet—possibly immersed in saltwater—in the past (Figure 10.18).

*Opportunity's* landing site had been chosen in part because the *Mars Odyssey* orbiter had detected a compound called hematite in the surface rocks there, a possible indicator of past water, and measurements made by the rover do indeed suggest that the rocks near the landing site have



been alternately underwater and dry for extended periods of time, possibly as a shallow lake alternately filled and evaporated repeatedly over the course of Martian history. If life ever did exist at this site, the sorts of rocks found there might have preserved a fossil record very well, but *Opportunity* was not equipped to carry out such studies.

*Spirit* ceased communications with Earth in 2010, more than 6 years into a mission originally expected to last 3–6 months. As of 2013, *Opportunity* is still operating and in communication with Earth. Mission controllers are delighted at the many important contributions the two rovers have made to planetary science.


NASA's *Phoenix* mission landed in the planet's north polar region in May 2008 (see Figure 10.4). Its objectives included determining if the Martian arctic is or was capable of supporting life, looking for ice or other evidence of water, and exploring the Martian polar climate. The spacecraft was not a rover, but instead contained a sophisticated

array of equipment to monitor weather conditions and collect and analyze the surrounding soil.

*Phoenix* confirmed the presence of subsurface water ice at the landing site (Figure 10.19) and found clay and carbonates in the soil, both indicators of a wet environment at some time in the past, although scientists do not know if the water resulted from seasonally melting ice or is an indicator of the much more distant past. Chemical analysis of the soil suggests that liquid or near-liquid water existed on or near the Martian surface as recently as 100–200 million years ago. Initially, the overall composition of the soil seemed Earth-like, but later analysis suggested some chemical differences that might make it less friendly to life, at least as we know it. The mission left unresolved the question of whether the soil samples scooped up by *Phoenix* contained any carbon-based organic compounds.


*Phoenix* touched down in late fall in the Martian northern hemisphere, and during its final weeks its sensors reported the first snow as winter closed in. The mission ended when diminishing sunlight and extreme low temperatures shut down the lander's power supply.

The most recent arrival on Mars is NASA's *Curiosity* rover, which soft-landed on the planet in August 2012. This car-sized robot is by far the most sophisticated lander NASA has ever



◀ **FIGURE 10.19** Martian

**Exploration** The *Phoenix* lander's robotic arm is shown here with a surface sample in its scoop, just before delivering it to an onboard miniature chemistry laboratory. The circular hardware at bottom is one of its solar panels. The inset shows one of the first trenches (about the size of this book) dug by *Phoenix* to a depth of about 8 centimeters. The white material near the top is almost surely ice, which melted soon after excavation. (NASA)






placed on the Martian surface. Mission goals include studying the Martian climate and geology, probing the past and present roles of water, determining whether the landing site (inside Gale Crater) has ever been conducive to microbial life, and evaluating the suitability of the planet for possible future human exploration.

*Curiosity*, shown in Figure 10.20, landed in a region thought to be an ancient streambed that may have held, in the distant past, flowing water up to a meter deep. The search for carbonate rocks, which would strongly support the idea of ancient rivers and oceans on Mars (see Section 10.5), is a key element of the mission. The first detailed scientific data were returned in December 2012 but, despite some early hints and much media speculation, no definitive detection of carbonates or organic compounds was possible. However, the analysis did reveal complex chemistry within the Martian soil, including water, sulfur, and chlorine-containing minerals, and clearly demonstrated the capabilities of the probe's science laboratory. The inset to Figure 10.20 shows holes bored into a surface rock by an onboard drill, allowing the first sample of subsurface Martian dust, presumably unaffected by weathering processes that may have altered the chemistry of the rock's outer layers, to be brought aboard for analysis.

Although the hoped-for carbonates have not yet (as of mid 2013) been found, mission scientists are confident that *Curiosity*, over its multi-year lifetime, will make many important contributions to our knowledge of the history and evolution of the Red Planet.

### CONCEPT Check

✓ Where has all the Martian water gone?



## 10.6 The Martian Atmosphere

Long before the arrival of the *Mariner* and *Viking* spacecraft, astronomers knew from Earth-based spectroscopy that the Martian atmosphere was quite thin and composed primarily of carbon dioxide. In 1964, *Mariner 4* confirmed these results, finding that the atmospheric pressure was only about 1/150 the pressure of Earth's atmosphere at sea level and that carbon dioxide made up at least 95 percent of the total atmosphere. With the arrival of *Viking*, more detailed measurements of the Martian atmosphere could be made. Its composition is now known to be 95.3 percent carbon dioxide, 2.7 percent nitrogen, 1.6 percent argon, 0.13 percent oxygen, 0.07 percent carbon monoxide, and about 0.03 percent water vapor. The level of water vapor is quite variable. Weather conditions encountered by *Mars Pathfinder* were quite similar to those found by *Viking 1*.

### Atmospheric Structure and Weather


As the *Viking* landers descended to the surface, they made measurements of the temperature and pressure at various heights. The results are shown in Figure 10.21. The Martian atmosphere contains a troposphere (the lowest-lying atmospheric zone, where convection and weather occur), which varies both from place to place and from season to season. ∞ (Sec. 7.2) The variability of the troposphere arises from the variability of the Martian surface temperature. At noon in the summertime, surface temperatures may reach 300 K. Atmospheric convection is strong, and the top of the troposphere can reach an altitude of 30 km. At night, the atmosphere retains little heat, and the temperature can drop by as much as 100 K. Convection then ceases and the troposphere vanishes.

On average, surface temperatures on Mars are about 50 K cooler than on Earth. The low early-morning temperatures often produce water-ice fog in the Martian canyons (Figure 10.22). Higher in the atmosphere, in the stratosphere, temperatures are low enough for carbon dioxide to solidify, giving rise to a high-level layer of carbon dioxide clouds and haze.

For most of the year, there is little day-to-day variation in the Martian weather: The Sun rises, the surface warms up, and light winds blow until sunset, when the temperature drops

#### ◀ FIGURE 10.20 *Curiosity* on Mars

This is a “self-portrait” of *Curiosity* on the floor of Gale crater; its sampling arm contains the camera, which was removed from the photo. The inset shows one of the first holes made in the Martian bedrock by *Curiosity*'s mini-drill; the hole is 1.6 cm across and 2 cm deep. (NASA/JPL)



◀ **FIGURE 10.21 Martian Atmosphere** Structure of the Martian atmosphere, as determined by *Viking* and *Mars Global Surveyor*. The troposphere, which rises to an altitude of about 30 km in the daytime, occasionally contains clouds of water ice or, more frequently, dust during the planetwide dust storms that occur each year. Above the troposphere lies the stratosphere. Note the absence of a higher temperature zone in the stratosphere, indicating the absence of an ozone layer.


## Atmospheric Evolution

Although there is some superficial similarity in composition between the atmospheres of Mars and Venus, the two planets obviously have quite different atmospheric histories—Mars’s “air” is over 10,000 times thinner than that on Venus. As with the other planets we have studied, we can ask *why* the Martian atmosphere is as it is.


Presumably, Mars acquired a secondary atmosphere *outgassed* from the planet’s interior quite early in its history, just as the other terrestrial worlds did. ∞ (Sec. 7.2) Around 4 billion years ago, as indicated by the runoff channels in the highlands, Mars may have had a fairly dense atmosphere, complete with blue skies, oceans, and rain. Even taking into account the larger distance from Mars to the Sun and the fact that the Sun was about 30 percent less luminous 4 billion years ago (see Chapter 22), planetary scientists estimate that the greenhouse effect from a Martian atmosphere a few times denser than Earth’s present atmosphere could have kept conditions fairly comfortable. A surface temperature above 0°C (the freezing point of water) seems quite possible.

Sometime during the next billion years, most of the Martian atmosphere disappeared. Possibly, some of it was expelled by impacts with large bodies in the early solar system, and a


again. Only in the southern summer does the daily routine change. Strong surface winds (without rain or snow) sweep up the dry dust, carry it high into the stratosphere, and eventually deposit it elsewhere on the planet. At its greatest fury, a Martian storm floods the atmosphere with dust, making the worst storm we could imagine on Earth’s Sahara Desert seem inconsequential by comparison. The dust can remain airborne for months at a time. The blown dust forms systems of sand dunes similar in appearance to those found on Earth.



(a)



(b)



▲ **FIGURE 10.22 Fog in the Canyons** (a) As the Sun’s light reaches and heats the canyon floor, it drives water vapor from the surface. When this vapor comes in contact with the colder air above the surface, it condenses again, and a temporary water-ice fog results, as seen here, near Mars’s northern polar cap. (b) Fog also shrouds the floor of the 200-km-wide Lowell Crater, imaged here by *Mars Global Surveyor* in the autumn of 2000. (NASA)

substantial part may have leaked away into space because of the planet's weak gravity. ∞ (More Precisely 8-1) However, most of the remainder probably became unstable and was lost in a kind of “reverse runaway greenhouse effect,” as we now describe. The following scenario, summarized in Figure 10.23, is accepted by many planetary scientists, although not by all—those who discount the evidence presented earlier for liquid water and a thick early atmosphere on Mars obviously require no explanation for their absence today.

The key mechanism controlling the level of carbon dioxide in a terrestrial planet's atmosphere is the absorption of carbon dioxide into the rocks that make up the planet's crust. The presence of liquid water greatly accelerates this process—carbon dioxide dissolves in the liquid water of the planet's rivers and lakes (and oceans, if any), ultimately reacting with surface material to form carbonate rocks. The result—at least in the absence of any opposing effect—is a continual depletion of atmospheric carbon dioxide.


On Venus, as we saw in Chapter 9, an opposing effect did exist, as the familiar greenhouse effect ran away to high temperatures and pressures. ∞ (Sec. 9.5) Carbon dioxide left the surface and entered the atmosphere as the temperature rose, resulting in the extreme conditions we now find on that planet today.

On Earth, a very different process acts to counteract the absorption of carbon dioxide into the surface. *Plate tectonics*

constantly recycles our planet's carbon dioxide, returning it to the atmosphere via volcanic activity. ∞ (Sec. 7.4) Eventually, these two competing processes come into balance, and the atmospheric concentration of carbon dioxide tends to remain roughly constant.

However, neither of these opposing processes operated on Mars. The planet was too cool for the greenhouse effect to run away. And, as we have seen, Mars's interior cooled faster than Earth's, and the planet apparently never developed large-scale plate motion. Even taking into consideration the large volcanoes discussed earlier in this chapter, Mars has had on average far less volcanism than Earth does, so the processes depleting carbon dioxide have been much more effective than those replenishing it, creating a “one-way street” in which the level of atmospheric carbon dioxide steadily declined.

As the Martian carbon dioxide was consumed and its greenhouse effect diminished, the planet cooled, causing still more carbon dioxide to leave the atmosphere. The reason for this is just as described above—lower temperatures allowed more carbon dioxide to be absorbed in the surface layers. The result was a runaway to lower temperatures and decreasing levels of carbon dioxide in the atmosphere—just the opposite the sequence of events on Venus. Calculations show that much of the Martian atmospheric carbon dioxide could have been depleted in this way in a relatively short period of time, perhaps




▲ **FIGURE 10.23 Atmospheric Change** (a) On Earth, geological and volcanic activity returns  $\text{CO}_2$  to the atmosphere, and a stable balance is struck. The result is a modest greenhouse effect. (b) On Mars, there is little volcanic activity, and the  $\text{CO}_2$  returned was insufficient to replenish the atmosphere. As a result, the Martian atmosphere thinned and cooled, losing most of its  $\text{CO}_2$  in as little as a few hundred million years. If life did arise on early Mars, it most likely had a rather brief window of time in which to develop.

as quickly as a few hundred million years (Figure 10.24), although some of it might have been replenished by volcanic activity, possibly extending the “comfortable” lifetime of the planet to a half-billion years or so. Much of the debate about the presence of liquid water on the Martian surface revolves around the time it took for the planet’s surface to freeze.


As the temperature continued to fall, water froze out of the atmosphere, lowering still further the level of atmospheric greenhouse gases and accelerating the cooling. (Recall from Section 9.5 that water vapor also contributes to the greenhouse effect.) Eventually, even carbon dioxide began to freeze out, particularly at the poles, and Mars reached the frigid state we see today—a cold, dry planet with most of its original complement of atmospheric gases now residing in or under the barren surface.

### CONCEPT Check

✓ What happened to the Martian atmosphere?



(a) Ancient Mars



(b) Today's Mars

## 10.7 Martian Internal Structure

The *Viking* landers carried seismometers to probe the internal structure of Mars. However, one failed to work, and the other was unable to clearly distinguish seismic activity from the buffeting of the Martian wind. As a result, no seismic studies of the Martian interior have yet been carried out. On the basis of studies of the stresses that occurred during the Tharsis uplift, astronomers estimate the thickness of the crust to be about 100 km.

During its visit to Mars in 1965, *Mariner 4* detected no planetary magnetic field, and for many years the most that could be said about the Martian magnetic field was that its strength was no more than a few thousandths the strength of Earth’s field (to the level of sensitivity of *Mariner’s* instruments). In 1997, *Mars Global Surveyor* detected a very weak Martian field, about 1/800 times that of Earth. However, this is probably a local anomaly, akin to the magnetic fluctuations detected by *Lunar Prospector* at certain locations on the surface of Earth’s Moon, and not a global field. ∞ (Sec. 8.7)

Because Mars rotates rapidly, the absence of a global magnetic field is taken to mean that the planet’s core is non-metallic, nonliquid, or both. ∞ (Sec. 7.5) The small size of Mars indicates that any radioactive (or other internal) heating of its interior would have been less effective at melting the planet than similar heating on Earth. The heat could reach the surface and escape more easily than on a larger planet such as Earth or Venus.

The evidence we noted earlier for ancient surface activity, especially volcanism, suggests that at least parts of the planet’s interior must have melted and possibly differentiated at some time in the past. But the lack of current activity, the absence of any significant magnetic field, the relatively low density ( $3900 \text{ kg/m}^3$ ), and an abnormally high abundance of iron at the surface all suggest that Mars never melted as extensively as did Earth. The latest data indicate that the Martian core has a diameter of about 2500 km, is composed largely of iron sulfide (a compound about twice as dense as surface rock), and is still at least partly molten.

The history of Mars appears to be that of a planet on which large-scale tectonic activity almost started, but was stifled by the planet’s rapidly cooling outer layers. The large upwelling of material that formed the Tharsis bulge might have developed into full-fledged plate tectonic motion on a larger, warmer planet, but the Martian mantle became too rigid and the crust too thick for that to occur. Instead, the upwelling continued to fire volcanic activity, almost up to the present day, but, geologically, much of the planet apparently died 2 billion years ago.

◀ **FIGURE 10.24 Martian Evolution** (a) Artist’s conception of Mars some 4 billion years ago, with a dwindling atmosphere and some lingering surface water. (b) A photo of Mars today. (Kees Veenenbos)

**CONCEPT Check**

✓ What is the principal reason for the lack of geological activity on Mars today?

**10.8 The Moons of Mars**

Unlike Earth's Moon, Mars's moons are tiny compared with their parent planet and orbit very close to it, relative to the planet's radius. Discovered by American astronomer Asaph Hall in 1877, the two Martian moons—Phobos (“fear”) and Deimos (“panic”)—are only a few tens of kilometers across. Their composition is quite unlike that of the planet. They are quite difficult to study from Earth because their proximity to Mars makes it hard to distinguish them from their much brighter parent. However, the many orbiters that have since visited the planet have studied both in great detail.


As shown in Figure 10.25, Phobos and Deimos are both quite irregularly shaped and heavily cratered. The larger of the two is Phobos (Figure 10.25a), which is about 28 km long and 20 km wide and is dominated by an enormous 10-km-wide crater named Stickney (after Angelina Stickney, Asaph Hall's wife, who encouraged him to persevere in his observations). The smaller Deimos (Figure 10.25b) is only 16 km long by 10 km wide. Its largest crater is 2.3 km in diameter. The fact that both moons have quite dark surfaces, reflecting no more than 6 percent of the light falling on them, contributes to the difficulty in observing them from Earth.

Phobos and Deimos move in circular, equatorial orbits, and they rotate synchronously (i.e., they each keep the same face permanently turned toward the planet). These


characteristics are direct consequences of the tidal influence of Mars. Both moons orbit Mars in the prograde sense—that is, in the same sense (counterclockwise, as seen from above the north celestial pole) as the planet orbits the Sun and rotates on its axis.

Phobos lies only 9378 km (less than three planetary radii) from the center of Mars and, as we saw earlier, has an orbital period of 7 hours and 39 minutes. This period is much less than a Martian day, so an observer standing on the Martian surface would see Phobos move “backward” across the Martian sky—that is, in a direction opposite that of the apparent daily motion of the Sun. Because the moon moves faster than the observer, it overtakes the planet's rotation, rising in the west and setting in the east, crossing the sky from horizon to horizon in about 5.5 hours. Deimos lies somewhat farther out, at 23,459 km, or slightly less than seven planetary radii, and orbits in 30 hours and 18 minutes. Because it completes its orbit in more than a Martian day, it moves “normally,” as seen from the ground (i.e., from east to west), taking almost 3 days to traverse the sky.

Astronomers have determined the masses of the two moons by measuring their gravitational effects on the many spacecraft that have now orbited the planet. [∞ \(Sec. 6.2\)](#) The density of the Martian moons is around  $2000 \text{ kg/m}^3$ , far less than that of any world we have yet encountered in our outward journey through the solar system. This is one reason that astronomers think it unlikely that Phobos and Deimos formed along with Mars. Instead, for many years it was thought more probable that they were asteroids that came too close to the planet and were slowed and captured by the outer fringes of the early Martian atmosphere.



(a)



(b)

▲ **FIGURE 10.25 Martian Moons** (a) A *Mars Express* photograph of the potato-shaped Phobos, not much larger than Manhattan Island. The prominent crater (called Stickney) at left is several kilometers across. (b) Like Phobos, the smaller moon, Deimos, has a composition unlike that of Mars. Both moons are probably captured asteroids. This close-up photograph of Deimos was taken by a *Viking* orbiter. Most of the boulders shown are about the size of a house. (ESA)

Recent measurements by Mars Global Surveyor and Mars Express now suggest that the composition of Phobos differs significantly from that of asteroids of comparable density, and its rather porous internal structure may be too weak for it to have survived the capture process. Instead, it may have formed after material ejected from Mars by a meteoritic impact was captured and held by the planet's gravity, a little like the collision hypothesis for the formation of Earth's Moon, but on a much smaller scale. ∞ (Sec. 8.8)

However it got there, Phobos, on its low-altitude Martian orbit, continues to interact with the planet's upper atmosphere. Its orbit is expected to decay, plunging the moon into the surface in just a few tens of millions of years.

### CONCEPT Check


✓ In what ways do Phobos and Deimos differ from Earth's Moon?

**The Big Question** For well over a century, people have wondered, Is there life on Mars? After years of careful observation, this intriguing world has yielded no hint of living creatures, intelligent or otherwise—not even bugs hiding beneath the dusty rocks. If Martians do exist, they must be microbes—small, hidden, barely eking out a living in a mostly dry, dusty, and chilly world. Perhaps future robots will find archeological evidence for life now extinct, or maybe they will find no organisms at all—dead or alive. Either way, the consequences for our place in the cosmos could be profound.


# Chapter Review

## SUMMARY

**1** Mars lies outside Earth's orbit and traverses the entire plane of the ecliptic, as seen from Earth. Mars is about half the radius and one-tenth the mass of Earth. It rotates at almost the same rate as Earth, and its axis of rotation is inclined to the ecliptic at almost the same angle as Earth's axis. Surface temperatures on Mars average about 50 K cooler than those on Earth. Otherwise, Martian weather is reminiscent of that on Earth, with dust storms, clouds, and fog.



**2** As a result of its axial tilt, Mars has daily and seasonal cycles much like those on our own planet, but they are more complex than those on Earth because of Mars's eccentric orbit. From Earth, the most obvious Martian surface features are the polar caps, which grow and diminish as the seasons change on Mars. The two polar caps on Mars each consist of a **seasonal cap** (p. 249), composed of carbon dioxide, which grows and shrinks, and a **residual cap** (p. 249), of water ice, which remains permanently frozen. The appearance of the planet also changes because of seasonal dust storms that obscure its surface.




**3** The Martian surface has vast plains, huge volcanoes, and deep channels and canyons. Mars's major surface feature is the Tharsis bulge, located on the planet's equator. This feature may have been caused by a "plume" of upwelling material in the youthful Martian mantle. Associated with the bulge are Olympus Mons, the largest known volcano in the solar system, and a huge crack, called the Valles Marineris, in the



planet's surface. The height of the Martian volcanoes is a direct consequence of Mars's low surface gravity. No evidence for recent or ongoing eruptions has been found. On the other side of Mars from Tharsis lies the Hellas basin, the site of a violent meteoritic impact early in the planet's history. There is a marked difference between the two Martian hemispheres. The northern hemisphere consists of rolling volcanic plains and lies several kilometers below the level of the heavily cratered southern hemisphere. The lack of craters in the north suggests that this region is younger. The cause of the north-south asymmetry is not known.

**4** There is strong evidence that Mars once had running water on its surface. **Runoff channels** (p. 245) are the remains of ancient Martian rivers, whereas **outflow channels** (p. 245) are the paths taken by flash floods that cascaded from the southern highlands into the northern plains. *Mars Global Surveyor* and *Mars Express* images also strongly suggest that liquid water once existed in great quantity on Mars, and the *Mars Exploration Rover* landers have returned direct evidence for a wet Martian past. The planet may have enjoyed a relatively brief, warm "Earth-like" phase early on in its evolution, with a thick atmosphere and rain, rivers, and lakes or even oceans.




**5** Today, much of the water on Mars is locked up in the polar caps and in the layer of permafrost lying under the Martian surface. *Viking* observations of fluidized ejecta surrounding impact craters indicated the presence of subsurface ice, and *Mars Odyssey* and *Mars Express* subsequently detected extensive ice deposits mixed with and



lying under the planet's surface layers. *Mars Explorer* has found numerous gullies in crater walls that appear to have been formed by running water, and some flows are known to have occurred within the past few years. Whether these flows imply liquid water or ice below the surface remains unclear.


**6** Like the atmosphere of Venus, Mars's atmosphere is composed primarily of carbon dioxide. However, unlike Venus's atmosphere, the cool Martian atmosphere has a density less than 1 percent that of Earth's. Mars may once have had a dense atmosphere, but it was lost, partly to space and partly to surface rocks and subsurface **permafrost** (p. 247) and polar caps. Even today, the thin atmosphere is slowly leaking away.

**7** Mars has an extremely weak magnetic field, which, together with the planet's rapid rotation, implies that its core



is nonmetallic, nonliquid, or both. The lack of current volcanism, the absence of any significant magnetic field, the planet's relatively low density, and a high abundance of surface iron all suggest that Mars never melted and differentiated as extensively as did Earth. Convection in the Martian interior seems to have been stifled 2 billion years ago by the planet's rapidly cooling and solidifying mantle.

**8** The Martian moons Phobos and Deimos are probably asteroids captured by Mars early in its history. Their densities are far less than that of any planet in the inner solar system. These moons may be representative of conditions in the early solar system.



**MasteringAstronomy**® For instructor-assigned homework go to [MasteringAstronomy.com](http://MasteringAstronomy.com).

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- Why is opposition the best time to see Mars from Earth? Why are some Martian oppositions better than others for viewing Mars?
- LO1** Imagine that you will be visiting the southern hemisphere of Mars during its summer. Describe the atmospheric conditions you might face.
- LO2** Describe the two Martian polar caps, their seasonal and permanent composition, and the differences between them.
- Why is Mars red?
- LO3** Describe the major large-scale surface features of Mars.
- Why were Martian volcanoes able to grow so large?
- Why couldn't you breathe on Mars?
- LO4 POS** What is the evidence that water once flowed on Mars? Is there liquid water on Mars today?
- LO5** Is there water on Mars today, in any form?
- POS** Why do some scientists think Mars once had an extensive ocean? Where was it located?
- LO6** Compare and contrast the evolution of the atmospheres of Mars, Venus, and Earth.
- LO7** What do measurements of Martian magnetism tell us about the planet's interior?
- POS** What is the evidence that Mars never melted as extensively as did Earth?
- Since Mars has an atmosphere, and it is composed mostly of a greenhouse gas, why isn't there a significant greenhouse effect to warm its surface?
- LO8** How were the masses of Mars's moons measured, and what did these measurements tell us about their origin?

## Conceptual Self-Test: Multiple Choice

- Compared with the Earth's orbit, the orbit of Mars (a) has the same eccentricity; (b) is more eccentric; (c) is less eccentric; (d) is smaller.
- As seen from Earth, Mars exhibits a retrograde loop about once every (a) week; (b) 6 months; (c) 2 years; (d) decade.
- Compared with Earth's diameter, the diameter of Mars is (a) significantly larger; (b) significantly smaller; (c) nearly the same size; (d) unknown.
- The lengths of the seasons on Mars can be determined by observing the planet's (a) tilt; (b) eccentricity; (c) polar caps; (d) moons.
- In terms of area, the extinct Martian volcano Olympus Mons is about the size of (a) Mt. Everest; (b) Colorado; (c) North America; (d) Earth's Moon.
- VIS** Figure 10.4 ("Mars Map") clearly shows (a) surface water and ice at northern latitudes; (b) a giant canyon

stretching all the way across the planet; (c) iron deposits in the mid-latitudes; (d) cratered terrain in the south.

7. **vis** The best evidence for the existence of liquid water on an ancient Mars is Figure (a) 10.11; (b) 10.13; (c) 10.14; (d) 10.16.
8. Compared with the atmosphere of Venus, the Martian atmosphere has (a) a significantly higher temperature; (b) significantly more carbon dioxide; (c) a significantly lower atmospheric pressure; (d) significantly more acidic compounds.
9. In comparison to the atmosphere of Venus, the vastly different atmospheric character of Mars is likely due to a/an (a) ineffective greenhouse effect; (b) reverse greenhouse effect; (c) absence of greenhouse gases that would hold in heat; (d) greater distance from the Sun.
10. The moons of Mars (a) are probably captured asteroids; (b) formed following a collision with Earth; (c) are the remnants of a larger moon; (d) formed simultaneously with Mars.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

1. ●● What is the maximum elongation of Earth, as seen from Mars? (For simplicity, assume circular orbits for both planets.)
2. ● Calculate the minimum and maximum angular diameters of the Sun, as seen from Mars.
3. ● What would you weigh on Mars?
4. ● What was the minimum size of a Martian surface feature resolvable during the 2003 opposition (see Figure 10.1) by an Earth-based telescope with an angular resolution of 0.05"?
5. ● The mass of the Martian atmosphere is about 1/150 the mass of Earth's atmosphere and is composed mainly (95 percent) of carbon dioxide. Taking the mass of Earth's atmosphere to be roughly  $5 \times 10^{18}$  kg, estimate the total mass of carbon dioxide in the atmosphere of Mars. Compare your answer with the mass of a seasonal polar cap, approximated as a circular sheet of frozen carbon dioxide ("dry ice," having a density of  $1600 \text{ kg/m}^3$ ) of diameter 3000 km and thickness 1 m.
6. ● How long would it take the wind in a Martian dust storm, moving at a speed of 150 km/h, to encircle the planet's equator?
7. ● The Hellas impact basin is roughly circular, 3000 km across, and 6 km deep. Taking the Martian crust to have a density of  $3000 \text{ kg/m}^3$ , estimate how much mass was blasted off the Martian surface when the basin formed. Compare your answer with the present total mass of the Martian atmosphere. (See Problem 5.)
8. ●● The outflow channel shown in Figure 10.9 is about 10 km across and 100 m deep. If it carried  $10^7$  metric tons ( $10^{10}$  kg) of water per second, as stated in the text, estimate the speed at which the water must have flowed.

## Activities

### Collaborative

1. If Mars is visible in the night sky, observe it with as large a telescope as is available to you; binoculars will not be of much use. Use your almanac (or go online) to find out which Martian season is occurring at the time of your observation, which hemisphere is tilted in Earth's direction, and what longitude is pointing toward Earth. Sketch what you see. Look carefully and take your time. Repeat this observation several times over the course of a night. Take turns making the sketches. Afterward, try to identify the various features you have seen by referring to known objects on Mars (e.g., as shown in Figure 10.2). You should also

be able to see the planet's rotation by watching the surface features move. Do the same the next night. Because Mars's rotation period is so similar to Earth's, you should see the same surface features again.

### Individual

1. Several months before opposition, Mars begins retrograde motion. Chart the planet's motion relative to the stars to determine when it stops moving eastward and begins moving toward the west. Notice the increase in Mars's brightness as it approaches opposition.





# Jupiter

## GIANT OF THE SOLAR SYSTEM

Beyond the orbit of Mars, the solar system is very different from our own backyard. The outer solar system presents us with a totally unfamiliar environment: huge gas balls, peculiar moons, complex ring systems, and a wide variety of physical and chemical phenomena, many of which are still only poorly understood. Although the jovian planets—Jupiter, Saturn, Uranus, and Neptune—differ from one another in many ways, we will find that they have much in common, too. As with the terrestrial planets, we learn from their differences as well as from their similarities.

Our study of these alien places begins with the jovian planet closest to Earth: Jupiter, the largest planet in the solar system. In mass, composition, and internal structure, it offers a model for the other jovian worlds.

**The Big Picture** Despite their many peculiar properties, other planets in our solar system might offer some hints and clues about our own. Weather patterns observed and deciphered in the atmospheres of the big jovian planets could be especially helpful. Astronomers are currently using telescopes and spacecraft to examine Jupiter's cloud heights, wind speeds, gas temperatures, and chemical composition. They seek not only to understand Jupiter itself, but also to gain a comparative understanding of Earth, most notably the changing climate on our much smaller world.

**LEFT:** Jupiter is one of the most fascinating objects in the solar system. This true-color mosaic, constructed from two dozen images taken by a camera onboard the *Cassini* spacecraft, is the most detailed portrait of Jupiter ever made, resolving features as small as 60 km across. Note the Great Red Spot at lower middle—a storm that has been under way for several hundred years. Everything seen here is a cloud, from the equatorial regions that show alternating light and dark belts, to the high-latitude areas that appear more mottled. (JPL)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Specify the ways in which Jupiter differs from the terrestrial planets in its physical and orbital properties.
- 2 Outline the processes responsible for the appearance of Jupiter's atmosphere.
- 3 Describe Jupiter's internal structure and composition, and explain how their properties are inferred from external measurements.
- 4 Summarize the characteristics of Jupiter's magnetosphere.
- 5 List the orbital and physical properties of the Galilean moons of Jupiter, and describe the appearance and interior structure of each.
- 6 Explain how tidal forces can produce enormous internal stresses in a jovian moon, and discuss some effects of those stresses.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive Figures, and self-guided tutorials.

## 11.1 Orbital and Physical Properties

Named after the most powerful god of the Roman pantheon, Jupiter is by far the largest planet in the solar system. Ancient astronomers could not have known the planet's true size, but their choice of names was apt.

### The View from Earth

Jupiter is the fifth planet from the Sun and the innermost jovian planet (Figure 11.1). It is the third-brightest object in the night sky (after the Moon and Venus), making it easy to locate and study. As in the case of Mars, Jupiter is brightest when it is near opposition. When this happens to occur close to perihelion, the planet can be up to  $50''$  across, and a lot of detail can be discerned through even a small telescope.

Figure 11.2(a) is a photograph of Jupiter, taken through a telescope on Earth. In contrast to the terrestrial worlds, Jupiter has many moons that vary greatly in size and other properties. The four largest, visible in this telescopic view (and, to a few people, with the naked eye), are known as the **Galilean moons**, after Galileo Galilei, who discovered them in 1610. [∞ \(Sec. 2.4\)](#) Figure 11.2(b) is a *Hubble Space Telescope* image of Jupiter taken during the opposition of December 1990. Notice both the alternating light and dark bands that cross the planet parallel to its equator and also the large oval at the lower right. These atmospheric features


are quite unlike anything found on the inner planets. Figure 11.2(c) is an up-close, true-color image of Jupiter's north polar region, taken by the *Cassini* spacecraft as it passed the planet in 2001 en route to Saturn.

### Mass and Radius

Since astronomers have been able to study the motion of the Galilean moons for quite some time, Jupiter's mass has long been known to high accuracy. It is  $1.9 \times 10^{27}$  kg, or 318 Earth masses—more than twice the mass of all the other planets combined. In the broadest sense, our solar system is a two-object system with a lot of additional debris. Nonetheless, as massive as Jupiter is, it is still only 1/1000 the mass of the Sun.

Knowing Jupiter's distance and angular size, we can easily determine the planet's radius, which turns out to be 71,500 km, or 11.2 Earth radii. More dramatically stated, more than 1400 Earths would be needed to equal the volume of Jupiter. From the planet's size and mass, we derive a density of  $1300 \text{ kg/m}^3$  for Jupiter. Here (as if we needed it) is yet another indicator that Jupiter is radically different from the terrestrial worlds: It is clear that, whatever Jupiter's composition, it cannot possibly be made up of the same material as the inner planets. (Recall from Chapter 7 that Earth's average density is  $5500 \text{ kg/m}^3$ ). [∞ \(Sec. 7.1\)](#)

In fact, theoretical studies of the planet's internal structure indicate that Jupiter must be composed primarily of hydrogen and helium. The enormous pressure in



▲ **FIGURE 11.1 Solar System Perspective** This is a variation on Figure 6.5—neither an overhead view nor an edge-on view of our solar system, but an oblique view from a distant perspective—illustrating the jovian planets relative to their terrestrial cousins. Jupiter orbits at a distance of 5.2 AU from the Sun, outside the asteroid belt but well inside the Kuiper belt.




(a)



(b)



(c)



◀ **FIGURE 11.2 Jupiter** (a) This photograph, made through a ground-based telescope, shows Jupiter and its four big Galilean moons. (b) A *Hubble Space Telescope* image captures Jupiter in true color as our human eyes would see it and reveals features as small as a few hundred kilometers across. (c) A *Cassini* spacecraft image of part of Jupiter, taken while this robot vehicle was on its way to Saturn, shows intricate clouds of different heights, thicknesses, and chemical composition. (AURA; NASA)

the planet's interior due to Jupiter's strong gravity greatly compresses these light gases, whose densities on Earth (at room temperature and sea level) are  $0.08$  and  $0.16 \text{ kg/m}^3$ , respectively, producing the relatively high average density we observe.


## Rotation Rate

As with other planets, we can attempt to determine Jupiter's rotation rate simply by timing a surface feature as it moves around the planet. However, in the case of Jupiter (and, indeed, all the gaseous outer planets), there is a catch: Jupiter has no solid surface. All we see are the features of clouds in the planet's upper atmosphere. With no solid surface to “tie them down,” different parts of Jupiter's atmosphere move independently of one another.


Visual observations and Doppler-shifted spectral lines indicate that the equatorial zones rotate a little faster (with a period of  $9^{\text{h}}50^{\text{m}}$ ) than the higher latitudes (with a period of  $9^{\text{h}}55^{\text{m}}$ ). Jupiter thus exhibits **differential rotation**—the rotation rate is not constant from one location to another. Differential rotation is not possible in solid objects like the terrestrial planets, but it is normal for fluid bodies such as Jupiter.

Observations of Jupiter's magnetosphere provide a more meaningful measurement of the rotation period. The planet's magnetic field is strong and emits radiation at radio wavelengths as charged particles accelerate in response to Jupiter's magnetic field. Careful studies show a periodicity of  $9^{\text{h}}55^{\text{m}}$  at these radio wavelengths. We assume that this measurement matches the rotation of the planet's interior, where the magnetic field arises. ∞ (Sec. 7.5) Thus, Jupiter's interior rotates at the same rate as the clouds at the planet's poles. The equatorial zones rotate more rapidly.


A rotation period of  $9^{\text{h}}55^{\text{m}}$  is fast for such a large object. In fact, Jupiter has the fastest rotation rate of any planet in the solar system, and this rapid spin has altered Jupiter's shape. As illustrated in Figure 11.3, a spinning object tends to flatten and develop a bulge around its midsection. ∞ (Sec. 6.6) The more loosely the object's matter is bound together, or the faster it spins, the larger the bulge becomes. In objects such as Jupiter, which are made up of gas or loosely packed matter, high spin rates can produce a quite pronounced bulge. Jupiter's equatorial



(a)



(b)

 **Interactive FIGURE 11.3 Rotational Flattening** All spinning objects tend to develop an equatorial bulge because rotation causes matter to push outward against the inward-pulling gravity. The size of the bulge depends on the mechanical strength of the matter and the rate of rotation.

radius (71,500 km) exceeds its polar radius (66,900 km) by about 6.5 percent.\*

But there is more to the story of Jupiter's shape. Jupiter's observed equatorial bulge also tells us something important about the planet's deep interior. Careful calculations indicate that Jupiter would be *more* flattened than it actually is if its core were composed of hydrogen and helium alone. To account for the planet's observed shape, we must assume that Jupiter has a dense, compact core, probably of rocky composition, about 5–10 times the mass of Earth. This is one of the few pieces of data we have on Jupiter's internal structure.

\*Earth also bulges slightly at the equator because of rotation. However, our planet is much more rigid than Jupiter, and the effect is much smaller—the equatorial diameter is only about 40 km larger than the distance from pole to pole, a tiny difference compared with Earth's full diameter of nearly 13,000 km. Relative to its overall dimensions, Earth is smoother and more spherical than a billiard ball.

## CONCEPT Check

- ✓ How do observations of a planet's magnetosphere allow astronomers to measure the rotation rate of the interior?

## 11.2 Jupiter's Atmosphere

Jupiter is visually dominated by two features: a series of ever-changing atmospheric bands arranged parallel to the equator and an oval atmospheric blob called the **Great Red Spot**, or, often, just the “Red Spot.” The bands of clouds, clearly visible in Figure 11.2, display many colors—pale yellows, light blues, deep browns, drab tans, and vivid reds, among others. Shown in more detail in Figure 11.4, a close-up photograph taken as *Voyager 1* sped past in 1979, the Red Spot is the largest of many features associated with Jupiter's weather. It seems to be a hurricane twice the size of planet Earth that has persisted for hundreds of years, the largest of numerous long-lived storm systems in the planet's atmosphere.


### Atmospheric Composition

Spectroscopic studies of sunlight reflected from Jupiter gave astronomers their first look at the planet's atmospheric composition. Radio, infrared, and ultraviolet observations provided more details later. The most abundant gas is molecular hydrogen ( $\text{H}_2$ , 86.1 percent by number of molecules), followed by helium (He, 13.8 percent). Together, these two gases make up over 99 percent of Jupiter's atmosphere. Small amounts of atmospheric methane ( $\text{CH}_4$ ), ammonia ( $\text{NH}_3$ ), and water vapor ( $\text{H}_2\text{O}$ ) are also found. Researchers think that hydrogen and helium in those same proportions make up the bulk of the planet's interior as well.

The abundance of hydrogen and helium on Jupiter is a direct consequence of the planet's strong gravity. Unlike the gravitational pull of the terrestrial planets, the gravity of the much more massive jovian planets is powerful enough to have retained even hydrogen. [∞ \(More Precisely 8-1\)](#) Little, if any, of Jupiter's original atmosphere has escaped since the planet formed 4.6 billion years ago.

### Atmospheric Bands

Astronomers generally describe Jupiter's banded appearance—and, to a lesser extent, the appearance of the other jovian worlds as well—as a series of bright **zones** and dark **belts** crossing the planet. These variations appear to be the result of *convective motion* in the planet's atmosphere. [∞ \(Sec. 7.2\)](#) *Voyager* sensors indicated that the light-colored zones lie above upward-moving convective currents in Jupiter's atmosphere. The dark belts are regions representing the other part of the convection cycle, during which



◀ **FIGURE 11.4 Jupiter's Red Spot**  
*Voyager 1* took this photograph of Jupiter's Great Red Spot (upper right) from a distance of about 100,000 km. Resolution is about 100 km. The arrows indicate direction of gas flow above, below, and inside the Red Spot. (NASA)


cause our weather on Earth. A major difference between Jupiter and Earth is that Jupiter's rapid rotation has wrapped these systems all the way around the planet, instead of forming localized circulating storms, as on our own world.

Observations made by the *Cassini* mission in 2000 during its Jupiter flyby have challenged this standard view, suggesting instead that upward convection is actually confined to the belts. For now, planetary scientists have no clear resolution to the apparent contradiction between the *Voyager* and the *Cassini* findings.

Underlying the bands is an apparently very stable pattern of eastward and westward wind flow, known as Jupiter's **zonal flow**. Figure 11.5 illustrates how the wind

material is generally sinking downward, as illustrated schematically in Figure 11.5.


Because of the upwelling material below them, the zones are regions of *high pressure*. The belts, conversely, are *low-pressure* regions. The belts and zones are Jupiter's equivalents of the familiar high- and low-pressure systems that




Cooler gas sinks into the atmosphere, creating darker bands atop lower-pressure regions.

Warm material rises and creates lighter-colored zones.

This is a real *Voyager* photo of Jupiter's clouds, showing its actual banded structure.



◀ **FIGURE 11.5 Jupiter's Convection** The colored bands in Jupiter's atmosphere are associated with vertical convective motion. As on Earth, winds tend to blow from high- to low-pressure regions. Jupiter's rapid rotation channels those winds into a global east-west flow pattern, as indicated by the three yellow-red arrows drawn atop the belts and zones. (NASA)



**Interactive FIGURE 11.6 Zonal Flow** This graph shows wind speed in Jupiter's atmosphere, measured relative to the planet's internal rotation rate. Alternations in wind direction are associated with the atmospheric band structure. By studying global circulation patterns in the atmospheres of the jovian worlds, scientists also learn about the basic processes driving Earth's atmosphere.

remains. The variations are not seasonal in nature: Having a low-eccentricity orbit and a rotation axis almost exactly perpendicular to its orbital plane, Jupiter has no seasons. (Sec. 1.4) Instead, the annual changes appear to be the result of dynamic motion in the planet's atmosphere.

## Atmospheric Structure and Color

None of the atmospheric gases listed earlier can, by itself, account for Jupiter's observed

direction alternates between adjacent bands as Jupiter's rotation deflects surface winds into eastward or westward streams. The interaction between convective motion in Jupiter's atmosphere and the planet's rapid rotation channels the largest convective eddies into the observed zonal pattern. Smaller eddies—like the Red Spot—cause localized irregularities in the zonal flow.


The connection between Jupiter's belts and zones and the zonal flow pattern is evident in Figure 11.6, which shows the wind speed at different planetary latitudes measured relative to the rotation of the planet's interior (determined from studies of Jupiter's magnetic field). As mentioned earlier, the equatorial regions of the atmosphere rotate faster than the planet; their average flow speed is some 85 m/s, or about 300 km/h, in the easterly direction. The speed of this equatorial flow is quite similar to that of the jet stream on Earth. At higher latitudes, there are alternating regions of westward and eastward flow, roughly symmetric about the equator, with the flow speed generally diminishing toward the poles. Near the poles, where the zonal flow disappears, the band structure vanishes also.

Because of the pressure difference between the two, the zones lie slightly higher in the atmosphere than do the belts. The associated temperature differences (the temperature increases as we descend into the atmosphere, as we will see next) and the resulting differences in chemical reactions are the basic reasons for the different colors of these jovian features. The zones and belts vary in both latitude and intensity during the year, although the general banded pattern

coloration. For example, frozen ammonia and water vapor would simply produce white clouds, not the many colors actually seen. Scientists suspect that the colors of the clouds are the result of complex chemical processes occurring in the planet's turbulent upper atmosphere, although the details are still not fully understood. When we observe Jupiter's colors, we are actually looking down to many different depths in the planet's atmosphere.

Based on the best available data and mathematical models, Figure 11.7 is a cross-sectional diagram of Jupiter's atmosphere. Since the planet lacks a solid surface to use as a reference level for measuring altitude, the top of the troposphere is conventionally taken to lie at 0 km. As on all planets, weather on Jupiter is the result of convection in the troposphere, so the clouds, which are associated with planetary weather systems, all lie at negative altitudes in the diagram. Just above the troposphere lies a thin, faint layer of haze created by *photochemical* reactions (reactions involving sunlight) similar to those that cause smog on Earth. The temperature at this level is about 110 K; it increases with altitude as the atmosphere absorbs solar ultraviolet radiation.

Jupiter's clouds are arranged in three main layers. Below the haze, at a depth of about 40 km (shown as -40 km in Figure 11.7), lies a layer of white, wispy clouds made up of ammonia ice. The temperature here is approximately 125–150 K; it increases quite rapidly with increasing depth. A few tens of kilometers below the ammonia clouds, the temperature is a little warmer—over 200 K—and the clouds are probably made up mostly of droplets or crystals of ammonium hydrosulfide,



▲ **FIGURE 11.7 Jupiter's Atmosphere** Models of the vertical structure of Jupiter's atmosphere show the planet's clouds arranged in three main layers, each with quite different colors and chemistry. The white regions are the tops of the upper ammonia clouds. The yellows, reds, and browns are associated with the second cloud layer, which is composed of ammonium hydrosulfide ice. The lowest (bluish) cloud layer is water ice. The blue curve shows how Jupiter's atmospheric temperature depends on altitude. (For comparison with Earth, see Figure 7.2.)

produced by reactions between ammonia and hydrogen sulfide in the planet's atmosphere. At deeper levels in the atmosphere, the ammonium hydrosulfide clouds give way to clouds of water ice or water vapor. This lowest cloud layer, which is not seen in visible-light images of Jupiter, lies some 80 km below the top of the troposphere.

Instead of being white (the color of ammonium hydro-sulfide on Earth), Jupiter's middle cloud layer is tawny in color. This is the level at which atmospheric chemistry begins to play a role in determining Jupiter's appearance. Many planetary scientists think that molecules containing the element sulfur, and perhaps even sulfur itself, are important in influencing the cloud colors—particularly the reds, browns, and yellows, all colors associated with sulfur or its compounds. It is also possible that compounds containing the element phosphorus contribute to the coloration.

Deciphering the detailed causes of Jupiter's distinctive colors is a difficult task. The cloud chemistry is complex and highly sensitive to small changes in atmospheric conditions, such as pressure, temperature, and chemical composition.


The atmosphere is in incessant, churning motion, causing conditions to change from place to place and from hour to hour. In addition, the energy that powers the reactions comes in many different forms: the planet's own interior heat, solar ultraviolet radiation, aurorae in the planet's magnetosphere, and lightning discharges within the clouds themselves. All of these factors combine to keep a complete explanation of Jupiter's appearance beyond our present grasp.

The preceding description of Jupiter's atmosphere, based largely on *Voyager* data, was put to the test in December 1995, when the *Galileo* atmospheric probe arrived at the planet. ∞ (Discovery 6-2) The probe survived for about an hour before being crushed by atmospheric pressure at an altitude of  $-150$  km (i.e., right at the bottom of Figure 11.7). Overall, *Galileo*'s findings on wind speed, temperature, and composition were in good agreement with the picture just presented. However, the probe's entry location was in Jupiter's equatorial zone and, as luck would have it, coincided with an atypical "hole" almost devoid of upper-level clouds (see Figure 11.8). The probe measured a temperature of 425 K at 150 km depth—a little higher than indicated in Figure 11.7, but consistent with the craft's having entered a clearing in Jupiter's cloud decks, where convective heat can more readily rise (and thus be detected). The probe also measured a slightly lower than expected water content, but that, too, may be normal for the hot, windy regions near Jupiter's equator.



▲ **FIGURE 11.8 Galileo's Entry Site** The arrow on this image shows where the *Galileo* atmospheric probe plunged into Jupiter's cloud deck on December 7, 1995. Until its demise, the probe took numerous weather measurements, transmitting those signals to the orbiting mother ship, which then relayed them to Earth. (NASA)





▲ **FIGURE 11.9 Red Spot Details** These *Voyager 2* close-up views of the Great Red Spot, taken 4 hours apart, show clearly the turbulent flow around its edges. The general direction of motion of the gas north of (above) the spot is westward (to the left), whereas gas south of the spot flows east. The spot itself rotates counterclockwise, suggesting that it is being “rolled” between the two oppositely directed flows. The colors have been exaggerated somewhat to enhance the contrast. (NASA)

The experts were somewhat surprised by the depth to which Jupiter’s winds continued. *Galileo*’s probe measured high wind speeds throughout its descent into the clouds, not just at the cloud tops, implying that heat deep within the planet, rather than sunlight, drives Jupiter’s weather patterns. Finally, complex organic molecules were sought, but not found. Some simple carbon-based molecules, such as ethane ( $C_2H_6$ ), were detected by one of the onboard spectrometers, but nothing suggesting prebiotic compounds (molecules that could combine to form the building blocks of life—see Section 28.1) or bacteria floating in the atmosphere was found. That same instrument also detected traces of phosphine ( $PH_3$ ), which may be a key coloring agent for Jupiter’s clouds.

## Weather on Jupiter



In addition to the zonal flow pattern, Jupiter has many “small-scale” weather patterns. The Great Red Spot (Figure 11.4) is a prime example. It was first reported by British scientist Robert Hooke in the mid-17th century, so we can be reasonably sure that it has existed continuously, in one form or another, for over 300 years. It may well be much older. *Voyager* observations showed the spot to be a region of swirling, circulating winds, rather like a whirlpool or a terrestrial hurricane—a persistent and vast atmospheric storm. The size of the spot varies, although it averages about twice the diameter of Earth. Its present dimensions are roughly 25,000 km by 15,000 km. The spot rotates around Jupiter at a rate similar to that of the planet’s interior, perhaps suggesting that the roots of the Great Red Spot lie far below the atmosphere.

The origin of the spot’s red color is uncertain, as is its source of energy, although it is generally supposed that the spot is somehow sustained by Jupiter’s large-scale atmospheric motion. Repeated observations show that the gas flow around the spot is counterclockwise, with a period of about 6 days. Turbulent eddies form and drift away from its edge. The spot’s center, however, remains quite tranquil in appearance, like the eye of a hurricane on Earth. The zonal motion north of the Great Red Spot is westward, whereas that to the south is eastward (see Figure 11.9), supporting the idea that the spot is confined and powered by the zonal flow. However, the details of how it is so confined are still unclear. Computer simulations of the complex fluid flows in Jupiter’s atmosphere only hint at answers.

Storms, which as a rule are much smaller than the Great Red Spot, may be quite common on Jupiter. Spacecraft photographs of the dark side of the planet reveal bright flashes resembling lightning. The *Voyager* mission discovered many smaller light- and dark-colored spots that are also apparently circulating storm systems. Note the **white ovals** in Figures 11.4 and 11.9, south of the spot. Like the spot itself, they rotate counterclockwise. Their high cloud tops give them their color. These particular white ovals are known to be at least 40 years old. Figure 11.10 shows a **brown oval**, a “hole” in the clouds that allows us to look down into Jupiter’s lower atmosphere. For unknown reasons, brown ovals appear only at latitudes around  $20^\circ N$ . Although not as long lived as the Great Red Spot, these systems can persist for many years or even decades.

Continuous monitoring of conditions on the outer planets has recently yielded important insights into the formation and evolution of large storm systems on the jovian worlds. In


The brown oval is nearly as large as Earth's diameter.




▲ **FIGURE 11.10 Brown Oval** This brown oval in Jupiter's northern hemisphere is actually a break in the upper cloud layer, allowing us to see deeper into the atmosphere, where the clouds are brown. (NASA)

the late 1990s, astronomers noted with interest the collision and merger of three relatively small white ovals in Jupiter's atmosphere (Figure 11.11a). For several years the resultant larger system remained a white oval, but in early 2006 it changed from white to brown to red, becoming in a matter of months a smaller version of the Great Red Spot! Figure 11.11(b) shows a *Hubble Space Telescope* image of the new red spot, known as "Junior," as well as a third, "baby" red spot that appeared in 2008 at almost the same latitude as the other two storms.


Scientists speculate that the red coloration of the Great Red Spot may be due to that storm's enormous size and strength, which lifts its cloud tops high above the surrounding clouds, where solar ultraviolet radiation causes chemical reactions producing the color. In that case, the reddening of the smaller spot may indicate that




(a)



(b)



(c)



▲ **FIGURE 11.11 Red Spot Junior** (a) Between 1997 and 2000, astronomers watched as three white ovals in Jupiter's southern hemisphere merged to form a single large storm. Each oval, captured here by the *Cassini* spacecraft cameras, is about half the size of Earth. (b) In early 2006 the white oval turned red, producing a second red spot! The color change may indicate that the storm is intensifying. (c) In mid-2008, the *Hubble* telescope recorded this sequence of images at monthly intervals (left to right), showing a "baby red spot" (arrows) approaching the Great Red Spot and being destroyed by it. Jupiter's storm systems allow terrestrial scientists to probe the complexities of atmospheric dynamics under conditions not found on Earth. (NASA)

## DISCOVERY 11-1

### A Cometary Impact

In July 1994 astronomers were granted a novel means of studying Jupiter's atmosphere and interior—the collision of a comet (called Shoemaker-Levy 9, after its discoverers) with the planet!

When it was discovered in March 1993, comet Shoemaker-Levy 9 appeared to have an odd “squashed” appearance. Higher resolution images such as that shown in part (a) on the facing page, revealed that the comet was really made up of several pieces, the largest no more than 1 km across. All the pieces were following the same orbit, but they were spread out along the comet's path, like a string of pearls 1 million kilometers long.

How could such an unusual object have originated? Tracing the orbit backward in time, researchers calculated that early in July 1992 the comet had approached within about 100,000 km of Jupiter. They realized that the objects shown in the figure were the fragments produced when a previously “normal” comet was captured by Jupiter and torn apart by its strong gravitational field. The data revealed an even more remarkable fact: On its next approach, roughly a year later, the comet would collide with Jupiter!

Between July 16 and July 22, 1994, fragments from Shoemaker-Levy 9 struck Jupiter's upper atmosphere, plowing into it at a speed of more than 60 km/s and causing a series of enormous explosions. Every major telescope on Earth, the *Hubble Space Telescope*, *Galileo* (which was only 1.5 AU from the planet at the time), and even *Voyager 2* were watching. Each impact created, for a period of a few minutes, a brilliant fireball hundreds of kilometers across with a temperature of many thousands of kelvins. The largest of the fireballs was bigger than planet Earth. The energy released in each explosion was comparable to a billion terrestrial nuclear detonations, rivaling in violence the prehistoric impact suspected of causing the extinction of the

dinosaurs on Earth 65 million years ago (see *Discovery 14-1*). One of the largest pieces of the comet, fragment G, produced the spectacular fireball shown in part (b).

The effects on the planet's atmosphere and the vibrations produced throughout Jupiter's interior were observable for days after the impact. The fallen material from the impacts spread slowly around Jupiter's bands and reached completely around the planet after 5 months. It took years for all the cometary matter to settle into Jupiter's interior.

As best we can determine, none of the cometary fragments breached the jovian clouds. Only *Galileo* had a direct view of the impacts on the back side of Jupiter, and in every case the explosions seemed to occur high in the atmosphere, above the uppermost cloud layer. Most of the dark material seen in the images (see part c) is probably pieces of the comet rather than parts of Jupiter. Water vapor was also detected spectroscopically, again apparently from the melted and vaporized comet—the composition of which resembled a “dirty snowball,” just as astronomers had long predicted (see Section 14.2).

When the Shoemaker-Levy 9 impacts were first observed in 1994, they were vindication for scientists (like Shoemaker) who had been claiming for years that collisions were important events in solar system evolution. However, they were still thought of as rare. Recent observations suggest that this is not the case. Thanks in large part to improving technology in the hands of amateur astronomers, Jupiter is now monitored more closely than ever before, and it is becoming clear that cometary impacts may be commonplace. The *HST* image in part (d) shows a 2009 impact comparable in violence to the 1994 events. Since then, three more impacts have been observed—two in 2010 and one in 2012. Scientists speculate that such collisions may become an important future source of information about the planet's interior.

that storm is intensifying and might even come to rival the Great Red Spot some day. Merger and growth may well be the mechanism by which large storms on the jovian planets form and strengthen. Small storms that approach too close to larger ones are swept up and absorbed. Figure 11.11(c) shows a time sequence of *HST* images of the Great Red Spot, the Junior Spot, and the 2008 “baby,” which was shredded within a few months by the swirling winds of the two larger storms.


Despite these many mysteries, we can offer at least a partial explanation for the longevity of storm systems on Jupiter. On Earth, a large storm, such as a hurricane, forms over the ocean and may survive for many days, but it dies quickly once it encounters land. Earth's continental landmasses disrupt the flow patterns that sustain the storm. Jupiter has no continents, so once a storm becomes established and reaches a size at which other storm systems cannot destroy it, apparently little affects it. The larger the system, the longer its lifetime.

### CONCEPT Check

- ✓ List some similarities and differences between Jupiter's belts, zones, and spots, on the one hand, and weather systems on Earth, on the other.

## 11.3 Internal Structure

Much of our knowledge of Jupiter's interior comes from theoretical modeling. Indeed, apart from data gained following the collision of a comet with Jupiter in 1994 (see *Discovery 11-1*), we have very little direct evidence of the planet's internal properties. Planetary scientists use all available bulk data on the planet—mass, radius, composition, rotation, temperature, etc.—to construct a model of the interior that agrees with observations. Modeling is an integral part of the scientific method, and our statements about Jupiter's structure are really statements about the model that best fits the observed facts. ∞ (Sec. 1.2) However, because the planet



consists largely of hydrogen and helium—two simple gases whose physics we think we understand well—we can be fairly confident that Jupiter’s internal structure is now understood.

### An Internal Energy Source

On the basis of Jupiter’s distance from the Sun, astronomers had expected to find the temperature of the cloud tops to be around 105 K. At that temperature, they reasoned, Jupiter would radiate back into space exactly the same amount of energy as it received from the Sun. When radio and infrared observations were first made of the planet, however, astronomers found that its blackbody spectrum corresponded to a temperature of 125 K instead. Subsequent measurements, including those made by *Voyager* and *Galileo*, have verified that finding. Although a difference of 20 K may seem small, recall from Chapter 3 that the energy emitted by a planet grows as the *fourth* power of the surface temperature (in Jupiter’s case, the temperature of the cloud tops). [∞ \(Sec. 3.4\)](#)

A planet at 125 K therefore radiates  $(125/105)^4$ , or about twice as much energy as a planet at 105 K radiates. Put another way, Jupiter actually emits about twice as much energy as it receives from the Sun. Thus, unlike any of the terrestrial planets, *Jupiter must have its own internal source of heat.*

What is responsible for Jupiter’s extra energy? It is not the decay of radioactive elements within the planet. That process must be occurring, as in Earth, but estimates of the total amount of energy released into Jupiter’s interior are far below the levels needed to account for the temperature we measure. [∞ \(Sec. 7.3\)](#) Nor is it nuclear fusion, the process that generates energy in the Sun. The temperature in Jupiter’s interior, high as it is, is still far too low for that (see *Discovery 11-2*). Instead, astronomers theorize that the source of Jupiter’s excess energy is the slow escape of gravitational energy released during the planet’s formation. As the planet took shape, some of its gravitational energy was converted into heat in the interior. That heat is still slowly leaking out through the planet’s heavy atmospheric blanket, resulting in the excess emission we observe.

## DISCOVERY 11-2

### Almost a Star?

Jupiter has a starlike composition—predominantly hydrogen and helium, with a trace of heavier elements. Did Jupiter ever come close to becoming a star itself? Might the solar system have formed as a double-star system? Probably not. Unlike a star, Jupiter is cold. Its central temperature is far too low to ignite the nuclear fires that power our Sun (see Section 16.6). Jupiter's mass would have to increase 80-fold before its central temperature would rise to the point where nuclear reactions could begin, converting Jupiter into a small, dim star.

Even so, it is interesting to note that, although Jupiter's present-day energy output is very small (by solar standards, at least), it must have been much greater in the distant past, while the planet was still contracting rapidly toward its present size. For a brief period—perhaps a few hundred million years—Jupiter might actually have been as bright as a faint star, although its brightness never came within a factor of 100 of the Sun's. Still, seen from Earth at that time, Jupiter would have been about 100 times brighter than the Moon!

What might have happened had our solar system formed as a double-star system? Conceivably, had Jupiter


been massive enough, its radiation might have produced severe temperature fluctuations on all the planets, perhaps to the point of making life on Earth impossible. Even if Jupiter's brightness were too low to cause us any problems, its gravitational pull (which would be 1/12 that of the Sun if its mass were 80 times its present value) might have made the establishment of stable, roughly circular planetary orbits in the inner solar system an improbable event, again to the detriment of life on Earth.

Curiously, in recent years astronomers have come to realize that, had Jupiter been too *small*, that also could have adversely affected the chances for life on our planet! As you learned in Chapter 6, Jupiter played a crucial role in clearing debris from the outer solar system during and after the period when the planets formed. [∞ \(Sec. 6.7\)](#) Had that not occurred, the meteoritic bombardment of our planet might have been too severe and too extended for complex life ever to have evolved. [∞ \(Sec. 8.5\)](#) Many stars near the Sun are now known to have Jupiter-sized planets orbiting them. It seems that the size of the “Jupiter,” or second-largest body, in a newborn planetary system may be a critical factor in determining the likelihood of the appearance of life there.

Despite the huge amounts of energy involved—Jupiter emits about  $4 \times 10^{17}$  watts more energy than it receives from the Sun—the loss is slight compared with the planet's total energy. On the basis of the planet's mass and temperature, as well as the rate at which thermal energy is leaving the planet, astronomers calculate that the average temperature of the interior of Jupiter decreases by only about a millionth of a kelvin per year. [∞ \(More Precisely 3-1\)](#)

### Jupiter's Deep Interior

Jupiter's clouds, with their complex chemistry, are probably less than 200 km thick. Below them, the temperature and pressure steadily increase as the atmosphere becomes the “interior” of the planet.



Both the temperature and the density of Jupiter's atmosphere increase with depth below the cloud cover. However, no “surface” of any kind exists anywhere inside. Instead, Jupiter's atmosphere just becomes denser and denser because of the pressure of the overlying layers. At a depth of a few thousand kilometers, the gas makes a gradual transition into the liquid state (see Figure 11.12). By a depth of about 20,000 km, the pressure is about 3 million times greater than atmospheric pressure on Earth. Under those conditions, the hot liquid hydrogen is compressed so much that it undergoes another transition, this time to a “metallic” state with properties in many ways similar to those of a liquid metal. Of particular importance for Jupiter's magnetic field (see Section 11.4) is that this metallic hydrogen is an excellent conductor of electricity.

As mentioned earlier, Jupiter's observed flattening requires that there be a relatively small (i.e., relatively small compared with the size of Jupiter), dense core at its center. On the basis of *Voyager* data, scientists once thought that the core might contain as much as 20 Earth masses of material.

**◀ FIGURE 11.12 Jupiter's Interior** Jupiter's internal structure, as deduced from spacecraft measurements and theoretical modeling. Pressure and temperature increase with depth, and the atmosphere gradually liquefies at a depth of a few thousand kilometers. Below 20,000 km, the hydrogen behaves like a liquid metal. At the center of the planet lies a large rocky core, somewhat terrestrial in composition, but much larger than any of the terrestrial planets.

However, following *Galileo's* arrival, it now appears that the core's mass could be as low as 5 Earth masses and perhaps even less. The precise composition of the core is unknown, but planetary scientists think that it contains much denser materials than the rest of the planet.

Current best estimates indicate that the core consists of “rocky” materials, similar to those found on the terrestrial worlds. (Note that the term *rocky* here refers to the *chemical composition* of the core, not to its physical state. At the high temperatures and pressures found deep in the jovian interiors, the core material bears little resemblance to rocks found on Earth's surface.) In fact, it now appears that all four jovian planets contain similarly large rocky cores and that the formation of such a large “terrestrial” planetary core may be a necessary stage in the process of building up a gas giant (see Section 15.2).

Because of the enormous pressure at the center of Jupiter—approximately 60 million times that on Earth's surface, or 12 times that at Earth's center—the core must be compressed to a very high density (perhaps twice the density of Earth's core). The jovian core is probably not much more than 20,000 km in diameter (still big enough for Earth to fit inside, with plenty of room left over), and the central temperature may be as high as 25,000 K.


### PROCESS OF SCIENCE Check

- ✓ How have astronomers determined the properties of Jupiter's core?

## 11.4 Jupiter's Magnetosphere

For decades, ground-based radio telescopes monitored radiation leaking from Jupiter's magnetosphere, but only when the *Pioneer* and *Voyager* spacecraft reconnoitered the planet in the mid-1970s did astronomers realize the full extent of its magnetic field. The *Galileo* probe spent many years orbiting within Jupiter's magnetosphere, returning a wealth of detailed information about its structure.


Jupiter, it turns out, is surrounded by a vast sea of energetic charged particles, mostly electrons and protons, somewhat similar to Earth's Van Allen belts, but much, much larger. The radio radiation detected on Earth is emitted when these particles are accelerated to very high speeds—close to the speed of light—by Jupiter's powerful magnetic field. This radiation is several thousand times more intense than that produced by Earth's magnetic field. The particles present a serious hazard to manned and unmanned space vehicles alike. Sensitive electronic equipment (not to mention even more sensitive human bodies) requires special protective shielding to operate for long in this hostile environment. *Galileo* was not expected to survive as long as it did.



▲ **FIGURE 11.13 Pioneer Mission** The *Pioneer 10* spacecraft (a forerunner of the *Voyager* missions) did not detect any solar particles while moving far behind Jupiter in 1976. Apparently, as sketched here, the tail of Jupiter's magnetosphere extends beyond the orbit of Saturn.

Direct measurements from spacecraft show Jupiter's magnetosphere to be almost 30 million kilometers across, roughly a million times more voluminous than Earth's magnetosphere and far larger than the entire Sun. As with Earth's, the size and shape of Jupiter's magnetosphere are determined by the interaction between the planet's magnetic field and the solar wind. Jupiter's magnetosphere has a long tail extending away from the Sun at least as far as Saturn's orbit (over 4 AU farther out from the Sun), as sketched in Figure 11.13. However, on the sunward side, the *magnetopause*—the boundary of Jupiter's magnetic influence on the solar wind—lies only 3 million kilometers from the planet. Near Jupiter's surface, the magnetic field channels particles from the magnetosphere into the upper atmosphere, forming aurorae vastly larger and more energetic than those observed on Earth (Figure 11.14). ∞ (Sec. 7.5)


The outer magnetosphere of Jupiter appears to be quite unstable, sometimes deflating in response to “gusts” in the solar wind and then reexpanding as the wind subsides. In the inner magnetosphere, Jupiter's rapid rotation has forced most of the charged particles into a flat *current sheet*, lying on the planet's magnetic equator, quite unlike the Van Allen belts surrounding Earth. ∞ (Sec. 7.5) The portion of the magnetosphere close to Jupiter is sketched



◀ **FIGURE 11.14 Aurorae on Jupiter** The main (underlying) image was taken by the *Hubble* telescope in visible (true-color) light, but the two insets at the poles were taken in the ultraviolet part of the spectrum. The oval-shaped aurorae, extending hundreds of kilometers above Jupiter's surface, result from charged particles escaping the jovian magnetosphere and colliding with the atmosphere, causing the gas to glow. (NASA)

in Figure 11.15. Notice that the planet's magnetic axis is not exactly aligned with its rotation axis, but is inclined to it at an angle of approximately  $10^\circ$ . Jupiter's magnetic field happens to be oriented opposite Earth's, with field lines running from north to south, rather than south to north as in the case of our own planet (see Figure 7.18).

Both ground- and space-based observations of the radiation emitted from Jupiter's magnetosphere imply that the *intrinsic* strength of the planet's magnetic field is nearly 20,000 times greater than Earth's. The existence of such a strong field further supports our theoretical model of Jupiter's internal structure. The conducting liquid interior that is thought to make up most of the planet should combine with Jupiter's rapid rotation to produce a large dynamo effect and a strong magnetic field, just as are observed. ∞ (Sec. 7.5)



▲ **FIGURE 11.15 Jupiter's Magnetosphere** Jupiter's inner magnetosphere is characterized by a flat current sheet (white dashed line) consisting of charged particles squeezed into the magnetic equatorial plane by the planet's rapid rotation. The plasma torus (green tube), a ring of charged particles associated with the moon Io, is discussed in Section 11.5. Compare this figure with Earth's magnetosphere in Figure 7.18.

**CONCEPT Check**


✓ Why is Jupiter's magnetosphere so much larger than Earth's?

**11.5 The Moons of Jupiter**

As of early 2013, Jupiter's official satellite count stands at 67. Table 11.1 presents some properties of the four largest members of Jupiter's moon system—the Galilean moons. The 63 small bodies not listed in the table all have diameters less than 300 km.

The Galilean moons are each comparable in size to Earth's Moon. ∞ (Sec. 2.4) Moving outward from Jupiter, the four are named Io, Europa, Ganymede, and Callisto, after the mythical attendants of the Roman god Jupiter. As illustrated in Figure 11.16, they move in nearly circular orbits about their parent planet. When the *Voyager 1* spacecraft passed close to the Galilean moons in 1979, it sent some remarkably detailed photographs back to Earth, allowing planetary scientists to discern fine surface features on each moon. ∞ (Discovery 6-2) More recently, in the late 1990s, the *Galileo* mission expanded our knowledge of these small, but complex, worlds still further. We will consider the Galilean satellites in more detail momentarily.

Within the orbit of Io lie four small satellites, all but one discovered by *Voyager* cameras. The largest of the four, Amalthea, is 260 km across and irregularly shaped. Discovered by American astronomer E. E. Barnard in 1892, it orbits at a distance of 181,000 km from Jupiter's center—only 110,000 km above the cloud tops. Like the Galilean moons, all four inner satellites have roughly circular, prograde orbits, and they rotate synchronously with their orbital motion due to Jupiter's strong tidal field.



**Interactive FIGURE 11.16 Galilean Moons** The orbits of Jupiter's Galilean moons, drawn to scale, as seen from above the planet's north pole. The four inserts show actual images of those moons, taken by the *Galileo* spacecraft and scaled here as they would appear from a distance of about 1 million kilometers. (NASA)

The remaining 59 small satellites lie well outside the Galilean moons. All have been discovered since the start of the 20th century, most since the late 1990s, thanks to painstaking observations using large Earth-based telescopes with specially designed instruments and software to scan large areas of the sky for very faint objects. Steadily improving technology means that small bodies once far too faint for Earth-based telescopes to see

**TABLE 11.1** The Major Moons of Jupiter\*

Name	Distance from Jupiter (km)	(planetary radii)	Orbital Period (days)	Size (longest diameter, km)	Mass** (Earth Moon masses)	Density (kg/m <sup>3</sup> )	(g/cm <sup>3</sup> )
Io	422,000	5.90	1.77	3640	1.22	3500	3.5
Europa	671,000	9.38	3.55	3130	0.65	3000	3.0
Ganymede	1,070,000	15.0	7.15	5270	2.02	1900	1.9
Callisto	1,880,000	26.3	16.7	4800	1.46	1900	1.9

\* This table does not include the 63 small moons described in the text. All of these small moons are less than 300 km in diameter (most are less than 10 km across).

\*\* Mass of Earth's Moon =  $7.4 \times 10^{22}$  kg =  $3.9 \times 10^{-5}$  Jupiter masses.





are now being detected and cataloged almost routinely. As a class, these outer moons are small—mostly less than a few tens of kilometers across—and move in eccentric, predominantly retrograde, orbits far from the planet. Their masses, and hence their densities, are unknown. However, their appearance and size suggest compositions more like asteroids or comets than their larger Galilean companions. Most astronomers think that these satellites did not form along with Jupiter, but instead are bodies that were captured by Jupiter’s strong gravitational field long after the planet and its larger inner moons originally formed.


## The Galilean Moons: A Model of the Inner Solar System

Jupiter’s Galilean moons have several interesting parallels with the terrestrial planets. Their orbits are direct (i.e., in the same sense as Jupiter’s rotation), are roughly circular, and lie close to Jupiter’s equatorial plane. They range in size from slightly smaller than Earth’s Moon (Europa) to slightly larger than Mercury (Ganymede). Figure 11.17 is a *Voyager 1* image of Io and Europa, with Jupiter providing a spectacular backdrop. Figure 11.18 shows the four Galilean moons to scale.

The similarity to the inner solar system continues with the fact that the moons’ densities decrease with increasing distance from Jupiter. ∞ (Sec. 6.4) Largely on the basis of detailed measurements made by *Galileo* of the moons’ gravitational fields, together with mathematical models of the interiors, researchers have built up fairly detailed pictures of each moon’s composition and internal structure



▲ **FIGURE 11.17 Jupiter, Up Close** *Voyager 1* took this photo of Jupiter with ruddy Io on the left and pearl-like Europa toward the right. Note the scale of objects here: Both Io and Europa are comparable in size to our Moon, and the Red Spot is roughly twice as big as Earth. (NASA)



*Europa and Ganymede may have layers of liquid water beneath their icy surfaces.*

▲ **FIGURE 11.18 Galilean Moon Interiors** Cutaway diagrams of the interior structure of the four Galilean satellites. Moving outward from Io to Callisto, densities steadily decrease as the composition shifts from rocky mantles and metallic cores in Io and Europa, to a thick icy crust and smaller core in Ganymede, to an almost uniform rock and ice mix in Callisto.

(Figure 11.18). The innermost two Galilean moons, Io and Europa, have thick rocky mantles, possibly similar to the crusts of the terrestrial planets, surrounding iron–iron sulfide cores. Io’s core accounts for about half that moon’s total radius. Europa has a water–ice outer shell between 100 and 200 km thick. The two outer moons, Ganymede and Callisto, are clearly deficient in rocky materials. Lighter materials, such as water and ice, may account for as much as half of their total mass. Ganymede appears to have a

relatively small metallic core topped by a rocky mantle and a thick icy outer shell. Callisto seems to be a largely undifferentiated mixture of rock and ice.

Many astronomers think that the formation of Jupiter and the Galilean satellites may in fact have mimicked, on a small scale, the formation of the Sun and the inner planets. For that reason, studies of the Galilean moon system could provide us with valuable insight into the processes that created our own world. We will return to this parallel in Chapter 15. So interested were mission planners in learning more about the Galilean moon system that the already highly successful *Galileo* mission was extended for 6 more years to allow for even more detailed study, particularly of Europa. ∞ (Discovery 6-2) The Galilean moons were scrutinized at resolutions as fine as a few meters during numerous extremely close passages by the spacecraft.

Not all the properties of the Galilean moons find analogs in the inner solar system, however. For example, all four Galilean satellites are locked into states of synchronous rotation by Jupiter's strong tidal field, so they all keep one face permanently pointing toward their parent planet. By contrast, of the terrestrial planets, only Mercury is strongly influenced by the Sun's tidal force, and even its orbit is not synchronous. ∞ (Sec. 8.4) Finally, inspection of Table 11.1 shows a remarkable coincidence in the orbital periods of the three inner Galilean moons: Their periods are almost exactly in the ratio 1:2:4 (and the fourth moon Callisto is

not too far from being the “8” in the sequence). This configuration may be the result of a complex, but poorly understood, three-body (or perhaps even four-body) resonance in the Galilean moon system, something not found among the terrestrial worlds.


## Io: The Most Active Moon

Io, the densest of the Galilean moons, is the most geologically active object in the entire solar system. Its mass and radius are fairly similar to those of Earth's Moon, but there the resemblance ends. Shown in Figure 11.19, Io's surface is a collage of reds, yellows, and blackish browns—resembling a giant pizza in the minds of some startled *Voyager* scientists. As the spacecraft sped past Io, it made an outstanding discovery: Io has active volcanoes! *Voyager 1* photographed eight erupting volcanoes. Six were still erupting when *Voyager 2* passed by 4 months later.


By the time *Galileo* arrived in 1995, several of the eruptions observed by *Voyager* had subsided. However, many new ones were seen—in fact, *Galileo* found that Io's surface features can change significantly in as little as a few weeks. In all, more than 80 active volcanoes have been identified on Io. The largest, called Loki (on the far side of Figure 11.20), is larger than the state of Maryland and emits more energy than all of Earth's volcanoes combined.




(a)



(b)



▲ **FIGURE 11.19** Io Jupiter's innermost moon, Io, is quite different in character from the other three Galilean satellites. Its surface is kept smooth and brightly colored by the moon's constant volcanism. The resolution of the *Galileo* photograph in (a) is about 7 km. In the more detailed *Voyager* image (b), features as small as 2 km across can be seen. (NASA)



**Interactive** **FIGURE 11.20** **Volcanoes on Io** The main image shows a *Galileo* view of Io, whose surface is kept smooth and brightly colored by constant volcanism, revealed here as dark, circular features. The left inset shows an umbrella-like eruption of one of Io's volcanoes as *Galileo* flew past this fascinating moon in 1997; the plume measures about 150 km high and 300 km across. The right inset shows another volcano, this one face-on, where surface features here are resolved to just a few kilometers. (NASA)


The top right inset in Figure 11.20 shows a volcano called Prometheus ejecting matter at speeds of up to 2 km/s to an altitude of about 150 km. These high-speed gases are quite unlike the (relatively) sluggish ooze that emanates from Earth's volcanoes. According to *Galileo*'s instruments, lava temperatures on Io generally range from 650 to 900 K, with the higher end of the range implying that at least some of the volcanism is similar to that found on Earth. However, temperatures as high as 2000 K—far hotter than any earthly volcano—have been measured at some locations. Mission scientists speculate that these “superhot” volcanoes may be similar to those that occurred on Earth more than 3 billion years ago.

The orange color immediately surrounding the volcanoes most likely results from sulfur compounds in the ejected material. In stark contrast to the surfaces of the other Galilean moons, Io's surface is neither cratered nor streaked (the circular features visible in Figures 11.19 and 11.20 are volcanoes), but is instead exceptionally smooth, mostly varying in altitude by less than about 1 km, although some volcanoes are several kilometers high. The smoothness is apparently the result of molten matter that constantly fills in any “dents and cracks.” This remarkable moon has the youngest surface of any known object in the solar system. Io also has a thin, temporary atmosphere made up primarily of sulfur dioxide, presumably the result of gases ejected by volcanic activity.

Io's volcanism has a major effect on Jupiter's magnetosphere. All the Galilean moons orbit within the magnetosphere and play some part in modifying its properties, but Io's influence is particularly marked. Although many of the charged particles in Jupiter's magnetosphere come from the solar wind, there is strong evidence that Io's volcanism is the primary source of heavy ions in the inner regions. Jupiter's magnetic field continually sweeps past Io, gathering up the particles its volcanoes spew into space and accelerating them to high speed. The result is the *Io plasma torus* (Figure 11.21; see also Figure 11.15), a doughnut-shaped region of energetic heavy ions that follows Io's orbital track, completely encircling Jupiter. (A plasma is a gas that has been heated to such high temperatures that all its atoms are ionized. A few neutral atoms have also been observed in the Io plasma torus.)

The plasma torus is quite easily detectable from Earth, but before *Voyager* its origin was unclear. *Galileo* made detailed studies of the plasma's dynamic and rapidly varying magnetic field. Spectroscopic analysis shows that sulfur is indeed one of the torus's major constituents, strongly implicating Io's volcanoes as its source. As a hazard to spacecraft—manned or unmanned—the plasma torus is formidable, with lethal radiation levels.

What causes such astounding volcanic activity on Io? The moon is far too small to have geological activity like that



▲ **FIGURE 11.21 Io Plasma Torus** Io's torus is the result of material ejected from Io's volcanoes and swept up by Jupiter's rapidly rotating magnetic field. Spectroscopic analysis indicates that the torus is made mainly of sodium and sulfur atoms and ions.

on Earth. Io should be long dead, like our own Moon. At one time, some scientists suggested that Jupiter's magnetosphere might be the culprit: Perhaps the (then-unknown) processes creating the plasma torus were somehow also stressing the moon. We now know that this is not the case. The real source of Io's energy is *gravity*—Jupiter's gravity. Io orbits very close to Jupiter—only 422,000 km, or 5.9 Jupiter radii, from the center of the planet. As a result, Jupiter's huge gravitational field exerts strong tidal forces on the moon. If Io were the only satellite in the Jupiter system, it would long ago have come into a state of synchronous rotation with the planet, just as our own Moon has with Earth, for the reasons discussed in Chapter 8. ∞ (Sec. 8.4) In that case, Io would move in a perfectly circular orbit, with one face permanently turned toward Jupiter, and the tidal bulge would be stationary with respect to the moon.


But Io is not alone. As it orbits, it is constantly tugged by the gravity of its nearest large neighbor, Europa. The tugs are small and not enough to cause any great tidal effect, but they are sufficient to make Io's orbit slightly noncircular, preventing the moon from settling into a precisely synchronous state. The reason for this effect is exactly the same as in the case of Mercury, also as discussed in Chapter 8. ∞ (Sec. 8.4) In a noncircular orbit, the moon's speed varies from place to place as it revolves around its planet, but its rate of rotation on its axis remains constant. Thus, it cannot keep one face always

turned toward Jupiter. Instead, as seen from Jupiter, Io rocks or “wobbles” slightly from side to side as it moves. The large (100 m) tidal bulge, however, always points directly toward Jupiter, so it moves back and forth across Io's surface as the moon wobbles. These conflicting forces result in enormous tidal stresses that continually flex and squeeze Io's interior.

Just as the repeated back-and-forth bending of a piece of wire can produce heat through friction, the ever-changing distortion of Io's interior constantly energizes the moon. This generation of large amounts of heat within Io ultimately causes huge jets of gas and molten rock to squirt out of the moon's surface. *Galileo's* sensors indicated extremely high temperatures in the outflowing material. It is likely that much of Io's interior is soft or molten, with only a thin solid crust overlying it. Researchers estimate that the total amount of heat generated within Io as a result of tidal flexing is about 100 million megawatts. This phenomenon makes Io one of the most fascinating objects in our solar system.

## Europa: Liquid Water Locked in Ice

Europa (Figure 11.22) is a world very different from Io. Lying outside Io's orbit, 671,000 km (9.4 Jupiter radii) from Jupiter, Europa showed relatively few craters on its surface in images taken by *Voyager*, suggesting geologic youth—perhaps just



**▲ FIGURE 11.22 Europa** (a) *Voyager 1* mosaic of Europa, showing resolution of about 5 km. (b) Europa's icy surface is only lightly cratered, indicating that some ongoing process must be obliterating impact craters soon after they form. The origin of the cracks crisscrossing the surface is uncertain. (c) A closer look from the *Galileo* spacecraft shows a smooth yet tangled surface, called Conamara Chaos, resembling the huge ice floes that cover Earth's polar regions. (d) This detailed (5-meter-resolution) *Galileo* image shows "pulled apart" terrain that suggests liquid water upwelling from the interior and freezing, filling in the gaps between separating surface ice sheets. (NASA)

a few million years. Recent activity has erased any scars of ancient meteoritic impacts. The dark areas are rocky deposits that may have come from the moon's interior or that may have been swept up by Europa as it moved in its orbit. Europa's surface also displays a vast network of lines crisscrossing bright, clear fields of water ice. Some of these linear "bands," or fractures, extend halfway around the satellite and resemble, in some ways, the pressure ridges that develop in ice floes on Earth's polar oceans.

On the basis of *Voyager* images, planetary scientists had theorized that Europa might be completely covered by an ocean of liquid water with its top frozen at the low temperatures prevailing so far from the Sun. In this view, the cracks in the surface are attributed to the tidal influence of Jupiter and the gravitational pulls of the other Galilean satellites, although these forces are considerably weaker than those powering Io's violent volcanic activity. However, other researchers had contended that Europa's fractured surface was instead related to some form of tectonic activity—involving ice rather than rock. High-resolution *Galileo* observations strongly support the former idea. Figure 11.22(c) is a *Galileo* image of this weird moon, showing what look like icebergs—flat chunks of ice that have been broken apart, moved several kilometers, and reassembled, perhaps by the action of water currents below. Mission scientists estimate that Europa's surface ice may be several kilometers thick, with a 100-km-deep liquid ocean below it.

Other detailed images of Europa's surface lend further support to this hypothesis. Figure 11.22(d) shows a region where Europa's icy crust appears to have been pulled apart and new material has filled in the gaps between the separating ice sheets. Elsewhere on the surface, *Galileo* found what appeared to be the icy equivalent of lava flows on Earth—regions where water apparently "erupted" through the surface and flowed for many kilometers before solidifying. The smooth "trenches" shown in Figure 11.22(d) strongly suggest local flooding of the terrain. The scarcity of impact craters on Europa implies that the processes responsible for these features did not stop long ago. Rather, they must be ongoing.

Further evidence comes from studies of Europa's magnetic field. Measurements made by *Galileo* on repeated flybys of the moon revealed that Europa has a weak magnetic field that constantly changes strength and direction. This finding is consistent with the idea that the field is generated by the action of Jupiter's magnetism on a shell of electrically conducting fluid about 100 km below Europa's surface—in other words, the salty layer of liquid water suggested by the surface observations. This discovery convinced quite a few skeptical scientists of the reality of Europa's ocean.

The likelihood that Europa has an extensive layer of liquid water below its surface ice opens up many interesting avenues of speculation about the possible development of life there. In the rest of the solar system, only Earth has liquid water on or near its surface, and most scientists agree that water played a key role in the appearance of life on Earth (see

Chapter 28). Europa may well contain more liquid water than exists on our entire planet! Of course, the existence of water does not *necessarily* imply the emergence of life. Europa, even with its liquid ocean, is still a hostile environment compared with Earth (although recent laboratory studies suggest that icy chemical reactions in Europa's dark, frigid depths may proceed much faster than had previously been thought possible). Nevertheless, the possibility—even a remote one—of life on Europa was an important motivating factor in the decision to extend the *Galileo* mission for 6 more years.

### PROCESS OF SCIENCE Check


✓ Why are scientists so interested in the existence of liquid water on Europa and Ganymede?

## Ganymede and Callisto: Fraternal Twins

The two outermost Galilean moons are Ganymede (at 1.1 million kilometers, or 15 planetary radii, from the center of Jupiter) and Callisto (at 1.9 million kilometers, or 26 Jupiter radii). The density of each is only about  $2000 \text{ kg/m}^3$ , suggesting that they harbor substantial amounts of ice throughout and are not just covered by thin icy or snowy surfaces. Ganymede, shown in Figure 11.23, is the largest moon in the solar system, exceeding not only Earth's Moon, but also the planet Mercury in size. It has many impact craters on its surface and patterns of dark and light markings that are reminiscent of the highlands and maria on Earth's own Moon. In fact, Ganymede's history has many parallels with that of the Moon (with water ice replacing lunar rock). The large, dark region clearly visible in Figure 11.23(a) is called Galileo Regio.

As on the inner planets, we can estimate ages on Ganymede by counting craters. We learn that the darker regions, such as Galileo Regio (marked in Figure 11.23a), are the oldest parts of Ganymede's surface. These regions are the original icy surface of the moon, just as the ancient highlands on our own Moon are its original crust. The surface darkens with age as micrometeorite dust slowly covers it. The light-colored parts of Ganymede are much less heavily cratered, so they must be younger. They are Ganymede's "maria" and probably formed in a manner similar to the way that maria on the Moon were created. ∞ (Sec. 8.5) Intense meteoritic bombardment caused liquid water—Ganymede's counterpart to our own Moon's molten lava—to upwell from the interior and flood the affected regions before solidifying.

Not all of Ganymede's surface features follow the lunar analogy. Ganymede has a system of grooves and ridges (shown in Figure 11.23c) that may have resulted from crustal tectonic motion, much as Earth's surface undergoes mountain building and faulting at plate boundaries. ∞ (Sec. 7.4) Ganymede's large size indicates that its original radioactivity probably helped heat and differentiate its interior, after which the moon cooled and the crust cracked. Ganymede seems to have had some early plate tectonic



▲ **FIGURE 11.23 Ganymede** (a) and (b) In these *Voyager 2* images of Ganymede, the dark regions are the oldest parts of the moon's surface and might represent its original icy crust. The largest dark region visible here, called Galileo Regio, spans some 3200 km. The lighter, younger regions are the result of flooding and freezing that occurred within a billion years or so of Ganymede's formation. The brightest-colored spots are recent impact craters. (c) Grooved terrain on Ganymede may have been caused by a process similar to plate tectonics on Earth. The area shown in this *Galileo* image spans about 50 km and reveals a multitude of ever-smaller ridges, valleys, and craters, down to the resolution of a football field. (NASA)

activity, but the process stopped about 3 billion years ago, when the cooling crust became too thick for such activity to continue. The *Galileo* data suggest that the surface of Ganymede may be older than was previously thought. With the improved resolution of that spacecraft's images (Figure 11.23c), some regions thought to have been smooth, and hence young, are now seen to be heavily splintered by fractures and thus probably very old.

In 1996, *Galileo* detected a weak magnetosphere surrounding Ganymede, making it the first moon in the solar system on which a magnetic field had been observed and implying that Ganymede has a modest iron-rich core. Ganymede's magnetic field is about 1 percent that of Earth. In December 2000, the magnetometer team reported fluctuations in the field strength similar to those near Europa, suggesting that Ganymede, too, may have liquid or perhaps "slushy" water under its surface. Recent observations of surface formations similar to those attributed to flowing water "lava" on Europa appear to support this view.


Callisto, shown in Figure 11.24, is in many ways similar in appearance to Ganymede, although it has more craters and fewer fault lines. Its most obvious feature is a huge series of concentric ridges surrounding each of two large basins. The larger of the two, on Callisto's Jupiter-facing side, is named Valhalla and measures some 3000 km across. It is clearly visible in the Figure. The ridges resemble the ripples made as a stone hits water, but on Callisto they probably resulted from a cataclysmic impact with an asteroid or comet. The upthrust

ice was partially melted, but it resolidified quickly, before the ripples had a chance to subside.

Today, both the ridges and the rest of the crust are frigid ice and show no obvious signs of geological activity (such as the grooved terrain on Ganymede). Apparently, Callisto froze before plate tectonic or other activity could start. The density of impact craters on the Valhalla basin indicates that it formed long ago, perhaps 4 billion years in the past. Yet, even on this frozen world, there are hints from *Galileo*'s magnetometers that there might be a thin layer of water, or more likely slush, deep below the surface.

Ganymede's internal differentiation indicates that the moon was largely molten at some time in the past; Callisto is undifferentiated and hence apparently never melted. Researchers are uncertain why two such similar bodies should have evolved so differently. Complicating matters further is Ganymede's magnetic field and possible subsurface liquid water, which suggest that the moon's interior may still be relatively warm. If that is so, then Ganymede's heating and differentiation must have happened quite recently—less than a billion years ago, based on recent estimates of how rapidly the moon's heat escapes into space.

Scientists have no clear explanation for how Ganymede could have evolved in this manner. Heating by meteoritic bombardment ended too early, and radioactivity probably could not have provided enough energy at such a late time. ∞ (Sec. 7.3) Some astronomers speculate that interactions among the inner moons, possibly related to the 1:2:4 near



▲ **FIGURE 11.24 Callisto** (a) Callisto resembles Ganymede in overall composition but is more heavily cratered. Extending nearly 1500 km from the Valhalla basin center, its concentric ridges formed when "ripples" from a large meteoritic impact froze before they could disperse completely. (b) This higher-resolution *Galileo* image of Callisto's equatorial region displays more clearly its heavy cratering. (NASA)


resonance mentioned earlier, may have been responsible. These interactions might have caused Ganymede's orbit to change significantly about a billion years ago, and prior tidal heating by Jupiter could have helped melt the moon's interior.

### CONCEPT Check

- ✓ What is the ultimate source of all the activity observed on Jupiter's Galilean satellites?

## 11.6 Jupiter's Ring

Yet another remarkable finding of the 1979 *Voyager* missions was the discovery of a faint ring of matter encircling Jupiter in the plane of the planet's equator (see Figure 11.25). The ring lies roughly 50,000 km above the top cloud layer of the planet, inside the orbit of the innermost moon. A thin sheet of material may extend all the way down to Jupiter's cloud tops, but most of the ring is confined within a region only a few thousand kilometers across. The outer edge of the ring is quite sharply defined. In the direction perpendicular to the equatorial plane, the ring is only a few tens of kilometers thick. The small, dark particles that make up the ring may be fragments chipped off by meteoritic impact from two small moons—Metis and Adrastea,



▲ **FIGURE 11.25 Jupiter's Ring** Jupiter's faint ring was captured nearly edge-on by *Voyager 2*'s cameras. Made of dark fragments of rock and dust possibly chipped off the innermost moons by meteorites, the ring was unknown before the two *Voyager* spacecraft arrived at the planet. (NASA)

discovered by *Voyager*—that lie very close to the ring itself. Despite differences in appearance and structure, Jupiter's ring can perhaps be best understood by studying the most famous planetary ring system—that of Saturn—so we will postpone further discussion of ring properties until the next chapter.

**The Big Question** Perhaps the most enduring question about Jupiter concerns the Red Spot, which is larger than Earth itself. What caused this huge feature to form in Jupiter's atmosphere, how has it continued swirling for hundreds of years, and why is it red-orange? Perhaps the spot is merely a gaseous vortex resembling a hurricane on Earth—yet it oddly remains in nearly the same location relative to its interior, unlike Earth's storms that move with weather patterns. Or maybe some underlying structure lurks beneath the storm, an atmospheric anomaly that somehow traps the turbulent gas above it. What's needed is a robot space probe that would fly directly into the Red Spot, radioing its secrets back to Earth.

# Chapter Review

## SUMMARY

1 Jupiter is the largest planet in the solar system. Its mass is more than twice the mass of all the other planets combined, although still only 1/1000 the mass of the Sun. Composed primarily of hydrogen and helium, Jupiter rotates rapidly, producing a pronounced equatorial bulge. The planet's flattened shape allows astronomers to infer the presence of a large rocky core in its interior. Unlike the terrestrial planets, Jupiter displays **differential rotation** (p. 267): the planet has no solid surface, and the rotation rate varies from



place to place in the atmosphere. Measurements of radio emission from Jupiter's magnetosphere provide a measure of the planet's interior rotation rate. Jupiter has many moons and a faint, dark ring extending down to the planet's cloud tops.




2 Jupiter's atmosphere consists of three main cloud layers, forming bands of bright **zones** (p. 268) and darker **belts** (p. 268) crossing the planet parallel to the equator. The bands are the result of convection in Jupiter's interior and the planet's rapid rotation. The lighter zones are the tops of upwelling, warm currents, and the darker belts are cooler regions where gas is sinking. Underlying them is a stable pattern of eastward or westward wind flow called



a **zonal flow** (p. 269). The wind direction alternates as we move north or south away from the equator. The colors we see are the result of chemical reactions, fueled by the planet's interior heat, solar ultraviolet radiation, auroral phenomena, and lightning. The main weather pattern on Jupiter is the **Great Red Spot** (p. 268), an Earth-sized hurricane that has been raging for at least three centuries. Smaller weather systems—**white ovals** (p. 272), **brown ovals** (p. 272), and new red spots—are also observed and can last for decades.



3 Jupiter's atmosphere becomes hotter and denser with depth, eventually becoming liquid. Interior pressures are so high that the hydrogen is “metallic” in nature near the center. The planet has a large “terrestrial” core about 10 times the mass of Earth. Jupiter radiates about twice as much energy into space as it receives from the Sun. The source of this energy is most likely heat released into the planet's interior when Jupiter formed 4.6 billion years ago, now slowly leaking from the surface.

4 The magnetosphere of Jupiter is about a million times more voluminous than Earth's magnetosphere, and the planet has a long magnetic “tail” extending away from the Sun to at least the distance of Saturn's orbit. Energetic particles spiral around magnetic field lines, accelerated by Jupiter's rotating magnetic field, producing intense radio radiation.



5 Jupiter and its system of moons resemble a small solar system. Sixty-three moons have been discovered so far. The outermost eight resemble asteroids and have retrograde orbits, suggesting that they may have been captured by Jupiter's gravity long after the planets and largest moons formed. Jupiter's four major moons are called the **Galilean moons** (p. 266), after their discoverer, Galileo Galilei. Their densities decrease with increasing distance from the planet. The innermost Galilean moon, Io, has active volcanoes and a smooth surface. Europa has a cracked, icy surface that probably conceals an ocean of liquid water, making the moon an interesting candidate for life in the solar system. Ganymede and Callisto have ancient, heavily cratered surfaces. Ganymede, the largest moon in the solar system, shows evidence of past geological activity, but now appears to be solid rock and ice, although recent evidence suggests that it, too, may have subsurface liquid water. Callisto apparently froze before tectonic activity could start there.

6 Io's volcanism is powered by the constant flexing of the moon by Jupiter's tidal forces. As Io orbits Jupiter, the moon “wobbles” because of the gravitational pull of Europa. The ever-changing distortion of its interior energizes Io, and geyserlike volcanoes keep its surface smooth with constant eruptions. Europa's fields of ice are nearly devoid of craters, but have extensive large-scale fractures, due most likely to the tidal influence of Jupiter, the gravitational effects of the other Galilean satellites, and the action of the underlying ocean.



**MasteringAstronomy**® For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information.

**LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- LO1** In terms of orbit and bulk properties, how does Jupiter compare to Earth?
- What is differential rotation, and how is it observed on Jupiter?
- POS** Describe some of the ways in which the *Voyager* and *Galileo* missions changed our perception of Jupiter.
- What is the Great Red Spot? What is known about the source of its energy?
- LO2** What is the cause of the colors in Jupiter's atmosphere?
- Why has Jupiter retained most of its original atmosphere?
- LO3 POS** Explain the theory that accounts for Jupiter's internal heat source.
- What is Jupiter thought to be like beneath its clouds? Why do we think this?
- LO4** What is responsible for Jupiter's enormous magnetic field?
- LO5** How does the density of the Galilean moons vary with increasing distance from Jupiter? Is there a trend to this variation? If so, why?
- LO6** What is the cause of Io's volcanic activity?
- What evidence do we have for liquid water below Europa's surface?
- How does the amount of cratering vary among the Galilean moons? Does it depend on their location? If so, why?
- POS** Why is there speculation that the Galilean moon Europa might be an abode for life?
- What might be the consequences of the discovery of life on Europa?

## Conceptual Self-Test: Multiple Choice

- Compared with Earth's orbit, the orbit of Jupiter is approximately (a) half as large; (b) twice as large; (c) 5 times larger; (d) 10 times larger.
- Compared with Earth's density, the density of Jupiter is (a) much greater; (b) much less; (c) about the same.
- The main constituent of Jupiter's atmosphere is (a) hydrogen; (b) helium; (c) ammonia; (d) carbon dioxide.
- VIS** Figure 11.6 ("Zonal Flow") shows that the most rapid westerly wind flows on Jupiter occur at (a) northern midlatitudes; (b) equatorial latitudes; (c) southern mid-latitudes; (d) polar latitudes.
- VIS** According to Figure 11.7 ("Jupiter's Atmosphere"), if ammonia and ammonium hydrosulfide ice were transparent to visible light, Jupiter would appear (a) bluish; (b) red; (c) tawny brown; (d) exactly as it does now.
- Jupiter's rocky core is (a) smaller than Earth's Moon; (b) comparable in size to Mars; (c) almost the same size as Venus; (d) larger than Earth.
- Jupiter's magnetosphere extends far into space, stretching (a) 1 AU; (b) 5 AU; (c) 10 AU; (d) 20 AU beyond the planet.
- The moon of Jupiter most similar in size to Earth's Moon is (a) Io; (b) Europa; (c) Ganymede; (d) Callisto.
- Io's surface appears very smooth because it (a) is continually resurfaced by volcanic activity; (b) is covered with ice; (c) has been shielded by Jupiter from meteorite impacts; (d) is liquid.
- The Galilean moons of Jupiter are sometimes described as a miniature inner solar system because (a) there are as many Galilean moons as there are terrestrial planets; (b) the moons have generally "terrestrial" composition; (c) the moons' densities decrease with increasing distance from Jupiter; (d) the moons all move on circular, synchronous orbits.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

- How does the force of gravity at Jupiter's cloud tops compare with the force of gravity at Earth's surface?
- What are the angular diameters of the orbits of Jupiter's four Galilean satellites, as seen from Earth at closest approach (assuming, for definiteness, that opposition occurs near perihelion)?
- Using the Figures given in the text, calculate how long it takes Jupiter's equatorial winds to circle the planet, relative to the interior.
- Calculate the rotational speed (in km/s) of a point on Jupiter's equator at the level of the cloud tops. Compare it with the orbital speed just above the cloud tops.
- Given Jupiter's age and current atmospheric temperature, what is the smallest possible mass the planet could have and still have retained its hydrogen atmosphere? [∞ \(More Precisely 8-1\)](#)
- If Jupiter had been just massive enough to fuse hydrogen (see *Discovery 11-2*), calculate what the planet's gravitational force on Earth would have been at closest approach, relative to the gravitational pull of the Sun. Assume circular orbits. Also, estimate what the magnitude of the planet's tidal effect on our planet would have been, again relative to that of the Sun.
- Calculate the ratio of Jupiter's mass to the total mass of the Galilean moons. Compare your answer with the ratio of Earth's mass to that of the Moon.
- Calculate the strength of Europa's gravitational pull on Io at closest approach, relative to Jupiter's gravitational attraction on Io.

## Activities


### Collaborative

- Jupiter's rapid (10 hour) rotation and ever-changing atmospheric features make it a highly dynamic telescopic object. Even a small telescope should reveal several of the planet's cloud belts, as well as its four brightest moons. Can you see the Great Red Spot? Make a series of sketches of the planet. You'll have to work fast because of the planet's rotation! Try to complete each sketch within 20–30 minutes, and prepare in advance a series of sheets of paper with a circle to represent the planet. Note the date and time of each sketch. By carefully timing some easily recognizable surface feature, such as the

Great Red Spot, measure Jupiter's rotation period. (You may have to come back on another night to be able to conveniently observe the chosen feature cross the entire planetary disk.) Repeat your observations a week or two later. Have any of the surface features, or their relation to one another, changed?

### Individual

- Consult a sky chart and find Jupiter in the night sky. It's an easy-to-find naked-eye object. Do any stars look as bright as Jupiter? What other differences do you notice between Jupiter and the stars?



# Saturn

## SPECTACULAR RINGS AND MYSTERIOUS MOONS

Saturn is one of the most beautiful and enchanting of all astronomical objects. Its rings are a breathtaking sight when viewed through even a small telescope, and they are probably the planet's best-known feature. Aside from its famous rings, though, Saturn presents us with another example of a jovian planet, allowing us to explore further the properties of these giant gaseous worlds.

Saturn is in many ways similar to its larger neighbor, Jupiter, in terms of composition, size, and structure, although its lower mass and greater distance from the Sun mean that Saturn's atmospheric colors are far less pronounced than those on Jupiter, and its weather patterns, although just as violent, are much harder to see. On the other hand, Saturn's rings and moons differ greatly from those of Jupiter. The comparison between these worlds provides us with invaluable insight into the structure and evolution of all the jovian planets.

**The Big Picture** Engineering feats of today's space age have allowed us to explore many of the planets and moons of our rich and varied solar system. Such thrilling exploration calls to mind earlier voyages of discovery, launched from Europe toward the Americas by Columbus, Magellan, Cortes, de Champlain, and many other seafarers who ventured forth across an unknown Atlantic and eventually around our world. Now, in the 21st century, robot spacecraft controlled by humans back on Earth act as modern spacefarers while exploring whole new, unknown worlds, and what they are finding is absolutely amazing.

**LEFT:** Saturn, a huge ball of lightweight gas surrounded by a spectacular ring system of orbiting rocky debris, is a planet much different from Earth. This image combines visible, infrared, and ultraviolet data acquired by cameras aboard the *Cassini* spacecraft as it cruised behind Saturn in 2005, revealing both the planet and its intricate rings in exquisite detail. We can also see, since this photo looks back toward the inner solar system, part of the bright Sun (bottom center) and a speck of reflected light from a dim and distant planet Earth (at the 8 o'clock position just outside the brightest ring). (JPL)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Summarize the orbital and physical properties of Saturn, and compare them with those of Jupiter.
- 2 Describe the composition and structure of Saturn's atmosphere and interior.
- 3 Explain why Saturn's internal heat source and magnetosphere differ from those of Jupiter.
- 4 Describe the structure and composition of Saturn's rings.
- 5 Define the Roche limit, and explain its relevance to the origin of Saturn's rings.
- 6 Summarize the general characteristics of Titan, and discuss the chemical processes in its atmosphere.
- 7 Outline some of the orbital and geological properties of Saturn's smaller moons.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.

## 12.1 Orbital and Physical Properties

Saturn was the outermost planet known to ancient astronomers. Named after the father of Jupiter in Roman mythology, Saturn orbits the Sun at almost twice the distance of Jupiter. The planet's sidereal orbital period of 29.4 Earth years was the longest natural unit of time known to the ancient world.

### Overall Properties

At opposition, when most of the Earth-based (or Earth-orbit-based) photographs of Saturn were taken, the planet is at its brightest and can lie within 8 AU of Earth. However, its great distance from the Sun still makes Saturn considerably fainter than either Jupiter or Mars. Saturn ranks behind Jupiter, the inner planets, and several of the brightest stars in the sky in terms of apparent brightness.


As with Jupiter, Saturn's many moons allowed an accurate determination of the planet's mass long before the arrival of spacecraft from Earth. Saturn's mass is  $5.7 \times 10^{26}$  kg, or 95 times the mass of Earth.  $\infty$  (Sec. 6.1) Although less than one-third the mass of Jupiter, Saturn is still an enormous body, at least by terrestrial standards.

From Saturn's distance and angular size, the planet's radius—and hence its average density—quickly follow. Saturn's equatorial radius is 60,000 km, or 9.5 Earth radii. The average density is  $700 \text{ kg/m}^3$ —less than the density of water ( $1000 \text{ kg/m}^3$ ). Here we have a planet that would float in the ocean—if Earth had one big enough! Saturn's low average density indicates that, like Jupiter, it is composed primarily of hydrogen and helium. Saturn's lower mass, however, results in lower interior pressure, so these gases are less compressed than in Jupiter's case.

#### ► FIGURE 12.1 Ring

**Orientation** Over time, Saturn's rings change their appearance to terrestrial observers as the tilted ring plane orbits the Sun. The roughly true-color images (inset) span a period of several years from 2003 (bottom) to nearly the present (top) and show how the rings change from our perspective on Earth, from almost edge-on to more nearly face-on. See also Figure 12.2 for a close-up image of its tilted ring system relative to Earth. (NASA)



At some times during Saturn's 29.5-year orbital period, the rings seem to disappear as Earth passes through their plane and we view them edge-on.



### Rotation Rate

Saturn, like Jupiter, rotates very rapidly and differentially.  $\infty$  (Sec. 11.1) The atmospheric rotation period, determined by tracking weather features observed in the planet's atmosphere, is  $10^{\text{h}14^{\text{m}}}$  at the equator and roughly  $10^{\text{h}40^{\text{m}}}$  at higher latitudes. However, the rotational period of the interior, obtained from *Cassini* measurements of magnetospheric outbursts, which should better trace the rotation of the planet's core, is  $10^{\text{h}46^{\text{m}}}$ , significantly longer than the surface values.

Curiously, the *Cassini* measurement is about 6 minutes longer than the corresponding result obtained (using similar means) by *Voyager* more than 20 years earlier. Scientists are uncertain as to the cause of this difference, although they do not think that Saturn's actual rotation rate has changed by as much as 1 percent during this relatively short time. Rather, it seems that the planet's magnetic field is not as good an indicator of interior rotation as was previously thought. Saturn's axis of rotation is significantly tilted with respect to the planet's orbital plane— $27^\circ$ , similar to that of both Earth and Mars. Researchers think that the interaction between the planet's magnetic field and the moon Enceladus (see Section 12.5) tends to slow the field's rotation, accounting for the observed difference in periods.



Because of Saturn's lower density, this rapid rotation makes Saturn even more flattened than Jupiter. In fact, Saturn is the “flattest” planet in the solar system, with a polar radius of just 54,000 km, about 10 percent less than the planet's equatorial radius. Careful calculations show that this degree of flattening, large as it is, is less than would be expected for a planet composed of hydrogen and helium alone. Accordingly, astronomers think that Saturn also has a rocky core, perhaps as much as 15 times the mass of Earth, or 1.5 times the mass of Jupiter's core.



## Rings

Saturn's best-known feature is its spectacular *ring system*. Because the rings lie in the planet's equatorial plane, their appearance (as seen from Earth) changes as Saturn orbits the Sun, as shown in Figure 12.1. As Saturn moves along its orbit, the angles at which the rings are illuminated and at which we view them vary. When the planet's north or south pole is tipped toward the Sun during Saturn's summer or winter, the highly reflective rings are at their brightest. During Saturn's spring and fall, the rings are close to being edge-on, both to the Sun and to us, so they seem to disappear altogether. The last two “ring crossings” occurred in 1996 and 2010.

One important deduction that we can make from this simple observation is that the rings are very thin. In fact, we now know that their thickness is only a few tens of meters, even though they are over 200,000 km in diameter.

### CONCEPT Check

- ✓ Why do some of the Earth-based images of Saturn in this chapter show the rings seen from above, whereas others show them from below?



## 12.2 Saturn's Atmosphere

Saturn is much less colorful than Jupiter. Figure 12.2 shows yellowish and tan cloud belts that parallel the equator, but these regions display less atmospheric structure than do the belts on Jupiter. No obvious large, long-lived “spots” or “ovals” adorn Saturn's cloud decks. Bands and storms do exist, but the color changes that distinguish them on Jupiter are largely absent on Saturn. This is the highest-resolution global image ever made of Saturn—a mosaic of more than 100 photographs taken in true color—showing great subtlety in the structure of its cloud deck.


### Composition and Coloration

Astronomers first observed methane in the spectrum of sunlight reflected from Saturn in the 1930s, about the same time that it was discovered on Jupiter. However, it was not until the early 1960s, when more sensitive observations became possible, that ammonia was finally detected on Saturn. In the planet's cold upper atmosphere, most ammonia is in the solid or liquid form, with relatively little of it present as a gas to absorb sunlight and create spectral lines. Astronomers finally made the first accurate determinations of Saturn's hydrogen and helium content in the late 1960s. These Earth-based measurements were later confirmed with the arrival of the *Pioneer* and *Voyager* spacecraft in the 1970s.

Saturn's atmosphere consists of molecular hydrogen ( $H_2$ , 92.4 percent), helium (He, 7.4 percent), methane ( $CH_4$ , 0.2 percent), and ammonia ( $NH_3$ , 0.02 percent). As on Jupiter, hydrogen and helium dominate—these most abundant elements never escaped from Saturn's atmosphere because of the planet's large mass and low temperature (see *More Precisely 8-1*). However, the fraction of helium on Saturn is far less than is observed on Jupiter (where, as we saw, helium accounts for nearly 14 percent of the atmosphere) or in the Sun.

It is extremely unlikely that the processes that created the outer planets preferentially stripped Saturn of nearly half its helium or that the missing helium somehow escaped from the planet while the lighter hydrogen remained behind. Instead, astronomers think that, at some time in Saturn's past, the heavier helium began to sink toward the center of the planet, reducing its abundance in the outer layers and leaving them relatively

◀ **FIGURE 12.2 Saturn** This image, acquired in 2005 by the *Cassini* spacecraft while approaching Saturn, is actually a mosaic of many images taken in true color. Note the bland coloration of the gas ball and the detail in the planet's rings. Resolution is 40 km. (NASA)



▲ **FIGURE 12.3 Saturn's Atmosphere** The vertical structure of Saturn's atmosphere contains several cloud layers, like Jupiter, but Saturn's weaker gravity results in thicker clouds and a more uniform appearance.

rich in hydrogen. We will return to the reasons for this differentiation and its consequences in a moment.

Figure 12.3 illustrates Saturn's atmospheric structure (recall the corresponding diagram for Jupiter, Figure 11.7). In many respects, Saturn's atmosphere is quite similar to Jupiter's, except that the temperature is a little lower because of Saturn's greater distance from the Sun and because its clouds are somewhat thicker. Since Saturn, like Jupiter, lacks a solid surface, we take the top of the troposphere as our reference level and set it to 0 km. The top of the visible clouds lies about 50 km below this level. As on Jupiter, the clouds are arranged in three distinct layers, composed (in order of increasing depth) of ammonia, ammonium hydrosulfide, and water ice. Above the clouds lies a layer of haze formed by the action of sunlight on Saturn's upper atmosphere.

The total thickness of the three cloud layers in Saturn's atmosphere is roughly 200 km, compared with about 80 km on Jupiter, and each layer is itself somewhat thicker than its counterpart on Jupiter. The reason for this difference is Saturn's weaker gravity (due to its lower mass). At the haze level, Jupiter's gravitational field is nearly two-and-a-half times stronger than Saturn's, so Jupiter's atmosphere

is pulled much more powerfully toward the center of the planet. Thus, Jupiter's atmosphere is compressed more than Saturn's, and the clouds are squeezed more closely together.

The colors of Saturn's cloud layers, as well as the planet's overall butterscotch hue, are due to the same basic cloud chemistry as on Jupiter. However, because Saturn's clouds are thicker, there are few holes and gaps in the top layer, so we rarely glimpse the more colorful levels below. Instead, we see different levels only in the topmost layer, which accounts for Saturn's rather uniform appearance.


## Weather

Saturn has atmospheric wind patterns that are in many ways reminiscent of those on Jupiter. There is an overall east-west zonal flow, which is apparently quite stable. Computer-enhanced images of the planet that bring out more cloud contrast (see Figure 12.4) clearly show the existence of bands, oval storm systems, and turbulent flow patterns looking very much like those seen on Jupiter. Scientists think that Saturn's bands and storms have essentially the same cause as does Jupiter's weather. Ultimately, the large-scale flows and small-scale storm systems are powered by convective motion in Saturn's interior and by the planet's rapid rotation.


The zonal flow on Saturn is considerably faster than on Jupiter and shows fewer east-west alternations, as can be seen from Figure 12.5 (see also Figure 11.6). The equatorial eastward jet stream, which reaches a speed of about 400 km/h on Jupiter, moves at a brisk 1500 km/h on Saturn and extends to much higher latitudes. Not until latitudes 40° north and south of the equator are the first westward flows found. Latitude 40° north also marks the strongest bands on Saturn and the most obvious ovals and turbulent eddies. Astronomers still do not fully understand the reasons for the differences between Jupiter's and Saturn's flow patterns.

In September 1990, amateur astronomers detected a large white spot in Saturn's northern hemisphere, just above the equator. A month later, when the *Hubble Space Telescope* imaged the phenomenon in more detail (Figure 12.6), the spot had developed into a band of clouds completely encircling the planet's equator. Astronomers think that the white coloration arose from crystals of ammonia ice formed when an upwelling plume of warm gas penetrated the cool upper cloud layers. Because the crystals were freshly formed, they had not yet been affected by the chemical reactions that color the planet's other clouds.

Such large spots are relatively rare on Saturn. They seem to appear roughly every 30 years, during the planet's northern summer. The previous one visible from Earth appeared in 1933, but it was smaller than the 1990 system and much shorter lived, lasting for only a few weeks. The turbulent flow patterns seen around the 1990 white spot had many similarities to the flow around Jupiter's Great Red Spot. Scientists hope that routine observations of such



▲ **FIGURE 12.4 Saturn's Cloud Structure** More structure is seen in Saturn's cloud cover when computer processing and artificial color are used to enhance the contrast of the image, as in these *Voyager* images of the entire gas ball and a smaller, magnified piece of it. (NASA)



▲ **FIGURE 12.5 Saturn's Zonal Flow** Winds on Saturn reach speeds even greater than those on Jupiter. As on Jupiter, the visible bands appear to be associated with variations in wind speed.


temporary phenomenon the jovian worlds will give them greater insight into the dynamics of planetary atmospheres.

Since *Cassini's* arrival at Saturn, researchers have been able to study the planet's storms in much more detail. Figure 12.7 shows the development of a particularly large and complex system that appeared in the northern hemisphere in late 2010. *Cassini's* detectors measure strong bursts of radio waves associated with it, most likely produced by intense lightning discharges deep below the cloud tops. The lightning is probably powered by convection and precipitation (water and ammonia "rain"), just as in thunderstorms on Earth, but the bursts are millions of times stronger than anything ever witnessed here at home.

Astronomers think that these large storms are rooted deep in Saturn's atmosphere—perhaps like Jupiter's Great Red Spot—and normally completely hidden below the upper cloud layers. Only occasionally does the storm flare up, producing a bright plume visible from outside. The largest storms can be quite long-lived—the system shown in Figure 12.7 is expected to persist at least into 2013. In addition, both *Hubble* and *Cassini* have observed many smaller (but still huge, by terrestrial standards!), shorter-lived storms on Saturn.

*Cassini* has also detected numerous small dark storms apparently associated with this and other large systems. These smaller storms seem to be "spun off" from the larger systems





▲ **FIGURE 12.6 Saturn Storms** Circulating and evolving cloud systems are evident on Saturn, imaged by the *Hubble Space Telescope* at approximately 2-hour intervals (left to right). (NASA)

and subsequently merge with the planet's zonal flow, perhaps providing a means for energy to flow from Saturn's warm interior into the cold atmosphere. However, the "big picture" connecting these large- and small-scale storm systems on Saturn with the Red Spot, white ovals, and the "Red Spot Junior" on Jupiter, remains to be worked out. ∞ (Sec. 11.2)

### CONCEPT Check

✓ **Why are atmospheric features on Saturn generally less vivid than those on Jupiter?**




▲ **FIGURE 12.7 Storm Alley** This huge storm on Saturn was observed by the *Cassini* spacecraft in 2011. While churning its way through the northern hemisphere, it seems to leave behind a "tail" wrapped around the planet. (NASA)

## 12.3 Saturn's Interior and Magnetosphere

Figure 12.8 depicts Saturn's internal structure. (Compare with Figure 11.12 for the case of Jupiter.) The picture was pieced together by planetary scientists using the same tools—spacecraft observations and theoretical modeling—that they employed to infer Jupiter's inner workings. Saturn has the same basic internal parts as Jupiter, but their relative proportions are somewhat different: Saturn's metallic hydrogen layer is thinner and its core is larger. Because of its lower mass, Saturn has a less extreme core temperature, density, and pressure than does Jupiter. The central pressure is around one-fifth of Jupiter's—about 2 to 3 times the pressure at the center of Earth.

### Internal Heating

Infrared measurements indicate that Saturn's surface (i.e., cloud-top) temperature is 97 K, substantially higher than the temperature at which Saturn would reradiate all the energy



▲ **FIGURE 12.8 Saturn's Interior** Saturn's internal structure, as deduced here from *Voyager* observations and computer modeling, can be compared with similar properties noted for Jupiter in Figure 11.12.



it receives from the Sun. In fact, Saturn radiates almost 3 times more energy than it absorbs. Thus, Saturn, like Jupiter, has an internal energy source. ∞ (Sec. 11.3) But the explanation behind Jupiter's excess energy—that the planet has a large reservoir of heat left over from its formation—doesn't work for Saturn. Smaller than Jupiter, Saturn must have cooled more rapidly—rapidly enough that its original supply of energy was used up long ago. What, then, is happening inside Saturn to produce this extra heat?

The explanation for the origin of Saturn's extra heat also explains the mystery of the planet's apparent helium deficit. At the temperatures and high pressures found in Jupiter's interior, liquid helium *dissolves* in liquid hydrogen. Inside Saturn, where the internal temperature is lower, the helium doesn't dissolve so easily and tends to form droplets instead. The phenomenon is familiar to cooks, who know that it is generally much easier to dissolve ingredients in hot liquids than in cold ones. Saturn probably started out with a fairly uniform solution of helium dissolved in hydrogen, but the helium tended to condense out of the surrounding hydrogen, much as water vapor condenses out of Earth's atmosphere to form a mist. The amount of helium condensation was greatest in the planet's cool outer layers, where the mist turned to rain about 2 billion years ago. A light shower of liquid helium has been falling through Saturn's interior ever since. This **helium precipitation** is responsible for depleting the outer layers of their helium content.

So we can account for the unusually low abundance of helium in Saturn's atmosphere: Much of it has rained down to lower levels. But what about the excess heating? The answer is simple: As the helium sinks toward the center, the planet's gravitational field compresses it and heats it up. The gravitational energy thus released is the source of Saturn's internal heat. In the distant future—in a billion years or so—the helium rain will stop, and Saturn will cool until its outermost layers radiate only as much energy as they receive from the Sun. When that happens, the temperature at Saturn's cloud tops will be 74 K. As Jupiter cools, it, too, may someday experience precipitate helium in its interior, causing its surface temperature to rise once again.

## Magnetospheric Activity

Saturn's electrically conducting interior and rapid rotation produce a strong magnetic field and an extensive magnetosphere. Probably because of the considerably smaller mass of Saturn's metallic hydrogen zone, the planet's basic magnetic field strength is only about one-twentieth that of Jupiter, or about a thousand times greater than that of Earth. The magnetic field at Saturn's cloud tops (roughly 10 Earth radii from the planet's center) is approximately the same as that at Earth's surface. *Voyager* measurements indicate that, unlike Jupiter's and Earth's magnetic axes, which are slightly tilted, Saturn's magnetic field is not inclined



▲ **FIGURE 12.9 Aurora on Saturn** An ultraviolet camera aboard the *Hubble Space Telescope* recorded this image of a remarkably symmetrical (orange) aurora on Saturn during a solar storm in 1998. (NASA)


with respect to its axis of rotation. Saturn's magnetic field, like Jupiter's, is oriented opposite that of Earth: that is, an Earth compass needle would point toward Saturn's south pole rather than its north. ∞ (Sec. 11.4) Figure 12.9 shows an aurora on Saturn, imaged in 1998 by the *Hubble Space Telescope*.

Saturn's magnetosphere extends about 1 million km toward the Sun and is large enough to contain the planet's ring system and the innermost 16 small moons. Saturn's largest moon, Titan, orbits about 1.2 million km from the planet, so it is found sometimes just inside the outer magnetosphere and sometimes just outside, depending on the intensity of the solar wind (which tends to push the sunward side of the magnetosphere closer to the planet). Because no major moons lie deep within Saturn's magnetosphere, the details of its structure are different from those of Jupiter's magnetosphere. For example, there is no equivalent of the Io plasma torus. ∞ (Sec. 11.5) Like Jupiter, Saturn emits radio waves, but as luck would have it, they are reflected from Earth's ionosphere (they lie in the AM band) and were not detected until the *Voyager* craft approached the planet.

## CONCEPT Check

✓ Where did Saturn's atmospheric helium go?


## 12.4 Saturn's Spectacular Ring System

The most obvious and well-known aspect of Saturn's appearance is, of course, its *planetary ring system*. Astronomers now know that all the jovian planets have rings, but Saturn's are by far the brightest, the most extensive, and the most beautiful.  (Sec. 11.6)


### The View from Earth

Galileo saw Saturn's rings first in 1610, but he did not recognize what he saw as a planet with a ring. At the resolution of his small telescope, the rings looked like bumps on the planet, or perhaps (he speculated) parts of a triple planet of some sort. Figure 12.10(a) and (b) shows two of Galileo's early sketches of Saturn. By 1616, Galileo had already realized that the “bumps” were not round, but rather elliptical in shape. In 1655, the Dutch astronomer Christian Huygens realized what the bumps were: a thin, flat ring, completely encircling the planet (Figure 12.10c).

In 1675, the French-Italian astronomer Giovanni Domenico Cassini discovered the first ring feature: a dark band about two-thirds of the way out from the inner edge. From Earth, the band looks like a gap in the ring (an observation that is not too far from the truth, although we now know that there is actually some ring material within it). This “gap”



▲ **FIGURE 12.10 Sketches of Saturn's Rings** Three artist's re-renderings of sketches of Saturn's rings, made (a) by Galileo in 1610, (b) by Galileo in 1616, and (c) by Huygens in 1655.



▲ **FIGURE 12.11 Saturn's Rings** Much fine structure, especially in the rings, appears in this image of Saturn taken with the *Cassini* spacecraft in 2005. The main ring features are marked and are shown here in false color in order to enhance contrast. The other rings listed in Table 12.1 are not visible here. (NASA)

is named the **Cassini division**, in honor of its discoverer. Careful observations from Earth show that the inner “ring” is in reality also composed of two rings. From the outside in, the three rings are known somewhat prosaically as the **A**, **B**, and **C rings**. The Cassini division lies between the A and B rings. The much narrower **Encke gap**, some 300 km wide, is found in the outer part of the A ring. These ring features are marked on Figure 12.11. No finer ring details are visible from our Earthly vantage point. Of the three main rings, the B ring is brightest, followed by the somewhat fainter A ring, and then by the almost translucent C ring. A more complete list of ring properties appears in Table 12.1. (The D, E, F, and G rings listed in the table are discussed later in this section.)

**TABLE 12.1** The Rings of Saturn

Ring	Inner Radius (km)	Outer Radius (km)	Width (km)
	(planetary radii)	(planetary radii)	
D	67,000	74,700	7700
C	74,700	92,000	17,300
B	92,000	117,500	25,500
Cassini division	117,500	122,300	4800
A	122,300	136,800	14,500
Encke gap*	133,400	133,700	300
F	140,300	140,400	100
G	165,800	173,800	8000
E	180,000	480,000	300,000

\*The Encke gap lies within the A ring.

## What Are Saturn's Rings?

A fairly obvious question—and one that perplexed the best scientists and mathematicians on Earth for almost two centuries—is “What are Saturn’s rings made of?” By the middle of the 19th century, various dynamic and thermodynamic arguments had conclusively proved that the rings could not be solid, liquid, or gas!

What is left? In 1857, after showing that a solid ring would become unstable and break up, Scottish physicist James Clerk Maxwell suggested that the rings are composed of a great number of small particles, all independently orbiting Saturn, like so many tiny moons. That inspired speculation was verified in 1895, when Lick Observatory astronomers measured the Doppler shift of sunlight reflected from the rings and showed that the velocities thus determined were exactly what would be expected from separate particles moving in circular orbits in accordance with Newton’s law of gravity. ∞ (Secs. 2.8, 3.5)

What sort of particles make up the rings? The fact that they reflect most (over 80 percent) of the sunlight striking them had long suggested to astronomers that they are made of ice, and infrared observations in the 1970s confirmed that water ice is indeed a prime constituent of the rings. Radar observations and later *Voyager* and *Cassini* studies of scattered sunlight showed that the diameters of

the particles range from fractions of a millimeter to tens of meters, with most particles being about the size (and composition) of a large snowball on Earth.


We now know that the rings are truly thin—only 10–15 m thick, according to *Cassini* measurements. Stars can occasionally be seen through them, like automobile headlights penetrating a snowstorm. Why are the rings so thin? The answer seems to be that collisions between ring particles tend to keep them all moving in circular orbits in a single plane. Any particle that tries to stray from this orderly motion finds itself in an orbit that soon runs into other ring particles. Over long periods, the ensuing jostling serves to keep all of the particles moving in circular, planar orbits. The asymmetric gravitational field of Saturn (a result of its flattened shape) sees to it that the rings lie in the planet’s equatorial plane.

## The Roche Limit


But why a ring of particles at all? What process produced the rings in the first place? To answer these questions, consider the fate of a small moon orbiting close to a massive planet such as Saturn. The moon is held together by internal forces—its own gravity, for example. As we bring our hypothetical moon closer to the planet, the tidal force on it increases. Recall from Chapter 7 that the effect of such a tidal force is to stretch the moon along the direction toward the planet—that is, to create a tidal bulge.

Recall also that the tidal force increases rapidly with decreasing distance from the planet. ∞ (Sec. 7.6) As the moon is brought closer to the planet, it reaches a point where the tidal force tending to stretch it out becomes *greater* than the internal forces holding it together. At that point, the moon is torn apart by the planet’s gravity, as shown in Figure 12.12. The pieces of the satellite then pursue their own individual orbits around that planet, eventually spreading all the way around it in the form of a ring.

For any given planet and any given moon, the critical distance inside of which the moon is destroyed is known as the *tidal stability limit*, or the **Roche limit**, after the 19th-century French



**Narrated FIGURE 12.12 Roche Limit** From top to bottom, these four frames illustrate how the tidal field of a planet first distorts (near top), and then destroys (at bottom), a moon that strays too close. Note that the distortion is exaggerated in the middle panels, and the moon’s infall does not happen directly; rather, its breakup occurs over the course of many orbits.



◀ **FIGURE 12.13**  
**Jovian Ring Systems**

All of the distances to the rings of Jupiter, Saturn, Uranus, and Neptune are expressed in planetary radii. The red line represents the Roche limit, and all the rings lie within (or very close to) this limit of their parent planets.

mathematician Edouard Roche, who first calculated it. As a handy rule of thumb, if our hypothetical moon is held together by its own gravity and its average density is comparable to that of the parent planet (both reasonably good approximations for Saturn's larger moons), then the Roche limit is roughly 2.4 times the radius of the planet. Thus, for Saturn, no moon can survive within a distance of 144,000 km of the planet's center, about 7000 km beyond the outer edge of the A ring. The main (A, B, C, D, and F) rings of Saturn occupy the region inside Saturn's Roche limit.

These considerations apply equally well to the other jovian worlds. Figure 12.13 shows the location of the ring system of each jovian planet relative to the planet's Roche limit. Given the approximations in our assumptions, we can conclude that all the major planetary rings are found within the Roche limit of their parent planet. Notice that, strictly speaking, the calculation of this limit applies only to low-density moons massive enough for their own gravity to be the dominant force binding them together. Sufficiently small moons (less than 10 km or so in diameter) can survive even within the Roche limit because they are held together mostly by interatomic (electromagnetic) forces, not by gravity.


## The Rings in Detail

Thus, as the two *Voyager* probes approached Saturn in 1980 and 1981, scientists on Earth were fairly confident that they understood the nature of the rings. However, there were many surprises in store. The *Voyager* flybys changed forever our

view of this spectacular region in our cosmic backyard, revealing the rings to be vastly more complex than astronomers had imagined. *Cassini's* 4-year tour of the Saturn system a quarter century later allowed much more extended and detailed study of many of the phenomena discovered by *Voyager* and yielded many new insights into this fascinating system.

As the *Voyager* probes approached Saturn, it became obvious that the main rings were actually composed of tens of thousands of narrow **ringlets**, shown (as seen by *Cassini*) in Figure 12.14. Although *Voyager* cameras did find several new gaps in the rings, the ringlets in the figure are actually not separated from one another by empty space. Instead, detailed studies reveal that the rings contain concentric regions of alternating high and low concentrations of ring particles. The ringlets are the high-density peaks. According to theory, the gravitational influence of Saturn's inner moons and the mutual gravitational attraction of the ring particles enables waves of matter to form and move in the plane of the rings, rather like ripples on the surface of a pond. The wave crests wrap all the way around the rings, forming tightly wound spiral patterns called *spiral density waves* that resemble grooves in a huge celestial phonograph record.

Although the ringlets are the result of spiral waves in the rings, the true gaps are not. The narrower gaps—roughly 20 of them—are thought to be kept clear by the action of small *moonlets* embedded in them. These moonlets are larger (perhaps 10 or 20 km in diameter) than the largest true ring particles, and they simply “sweep up” ring material through collisions as they go. However, despite careful



**Interactive FIGURE 12.14 Saturn's Rings, Up Close** *Cassini* took this true-color image of Saturn's dazzling ring structure just before flying through the planet's tenuous outer rings. Earth is superposed at right, to proper scale, for a size comparison. The inset at left is an overhead view of a portion of the B ring, showing the ringlet structure in even more detail; in fact the resolution here is an incredible 4 km. (NASA)

searches of *Voyager* and *Cassini* images, only two of these moonlets have so far been found. The 30-km-wide moonlet Pan, discovered in archival *Voyager* data in 1991, orbits in the Encke gap (marked in Figure 12.11). The 7-km Daphnis, imaged by *Cassini* in 2005, resides in the narrow Keeler gap, close to the outer edge of the A ring.

Astronomers have found indirect evidence for embedded moonlets in the form of the scalloped “wakes” they leave behind them in the rings, but no other direct sightings have occurred. Despite their elusiveness, moonlets are nevertheless regarded as the best explanation for the small gaps and associated fine structure in the rings. In 2006 *Cassini* found evidence for a new class of intermediate-sized (100-m diameter) moonlets, which may play an important role in clearing some small or partial gaps.

*Voyager 2* found a series of faint rings, now known collectively as the **D ring**, inside the inner edge of the C ring, stretching down almost to Saturn's cloud tops. The D ring contains relatively few particles and is so dark that it is completely invisible from Earth. Two other faint rings, discovered by *Pioneer 11* and *Voyager 1*, respectively, lie well outside the main ring structure. Both the **G ring** and the **E ring**, are faint and diffuse, more like Jupiter's ring than the main A, B, and C rings of the Saturn system. ∞ (Sec. 11.6) The E ring appears to be associated with volcanism on the moon Enceladus (see Section 12.5).


Figure 12.15 is a *Cassini* image showing the rings from a perspective never before seen—behind the planet looking back toward the eclipsed Sun. Just as diffuse airborne dust is most easily seen against the light streaming in through a sunlit window, the planet's faint rings show up clearly in this remarkable back-lit view. In addition, this image reveals several additional faint rings, some of them also associated with the orbits of various small moons.

The *Voyager 2* cameras revealed one other completely unexpected feature. A series of dark radial “spokes” formed on the B ring, moved around the planet for about one ring orbital period, and then disappeared (Figure 12.16). Careful scrutiny of these peculiar drifters showed that they were composed of very fine (micron-sized) dust hovering a few tens of meters *above* the plane of the rings. Scientists think that this dust was held in place by electromagnetic forces generated in the ring plane, perhaps resulting from collisions among particles there or interactions with the planet's magnetic field. The spokes faded as the ring revolved.

Astronomers had expected that the creation and dissolution of such spokes would be regular occurrences in the Saturn ring system, but during the first year of *Cassini*'s tour, none were seen. However, as with the planet's faint rings, lighting and viewing conditions seem to be critical to the spokes' visibility. As the planet's orientation changed relative to the Sun, spokes were finally observed in late 2005. Spoke activity is expected to be common for the remainder of the *Cassini* mission, and researchers hope that repeated observations will allow them to understand this peculiar phenomenon.

## Orbital Resonances and Shepherd Satellites


*Voyager* images showed that the largest gap in the rings, the Cassini division, is not completely empty of matter. In fact, as can be seen in Figure 12.14, the division contains a series of faint ringlets and gaps (and, presumably, embedded moonlets, too). The overall concentration of ring particles in the division as a whole is, however, much lower than in the A and B rings. The diffuse nature of the division causes it to appear bright in Figure 12.15. Although its small



▲ **FIGURE 12.15 Back-Lit Rings** *Cassini* took this spectacular image of Saturn’s rings as it passed through Saturn’s shadow. The normally hard to see F, G, and E outer rings are clearly visible in this contrast-enhanced image. The inset shows the moon Enceladus orbiting within the E ring; its eruptions likely give rise to the ring’s icy particles. (NASA)

internal gaps probably result from embedded satellites, the division itself does not. Instead, it owes its existence to another solar system *resonance*, this time involving particles orbiting in the division, on the one hand, and Saturn’s innermost major moon, Mimas, on the other. ∞ (Sec. 8.4)

A ring particle moving in an orbit within the Cassini division has an orbital period exactly half that of Mimas. Particles in the division thus complete exactly two orbits around Saturn in the time taken for Mimas to orbit once—a configuration known as a 2:1 resonance. Applying Kepler’s



third law (recast to refer not to the planets, but to Saturn’s moons), we can show that this 2:1 resonance with Mimas corresponds to a radius of 117,000 km, the inner edge of the division. ∞ (More Precisely 2-2)

The effect of this resonance is that particles in the division receive a gravitational tug from Mimas at exactly the same location in their orbit every other time around. Successive tugs reinforce one another, and the initially circular trajectories of the ring particles soon get stretched out into ellipses. In their new orbits, these particles collide with other particles and eventually find their way into new circular orbits at other radii. The net effect is that the number of ring particles in the Cassini division is greatly reduced.

Particles in “nonresonant” orbits (i.e., at radii whose orbital period is not simply related to the period of Mimas) also are acted upon by Mimas’s gravitational pull. But the times when the force is greatest are spread uniformly around the orbit, and the tugs cancel out. It’s a little like pushing a child on a swing: Pushing at the same point in the swing’s motion each time produces much better results than do random shoves. Thus, Mimas (or any other moon) has a large effect on the ring at those radii at which a resonance exists and little or no effect elsewhere.

◀ **FIGURE 12.16 Spokes in the Rings** Saturn’s B ring showed a series of dark temporary “spokes” as *Voyager 2* flew by at a distance of about 4 million km. The spokes are caused by small particles suspended just above the ring plane. *Cassini*, too, has seen spokes, although (so far) they have not been as prominent as those seen by *Voyager* 25 years earlier. (NASA)

Table 12.2 The Major Moons of Saturn\*

Name	Distance from Saturn		Orbit Period (days)	Size (longest diameter, km)	Mass** (Earth moon masses)	Density	
	(km)	(planetary radii)				(kg/m <sup>3</sup> )	(g/cm <sup>3</sup> )
Mimas	186,000	3.08	0.94	398	0.00051	1100	1.1
Enceladus	238,000	3.95	1.37	498	0.00099	1100	1.1
Tethys	295,000	4.89	1.89	1060	0.0085	1000	1.0
Dione	377,000	6.26	2.74	1120	0.014	1400	1.4
Rhea	527,000	8.74	4.52	1530	0.032	1200	1.2
Titan	1,220,000	20.3	16.0	5150	1.83	1900	1.9
Hyperion	1,480,000	24.6	21.3	370			
Iapetus	3,560,000	59.1	79.3	1440	0.022	1000	1.0


\* Moons larger than 300 km in diameter only.

\*\* Mass of Earth's Moon =  $7.4 \times 10^{22}$  kg =  $1.3 \times 10^{-4}$  Saturn masses.

We now know that resonances between ring particles and moons play an important role in shaping the fine structure of Saturn's rings. For example, the sharp outer edge of the A ring is thought to be governed by a 3:2 resonance with Mimas (three ring orbits in two Mimas orbital periods). Most theories of planetary rings predict that the ring system should spread out with time, basically because of collisions among ring particles. Instead, the A ring's outer edge is "patrolled" by a small satellite named Atlas,

held in place by the gravity of Mimas, that prevents ring particles from diffusing away. Compare Tables 12.1 and 12.2 and see if you can identify other resonant connections between Saturn's moons or between the moons and the rings. (You should be able to find quite a few—Saturn's ring system is a complex place!)

Outside the A ring lies perhaps the strangest ring of all. The faint, narrow **F ring** (shown in Figure 12.17) was discovered by *Pioneer 11* in 1979, but its full complexity



▲ **FIGURE 12.17 Shepherd Moon** (a) Saturn's narrow F ring appears to contain kinks and braids, making it unlike any of Saturn's other rings. Its thinness is caused by two shepherd satellites that orbit near the ring—one a few hundred kilometers inside the ring and the other a similar distance outside it. (b) One of the potato-shaped shepherd satellites (Prometheus, here roughly 100 km across) can be seen in this enlarged view. (NASA)



became evident only when *Voyager 1* took a closer look. Unlike the inner major rings, the F ring is narrow—less than a hundred kilometers wide. It lies just inside Saturn’s Roche limit, separated from the A ring by about 3500 km. Its narrowness by itself is unusual, as is its slightly eccentric shape, but the F ring’s oddest feature is its irregular, “kinked” structure, making it look as though it is made up of several separate strands braided together! This remarkable appearance sent dynamicists scrambling in search of an explanation. It now seems as though the ring’s intricate structure, as well as its thinness, arise from the influence of two small moons, known as **shepherd satellites**, that orbit on either side of it (Figure 12.17b).

Two small, dark satellites, each little more than 100 km in diameter and called Prometheus and Pandora, orbit about 1000 km on either side of the F ring. Their gravitational influence on the F-ring particles keeps the ring tightly confined in its narrow orbit. Any particle straying too far out of the F ring is gently guided back into the fold by one or the other of the shepherd moons. (The moon Atlas confines the A ring in a somewhat similar way.)

However, the details of how Prometheus and Pandora produce the braids in the F ring, and why the two moons are there at all, in such similar orbits, remain unclear. Scientists theorize that there may be many more unseen small moonlets orbiting near the F ring and affecting its appearance. There is evidence that other eccentric rings found in the gaps in the A, B, and C rings may also result from the effects of shepherd moonlets. Beyond the F ring, the G ring apparently lacks ringlets and peculiar internal structure, but its relative narrowness and sharp edges suggest the presence of shepherd satellites, although so far none has been found. *Cassini* discovered that the G ring has regions of enhanced brightness, known as *arcs*, that move in resonance with Mimas and likely harbor confining moonlets. One of these moonlets was discovered by *Cassini* in 2009. *Cassini* has also detected faint arcs associated with other known moons. The arcs are most likely the result of micrometeoroid impacts ejecting material from those small bodies.

## Origin of the Rings

Two possible origins have been suggested for Saturn’s rings. Astronomers estimate that the total mass of ring material is no more than  $10^{15}$  tons—enough to make a satellite about 250 km in diameter. If such a satellite strayed inside Saturn’s Roche limit or was destroyed (perhaps by a collision) near that radius, a ring could have resulted. An alternative view is that the rings represent material left over from Saturn’s formation stage 4.6 billion years ago. In this scenario, Saturn’s tidal field prevented any moon from forming inside the Roche limit, so the material has remained a ring ever since. Which view is correct?

All the dynamic activity observed in Saturn’s rings suggests to many researchers that the rings must be quite

young—perhaps no more than 50 million years old, or 100 times younger than the solar system. There is just too much going on, the argument goes, for the rings to have remained stable for billions of years, so they probably aren’t left over from the planet’s formative stages. If this is so, then either the rings are continuously replenished, perhaps by fragments of Saturn’s moons chipped off by meteorites or by activity on the larger moons (see Section 12.5), or they are the result of a relatively recent, possibly catastrophic, event in the planet’s system—a small moon that may have been hit by a large comet or even by another moon.


Recently, *Cassini* has complicated this discussion by finding that different rings may have different ages and may even have formed in different ways. The *Cassini* data, coupled with theoretical simulations, suggest that the B ring might in fact be billions of years old but manages to preserve its youthful appearance by continually clumping and recycling its material, exposing fresh, bright water ice.

Astronomers prefer not to invoke catastrophic events to explain specific phenomena, but the more we learn of the universe, the more we realize that catastrophe probably plays an important role. For now, the details of the formation of Saturn’s ring system simply aren’t well understood.

### CONCEPT Check

- ✓ What do the Roche limit and orbital resonances have to do with planetary rings?

## 12.5 The Moons of Saturn

Saturn has the most extensive, and in many ways the most complex, system of natural satellites of all the planets. The planet’s eight largest moons, all more than 300 km in diameter, are listed in Table 12.2. Observations of sunlight reflected from them suggest that most are covered with snow and ice. Many of them are probably made almost entirely of water ice. Even so, they are a curious and varied lot, and many aspects of their structure and history are still not well understood. Most of our detailed knowledge of these moons comes from the *Pioneer* and *Voyager* flybys in the late 1970s and early 1980s, and from *Cassini*, which is currently orbiting the planet.  (Discovery 6-2)

The moons fall into three fairly natural groups. First, there are many “small” moons—irregularly shaped chunks of ice, all less than 400 km across—that exhibit a bewildering variety of complex and fascinating motion. Only the largest, Hyperion, is listed in Table 12.2. Second, there are six “medium-sized” moons—spherical bodies with diameters ranging from about 400 to 1500 km—that

offer clues to the past and present state of the environment of Saturn while presenting many puzzles regarding their own appearance and history. Finally, there is Saturn's single "large" moon—Titan—which, at 5150 km in diameter, is the second-largest satellite in the solar system. (Jupiter's Ganymede is a little bigger.) Titan has an atmosphere denser than Earth's and (some scientists think) surface conditions that may be conducive to life. Notice, incidentally, that Jupiter has no "medium-sized" moons, as just defined: The Galilean satellites are large, like Titan, and all of Jupiter's other satellites are small—no more than 300 km in diameter.


As in the case of Jupiter, improving observational techniques in recent years have led to an explosion in the number of known moons orbiting Saturn—to at least 62 as of mid-2013. Like the many small satellites of Jupiter, these moons are all very faint (and hence small) and revolve around Saturn quite far from the planet on rather inclined, often retrograde, orbits, much as Saturn's other "small" moons do. Most likely, they are chunks of debris captured from interplanetary space after close encounters with Saturn.

## Titan: A Moon with an Atmosphere

The largest and most intriguing of all Saturn's moons, Titan, was discovered by Christian Huygens in 1655. Even through a large Earth-based optical telescope, this moon is visible only as a barely resolved reddish disk. However, long before the *Voyager* or *Cassini* missions, astronomers already knew from spectroscopic observations that the moon's reddish coloration was caused by something quite special—an atmosphere. So eager were mission planners to obtain a closer look that they programmed *Voyager 1* to pass very close to Titan, even though it meant that the spacecraft could not then use Saturn's gravity to continue on to Uranus and Neptune. Instead, *Voyager 1* left the Saturn system on a path taking the craft out of the solar system well above the plane of the ecliptic. [∞ \(Discovery 6-1\)](#)

A *Voyager 1* image of Titan is shown in Figure 12.18(a). Unfortunately, despite the spacecraft's close pass, the moon's

surface remained a mystery. A thick, uniform layer of haze (similar to the photochemical smog created by chemical reactions powered by light found over many cities on Earth) that envelops the moon completely obscured the spacecraft's view. Still, *Voyager 1* was able to provide mission specialists with detailed atmospheric data. Figure 12.18(b) shows one of the best Earth-based views of the moon, taken in infrared



► **FIGURE 12.18 Titan** (a) Larger than the planet Mercury and roughly half the size of Earth, Titan was photographed in visible light from only 4000 km away as *Voyager 1* passed by in 1980. Only Titan's upper cloud deck can be seen here. The inset shows a contrast-enhanced, true-color image of the haze layers in Titan's upper atmosphere, taken by the *Cassini* spacecraft in 2005. (b) In the infrared, as captured with the adaptive-optics system on the Canada-France-Hawaii telescope on Mauna Kea, some large-scale surface features can be seen. The bright regions are thought to be highlands, possibly covered with frozen methane. The brightest area is nearly 4000 km across—about the size of Australia. (NASA; CFHT)


## DISCOVERY 12-1

### Dancing Among Saturn's Moons

In Chapter 6 we described how mission planners routinely use “gravitational slingshots” to accelerate and redirect a spacecraft’s trajectory as it speeds toward its target. ∞ (Discovery 6-1) Many of NASA’s planetary missions have relied heavily on this expertise. ∞ (Discovery 6-2) The *Galileo* and *Cassini* probes also made heavy use of gravity assists to control the spacecrafts’ intricate orbits within the moon systems of Jupiter and Saturn. Here we describe how *Cassini* has managed to make repeated close approaches to many of the moons in the Saturn system—the key to the extraordinary data now streaming back to scientists on Earth.

The first figure illustrates how a spacecraft can be moved onto a higher or lower orbit depending on the details of its interaction with a moon. Passing behind the moon, the spacecraft acquires some of the moon’s energy and speeds up, moving farther from the planet. Conversely, passage in front of the moon slows the spacecraft down, reducing its orbital semimajor axis. In either case, the *direction* of the spacecraft’s trajectory is changed, setting it on course for its next destination. In the case of *Cassini* and Saturn, the moon responsible for the lion’s share of the close encounters is Titan—very convenient, as a close study of Titan is one of the mission’s prime objectives.

By carefully combining a series of close encounters with Titan and other moons, *Cassini*’s controllers sent the probe throughout the moon system, allowing it to visit many of the moons multiple times. The second figure shows a possible trajectory computed before launch. Each orbit is the result of a carefully computed interaction with Titan or some other moon during the previous pass. In fact, this diagram differs in detail from *Cassini*’s actual flight path—small errors in the exact distance from each moon at one pass cause large changes in the




next encounter, and the actual trajectory was revised several times during the mission.

The *Cassini* mission was originally scheduled to end in 2008, 4 years after its arrival at Saturn, but the gravity-assisted maneuver that placed the probe in orbit around the planet was so accurate that almost no fuel had to be used to fine-tune the trajectory. As a result, the highly successful mission—now named *Cassini Equinox*, after the ring-plane crossing in August 2009—has been extended until 2017. Even as *Cassini*’s fuel supply dwindles, mission controllers should still be able to adjust its trajectory, using mainly Titan’s gravity, to reach the desired parts of the Saturn system.

Eventually, though, the maneuvering fuel will be nearly gone and the mission will have to be brought to a close. Mission planners are undecided as to the best way to end the

*Cassini* tour. They could place *Cassini* into a wide stable orbit, taking measurements for years to come until the probe’s nuclear fuel cells are depleted. Or the probe could crash into Saturn, just as *Galileo* did with Jupiter. Even crash-landing on Titan has been considered, although that would risk polluting the moon with nuclear material. A final option is to send *Cassini* flying through the main rings of Saturn, taking fascinating photographs and amassing data until it “hits something big.” Stay tuned!



light to penetrate the smog. A detailed, close-up look at Titan is a key part of the ongoing *Cassini* mission. As described in *Discovery 12-1*, the spacecraft's carefully choreographed trajectory includes dozens of additional visits to the moon by the end of its extended mission.

Titan's atmosphere is thicker and denser than Earth's, and it is certainly far more substantial than that of any other moon. Prior to *Voyager 1*'s arrival in 1980, only methane and a few other simple hydrocarbons had been conclusively detected on Titan (*hydrocarbons* are molecules consisting solely of hydrogen and carbon atoms; methane,  $\text{CH}_4$ , is the simplest). Radio and infrared observations made by *Voyager 1* and *Cassini* showed that the atmosphere is actually made up mostly of nitrogen ( $\text{N}_2$ , roughly 98 percent) and methane ( $\text{CH}_4$ , about 2 percent), with traces of argon and other gases.

Titan's atmosphere seems to act like a gigantic chemical factory. Powered by the energy of sunlight, the atmosphere is undergoing a complex series of chemical reactions that maintain steady (but trace) levels of hydrogen gas ( $\text{H}_2$ ), the hydrocarbons ethane ( $\text{C}_2\text{H}_6$ ) and propane ( $\text{C}_3\text{H}_8$ ), and carbon monoxide ( $\text{CO}$ ), ultimately resulting in the observed


smog. Spectrometers aboard *Cassini* have detected organic molecules below the haze layers. The upper atmosphere is thick with aerosol haze (droplets so small that they remain suspended in the atmosphere), and the unseen surface appears to be covered with organic material that has settled down from the clouds.

Figure 12.19 shows the probable structure of Titan's atmosphere. The diagram was drawn following the *Voyager 1* flybys, but it is largely consistent with the later data returned by *Cassini*. Despite the moon's low mass (a little less than twice that of Earth's Moon), and hence its low surface gravity (one-seventh of Earth's), the atmospheric pressure at ground level is 60 percent greater than on Earth. Titan's atmosphere contains about 10 times more gas than Earth's atmosphere. Titan's surface temperature is a frigid 94 K, roughly what we would expect on the basis of that moon's distance from the Sun. At the temperatures typical of the lower atmosphere, methane and ethane behave like water on Earth, raising the possibility of methane rain, snow, and fog and ethane rivers, lakes, and oceans! At higher levels in the atmosphere, the temperature rises, the result of photochemical absorption of solar radiation.


Because of Titan's weaker gravitational pull, the atmosphere extends some 10 times farther into space than does our own. The top of the main haze layer lies about 200 km above the surface, although there are additional layers, seen primarily through their absorption of ultraviolet radiation, at 300 km and 400 km. (See the inset in Figure 12.19). Below the haze the atmosphere is reasonably clear, although rather gloomy, because so little sunlight gets through. *Cassini* detected low-lying methane clouds at roughly the altitudes predicted by the model, although the clouds were less common than scientists had expected.

Why does Titan have such a thick atmosphere, when similar moons of Jupiter, such as Ganymede and Callisto, have none? The answer lies largely in Titan's low surface temperature, which makes it easier for Titan to retain an atmosphere. [More Precisely 8-1](#) Also, at such low temperatures, methane and ammonia, both of which were present in abundance at early times, would have been readily absorbed into Titan's icy surface. As a result, Titan was initially laden with much more methane and ammonia gas than was either Ganymede or Callisto.

As Titan's internal radioactivity warmed the moon, the ice released the trapped gases, forming a thick methane-ammonia atmosphere. Sunlight split the ammonia into hydrogen, which escaped into space, and nitrogen, which



◀ **FIGURE 12.19 Titan's Atmosphere** The structure of Titan's atmosphere, as deduced from *Voyager 1* observations. The solid blue line represents temperature at different altitudes. The inset shows the haze layers in Titan's upper atmosphere, depicted in false-color green above Titan's orange surface in this *Voyager 1* image. (NASA)



◀ **FIGURE 12.20 Titan Revealed** *Cassini's* telescopes captured this infrared, false-color view of Titan's surface in late 2004. The semicircular area near the center may be an old impact basin and the dark linear feature to its northwest perhaps mountain ranges caused by ancient tectonic activity. The inset shows a circular surface feature thought to be an icy volcano, further suggesting some geological activity on this icy moon's surface. Resolution of the larger image is 25 km; that of the inset is 10 times better. (NASA/ESA)





Figure 12.20 shows how *Cassini's* infrared instruments can penetrate the moon's atmosphere, revealing details of the surface. The image shows light and dark regions near the center of the field of view, thought to be icy plateaus, apparently smeared with hydrocarbon tar. Ridges and cracks on the moon's surface suggest that geological activity, in the form of "titanquakes," may be common. The rather blurred boundaries between the light and dark regions, the peculiar surface markings in the light-colored region, and the absence of extensive cratering suggest that some sort of erosion is occurring, perhaps as a result of wind or volcanic activity. Radar imaging reveals few large (10 to 100 km diameter) craters on the moon's surface, but there do not seem to be as many small craters as would be expected given Titan's location in Saturn's congested ring plane. The inset in the figure shows what may be an icy volcano, supporting the view that the moon's surface is geologically active.

remained in the atmosphere. The methane, which is more tightly bound and so was less easily broken apart, survived intact. Together with argon outgassed from Titan's interior, these gases form the basis of the atmosphere we see on Titan today. Detailed studies by *Cassini* scientists of the composition of Titan's atmosphere suggest that the atmosphere is steadily escaping, in large part due to the constant buffeting by Saturn's harsh magnetosphere, and may have been much thicker—perhaps 5–10 times denser—in the distant past.


## Titan's Surface and Interior Structure

*Cassini's* observations have refined our models of Titan's atmosphere, but they have revolutionized our knowledge of the moon's surface and interior. Under the frigid conditions found on Titan, as on Ganymede and Callisto, water ice plays the role of rock on Earth, and liquid water the role of lava. ∞ (Sec. 11.5) Before *Cassini* came on the scene, speculation about what might be found on Titan's surface ran the gamut from oceans of liquid methane or ethane to icy valleys laden with hydrocarbon sludge.

In January 2005, the *Huygens* probe, transported to Saturn by *Cassini* and released 3 weeks earlier, arrived at Titan and parachuted through the thick atmosphere to the moon's surface. Figure 12.21(a) shows an intriguing image radioed back from *Huygens* during its descent. It appears to show a network of drainage channels leading to a shoreline, but this interpretation remains uncertain. The lack of clouds just described meant that the methane rain supposedly responsible for the channels might not be as widespread as initially thought. Current models suggest that the rain may be a seasonal phenomenon, falling in winter and evaporating in summer. *Huygens* reached Titan near the end of winter at its landing site. The probe landed on solid ground, and for the next hour transmitted images and instrument readings to *Cassini* as it passed overhead. The view from the surface (Figure 12.21b) reveals a hazy view of an icy landscape. The "rocks" in the foreground are a few centimeters across and show evidence of erosion by liquid of some sort. Later detailed analysis of the data indicated that the lander skidded to a stop on a slushy surface covered with a thin solid layer—like frozen snow on Earth, except that it was most probably composed of ethane.





(a)



(b)

◀ **FIGURE 12.21****Titan's Surface**

(a) This photograph of the surface was taken from an altitude of 8 km as the *Huygens* probe descended. It shows a network of dark channels reminiscent of streams or rivers (at center) draining from the light-shaded uplifted terrain into darker, low-lying regions (at bottom). (b) *Huygens*'s view of its landing site, in approximately true color. The foreground icy "rocks" are only a few centimeters across. (NASA/ESA)



◀ **FIGURE 12.22 Titan's Lakes** Radar aboard *Cassini* detected many smooth regions (colored dark blue in this false-colored radio image), thought to be lakes of liquid methane, near Titan's north pole. The largest features are much smaller than any of the Great Lakes on Earth. (NASA/ESA)

In 2003, radio astronomers using the Arecibo telescope reported the detection of liquid hydrocarbon lakes on Titan's surface. ∞ (Sec. 5.5) In 2007, *Cassini* mission specialists confirmed this finding, presenting radar images showing numerous lakes, some many tens of kilometers in length, near Titan's north polar regions, at that time the coolest part of the moon. Figure 12.22 shows an example of these new images. As with the *Magellan* radar images of Venus, the darkest regions are extremely smooth, implying that they are composed of liquid; the shapes of the regions also strongly suggest liquid bodies.

Subsequently, *Cassini* confirmed the presence of liquid ethane in the lakes, although their exact composition remains uncertain. Methane is surely present: The methane rain feeds the lakes, but it is much more volatile than ethane under the conditions found on Titan, so it probably evaporates rapidly, leaving heavier hydrocarbons behind. Computer models indicate that the lakes are mostly ethane (75 percent), with methane (10 percent) and propane

(7 percent) making up much of the remainder. No waves have been detected on the lakes, hinting that they may also contain heavier tar-like hydrocarbons, increasing their viscosity. The number of lakes observed near Titan's south pole is significantly smaller than the number seen in the north, suggesting that the lakes grow during the rainy winter season.

Scientists hope that repeated observations on future Cassini passes will reveal changes in the size and structure of the lakes as spring on Titan gives way to summer. Exploration of the lakes by *Cassini* and future missions may afford scientists the opportunity to study the kind of chemistry thought to have occurred billions of years ago on Earth—the prebiotic chemical reactions that eventually led to life on our own planet (see Chapter 28).


Astronomers had also long expected that Titan's internal composition and structure would be similar to those of Ganymede and Callisto, because all three moons have quite similar masses, radii, and, hence, average densities. (Titan's density is  $1900 \text{ kg/m}^3$ . ∞ (Sec. 11.5) Titan contains a rocky core surrounded by a thick mantle of water ice. Each pass of *Cassini* allows scientists to probe the gravity of Titan, and repeated passes, coupled with knowledge of the properties of Titan's likely constituents, have resulted in the construction of some remarkably detailed models of the moon's interior. Figure 12.23 shows a recent such model. It indeed shows a rock and ice core and an icy mantle, but, intriguingly, also predicts the presence of a thick layer of liquid water a few tens of kilometers below the surface. Thus, Titan joins Europa, Ganymede, and Earth on the list of solar system objects containing large bodies of liquid water, with all that that implies for the prospects of life developing there. ∞ (Sec. 11.5)

### PROCESS OF SCIENCE Check

✓ Why are planetary scientists so interested in Titan?

## Saturn's Medium-Sized Moons

Saturn's complement of midsized moons consists (in order of increasing distance from the planet) of Mimas (at 3.1 planetary radii), Enceladus (4.0), Tethys (4.9), Dione (6.3), Rhea (8.7), and Iapetus (59.1). These moons are shown, to proper scale, in Figure 12.24. All six were known from Earth-based observations long before the Space Age. The inner five move on nearly circular trajectories, and all are tidally locked into synchronous rotation (so that one side always faces the planet) by Saturn's gravity. They therefore all have permanently "leading" and "trailing" faces as they move in their orbits, a fact that is important in understanding their often asymmetrical surface markings.




▲ **FIGURE 12.23 Titan's Interior** Based on measurements of Titan's gravitational field during numerous flybys, Titan's interior appears to be largely a rock-ice mixture. Most intriguing is the subsurface layer of liquid water, similar to that hypothesized on Jupiter's Europa and Ganymede.

Unlike the densities of the Galilean satellites of Jupiter, the densities of these six moons do not show any correlation with distance from Saturn. The densities of Saturn's midsized moons are all between  $1000$  and  $1400 \text{ kg/m}^3$ , implying that nearness to the central planetary heat source was a less important influence during their formation than it was in the Jupiter system. ∞ (Sec. 11.5) Scientists think that the midsized moons are composed largely of rock and water ice, like Titan. Their densities are lower than Titan's primarily because their smaller masses produce less compression of their interiors. All show heavy cratering, indicating the cluttered and violent planetary environment in the early solar system as fragments collided to form the outer planets and their moons. ∞ (Sec. 6.6)

The largest of the six, Rhea, has a mass only one-thirtieth that of Earth's Moon, and its icy surface is highly reflective and heavily cratered. At the low temperatures found on the surface of this moon, water ice is very hard and behaves rather like rock on the inner planets. For that reason, Rhea's surface craters look very much like craters on the Moon or Mercury. The density of craters is similar to that in the lunar highlands, indicating that the surface is old, and there is no evidence of extensive geological activity.

Prior to *Cassini*'s arrival, Rhea's main riddle was the presence of so-called *wispy terrain*—prominent light-colored streaks—on its trailing side (the right side of the image in Figure 12.24). The leading face, by contrast, shows no such markings, only craters. Astronomers thought that the wisps might have been caused by some event in the distant past during which water was somehow released from the interior and condensed on the surface. However, *Cassini* images reveal that the markings are in fact bright complexes of ice cliffs created by *tectonic fractures*, where stresses in the moon's icy interior as it cooled and solidified caused the surface layers to crack and



▲ **FIGURE 12.24 Saturn's Midsized Moons** Saturn's six medium-sized satellites as seen by the *Cassini* spacecraft and compared to Earth's Moon, all to scale. All are heavily cratered and all are shown here in natural color. (UC/Lick Observatory; NASA)

buckle. ∞ (Sec. 10.4) Any similar features on the leading side have presumably been obliterated by cratering, which should be more frequent on the satellite's forward-facing surface.


Inside Rhea's orbit lie the orbits of Tethys and Dione. These two moons are comparable to each other in size and have masses somewhat less than half the mass of Rhea. Like Rhea, they have reflective surfaces that are heavily cratered, but each shows signs of surface activity, too. Dione's trailing face (at the left of the moon in Figure 12.24) has prominent bright streaks, which *Cassini* revealed to be ice cliffs, as just described on Rhea. The cliffs cut across many craters, showing them to be considerably more recent than the period of heaviest bombardment. ∞ (Sec. 8.5) Dione also has "maria" of sorts, where flooding appears to have obliterated the older craters. The cracks on Tethys (upper left in Figure 12.24) may also be tectonic fractures, or could possibly be the result of a violent impact early in the moon's history (a large impact basin lies on the far side of the moon in the view shown here).

The innermost, and smallest, medium-sized moon is Mimas. Despite its low mass—only 1 percent the mass of Rhea—its closeness to the rings causes resonant interactions with the ring particles, resulting most notably in the Cassini division, as we have already seen. Possibly because of its proximity to the rings, Mimas is heavily cratered. The moon's chief surface feature is an enormous crater, called Herschel, on the leading face (at center in Figure 12.24). The diameter of this crater is almost one-third that of the moon itself. The impact that formed Herschel must have come very close to destroying Mimas completely. It is quite possible that the debris produced by such impacts is responsible for creating or maintaining the spectacular rings we see.

Enceladus, shown in Figure 12.25, orbits just outside Mimas. Its size, mass, composition, and orbit are so similar to those of Mimas that one might guess that the two moons would also be similar to each other in appearance and history. However, this is not so. Enceladus is so bright and shiny—it reflects virtually 100 percent of the sunlight



► **FIGURE 12.25 Enceladus** In this puzzling image taken by the *Cassini* spacecraft, Saturn's tiny ice-covered moon Enceladus shows evidence of youthful terrain in the south where craters are mostly absent. The long blue "tiger stripe" streaks (about 1 km wide) are fractures in the ice through which gas escapes to form a thin but real atmosphere. The inset shows some of the jets that are launched from geysers on the limb of this peculiar moon. (*JPL*)



falling on it—that astronomers think that its surface must be completely coated with fine crystals of pure ice, the icy “ash” of water “volcanoes” formed when liquid water emerges under pressure from the moon’s interior.

*Voyager* found that the moon bears visible evidence of large-scale volcanic activity of some sort. Much of its surface is devoid of impact craters, which seem to have been erased by what look like lava flows, except that the “lava” is water, temporarily liquefied during recent internal upheavals and now frozen again. Flybys of Enceladus by *Cassini* in 2008 and 2009 have detected what appears to the Earth-like plate tectonics near the moon’s south pole. The “tiger stripe” fractures visible in Figure 12.25 may be akin to spreading sites on Earth, such as the mid-Atlantic ridge. ∞ (Sec. 7.4) Arguing that the processes involved may actually be more similar to the geothermal activity found in many volcanic regions on Earth, some astronomers prefer to describe these features as *geysers*, rather than volcanoes. Similar activity has been found on Neptune’s moon Triton (see Section 13.5).


Long before *Cassini*’s arrival, the apparent association of Enceladus with the nearby thin cloud of small, reflective particles making up Saturn’s E ring provided strong circumstantial evidence that the moon is responsible for the ring. The E ring is known to be densest near Enceladus. Calculations indicate that the ring is unstable because of the disruptive effects of the solar wind, supporting the view that volcanism on Enceladus continually supplies new particles to maintain the ring.

*Cassini* confirmed much of the speculation about Enceladus’s internal activity and its connection with the E ring, finding evidence for every stage of the scenario just outlined. The probe detected icy jets emerging from geysers near the moon’s south pole (see the inset to Figure 12.25) and a transient water-vapor atmosphere surrounding the moon, densest around the south pole. *Cassini* also found a large increase in the density of E-ring particles near Enceladus as the atmosphere continually escapes from the moon’s

weak gravity. The geysers steadily replenish the atmosphere as it escapes to form the E ring. Interestingly, the jets have been found to contain both salt and organic material, fueling speculation about the possibility of a salty ocean just below Enceladus’s icy crust, as well as the likelihood of finding life there.

Why is there so much activity on such a small moon? The best explanation seems to be that the internal heating is the result of tidal stresses, much like those driving the volcanism on Io. ∞ (Sec. 11.5) Enceladus appears to be locked into an orbital resonance with some of the other moons, causing its orbit to be slightly nonspherical. Even though Saturn’s tidal force on Enceladus is only one-quarter of the force exerted by Jupiter on Io, the departure from perfect synchronism may be enough to cause the activity observed.

Why is the activity concentrated on the *south* pole? Instead of being one of the coolest spots on the moon, as one might expect based on the small amount of sunlight reaching the surface there, the south pole of Enceladus is actually several kelvins *warmer* than the equator! One possibility is that upwelling material at a warm spot on the moon may have caused the entire moon to “roll over” to place the low-density



◀ **FIGURE 12.26 Saturn's Diffuse Ring** This artist's conception shows a nearly invisible ring well beyond Saturn and its usual family of majestic rings; the planet itself appears as just a small dot at the center, which is then enlarged in the circular inset. Composed of thinly dispersed dust particles that radiate heat, the ring can be detected only in the infrared part of the spectrum—which is why it escaped detection until 2009. (SST/JPL)

warm region on the rotation axis—that is, at the pole—in much the same way as a spinning bowling ball will tend to rotate so as to place the (low-density) holes on the spin axis.

The outermost midsize moon is Iapetus, which orbits Saturn on a somewhat eccentric, inclined orbit with a semimajor axis of 3.6 million kilometers. Its mass is about three-quarters that of Rhea. Iapetus is a two-faced moon. The dark, leading face (at the bottom in Figure 12.24) reflects only about 3 percent of the sunlight reaching it, whereas the icy trailing side reflects 50 percent. Similar dark deposits seen elsewhere in the solar system are thought to be organic (containing carbon) in nature; they can be produced by the action of solar radiation on hydrocarbon (e.g., methane) ice.


Astronomers have long puzzled over how the dark markings could adorn only one side of Iapetus, as there seemed to be no nearby material for it to sweep up as it orbited the planet. However, in 2009 the *Spitzer Space Telescope* discovered a new huge, yet very diffuse ring more than 6 million km from Saturn, invisible in optical light but quite prominent in the infrared. ∞ (Sec. 5.7) As illustrated in Figure 12.26, Iapetus lies at the inner edge of this ring, and the steady accumulation of ring particles over billions of years could naturally account for the moon's asymmetric appearance.

Iapetus's other prominent surface feature is a giant 20-km-high, 1400-km-long ridge spanning half of the moon's circumference. Discovered by *Cassini* in 2005, it is clearly visible cutting across the bottom third of the moon in Figure 12.24. It is unique in the solar system, and so far defies explanation.

## The Small Satellites

Finally we come to Saturn's many small moons. Their masses are poorly known (they are inferred mainly from their gravitational effects on the rings), but they are thought to be similar in composition to the small moons of Jupiter. Most are less than a few tens of kilometers across. The largest, Hyperion and Phoebe, were discovered in the 19th century, in 1848 and 1898, respectively. The others have all been discovered since the latter part of the 20th century. Only the moons in or near the rings themselves were actually discovered by the *Voyager* and *Cassini* spacecrafts. These tiny bodies play no significant role in the “big picture” of the Saturn system, but their interactions with the rings and intricate dynamics have intrigued astronomers for decades.


Just 10,000 km beyond the F ring lie the so-called *co-orbital satellites* Janus and Epimetheus. As the name implies, these two satellites “share” an orbit, but in a very strange way. At any given instant, both moons are in circular orbits about Saturn, but one of them has a slightly smaller orbital radius than the other. Each satellite obeys Kepler's laws, so the inner satellite orbits slightly faster than the outer one and slowly catches up to it. The inner moon takes about 4 Earth years to “lap” the outer one. As the inner satellite gains ground on the outer one, a strange thing happens: As illustrated in Figure 12.27, when the two get close enough to begin to feel each other's weak gravity, they switch orbits—the new inner moon (which used to be the outer one) begins to pull away from its companion, and the whole process begins again! No one knows how the co-orbital satellites came



◀ **FIGURE 12.27 Orbit-Sharing Satellites** Saturn's co-orbital satellites Janus and Epimetheus play a never-ending game of tag while moving around the planet in their orbits. Their peculiar motions are depicted here by the labeled points that represent the locations of the two moons at a few successive times. The whole process then repeats, apparently forever.

to be engaged in this curious dance. Possibly they are portions of a single moon that broke up, perhaps after a meteoritic impact, leaving the two pieces in almost the same orbit.

In fact, many of the other small moons also share orbits, this time with larger moons. Telesto and Calypso have orbits that are synchronized with the orbit of Tethys, so that the two smaller moons always remain fixed relative to the larger moon, lying precisely  $60^\circ$  ahead of and  $60^\circ$  behind it as it travels around Saturn (see Figure 12.28). The moon Helene is similarly tied to Dione. These  $60^\circ$  points are known as **Lagrangian points**, after the French mathematician Joseph Louis Lagrange, who first studied them. Later we will see further examples of this special 1:1 orbital resonance in the motion of some asteroids about the Sun, trapped in the Lagrangian points of Jupiter's orbit.



▲ **FIGURE 12.28 Synchronous Orbits** The orbits of the moons Telesto and Calypso are tied to the motion of the moon Tethys. The combined gravitational pulls of Saturn and Tethys keep the small moons exactly  $60^\circ$  ahead and behind the larger moon at all times, so all three moons share an orbit and never change their relative positions.

### CONCEPT Check

- ✓ Why do Saturn's midsize moons show asymmetric surface markings?

**The Big Question** Looking at Saturn, whether in the spectacular images of this chapter or through a small telescope in someone's backyard, we have to wonder, Why does this planet have such awesome rings around it? Perhaps the loose rock and ice scattered throughout the rings is trash left over from the early days of the solar system—formative matter that never managed to coalesce into moons close to the planet. Or perhaps some of the moons did form early on, yet were summarily dismembered, owing to their flimsy material and proximity to the planet—literally torn apart by fierce gravitational tides. The *Cassini-Huygens* mission is now actively trying to address this deep astronomical puzzle.

# Chapter Review

## SUMMARY


1 Saturn was the outermost planet known to ancient astronomers. Its rings and moons were not discovered until after the invention of the telescope. The rings lie in the planet's equatorial plane, which is tilted at  $27^\circ$  with respect to the planet's orbit, so their appearance from Earth changes as Saturn orbits the Sun. Saturn is smaller than Jupiter, but still much larger than any of the terrestrial worlds. Like Jupiter, Saturn rotates rapidly, producing a pronounced flattening, and displays differential rotation. Strong radio emission from the planet's magnetosphere allows the rotation rate of the interior to be determined.



2 As on Jupiter, weather systems are seen on Saturn, although they are less distinct. Large storms are occasionally seen. Saturn has weaker gravity and a more extended atmosphere than Jupiter. The planet's overall butterscotch hue is due to cloud chemistry similar to that occurring in Jupiter's atmosphere. Saturn, like Jupiter, has bands, ovals, and turbulent flow patterns powered by convective motion in the interior. *Cassini* imaged the planet's south polar vortex.



3 Again like Jupiter, Saturn emits far more radiation into space than it receives from the Sun. Unlike Jupiter's, Saturn's excess energy emission is the result of **helium precipitation** (p. 297) in the planet's interior, where helium liquefies and forms droplets that then fall toward the center of the planet. This process is also responsible for Saturn's observed helium deficit. Saturn's interior is theoretically similar to that of Jupiter, but with a thinner layer of metallic hydrogen and a larger core. Its lower mass gives Saturn a less extreme core temperature, density, and pressure than Jupiter's core has. Saturn's conducting interior and rapid rotation produce a strong magnetic field and an extensive magnetosphere that contains the planet's ring system and many of the innermost moons.




4 From Earth, the main visible features of the rings are the **A, B, and C rings** (p. 298), the **Cassini division** (p. 298), and the **Encke gap** (p. 298). The rings are made up of trillions of icy particles ranging in size from dust grains to boulders. Their total mass is comparable to that of a small moon. Both divisions are dark because they are almost empty of ring particles. The main rings contain tens of thousands of narrow **ringlets** (p. 300). Interactions between the ring particles and the planet's inner moons are responsible for much of the fine structure observed. The narrow **F ring** (p. 303) lies just outside




the A ring and has a kinked, braided structure, apparently caused by two small **shepherd satellites** (p. 304) that orbit close to the ring and prevent it from breaking up. Beyond the F ring is the faint, narrow **G ring** (p. 301), whose sharp edges and bright arcs suggest a shepherd moonlet, although none has been found. The faint **D ring** (p. 301), lies between the C ring and Saturn's cloud layer. The diffuse **E ring** (p. 301) is associated with the moon Enceladus.

5 The **Roche limit** (p. 299) of a planet is the distance within which the planet's tidal field would overwhelm the internal gravity of a moon, tearing it apart and forming a ring. All known planetary ring systems lie inside their parent planets' Roche limits. Planetary rings may have lifetimes of only a few tens of millions of years. If so, the fact that we see rings around all four jovian planets means that they must constantly be reformed or replenished, perhaps by material chipped off moons by meteoritic impact or by the tidal destruction of entire moons.



6 Saturn's single large moon Titan is the second-largest moon in the solar system. Its thick atmosphere obscures the moon's surface and may be the site of complex cloud and surface chemistry. The moon's surface is so cold that water behaves like rock and liquid methane flows like water. Sensors aboard *Cassini* have allowed mission scientists to map the moon's surface for the first time, revealing evidence for ongoing erosion and volcanic activity. The *Huygens* probe landed on the icy surface and photographed what may be channels carved by flowing methane. The existence of Titan's atmosphere is a direct consequence of the cold conditions that prevailed at the time of the moon's formation.



7 The medium-sized moons of Saturn are made up predominantly of rock and water ice. They show a wide variety of surface terrains, are heavily cratered, and are tidally locked into synchronous orbits by the planet's gravity. The innermost midsized moon Mimas exerts an influence over the structure of the rings. The Cassini division is the result of resonance between ring particles there and Mimas. The moon Iapetus has an equatorial ridge and a marked contrast between its leading and trailing faces, and Enceladus has a highly reflective appearance, the result of water "volcanoes" on its surface. Rhea and Dione have extensive ice cliffs on their surfaces, the result of cracking of the outer layers as the moons cooled. Saturn's small moons exhibit a wide variety of complex motion. Several moons "share" orbits, in some cases lying at the **Lagrangian points** (p. 314)  $60^\circ$  ahead of and  $60^\circ$  behind the orbit of a larger moon.



**MasteringAstronomy**<sup>®</sup> For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- LO1** Why does Saturn have a less varied appearance than Jupiter? What does its shape tell us about its deep interior?
- Seen from Earth, Saturn's rings sometimes appear broad and brilliant, but at other times seem to disappear. Why?
- Compare and contrast the atmospheres and weather systems of Saturn and Jupiter, and tell how the differences affect each planet's appearance.
- LO2** Compare the thicknesses of Saturn's various layers (clouds, molecular hydrogen, metallic hydrogen, and core) with the equivalent layers in Jupiter. Why do the thicknesses differ?
- LO3** What mechanism is responsible for the relative absence of helium in Saturn's atmosphere, compared with Jupiter's atmosphere?
- LO4 POS** When were Saturn's rings discovered? When did astronomers realize what they were?
- LO5** What would happen to a satellite if it came too close to Saturn?
- POS** What evidence supports the idea that a relatively recent catastrophic event was responsible for Saturn's rings?
- What effect does Mimas have on Saturn's rings?
- What are shepherd satellites?
- LO6** Compare and contrast Titan with Jupiter's Galilean moons.
- Why does Titan have a dense atmosphere, whereas other large moons in the solar system don't?
- POS** What is the evidence for geological activity on Enceladus?
- LO7** What is the connection between Enceladus and Saturn's rings?
- Describe the behavior of Saturn's co-orbital satellites.

## Conceptual Self-Test: Multiple Choice

- VIS** From Figure 12.1 ("Ring Orientation"), the next time Saturn's rings will appear roughly edge-on as seen from Earth will be around (a) 2018; (b) 2022; (c) 2025; (d) 2035.
- Compared with the time it takes Jupiter to orbit the Sun once, the time it takes Saturn, which is twice as far away, to orbit the Sun is (a) significantly less than twice as long; (b) about twice as long; (c) significantly more than twice as long.
- Saturn's cloud layers are much thicker than those of Jupiter because Saturn has (a) more moons; (b) lower density; (c) a weaker magnetic field; (d) weaker surface gravity.
- VIS** According to Figure 12.5 ("Saturn's Zonal Flow"), the winds on Saturn are fastest at (a) the north pole; (b) 50° N latitude; (c) the equator; (d) 50° S latitude.
- Saturn's icy, rocky core is roughly (a) half the mass of; (b) the same mass as; (c) twice as massive as; (d) 10 times more massive than planet Earth.
- Of the following, which are most like the particles found in Saturn's rings? (a) house-sized rocky boulders; (b) grains of silicate sand; (c) asteroids from the asteroid belt; (d) fist-sized snowballs.
- A moon placed at a planet's Roche limit will (a) change color; (b) break into smaller pieces; (c) develop a magnetic field; (d) flatten into a disk.
- The atmospheric pressure at the surface of Titan is (a) less than; (b) about the same as; (c) about one-and-a-half times greater than; (d) about 16 times greater than the atmospheric pressure at Earth's surface.
- A tidally locked moon of Saturn (a) always presents the same face to the planet; (b) does not rotate; (c) always stays above the same point on the planet's surface; (d) maintains a constant distance from all the other moons.
- The moons Telesto and Calypso, orbiting at the Lagrangian points of Saturn and the moon Tethys (a) orbit twice as far from Saturn as does Tethys; (b) orbit closer to Saturn than does Tethys; (c) always stay the same distance apart; (d) always stay between Saturn and the Sun.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

1. ● What is the angular diameter of Saturn's A ring, as seen from Earth at closest approach?
2. ● What is the size of the smallest feature visible in Saturn's rings, as seen from Earth at closest approach with a resolution of  $0.05''$ ?
3. ●● What would be the mass of Saturn if it were composed entirely of hydrogen at a density of  $0.08 \text{ kg/m}^3$ , the density of hydrogen at sea level on Earth? Assume for simplicity that Saturn is spherical. Compare your answer with Saturn's actual mass and with the mass of Earth.
4. ● How long does it take for Saturn's equatorial flow, moving at  $1500 \text{ km/h}$ , to encircle the planet? Compare your answer with the wind-circulation time on Jupiter.
5. ● On the basis of the data given in Sections 12.1 and 12.3 (Figure 12.8), estimate the average density of Saturn's core.
6. ●● The text states that the total mass of material in Saturn's rings is about  $10^{15}$  tons ( $10^{18}$  kg). Suppose the average ring particle is  $6 \text{ cm}$  in radius (the size of a large snowball) and has a density of  $1000 \text{ kg/m}^3$ . How many ring particles are there?
7. ● What is the orbital speed of ring particles at the inner edge of the B ring, in kilometers per second? Compare your answer with the speed of a satellite in low Earth orbit ( $500 \text{ km}$  altitude, say). Why are these speeds so different?
8. ●● Assuming a spherical shape and a uniform density of  $2000 \text{ kg/m}^3$ , calculate how small an icy moon would have to be before a  $40 \text{ m/s}$  (about  $90 \text{ mph}$ ) fastball could escape.

## Activities


### Collaborative

1. Saturn moves more slowly among the stars than does any other visible planet. How many degrees per year does it move? Look in an almanac to see where the planet is now. What constellation is it in now? Can you see any atmospheric features? While looking at Saturn through a telescope, can you see any of its moons? They line up with the rings. How many can you see? Can you identify them using the almanac? Titan is often the farthest moon out and is always the brightest. By observing it a couple of times per night over a period of 2–3 weeks, can

you determine the radius (relative to Saturn) and period of its orbit?

### Individual

1. Binoculars may not reveal the rings of Saturn, but most small telescopes will. Use a telescope to look at Saturn. Does Saturn appear flattened? Examine the rings, and sketch what you see. How are they tilted? Can you see a dark line in the rings? This is the Cassini division. It once was thought to be a gap in the rings, but *Voyager* found that it is filled with tiny ringlets. Can you see the shadow of the rings on Saturn?



# Uranus and Neptune

## THE OUTER WORLDS OF THE SOLAR SYSTEM

The two outermost planets were unknown to the ancients and were discovered by telescopic observations: Uranus in 1781 and Neptune in 1846. Uranus and Neptune have similar bulk properties, so it is natural to consider them together; they are part of the jovian family of planets.

Yet as we study the properties of Uranus and Neptune in more detail, we find important differences between the outer and inner jovian worlds. The two outermost jovian planets are the smallest and least massive of the four, and their internal structures and the details of their atmospheric composition differ significantly from those of their larger jovian cousins. Their moons and rings, too, deviate from those found around Jupiter and Saturn. These differences are not mere anomalies; rather, they have much to tell us about the environment in which the outer planets formed and evolved.

**The Big Picture** The outer planets likely played important roles in the origin and evolution of our solar system. However, Uranus and Neptune reside at truly remote parts of our planetary family, and it's not easy to reach them even with robots. They have so far been explored only briefly by a few spacecraft speeding by on their way toward interstellar space. Given other priorities of the U.S. space program, it will likely be many years before robotic devices revisit these distant, mysterious worlds.

**LEFT:** Uranus, the seventh planet out from the Sun, is often viewed as bland, featureless, and boring. However, this new image reveals a surprising amount of activity within its gas-enshrouded clouds that display bands reminiscent of those on Jupiter and Saturn. Among the finest, highest-resolution photos ever taken of Uranus, it also shows circulating clouds, massive hurricanes, and unusual convective features near its north pole (at right), as well as a very thin ring. The image is actually an infrared view of radiation from the Sun reflected from the planet and captured by a new camera on an exceptionally clear night at the Keck Observatory. (L. Sromovsky and P. Fry)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Describe how both chance and calculation played major roles in the discoveries of the outer planets.
- 2 Summarize the similarities and differences between Uranus and Neptune, and compare these planets with the other two jovian worlds.
- 3 Describe what is known about the interiors of Uranus and Neptune.
- 4 Explain what the moons of the outer planets tell us about their past.
- 5 Contrast the rings of Uranus and Neptune with those of Jupiter and Saturn.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.



## 13.1 The Discoveries of Uranus and Neptune

The two outermost planets, Uranus and Neptune, were unknown to ancient astronomers. Both were discovered after the dawn of the modern scientific age, and both discoveries are testaments to the power of two pillars of modern science—improved *technology* and mathematical *modeling*. ∞ (Sec. 1.2)

### Uranus

The planet Uranus was discovered by British astronomer William Herschel in 1781. Herschel was engaged in charting the faint stars in the sky when he came across an odd-looking object that he described as “a curious either nebulous star or perhaps a comet.” Repeated observations showed that it was neither. The object appeared as a disk in Herschel’s 6-inch telescope and moved relative to the stars, but it traveled too slowly to be a comet. Herschel soon realized that he had found the seventh planet in the solar system.


This was the first new planet discovered in well over 2000 years, and the event caused quite a stir at the time. The story goes that Herschel’s first instinct was to name the new planet “Sidus Georgium” (Latin for “George’s star”), after his king, George III of England. The world was saved from a planet named George by the wise advice of another astronomer, Johann Bode, who suggested instead that the tradition of using names from Greco–Roman mythology be continued and that the planet be named Uranus, after the father of Saturn.

Uranus is in fact just barely visible to the naked eye if you know exactly where to look. At opposition, it has a maximum angular diameter of 4.1” and shines just above the unaided eye’s threshold of visibility. It looks like a faint, undistinguished star. No wonder it went unnoticed by the ancients. Even today, few astronomers have seen it without a telescope.

Through most large Earth-based optical telescopes (Figure 13.1), Uranus appears hardly more than a tiny pale greenish disk. With the flyby of *Voyager 2* in 1986, our knowledge of Uranus increased dramatically, although close-up images of the planet still showed virtually no surface detail (Figure 13.2). Not until the chapter-opening photo on page 318 was acquired under ideal observing conditions and with state-of-the-art technology did Uranus reveal bands and spots reminiscent of those on all the other jovian worlds.

### Neptune

Following the discovery of Uranus, astronomers set about charting its orbit and quickly discovered a small discrepancy between the planet’s predicted position and where they actually observed it. Try as they might, astronomers could not find an elliptical orbit that fit the planet’s trajectory to within the accuracy of their measurements. Half a century



▲ **FIGURE 13.1 Uranus from Earth** Details are virtually invisible on photographs of Uranus made with most Earth-based telescopes. (Arrows point to three of the planet’s moons.) (UC/Lick Observatory)

after Uranus’s discovery, the discrepancy had grown to a quarter of an arc minute, far too big to be explained away as observational error.

The logical conclusion was that an unknown body must be exerting a gravitational force on Uranus—much weaker



▲ **FIGURE 13.2 Uranus, Close Up** This image of Uranus, taken from a distance of about 1 million kilometers, was sent back to Earth by the *Voyager 2* spacecraft as it whizzed past the giant planet at 10 times the speed of a rifle bullet. Color is natural and the upper atmosphere is nearly featureless, except for a few wispy clouds in the northern hemisphere. (NASA)

than that of the Sun, but still measurable. But what body could that be? Astronomers realized that there had to be *another* planet in the solar system perturbing Uranus's motion.

In the 1840s, two mathematicians independently solved the difficult problem of determining the new planet's mass and orbit. A British astronomer, John Adams, reached the solution in September 1845; in June of the following year, the French mathematician Urbain Leverrier came up with essentially the same answer. British astronomers seeking the new planet found nothing during the summer of 1846. In September, a German astronomer named Johann Galle began his own search from the Berlin Observatory, using a newly completed set of more accurate sky charts. He found the new planet within one or two degrees of the predicted position—on his first attempt. After some wrangling over names and credits, the new planet was named Neptune, and Adams and Leverrier (but not Galle!) are now jointly credited with its discovery.

With an orbital period of 163.7 Earth years, Neptune is only now completing one revolution since its discovery. Unlike Uranus, distant Neptune cannot be seen with the naked eye, although it can be seen with a small telescope—in fact, according to his notes, Galileo might actually have seen Neptune, although he had no idea what it really was at the time. Through a large telescope, Neptune appears as a bluish disk, with a maximum angular diameter of  $2.4''$  at opposition.

Figure 13.3 shows a long Earth-based exposure of Neptune and its largest moon, Triton. Neptune is so distant that surface features on the planet are virtually impossible to discern. Even under the best observational conditions, only a few markings can be seen. These features are suggestive of multicolored cloud bands, with light bluish hues seeming



▲ **FIGURE 13.3 Neptune from Earth** Neptune and two of its moons, Triton (left arrow) and Nereid (right), imaged with a large Earth-based telescope. (UC/Lick Observatory)

to dominate. With *Voyager 2*'s arrival, much more detail emerged, as shown in Figure 13.4. Superficially, at least, Neptune resembles a blue-tinted Jupiter, with atmospheric bands and spots clearly evident.

### PROCESS OF SCIENCE Check

- ✓ How did observations of the orbit of Uranus lead to the discovery of Neptune?



(a)



(b)

◀ **FIGURE 13.4 Neptune, Close Up** (a) Neptune as seen in natural color by *Voyager 2*, from a distance of roughly 1 million kilometers. (b) A closer view, resolved to about 10 km, shows cloud streaks ranging in width from 50 km to 200 km. (NASA)

## 13.2 Orbital and Physical Properties

With orbital semimajor axes of 19.2 and 30.1 AU, respectively, and orbital periods on the order of a century, Uranus and Neptune lie in the outer reaches of the Sun's planetary family. Their orbits lie just inside the Kuiper belt, for which (see Chapters 14 and 15) Uranus and Neptune are largely responsible.


Figure 13.5 shows Uranus and Neptune to scale, along with Earth for comparison. The two giant planets are quite similar to one another in their bulk properties. The radius of Uranus is 4.0 times that of Earth, that of Neptune 3.9 Earth radii. Their masses (first determined from terrestrial observations of their larger moons and later refined by *Voyager 2*) are 14.5 Earth masses for Uranus and 17.1 Earth masses for Neptune. Uranus's average density is  $1300 \text{ kg/m}^3$ , and Neptune's is  $1600 \text{ kg/m}^3$ . These densities imply that large rocky cores constitute a greater fraction of the planets' masses than do the cores of either Jupiter or Saturn. The cores themselves are probably comparable in size, mass, and composition to those of the two larger giants.


Like the other jovian planets, Uranus has a short rotation period. Earth-based observations of the Doppler shifts in spectral lines first indicated that Uranus's "day" was between 10 and 20 hours long. The precise value of the planet's rotation period—accurately determined when *Voyager 2* timed radio signals associated with Uranus's magnetosphere—is now known to be 17.2 hours. Again,

as with Jupiter and Saturn, the planet's atmosphere rotates differentially. However, Uranus's atmosphere actually rotates *faster* at the poles (where the period is 14.2 hours) than near the equator (where the period is 16.5 hours).


Each planet in our solar system seems to have some outstanding peculiarity, and Uranus is no exception. Unlike all the other planets, whose spin axes are roughly perpendicular to the plane of the ecliptic, Uranus's axis of rotation lies almost within that plane— $98^\circ$  from the perpendicular, to be precise. (Because the north pole lies below the ecliptic plane, the rotation of Uranus, like that of Venus, is classified as retrograde.) We might say that, relative to the other planets, Uranus lies tipped over on its side. As a result, the north (spin) pole of Uranus, at some time in its orbit, points almost directly toward the Sun.\* Half a Uranus year later, its south pole faces the Sun, as illustrated in Figure 13.6. When *Voyager 2* encountered the planet in 1986, the north pole happened to be pointing nearly at the Sun, so it was mid-summer in the northern hemisphere.

The strange orientation of Uranus's rotation axis produces some extreme seasonal effects. Starting at the height of northern summer, when the north pole points closest to the Sun, an observer near that pole would find the Sun would never set. Rather, it would appear to move in a small circle in the sky around the planet's north celestial pole as the planet

\*As in Chapter 9, we adopt the convention that a planet's rotation is always counterclockwise as seen from above the north pole (i.e., planets always rotate from west to east).  (Sec. 9.2).



◀ **FIGURE 13.5 Jovian Planets** Jupiter, Saturn, Uranus, and Neptune, showing their relative sizes compared to Earth. Uranus and Neptune are quite similar in their bulk properties, each one probably having a core about 10 times more massive than Earth. Jupiter and Saturn are both much larger, but their rocky cores are probably comparable in mass to those of Uranus and Neptune. (NASA)



▲ **FIGURE 13.6 Seasons on Uranus** Because of Uranus's axial tilt of  $98^\circ$ , the planet experiences the most extreme seasons known in the solar system.

rotated, completing one circuit (counterclockwise) every 17 hours. ∞ (Sec. 1.3) Over time, as Uranus moved along its orbit and its rotation axis pointed farther and farther from the Sun, the circle would gradually increase in size, with the Sun dipping slightly lower in the sky each day. Eventually, the Sun would begin to set and rise again in a daily cycle, and the nights would grow progressively longer with each passing day. Twenty-one Earth years after the summer solstice, the autumnal equinox would occur, with day and night each 8.5 hours long.

The days would continue to shorten, until one day the Sun would fail to rise at all. The ensuing period of total darkness would be equal in length to the earlier period of constant daylight, plunging the northern hemisphere into the depths of winter. Eventually, the Sun would rise again; the days would lengthen through the vernal equinox and beyond, and in time the observer would again experience a long summer of uninterrupted (though dim) sunshine. From the point of view of an observer on the equator, by contrast, summer and winter would be almost equally cold seasons, with the Sun never rising far above the horizon. Spring and fall would be the warmest times of year, with the Sun passing almost overhead each day.

Astronomers are unsure why Uranus is tilted in this way—the other planets all have rotation axes lying well out of the ecliptic plane. For many years the leading explanation has been that a single catastrophic event during the late formative stages of the solar system, such as a glancing collision between the planet and another body a few times

more massive than Earth, abruptly altered Uranus's spin. ∞ (Sec. 6.7) However, recent computer simulations suggest that, if that were the case, the planet's moon system would not have been able to "keep up" with the sudden change, and the moons would not have the prograde, equatorial orbits now observed. Instead, it seems more likely that the shift in the planet's axis occurred as a result of two or more smaller (but still substantial) impacts gentle enough to allow the moons to retain their orbits. The problem is that this runs counter to the established view that such encounters were rare in the outer solar system. The resolution may force astronomers to rethink some details of the condensation theory (described in Chapter 6).

Neptune's clouds show more variety and contrast than do those of Uranus, and Earth-based astronomers studying them determined a rotation rate for Neptune even before *Voyager 2*'s flyby in 1989. The average rotation period of Neptune's atmosphere is 17.3 hours (quite similar to that of Uranus). Measurements of Neptune's radio emissions by *Voyager 2* showed that the magnetic field of the planet, and presumably also its interior, rotates once every 16.1 hours. Thus, Neptune is unique among the jovian worlds in that its atmosphere rotates *more slowly* than its interior. Neptune's axis of rotation is inclined  $29.6^\circ$  to a line perpendicular to the planet's orbital plane, quite similar to the  $27^\circ$  tilt of Saturn.

### CONCEPT Check

✓ What is unusual about the rotation of Uranus?

## 13.3 The Atmospheres of Uranus and Neptune

### Composition

Spectroscopic studies of sunlight reflected from Uranus's and Neptune's dense clouds indicate that the two planets' outer atmospheres (the parts we actually measure spectroscopically) are quite similar to the atmospheres of Jupiter and Saturn. The most abundant element is molecular hydrogen ( $H_2$ , 84 percent), followed by helium (He, about 14 percent) and methane ( $CH_4$ ), which is more abundant on Neptune (about 3 percent) than on Uranus (2 percent). Ammonia ( $NH_3$ ), which plays such an important role in the Jupiter and Saturn systems, is not present in any significant quantity in the outermost jovian worlds.

The abundances of gaseous ammonia and methane vary systematically among the jovian planets. Jupiter has much more gaseous ammonia than methane, but moving outward from the Sun, we find that the more distant planets have steadily decreasing amounts of ammonia and relatively greater amounts of methane. The reason for this variation is temperature. Ammonia gas freezes into ammonia ice crystals at about 70 K. This temperature is cooler than the cloud-top temperatures of Jupiter and Saturn, but warmer than those of Uranus (58 K) and Neptune (59 K). Thus, the outermost jovian planets have little or no *gaseous* ammonia in their atmospheres, so their spectra (which record atmospheric gases only) show only traces of ammonia.

The increasing amounts of methane are largely responsible for the outer jovian planets' blue coloration. Methane absorbs long-wavelength red light quite efficiently, so sunlight reflected from the planets' atmospheres is deficient in red and yellow photons and appears


bluish-green or blue. As the concentration of methane increases, the reflected light should appear bluer—just the trend that is observed: Uranus, with less methane, looks bluish-green, whereas Neptune, with more, looks distinctly blue.

### Weather

*Voyager 2* detected just a few cloud features in Uranus's atmosphere (Figure 13.2), and even those became visible only after extensive computer enhancement. Figure 13.7 shows a series of *Hubble Space Telescope* views of the planet. Parts (a) through (c) are heavily processed optical images that show the progress of a pair of bright clouds around the planet. Part (d) shows a false-color, near-infrared rendition of Uranus. The colors in this image generally indicate the depth to which we can see into the atmosphere. Blue-green regions are clear atmospheric regions where astronomers can study conditions down to the lower cloud levels. Yellow-gray colors show sunlight reflecting from higher cloud layers or from atmospheric haze. Orange-red colors, such as the prominent "spots" on the south (right) edge of this image, indicate very high clouds, much like the wispy, white cirrus clouds often seen at high altitudes in Earth's atmosphere. Like cirrus clouds on Earth, these Uranian clouds are made up predominantly of ice crystals, formed in the planet's cold upper atmosphere.

Uranus apparently lacks any significant internal heat source, and because the planet has a low surface temperature, its clouds are found only at low-lying, warmer levels in

▼ **FIGURE 13.7 Uranus's Rotation** (a), (b), and (c) These computer-enhanced *Hubble Space Telescope* images, taken at roughly 4-hour intervals, show the motion of a pair of bright clouds (labeled A and B) in Uranus's southern hemisphere. (The numbers at the top give the time of each photo.) (d) An infrared image of Uranus shows that planet's ring system, as well as a number of clouds (pink and red regions) in the upper atmosphere. (NASA)




the atmosphere. The absence of high-level clouds means that we must look deep into the planet's atmosphere to see any structure, so the bands and spots that characterize flow patterns on the other jovian worlds are largely "washed out" (in visible light) on Uranus by intervening stratospheric haze. Take another look at the chapter-opening image of Uranus showing its atmospheric structure more clearly. Note the mottled structure near the north pole (at right), which scientists think may be a polar vortex similar to the one seen on Venus. ∞(Sec. 9.5)

From computer-processed images such as those shown in Figure 13.7, astronomers have learned that Uranus's atmospheric clouds and flow patterns move around the planet in the same sense as the planet's rotation, with wind speeds ranging from 200 to 500 km/h. In fact, tracking these clouds allowed Uranus's differential rotation, mentioned earlier, to be measured. Despite the odd angles at which sunlight may strike the surface, the planet's rapid rotation still channels the wind flow into bands reminiscent of those found on Jupiter and Saturn. Even though the predominant flow is in the east–west direction, the atmosphere seems to be quite efficient at transporting energy between the northern and southern hemispheres. For example, during the *Voyager 2* flyby in 1986, although much of the south

was in total darkness, the temperature there was only a few kelvins less than in the north.

Neptune's clouds and band structure are much more easily seen than Uranus's. Although Neptune lies at a greater distance from the Sun, the planet's upper atmosphere is actually slightly warmer than that of Uranus. Like Jupiter and Saturn, but unlike Uranus, Neptune has an internal energy source—in fact, Neptune radiates 2.7 times more heat than it receives from the Sun. The cause of this heating is still uncertain. Some scientists have suggested that Neptune's excess methane has helped "insulate" the planet, tending to maintain its initially high internal temperature. If that is so, then the source of Neptune's internal heat is the same as Jupiter's: energy left over from the planet's formation. ∞ (Sec. 11.3) The combination of extra heat and less haze may be responsible for the greater visibility of Neptune's atmospheric features (see Figure 13.8), as its cloud layers lie at higher levels in the atmosphere than do those of Uranus.


Neptune sports several storm systems similar in appearance to those seen on Jupiter (and assumed to be produced and sustained by the same basic processes). The largest such storm, known simply as the **Great Dark Spot**, is shown in Figure 13.8(a). ∞ (Sec. 11.2) Discovered by *Voyager 2* in



(a)

◀ **FIGURE 13.8 Neptune's Dark Spot**

(a) Close-up views, taken by *Voyager 2* of the Great Dark Spot of Neptune, show a large storm system in the planet's atmosphere, possibly similar in structure to Jupiter's Great Red Spot. The entire dark spot is roughly the size of Earth. (b) These *Hubble Space Telescope* views of Neptune were taken years apart (as marked). Some cloud features (mostly methane ice crystals) are tinted pink here because they were imaged in the infrared, but they are really white in visible light. Note that the Great Dark Spot has disappeared in recent years, for unknown reasons. (NASA)



(b)

1989, the spot was about the size of Earth, was located near the planet's equator, and exhibited many of the same general characteristics as the Great Red Spot on Jupiter. The flow around it was counterclockwise, as with the Red Spot, and there appeared to be turbulence where the winds associated with the Great Dark Spot interacted with the zonal flow to its north and south. The flow around this and other dark spots may drive updrafts to high altitudes, where methane crystallizes out of the atmosphere to form high-lying cirrus clouds—those visible in Figure 13.8(a) lie some 50 km above the main cloud tops. Astronomers did not have long to study the Dark Spot's properties, however: As shown in Figure 13.8(b), when the *Hubble Space Telescope* viewed Neptune after the mid-1990s, the spot had vanished, although several new storms (bright spots) had appeared.

Infrared views such as those shown in Figure 13.8(b) reveal Neptune's dynamic weather patterns. The planet's weather can change in as little as a few rotation periods, and winds blow at speeds in excess of 1500 km/h—almost half the speed of sound in Neptune's upper atmosphere—with storms the size of Earth more the rule than the exception. The planet's stormy disposition is well established, but very difficult to understand. On Earth, weather systems are driven by the heat of the Sun. However, Neptune lies far from the Sun, in the outer solar system, and the Sun's heating effect is minuscule—nearly a thousand times less than at Earth. How can Neptune be so cold, yet so active?

Intriguingly, the three images of Neptune shown in Figure 13.8(b) reveal that the planet's southern hemisphere has brightened significantly over the period shown. Apparently, despite the Sun's faint heating, the planet is responding to the increase in solar energy as its southern half slowly moves from spring into summer. Infrared VLT observations indicate that the south polar region, which has been sunlit for the past three decades, is about 10 K warmer than the rest of the planet.

### CONCEPT Check

✓ Why are planetary scientists puzzled by the strong winds and rapidly changing storm systems on Neptune?

## 13.4 Magnetospheres and Internal Structure

*Voyager 2* found that both Uranus and Neptune have fairly strong internal magnetic fields—about a hundred times stronger than Earth's field and one-tenth as strong as Saturn's. However, because Uranus and Neptune are so much larger than Earth, the magnetic fields at the cloud tops—spread out over far larger volumes than is the field on Earth—are actually comparable in strength to Earth's field. Uranus and Neptune each have substantial magnetospheres,


populated largely by electrons and protons either captured from the solar wind or created from ionized hydrogen gas escaping from the planets themselves.

When *Voyager 2* arrived at Uranus, it discovered that the planet's magnetic field was tilted at about 60° to the axis of rotation. On Earth, such a tilt would put the north magnetic pole somewhere in the Caribbean. Furthermore, on Uranus, the magnetic field lines are *not* centered on the planet. It is as though Uranus's field were due to a bar magnet that is tilted with respect to the planet's rotation axis and displaced from the center by about one-third the radius of the planet. Figure 13.9 shows the magnetic field structures of the four jovian planets, with Earth's also shown for comparison. The locations and orientations of the bar magnets represent the observed planetary fields, and the sizes of the bars indicate magnetic field strength.


Because dynamo theories generally predict that a planet's magnetic axis should be roughly aligned with its rotation axis—as on Earth, Jupiter, Saturn, and the Sun—the misalignment on Uranus suggested to some researchers that perhaps the planet's field had been caught in the act of reversing. ∞ (Sec. 7.5) Another possibility was that the oddly tilted field was in some way related to the planet's axial tilt—perhaps one catastrophic collision skewed both axes at the same time. Those ideas evaporated in 1989 when *Voyager 2* found that Neptune's field is also inclined to the planet's axis of rotation, at an angle of 46° (see Figure 13.9d), and also substantially offset from the center of the planet. It now appears that the internal structures of Uranus and Neptune are different from those of Jupiter and Saturn, and this difference changes how the former planets' magnetic fields are generated.

Theoretical models indicate that Uranus and Neptune have rocky cores similar to those found in Jupiter and Saturn—about the size of Earth and perhaps 10 times more massive. However, the pressure outside the cores of Uranus and Neptune (unlike the pressure within Jupiter and Saturn) is too low to force hydrogen into the metallic state, so hydrogen stays in its molecular form all the way into the planets' cores. Astronomers theorize that deep below the cloud layers, Uranus and Neptune may have high-density, “slushy” interiors containing thick layers of water clouds. It is also possible that much of the planets' ammonia is dissolved in the hypothetical water, accounting for the absence of ammonia at higher levels. Such an ammonia solution would provide a thick, electrically conducting ionic layer that could conceivably explain the planets' misaligned magnetic fields if the circulating electrical currents that generate the fields occur mainly in regions far from the planets' centers and rotation axes.

At present, we simply don't know enough about the interiors of Uranus and Neptune to assess the correctness of this picture. Our current state of knowledge is summarized in Figure 13.10, which compares the internal structures of the four jovian worlds.



**Interactive FIGURE 13.9 Jovian Magnetic Fields** Comparison of the magnetic field strengths, orientations, and offsets in the four jovian planets. The planets are drawn to scale, and in each case the magnetic field arises from an imaginary bar magnet. The size and location of each magnet represent the strength and orientation of the planetary field.



**Interactive FIGURE 13.10 Jovian Interiors** A comparison of the interior structures of the four jovian planets. (a) The planets drawn to scale. (b) The relative proportions of the various internal zones.



**CONCEPT Check**

✓ What is odd about the magnetic fields of Uranus and Neptune?

## 13.5 The Moon Systems of Uranus and Neptune

Like Jupiter and Saturn, both Uranus and Neptune have extensive moon systems, each consisting of a few large moons, long known from ground-based observations, and many smaller moonlets, discovered by *Voyager 2* or recently detected from Earth.

### Uranus's Moons

As of 2013, 27 moons are known to orbit Uranus. The properties of those more than 300 km in diameter are listed in Table 13.1.

William Herschel discovered and named Titania and Oberon, the two largest of Uranus's five major moons, in 1789. British astronomer William Lassell found Ariel and Umbriel, the next-largest moons, in 1851. Gerard Kuiper found Miranda, the smallest, in 1948. In order of increasing distance from the planet, they are Miranda (at 5.1 planetary radii), Ariel (7.5), Umbriel (10.4), Titania (17.1), and Oberon (22.8). Ten smaller moons discovered by *Voyager 2* all lie inside the orbit of Miranda. Many of them are intimately related to the Uranian ring system. All of these moons revolve in the planet's skewed equatorial plane, almost perpendicular to the ecliptic, in circular, tidally locked orbits, sharing their parent's extreme seasons.

Of the remaining 22 moons, one, orbiting close to the planet, was found after careful reanalysis of *Voyager 2*

images. All the rest were discovered via systematic ground-based searches made since 1997, with techniques similar to those that have been so successful in identifying new moons of Jupiter and Saturn. ∞ (Sec. 12.5) These small bodies orbit far from Uranus, mostly on retrograde, highly inclined orbits. Like the outer moons of Jupiter and Saturn, and like Phobos and Deimos of Mars, each is thought to be interplanetary debris captured following a glancing encounter with the planet's atmosphere.

The five largest Uranian moons are similar in many respects to the six midsize moons of Saturn. ∞ (Sec. 12.5) Their densities lie in the range from 1100 to 1700 kg/m<sup>3</sup>, suggesting a composition of ice and rock, like Saturn's moons, and their diameters range from 1600 km for Titania and Oberon, to 1200 km for Umbriel and Ariel, to 480 km for Miranda. Uranus has no moons comparable to the Galilean satellites of Jupiter or to Saturn's single large moon, Titan. Figure 13.11 shows Uranus's five large moons to scale, along with Earth's Moon and Neptune's two midsize moons (named Proteus and Nereid) for comparison.

The outermost of the five moons, Titania and Oberon, are heavily cratered and show little indication of geological activity. Their overall appearance (and quite possibly their history) is comparable to that of Saturn's moon Rhea, except that they lack Rhea's wispy streaks. Also, like all Uranian moons, they are considerably less reflective than Saturn's satellites, suggesting that their icy surfaces are quite dirty.

One possible reason for the lesser reflectivity may simply be that the planetary environment in the vicinity of Uranus and Neptune contains more small "sooty" particles than do the parts of the solar system that are closer to the Sun. An alternative explanation, now considered more likely


**TABLE 13.1** The Major Moons of Uranus and Neptune\*

Name	Distance from Planet		Orbital Period (days)	Size (longest diameter, km)	Mass** (Earth Moon masses)	Density	
	(km)	(planetary radii)				(kg/m <sup>3</sup> )	(g/cm <sup>3</sup> )
Miranda (U)	130,000	5.08	1.41	480	0.00090	1100	1.1
Ariel (U)	191,000	7.48	2.52	1160	0.018	1600	1.6
Umbriel (U)	266,000	10.4	4.14	1170	0.016	1400	1.4
Titania (U)	436,000	17.1	8.71	1580	0.048	1700	1.7
Oberon (U)	583,000	22.8	13.5	1520	0.041	1600	1.6
Proteus (N)	118,000	4.75	1.12	440			
Triton (N)	355,000	14.3	-5.88 <sup>†</sup>	2710	0.292	2100	2.1
Nereid (N)	5,510,000	223	360	340	0.0000034	1200	1.2

\* Only moons larger than 300 km in diameter are listed.

\*\* Mass of Earth's Moon =  $7.4 \times 10^{22}$  kg =  $8.5 \times 10^{-4}$  Uranus mass =  $7.3 \times 10^{-4}$  Neptune mass.

<sup>†</sup> Retrograde orbit.



▲ **FIGURE 13.11 Moons of Uranus and Neptune** The five largest moons of Uranus and two midsize moons of Neptune (Proteus and Nereid) are shown to scale, with part of Earth's Moon (also to scale) for comparison. The smallest details visible on these moons are about 15 km across. (NASA; *Lick Observatory*)


by many planetary scientists, cites the effects of radiation and high-energy particles that strike the surfaces of these moons. The impacts tend to break up the molecules on the moons' surfaces, eventually leading to chemical reactions that slowly build up a layer of dark, organic material. This **radiation darkening** is thought to contribute to the generally darker coloration of many of the moons and rings in the outer solar system. In either case, the longer a moon has been inactive and untouched by meteoritic impact, the darker its surface should be.

The darkest of the moons of Uranus is Umbriel. This moon displays little evidence of any past surface activity; its only mark of distinction is a bright spot about 30 km across, of unknown origin, in its northern hemisphere. By contrast, Ariel, similar in size to Umbriel, but closer to Uranus, does appear to have undergone some activity in the past. Ariel shows signs of resurfacing in places and exhibits surface cracks a little like those seen on another of Saturn's moons, Tethys. However, unlike Tethys, whose cracks are probably due to meteoritic impact, Ariel's activity likely occurred when internal forces and external tidal stresses (due to

the gravitational pull of Uranus) distorted the moon and cracked its surface.

Strangest of all of Uranus's icy moons is Miranda, shown in Figure 13.12. Before the *Voyager 2* encounter, astronomers expected that Miranda would resemble Mimas, the moon of Saturn whose size and location it most closely approximates. However, instead of being a relatively uninteresting, cratered, geologically inactive world, Miranda displays a wide range of surface terrains, including ridges, valleys, large oval faults, and many other tortuous geological features.

To explain why Miranda seems to combine so many different types of surface features, some researchers have hypothesized that this baffling object has been catastrophically disrupted several times (by internal or external processes), with the pieces falling back together in a chaotic, jumbled way. Certainly, the frequency of large craters on the outer moons suggests that destructive impacts may once have been quite common in the Uranian system. It will be a long time, though, before we can obtain more detailed information to test this theory.



▲ **FIGURE 13.12 Miranda** The asteroid-sized innermost moon of Uranus, photographed by *Voyager 2*. Miranda has a fractured surface suggestive of a violent past, but the cause of the grooves and cracks is currently unknown. Resolution in the inset is a remarkable 2 km. The long “canyon” near the bottom of the inset is nearly 20 km deep. (NASA)

## Neptune’s Moons

From Earth, we can see only 2 moons orbiting Neptune. William Lassell discovered Triton, the inner moon, in 1846. The moon Nereid was located by Gerard Kuiper in 1949. *Voyager 2* discovered 6 additional moons, all less than a few hundred kilometers across and all lying within Nereid’s orbit. Five more small moons, on wide, eccentric orbits, have been discovered by ground-based surveys since 2002, for a total of 13. The planet’s 3 moons larger than 300 km in diameter are listed in Table 13.1. Proteus and Nereid, Neptune’s only midsize moons (by our previous definition) are shown in Figure 13.11. ∞ (Sec. 12.5)



In its moons, we find Neptune’s contribution to our list of solar system peculiarities. Unlike the other jovian worlds, Neptune has no regular moon system—that is, no moons on roughly circular, equatorial, prograde orbits. The largest moon, Triton, is 2700 km in diameter and occupies a circular retrograde orbit 355,000 km (14.3 planetary radii) from the planet, inclined at about 20° to Neptune’s equatorial plane. Triton is the only large moon in our solar system to have a retrograde orbit. The other moon visible from Earth, Nereid, is only 340 km across. This moon orbits Neptune in the prograde sense, but on an elongated trajectory that brings it as close as 1.4 million km to the planet and as far away as 9.7 million km. Nereid is probably similar in both size and composition to Neptune’s small inner moons.

*Voyager 2* approached to within 24,000 km of Triton’s surface, providing us with virtually all that we now know

about that distant, icy world. Astronomers redetermined the moon’s radius (correcting it downward by about 20 percent) and measured its mass for the first time. Along with Saturn’s Titan and the four Galilean moons of Jupiter, Triton is one of the six large moons in the outer solar system. Triton is the smallest of them, with about half the mass of the next smallest, Jupiter’s Europa.

Lying 4.5 billion km from the Sun, and with a fairly reflective surface, Triton has a surface temperature of just 37 K. It has a tenuous nitrogen atmosphere, perhaps a hundred thousand times thinner than Earth’s, and a surface that most likely consists primarily of water ice. A *Voyager 2* mosaic of Triton’s south polar region is shown in Figure 13.13. The moon’s low temperatures produce a layer of nitrogen frost that forms and evaporates over the polar caps, a little like the carbon dioxide frost responsible for the seasonal caps on Mars. The frost is visible as the orange region at the right of the figure.

Overall, Triton exhibits a marked lack of cratering, presumably indicating that surface activity has obliterated the evidence of most impacts. There are many other signs of an active past. For example, Triton’s face is scarred by large fissures similar to those seen on Ganymede, and Triton’s odd cantaloupe-like terrain may indicate repeated faulting and deformation over the moon’s lifetime. In addition, Triton has numerous frozen “lakes” of water ice (Figure 13.14), which may be volcanic in origin. The basic process may be similar to the water volcanism observed on Saturn’s moon Enceladus. ∞ (Sec. 12.5)



▲ **FIGURE 13.13 Triton** The south polar region of Triton, showing a variety of terrains ranging from deep ridges and gashes to what appear to be lakes of frozen water, all indicative of past surface activity. The orange region at lower right is nitrogen frost, forming the moon's polar cap. The long black streaks at bottom left were probably formed by geysers of liquid nitrogen on the surface. (NASA)



Triton's surface activity is not just a thing of the past. As *Voyager 2* passed the moon, its cameras detected two great jets of nitrogen gas erupting from below the surface and rising several kilometers above it. It is thought that these “geysers” form when liquid nitrogen below Triton's surface is heated and vaporized by some internal energy source or perhaps even by the Sun's feeble light. Vaporization produces high pressure, which forces the gas through fissures in the crust, creating the displays *Voyager 2* saw. Scientists conjecture that nitrogen geysers may be common on Triton and are perhaps responsible for much of the moon's thin atmosphere. The long black streaks at the bottom left of Figure 13.13 may have formed when geysers carried dark carbon-rich material from the moon's interior to the surface. Winds in Triton's thin atmosphere may also play a role in spreading the material over the surface.

The event or events that placed Triton on a retrograde orbit and Nereid on such an eccentric path are unknown, but they are the subject of considerable speculation. Triton's peculiar orbit and surface features suggest to some astronomers that the moon did not form as part of the Neptunian system, but instead was captured, perhaps not too long ago, astronomically speaking—maybe even as little as a few hundred million years. Other astronomers, basing their views on Triton's chemical composition, maintain that the moon formed “normally,” but was later kicked into

its abnormal orbit by some catastrophic event, such as an interaction with another, similar-sized body.

The surface deformations on Triton certainly suggest fairly violent and relatively recent events in the moon's past. However, they were most likely caused by the tidal stresses produced in Triton as Neptune's gravity made the moon's orbit more circular and synchronized its spin, and they give little indication of the processes responsible for the orbit.

Whatever its past, Triton's future seems clear. Because of its retrograde orbit, the tidal bulge Triton raises on Neptune tends to make the moon spiral *toward* the planet rather than away from it (as our Moon moves away from Earth). ∞ (Sec. 7.6) Thus, Triton is doomed to be torn apart by Neptune's tidal gravitational field, probably in no more than 100 million years or so, the time required for the moon's inward spiral to bring it inside Neptune's Roche limit. ∞ (Sec. 12.4) The shredded moon will form a new ring



▲ **FIGURE 13.14 Water Ice on Triton** This roughly circular lakelike feature on Triton may have been caused by the eruption of an ice volcano. The water “lava” has since solidified, leaving a smooth surface. The absence of craters implies that this eruption was a relatively recent event. The inset at bottom is a computer-generated view along Triton's surface, illustrating the topography of the area. (NASA)

around the planet (see Figure 12.12). By that time, it is conceivable that large portions of Saturn's ring system may have disappeared, so Neptune will then be the only planet in the solar system with spectacular rings!  $\infty$  (Sec. 12.4)

### CONCEPT Check


✓ Why is Triton much less heavily cratered than the other moons of Uranus and Neptune?

## 13.6 The Rings of the Outermost Jovian Planets

All the jovian planets have rings. However, just as the ring system of Saturn differs greatly from that of Jupiter, the ring systems of Uranus and Neptune also differ both from one another and from those of the two larger jovian worlds.

### The Rings of Uranus


The ring system surrounding Uranus was discovered in 1977, when astronomers observed a **stellar occultation**: The rings passed in front of a bright star, momentarily dimming the star's light (Figure 13.15). Such an alignment happens a few times per decade and allows astronomers to measure planetary structures that are too small and faint to be detected directly. The 1977 observation was actually aimed at studying the planet's atmosphere by watching how it absorbed starlight. However, 40 minutes before and after Uranus itself occulted (passed in front of) the star, the flickering starlight revealed the presence of a set of rings. The discovery was particularly exciting because, at the time, only Saturn was known to have rings. Jupiter's rings went unseen until *Voyager 1* arrived there in 1979, and those of Neptune were unambiguously detected only in 1989, by *Voyager 2*.



▲ **FIGURE 13.15 Occultation of Starlight** By carefully watching the dimming of distant starlight as a planet crosses the line of sight, astronomers can infer fine details about that planet. The rings of Uranus were discovered with this technique.

The ground-based observations revealed the presence of a total of nine thin rings around Uranus. The main rings, in order of increasing radius, are named Alpha, Beta, Gamma, Delta, and Epsilon, and they range from 44,000 to 51,000 km from the planet's center. All lie within the Roche limit of Uranus, which is about 62,000 km from the planet's center. A fainter ring, known as the Eta ring, lies between the Beta and Gamma rings, and three other faint rings, known as 4, 5, and 6, lie between the Alpha ring and the planet itself. In 1986, *Voyager 2* discovered two more even fainter rings, one between Delta and Epsilon and one between ring 6 and Uranus. The main rings are shown in Figure 13.16. More details on the rings are given in Table 13.2.

The rings of Uranus are quite different from those of Saturn. Whereas Saturn's rings are bright and wide, with relatively narrow gaps between them, the rings of Uranus are dark, narrow, and widely spaced. With the exception of the Epsilon ring and the diffuse innermost ring, the



**▲ FIGURE 13.16 Uranus's Rings** The main rings of Uranus, as imaged by *Voyager 2*. All nine of the rings known before the spacecraft's arrival can be seen in this photo. The two rings discovered by *Voyager 2* are too faint to be seen here. The inset at top shows a close-up of the Epsilon ring, revealing some internal structure. The width of this ring averages 30 km; special image processing has magnified the resolution in the inset to about 100 m—the size of a football field. (NASA)

**TABLE 13.2 The Rings of Uranus**


Ring	Inner Radius (km)	(planetary radii)	Outer Radius* (km)	(planetary radii)	Width (km)
1986U2R	37,000	1.45	39,500	1.55	2500
6	41,800	1.64			2
5	42,200	1.65			2
4	42,600	1.67			3
Alpha	44,700	1.75			4–10
Beta	45,700	1.79			5–11
Eta	47,200	1.83			2
Gamma	47,600	1.86			1–4
Delta	48,300	1.90			3–7
1986U1R	50,000	1.96			2
Epsilon	51,200	2.00			20–100

\* Most of Uranus's rings are so thin that there is little difference between their inner and outer radii.

rings of Uranus are all less than about 10 km wide, and the spacing between them ranges from a few hundred to about a thousand kilometers. However, like Saturn's rings, all Uranus's rings are less than a few tens of meters thick (that is, measured in the direction perpendicular to the ring plane).

The density of particles within Uranus's rings is comparable to that found in Saturn's A and B rings. The particles that make up Saturn's rings range in size from dust grains to large boulders, but in the case of Uranus, the particles show a much smaller spread—few, if any, are smaller than a centimeter or so in diameter. The ring particles are also considerably less reflective than Saturn's ring particles, possibly because they are covered with the same dark material as Uranus's moons. The Epsilon ring (shown in detail in the inset for Figure 13.16) exhibits properties a little like those of Saturn's F ring. It has a slight eccentricity of 0.008 and is of variable width, although no braids were found in it. It also appears to be composed of ringlets.

Like the F ring of Saturn, Uranus's narrow rings require *shepherd satellites* to keep them from diffusing. ∞ (Sec. 12.4) In fact, the theory of shepherd satellites was first worked out to explain the rings of Uranus, which had been detected by stellar occultation even before *Voyager 2*'s encounter with Saturn. Thus, the existence of the F ring did not come as quite such a surprise as it might have otherwise! Presumably, many of the small inner satellites of Uranus play some role in governing the




▲ **FIGURE 13.17 Uranian Shepherd Moons** These two small moons, named Cordelia (U7) and Ophelia (U8), were discovered by *Voyager 2* in 1986. They shepherd Uranus's Epsilon ring, keeping it from diffusing away. (NASA)

appearance of the rings. *Voyager 2* detected Cordelia and Ophelia, the shepherds of the Epsilon ring (see Figure 13.17). Many other, undetected, shepherd satellites must also exist.

## The Rings of Neptune

As shown in Figure 13.18 and presented in more detail in Table 13.3, Neptune is surrounded by five dark rings. Three are quite narrow, like the rings of Uranus; the other two are broad and diffuse, more like Jupiter's ring. The dark coloration probably results from radiation darkening, as discussed earlier in the context of the moons of Uranus. All the rings lie within Neptune's Roche limit. The outermost (Adams) ring is noticeably clumped in places. From Earth, we see not a complete ring, but only partial arcs—the unseen parts of the ring are simply too thin (unclumped) to be detected. The connection between the rings and the planet's small inner satellites has not yet been firmly established, but many astronomers think that the clumping is caused by shepherd satellites.

Although all the jovian worlds have ring systems, the rings themselves differ widely from planet to planet. Is there some “standard” way in which rings form around a planet? Also, is there a standard manner in which ring systems evolve? Or do the processes of ring formation and evolution depend entirely on the particular planet in question? If, as now appears to be the case, ring systems are relatively



▲ **FIGURE 13.18 Neptune's Faint Rings** In this long-exposure image, Neptune (center) is heavily overexposed and has been artificially obscured (by an instrument) to make the rings easier to see. One of the two fainter rings lies between the inner bright ring (Leverrier) and the planet. The others lie between the Leverrier ring and the outer bright (Adams) ring. (NASA)

short-lived, their formation must be a fairly common event. Otherwise, we would not expect to find rings around all four jovian planets at once. There are many indications that the individual planetary environment plays an important role in determining a ring system's appearance and longevity. Although many aspects of ring formation and evolution are now understood, it must be admitted that no comprehensive theory yet exists.

### CONCEPT Check

- ✓ **What does the Epsilon ring of Uranus have in common with the F ring of Saturn?**

**TABLE 13.3** The Rings of Neptune

Ring	Inner Radius (km)	Outer Radius* (km)	Width (km)
	(planetary radii)	(planetary radii)	(planetary radii)
Galle (1989N3R)	40,900	42,900	2000
Leverrier (1989N2R)	53,200	53,200	100
Lassell (1989N4R)**	53,200	57,200	4000
Arago (1989N4R)**	57,200	57,200	100
Adams (1989N1R)	62,900	62,900	50

\* Three of Neptune's rings are so thin that there is little difference between their inner and outer radii.

\*\* Lassell and Arago were originally identified as a single ring.

**The Big Question** All four jovian planets are basically huge gas balls surrounding smaller rocky balls deep down inside. It's cold at the top of the cloud deck and hot near the interior surface—conditions unlikely to support life. But a basic question lingers in the minds of many astronomers: Could buoyant, floating life possibly exist at intermediate altitudes where temperatures are more moderate? If so, those life forms must be small: If they got too big they would rise into the interior and cook, yet if they were too small, they would float to the top of the clouds and freeze.


# Chapter Review

## SUMMARY


1 The outer planets Uranus and Neptune were unknown to ancient astronomers. Uranus was discovered in the 18th century, by chance. Neptune was discovered after mathematical calculations of Uranus's slightly non-Keplerian orbit revealed the presence of an eighth planet. At opposition, Uranus is barely visible to the unaided eye. Through a telescope, the planet appears as a pale green disk. Neptune cannot be seen with the naked eye, but a telescope shows it as a tiny bluish disk. Today, we know the giant planets Uranus and Neptune mainly through data taken by *Voyager 2*.



2 The masses of the outer planets are determined from measurements of their orbiting moons. The radii of Uranus and Neptune were relatively poorly known until the *Voyager 2* flybys in the 1980s. Uranus and Neptune have similar bulk properties; they are smaller, less massive, and denser than Jupiter or Saturn. For unknown reasons, Uranus's spin axis lies nearly in the plane of the ecliptic, leading to extreme seasonal variations in solar heating on the planet as it orbits the Sun. Surface features are barely discernible on Uranus, but computer-enhanced images from *Voyager 2* revealed atmospheric clouds and flow patterns moving beneath the planet's haze. Neptune, although farther away from us, has atmospheric features that are clearer because of warmer temperatures and less haze. The **Great Dark Spot** (p. 325) on Neptune had many similarities to Jupiter's Red Spot, but disappeared in 1994.




3 The relatively high densities of Uranus and Neptune imply large, rocky cores making up greater fractions of the planets' masses than in either Jupiter or Saturn. Unlike the other jovian planets, Uranus has no excess heat emission. The source of Neptune's excess energy, like that




of Jupiter's, is most likely heat left over from the planet's formation. Both Uranus and Neptune have substantial magnetospheres. *Voyager 2* discovered that the magnetic fields of the two planets are tilted at large angles to the planets' rotation axes. The reason for these large tilts is not known.

4 All but two of Uranus's moons revolve in the planet's equatorial plane, almost perpendicular to the ecliptic, in circular synchronous orbits. Like the moons of Saturn, the medium-sized moons of Uranus are made up predominantly of rock and water ice. Many of them are heavily cratered and in some cases must have come close to being destroyed by the meteoritic impacts whose craters we now see. The strange moon Miranda has geological features that suggest repeated violent impacts in the past. Neptune's large moon Triton has a fractured surface of water ice and a thin atmosphere of nitrogen, probably produced by nitrogen "geysers" on its surface. Triton is the only large moon in the solar system to have a retrograde orbit around its parent planet. This orbit is unstable and will eventually cause Triton to be torn apart by Neptune's gravity.



5 Uranus has a series of dark, narrow rings, first detected from Earth by **stellar occultation** (p. 332)—their obscuration of the light received from background stars. Shepherd satellites are responsible for the rings' thinness. Neptune has three narrow rings like Uranus's and one broad ring like Jupiter's. The four were discovered by *Voyager 2*. The dark coloration of both the rings and the moons of the outer giant planets may be due to **radiation darkening** (p. 329), whereby exposure to solar high-energy radiation slowly causes a dark hydrocarbon layer to build up on a body's icy surface.





**MasteringAstronomy**® For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- LO1 POS** Why did astronomers suspect an eighth planet beyond Uranus?
- POS** How did astronomers determine where to look for Neptune?
- How did Uranus come to be spinning “on its side”?
- LO2** How and why do the overall colors and appearance of Uranus and Neptune differ from those of Jupiter and Saturn?
- Why are storms and other atmospheric features more easily seen on Neptune than on Uranus?
- LO3** How are the interiors of Uranus and Neptune thought to differ from those of Jupiter and Saturn?
- How do the magnetic fields of Uranus and Neptune compare with that of Earth?
- Describe a day on Titania.
- LO4** What is unique about Miranda? Give a possible explanation.
- How does Neptune’s moon system differ from those of the other jovian worlds? What do these differences suggest about the origin of Neptune’s moon system?
- What causes Triton’s geysers?
- POS** How were the rings of Uranus and Neptune discovered?
- LO5** The rings of Uranus are dark, narrow, and widely spaced. Which of these properties makes them different from the rings of Saturn?
- How do the rings of Neptune differ from those of Uranus and Saturn?
- POS** Why was the discovery of Uranus in 1781 so surprising? Might there be similar surprises in store for today’s astronomers?

## Conceptual Self-Test: Multiple Choice

- The discovery of new planets mostly requires (a) complex calculations and large supercomputers; (b) the patient use of improving technology; (c) an astronomy degree from a large university; (d) pure luck.
- Uranus was discovered about the same time as (a) Columbus reached North America; (b) the U.S. Declaration of Independence; (c) the American Civil War; (d) the Great Depression in the United States.
- Compared with Uranus, the planet Neptune is (a) much smaller; (b) much larger; (c) roughly the same size; (d) tilted on its side.
- The jovian planets with the largest diameters also tend to (a) have the slowest rotation rates; (b) move most slowly in their orbit around the Sun; (c) have the fewest moons; (d) have magnetic field axes most closely aligned with their axes of rotation.
- The five largest moons of Uranus (a) all orbit in the ecliptic plane; (b) can never come between Uranus and the Sun; (c) all orbit directly above the planet’s equator; (d) all have significantly eccentric orbits.
- Moons that show few craters probably (a) are captured asteroids; (b) have been shielded from impacts by their host planet; (c) have had their smaller craters obliterated by larger impacts; (d) have warm interiors.
- A gas giant planet orbiting a distant star would be expected to have (a) a ring system like that of Saturn; (b) a density less than water; (c) many large moons orbiting in different directions; (d) evidence for hydrogen in its spectrum.
- VIS** Uranus’s rings were discovered by the occultation of starlight, as shown in Figure 13.15 (“Occultation of Starlight”). If Uranus were moving more rapidly relative to Earth, the graph in that figure would appear (a) more compressed horizontally; (b) the same; (c) more stretched out horizontally.
- The discovery of a moon orbiting a planet allows astronomers to measure (a) the planet’s mass; (b) the moon’s mass and density; (c) the planet’s ring structure; (d) the planet’s cratering history.
- The solar system object most similar to Neptune is (a) Earth; (b) Jupiter; (c) Saturn; (d) Uranus.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

- What is the angular diameter of the Sun, as seen from Uranus? Compare your answer with the angular diameter of Titania, as seen from the planet's cloud tops. Would you expect solar eclipses to occur on Uranus?
- What is the gravitational force exerted on Uranus by Neptune, at closest approach? Compare your answer with the Sun's gravitational force on Uranus.
- Estimate the speed of cloud A in Figure 13.7, assuming that it lies near the equator. Is your estimate consistent with the rotation speed of the planet?
- If the core of Uranus has a radius twice that of planet Earth and an average density of  $8000 \text{ kg/m}^3$ , what is the mass of Uranus outside the core? What fraction of the planet's total mass is core?
- Add up the masses of all the moons of Uranus and Neptune. (Neglect the masses of the small moons—they contribute little to the result.) How does this sum compare with the mass of Earth's Moon?
- From Wien's law, at what wavelength does Uranus's thermal emission peak? In what part of the electromagnetic spectrum does this wavelength lie? [∞ \(More Precisely 3-2\)](#)
- On the basis of the earlier discussion of planetary atmospheres, would you expect Triton to have retained a nitrogen atmosphere? [∞ \(More Precisely 8-1\)](#)
- How close is Triton to Neptune's Roche limit?

## Activities

### Collaborative

- The search for Neptune requires a determined effort! A telescope is best for the search, but high-powered binoculars mounted on a steady support will also do. Comparing Uranus and Neptune, which planet appears bluer? Through a telescope, does either planet appear as a disk, or do they look more like points of light? Repeat your observations over a period of days or weeks. Can you detect any movement of either planet relative to the background stars in your field of view?

### Individual

- Consult a sky chart online or in a magazine, and locate Uranus in the night sky. It may be barely visible to the naked eye, but binoculars will make the search much easier. (Hint: Uranus shines more steadily than the background stars.) Can you detect the planet's color with your eyes alone? Through binoculars?



# Solar System Debris

## KEYS TO OUR ORIGIN

According to the current definition, there are only eight planets in the solar system. But hundreds of thousands of other celestial bodies are also known to revolve around the Sun. These minor bodies—the asteroids, comets, Kuiper belt objects, and meteoroids—are small and of negligible mass compared with the planets and their major moons. On the basis of statistics, astronomers estimate that there are more than a billion such objects still to be discovered. Yet each is a separate world, with its own story to tell about the early solar system.

These small bodies may seem to be only rocky and icy “debris,” but more than the planets themselves, they hold a record of the formative stages of our planetary system. Many are nearly pristine, unevolved bodies with much to teach us about our local origins.

**The Big Picture** Only within the past few decades have scientists taken seriously the idea that life on Earth has been disrupted over the course of billions of years by asteroid and comet impacts. It’s very much in our own interest to monitor such stray objects that occasionally glide by Earth—and sometimes hit us! Debris in the solar system might be the key to understanding the birth of our planet and even of life—but they also might well determine our fate.

**LEFT:** Comets are sometimes called “dirty snowballs,” since they are composed of ice and snow often contaminated with dusty debris. This spectacular photograph of Comet McNaught was captured from Mount Paranal in Chile, looking out over the Pacific Ocean in 2007. Its colorful tail stretched nearly a quarter of the way across the sky. Other fragile objects like this one are now heading our way while rounding the Sun, and some of them might also melt, vaporize, and break apart while creating awesome displays in the skies above. (S. Deiries/ESO)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Describe the orbital properties of the major groups of asteroids.
- 2 Summarize the composition and physical properties of a typical asteroid.
- 3 Detail the composition and structure of a typical comet, and explain the formation and appearance of its tail.
- 4 List the key characteristics of cometary orbits, and say what they tell us about the probable origin of comets.
- 5 Describe the solar system beyond Neptune, and explain why astronomers no longer regard Pluto as a planet.
- 6 Distinguish among the terms *meteor*, *meteoroid*, and *meteorite*.
- 7 Summarize the orbital and physical properties of meteoroids, and explain what these properties suggest about the probable origin of meteoroids.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.

## 14.1 Asteroids

**Asteroids** are relatively small, predominantly rocky objects that revolve around the Sun. Their name literally means “star-like bodies,” but asteroids are definitely not stars. They are too small even to be classified as planets. Astronomers often refer to them as “minor planets” or, sometimes, “planetoids.”

Asteroids differ from planets in both their orbits and their size. As illustrated in Figure 14.1, they generally move on somewhat eccentric trajectories between Mars and Jupiter, unlike the almost circular paths of the major planets. Few asteroids are larger than 300 km in diameter, our cutoff for a “midsize” moon, and most are far smaller—as small as a tenth of a kilometer across. The largest known asteroid, Ceres (inset to Figure 14.1), is just 1/10,000 the mass of Earth and measures only 940 km across. Taken together, the known asteroids amount to less than the mass of the Moon, so they do not contribute significantly to the total mass of the solar system.

### Orbital Properties


European astronomers discovered the first asteroids early in the 19th century as they searched the sky for an additional planet orbiting between Mars and Jupiter. Italian astronomer

Giuseppe Piazzi was the first to discover an asteroid. He detected Ceres in 1801 and measured its orbital semimajor axis to be 2.8 AU. Within a few years, three more asteroids—Pallas (2.8 AU), Juno (2.7 AU), and Vesta (3.4 AU)—were discovered.

By the start of the 20th century, astronomers had cataloged several hundred asteroids with well-determined orbits. Now, at the end of the first decade of the 21st century, the list has grown to over 350,000. The total number of known asteroids (including those whose orbits are not yet known with sufficient accuracy to make them “official”) now exceeds 600,000. The vast majority of these bodies are found in a region of the solar system known as the **asteroid belt**, located between 2.1 and 3.3 AU from the Sun—roughly midway between the orbits of Mars (1.5 AU) and Jupiter (5.2 AU). All but a handful (about 30) of the known asteroids revolve about the Sun in prograde orbits, in the same sense as the planets.

The compact concentration of asteroids in a well-defined belt has long suggested to astronomers that they are either the fragments of a planet broken up long ago or primal rocks that never managed to accumulate into a genuine planet. On the basis of the best evidence currently available, researchers strongly favor the latter view. There is far too little mass in the belt to constitute a planet, and the marked chemical differences among individual asteroids indicate that they could

not all have originated in a single body. Instead, as described in Chapter 6, Jupiter’s strong gravitational field continuously disturbs the motions of these primitive chunks, preventing them from aggregating into a larger body. [∞ \(Sec. 6.7\)](#)



**MA** **Narrated Interactive FIGURE 14.1 Inner Solar System** (a) The main asteroid belt, along with the orbits of Earth, Mars, and Jupiter (drawn obliquely, that is neither face-on nor edge-on). Note the Trojan asteroids at two locations in Jupiter’s orbit. Some Apollo (Earth-crossing) and Amor (Mars-crossing) orbits are shown. (b) Little surface detail is evident on the largest asteroid known, the dwarf planet Ceres, although image processing reveals what seems to be a large impact crater some 250 km across near the center of the frame. (NASA/SWRI)

## Physical Properties

With only a few exceptions (such as Ceres, shown in Figure 14.1), asteroids are too small to be resolved by Earth-based telescopes, so astronomers must rely on indirect methods to find their sizes, shapes, and composition. Consequently, only a few of their physical and chemical properties are accurately known. To the extent that astronomers can determine their compositions, asteroids have been found to differ not only from the eight known planets and their many moons, but also among themselves.

Asteroids are classified by their spectroscopic properties. The darkest, or least reflective, asteroids contain a large fraction of carbon in their makeup. These asteroids are known as *C-type* (or *carbonaceous*) asteroids. The more reflective *S-type* asteroids contain silicate, or rocky, material. Generally speaking, S-type asteroids predominate in the inner portions of the asteroid belt, and the fraction of C-type bodies steadily increases as we move outward. Overall, about 15 percent of all asteroids are S-type, 75 percent are C-type, and 10 percent are other types (mainly the *M-type* asteroids, containing large fractions of nickel and iron). Many planetary scientists think that the carbonaceous asteroids consist of very primitive material representative of the earliest stages of the solar system. Carbonaceous asteroids have not been subject to significant heating or undergone chemical evolution since they first formed 4.6 billion years ago.

In most cases, astronomers estimate the sizes of asteroids from the amount of sunlight they reflect and the amount of heat they radiate. These observations are difficult, but size measurements have been obtained in this way for a few thousand asteroids. On rare occasions, astronomers witness an asteroid occulting a star, allowing them to determine the asteroid's size and shape with great


accuracy. The largest asteroids are roughly spherical, but the smaller ones can be highly irregular.

The three largest asteroids—Ceres, Pallas, and Vesta—have diameters of 940 km, 580 km, and 540 km, respectively. Only 15 asteroids are more than 300 km across, and most are much smaller. Almost assuredly, many hundreds of thousands more await discovery. However, observers estimate that they are mostly very small. Probably 99 percent of all asteroids larger than 100 km are known and cataloged, and at least 50 percent of asteroids larger than 10 km are accounted for. Although the vast majority of asteroids are probably less than a few kilometers across, most of the *mass* in the asteroid belt resides in objects greater than a few tens of kilometers in diameter.

Vesta is unique among asteroids in that, despite its small size, it appears to have undergone *volcanism* in its distant past. On the basis of their orbits and their overall spectral similarities to Vesta, numerous meteorites (Section 14.3) found on Earth are thought to have been chipped off that asteroid following collisions with other members of the asteroid belt. Remarkably, these meteorites have compositions similar to that of terrestrial basalt, indicating that they were subject to ancient volcanic activity. As discussed below, NASA's *Dawn* mission has recently provided important insights into Vesta's internal structure and history.

## Asteroid Observations from Space

The first close-up views of asteroids were provided by the Jupiter probe *Galileo*, which, on its roundabout path to the giant planet, passed twice through the asteroid belt, making close encounters with the S-type asteroids Gaspra (in 1991) and Ida (1993). [Discovery 6-2](#) Gaspra and Ida (Figure 14.2a,b) are irregularly shaped bodies with maximum diameters of about 20 km and 60 km, respectively.




**▲ FIGURE 14.2 Asteroids, Close-up** (a) The S-type asteroid Gaspra, as seen from a distance of 1600 km by the space probe *Galileo*. (b) The S-type asteroid Ida, photographed by *Galileo* from a distance of 3400 km. Note Ida's moon, Dactyl, at right. (c) The C-type asteroid Mathilde, imaged by the *NEAR* spacecraft on its way to the near-Earth asteroid Eros. Note the huge craters on this low-density rock. Resolution for all these photos is about 100 m. (NASA; JHU)

They are pitted with craters ranging in size from a few hundred meters to 2 km across and are covered with layers of dust of variable thickness. Based on the extent of cratering on their surfaces, Ida is estimated to be about a billion years old—far older than Gaspra, which is a mere 200 million years. Both are thought to be fragments of larger objects that broke up into smaller pieces following violent collisions long ago.

To the surprise of most mission scientists, close inspection of the Ida image (Figure 14.2b) revealed the presence of a tiny moon, now named Dactyl, just 1.5 km across, orbiting the asteroid at a distance of about 90 km. Given the relative congestion of the asteroid belt, scientists think that collisions between asteroids may be quite common. The more violent impacts are likely sources of interplanetary dust and smaller asteroids; the less violent ones may result in bound systems and moons such as Dactyl. By studying the *Galileo* images, astronomers were able to approximate Dactyl's orbit around Ida and hence (using Newton's law of gravity) estimate Ida's mass at about  $5\text{--}10 \times 10^{16}$  kg. [∞ \(More Precisely 2-2\)](#) This in turn allowed them to measure Ida's density as  $2200\text{--}2900$  kg/m<sup>3</sup>, a range consistent with its rocky, S-type classification.

In 1997, the *Near Earth Asteroid Rendezvous (NEAR)* spacecraft visited the C-type asteroid Mathilde on its way to the mission's main target: the S-type asteroid Eros. Shown in Figure 14.2(c), Mathilde is about 60 km across. By sensing its gravitational pull, *NEAR* measured Mathilde's mass at about  $10^{17}$  kg, implying a density of just  $1400$  kg/m<sup>3</sup>. To account for this low density, scientists think that Mathilde, like many smaller asteroids, is more like a loosely bound “rubble pile” than solid rock. The relatively soft internal consistency may also explain the unexpectedly large size of many of the craters on Mathilde's surface. A solid object would probably have shattered after an impact violent enough to cause such large craters. However, like a crumple zone in a car, Mathilde's porous interior could have absorbed and dissipated the impactor's energy, allowing the asteroid to survive the event.

Upon arrival at Eros on February 14, 2000, *NEAR* (by then renamed *NEAR Shoemaker*) went into orbit around the asteroid. For 1 year, the spacecraft sent back high-resolution images of Eros (Figure 14.3) and made detailed measurements of its size and shape ( $34 \times 11 \times 11$  km), as well as its gravitational and magnetic fields, composition, and structure. The craft's various sensors revealed Eros to be a heavily cratered, solid body of mass  $7 \times 10^{15}$  kg and roughly uniform density around  $2700$  kg/m<sup>3</sup>. The asteroid's interior is extensively fractured due to innumerable impacts in the past. The measurements are consistent with Eros being a primitive, unevolved sample of material from the early solar system.



▲ **FIGURE 14.3 Asteroid Eros** A mosaic of detailed images of the asteroid Eros, as seen by the *NEAR Shoemaker* spacecraft that actually landed on this asteroid, displays craters of all sizes, ranging from 50 m (the resolution of the image) to 5 km. The inset shows a close-up image of a “younger” section of the surface, where loose material from recent impacts has apparently filled in and erased all trace of older craters. (*JHU/NASA*)

In July 2011, NASA's *Dawn* probe entered orbit around Vesta, the third largest and second most massive asteroid in the solar system. Armed with cameras to map the surface and instruments to probe the composition of both the surface and the interior, *Dawn* remained at Vesta for more than a year, moving through a series of orbits that brought it within 200 km of the surface.

Vesta's mass of  $2.6 \times 10^{20}$  kg and mean radius of 540 km imply a density of  $3500$  kg/m<sup>3</sup>, considerably higher than most other asteroids. *Dawn* found that Vesta's interior has a differentiated structure with a crust, mantle, and a 200-km-wide iron core and that the asteroid may have been entirely molten in the distant past. Many scientists think that Vesta is a genuine protoplanet dating back to the formative stages of the solar system—the only one known to have survived to the present day. [∞ \(Sec. 6.6\)](#)

Vesta's most striking surface features are a set of deep troughs girdling the asteroid's equator (Figure 14.4a) and a collection of large impact craters in the southern hemisphere, the largest forming a giant circular basin at

the asteroid's south pole (Figure 14.4b). At the center of the south polar basin lies one of the largest mountains in the solar system, rising some 22 km above its surroundings—three times the height of Mount Everest. Cratering age estimates indicate that the southern hemisphere is 1–2 billion years old, much younger than the north, and the south polar region is youngest of all—perhaps as little as 1 billion years. The polar basin and equatorial troughs probably formed when a collision with another large body devastated the asteroid and fractured its interior. The impact may have ejected many of the meteorites associated with

Vesta that have been found on Earth. The asteroid belt is a violent place!

*Dawn* departed Vesta in September 2012, heading for its next destination, the dwarf planet Ceres, where it should arrive early in 2015.

### CONCEPT Check


- ✓ Describe some basic similarities and differences between asteroids and the inner planets.

## Earth-Crossing Asteroids


The orbits of most asteroids have eccentricities lying in the range from 0.05 to 0.3, ensuring that they always remain between the orbits of Mars and Jupiter. Very few asteroids have eccentricities greater than 0.4. Those that do are of particular interest to us, however, as their paths may intersect Earth's orbit, leading to the possibility of a collision with our planet. These bodies are collectively known as **Earth-crossing asteroids**. Those stray asteroids having highly elliptical orbits or orbits that do not lie in the main asteroid belt have probably been influenced by nearby Mars and, especially, Jupiter. The gravitational fields of those two planets can disturb normal asteroid orbits, deflecting them into the inner solar system. Earth-crossing asteroids are termed *Apollo asteroids* (after the first known Earth-crossing asteroid, Apollo) if their orbital semimajor axes exceed 1 AU and *Aten asteroids* otherwise. Asteroids whose orbits cross only the orbit of Mars are known as *Amor asteroids* (see Figure 14.1).

As of early 2013, some 10,000 Earth-crossing asteroids are known. Most have been discovered since the late 1990s, when systematic searches for such objects began. More than 1400 Earth crossers are officially designated “potentially hazardous,” meaning that they are more than about 150 m in diameter (three times the size of the impactor responsible for the Barringer crater shown in Figure 8.18) and move in orbits that could bring them within 0.05 AU (7.5 million kilometers) of our planet.

From a human perspective, perhaps the most important consequence of the existence of Earth-crossers is the very real possibility of an actual collision with Earth. For example, the 3-km-wide asteroid 4179 Toutatis (Figure 14.5) missed our planet by “only” 1.5 million kilometers in 2004—a close call by cosmic standards. Two years earlier, the “nonhazardous,” but still formidable, asteroid 2002 MN, some 100 m across, came much closer, missing us by a mere 120,000 km (less than one-third of the distance to the Moon). It was detected 3 days after it passed our planet! In a widely publicized recent near-miss, the 50-m wide object 2012 DA14 skirted Earth by a mere 28,000 km in February 2013. Coincidentally, the day before, an unrelated, much smaller—and previously completely unknown—20-m meteoroid entered the atmosphere and exploded over southern Russia, releasing as much energy




(a)



(b)

▲ **FIGURE 14.4 Asteroid Vesta** NASA's *Dawn* spacecraft snapped these shots of Vesta, the solar system's second-largest asteroid in 2011. (a) Note the deep grooves that span much of this rocky body, nearly all the way around its equator. (b) The towering mountain near the bottom of this image is more than twice the height of Mount Everest on Earth. (JPL)





▲ **FIGURE 14.5 Asteroid Toutatis** This is a recently released image of asteroid Toutatis 4179, which is about the size of a small city. It was taken by the *Chang'e 2* spacecraft, which had been orbiting the Moon until 2012 when it was retasked to flyby Toutatis. This photograph represents a remarkable achievement for the new Chinese Space Agency, adding China to the other nations (U.S., Russia, and Europe) capable of exploring deep space. (*Xinhuashidian*)

as half a dozen modern nuclear bombs and causing widespread surface damage.

All told, between 2002 and 2012, more than 250 potentially hazardous asteroids (that we know of) passed within 0.05 AU of our planet; 4179 Toutatis was the largest. A similar number are expected to pass within this distance between 2013 and 2023.

None of the currently known potentially hazardous asteroids are expected to impact Earth during the next century—the closest predicted near miss will occur in April, 2029, when the 350-m asteroid 2004 MN4 (also known as Apophis) will pass 30,000 km above our planet's surface. That said, calculations imply that most Earth-crossing asteroids *will* in fact eventually collide with Earth. On average, during any given million-year period, our planet is struck by about three asteroids. Because Earth is largely covered with water, two of those impacts are likely to occur in the ocean and only one on land. Several dozen large land basins and eroded craters on our planet are suspected to be sites of ancient asteroid collisions (see, for example, Figure 14.27, later in the chapter). The many large impact craters on the Moon, Venus, and Mars are direct evidence of similar events on other worlds (see also *Discovery 11-1*).

Most known Earth-crossing asteroids are relatively small—less than about 1 km in diameter (although, as we have seen, larger ones are known, and one 10 km in diameter has been identified). Even so, a visit of even a kilometer-sized asteroid to Earth could be catastrophic by human standards. Such an object packs enough energy to

devastate an area some 100 km in diameter. The explosive power would be equivalent to about a million 1-megaton nuclear bombs—a hundred times more than all the nuclear weapons currently in existence on Earth. A fatal blast wave (the shock from the explosion, spreading rapidly outward from the site of the impact) and a possible accompanying tsunami (tidal wave) from an ocean impact would doubtless affect a much larger area still. Should an asteroid hit our planet hard enough, it might even cause the extinction of entire species—indeed, many scientists think that the extinction of the dinosaurs was the result of just such an impact (see *Discovery 14-1*).


Some astronomers take the prospect of an asteroid impact sufficiently seriously that they maintain an “asteroid watch”—an effort to catalog and monitor all Earth-crossing asteroids in order to maximize our warning time of any impending collision. Several large, dedicated telescopes now scan the skies for faint objects in our neighborhood. Currently, our options in the event that an impending impactor is seen are very limited—science fiction movies aside, we could neither destroy nor deflect an asteroid just a few days away from our planet. However, scientists are confident that, given enough warning, a small “nudge” from thrusters placed on the impactor's surface years ahead of the collision could shift its orbit by just enough to miss us.

## Orbital Resonances

Although most asteroids orbit in the main belt, between about 2 and 3 AU from the Sun, an additional class of asteroids, called the **Trojan asteroids**,\* orbits at the distance of Jupiter. Several hundred such asteroids are now known. They are locked into a 1:1 orbital resonance with Jupiter by that planet's strong gravity, just as some of the small moons of Saturn share orbits with the medium-sized moons Tethys and Dione, as described in Chapter 12. ∞ (Sec. 12.5)

Calculations first performed by the French mathematician Joseph Louis Lagrange in 1772 show that there are exactly five places in the solar system where a small body can orbit the Sun in synchrony with Jupiter, subject to the combined gravitational influence of both large bodies. (Lagrange in fact demonstrated that five such points exist for any planet.) These places are now known as the *Lagrangian points* of the planet's orbit. As illustrated in Figure 14.6, three of the points (referred to as  $L_1$ ,  $L_2$ , and  $L_3$ ) lie on the line joining Jupiter and the Sun (or its extension in either direction). The other two— $L_4$  and  $L_5$ —are located on Jupiter's orbit, exactly  $60^\circ$  ahead of and  $60^\circ$  behind the planet. All five Lagrangian points revolve around the Sun at the same rate as Jupiter.

\*The first few hundred asteroids discovered were named after characters in Greek mythology. The first asteroid found in this orbit was called Achilles. As more asteroids were found sharing this orbit, they became known as the Trojans.



▲ **FIGURE 14.6 Lagrangian Points** The Lagrangian points of the Jupiter–Sun system, where a third body could orbit in synchrony with Jupiter on a circular trajectory. Only the  $L_4$  and  $L_5$  points are stable. They are the locations of the Trojan asteroids (see Figure 14.1).

In principle, an asteroid at any of the Lagrangian points will circle the Sun in lockstep with Jupiter, always maintaining the same position relative to the planet. However, the three Lagrangian points that are in line with Jupiter and the Sun are known to be *unstable*—a body displaced, however slightly, from any of those points will drift slowly away from it, not back toward it. Since matter in the solar system is constantly subjected to small perturbations—by the planets, the asteroids, even the solar wind—matter does not accumulate in these regions. Thus, no asteroids orbit near the  $L_1$ ,  $L_2$ , or  $L_3$  point of Jupiter’s orbit.

The other two Lagrangian points,  $L_4$  and  $L_5$ , are both *stable*—matter placed near them tends to remain in their vicinity. Consequently, asteroids tend to accumulate near these points (see Figure 14.1). For unknown reasons, Trojan asteroids tend to be found near Jupiter’s leading ( $L_4$ , or

*eastern*, as it lies to the east of Jupiter in the sky) Lagrangian point, rather than the trailing ( $L_5$ ) point. Recently, a few small asteroids have been found similarly trapped in the Lagrangian points of Venus, Earth, and Mars.


The main asteroid belt also has structure due to resonances—not as obvious as the Trojan orbits or the prominent gaps and ringlets in Saturn’s ring system, but observable nonetheless. Careful study reveals that asteroids whose orbital periods are simple fractions ( $\frac{1}{2}$ ,  $\frac{1}{3}$ ,  $\frac{2}{5}$ , etc.) of Jupiter’s period are conspicuously absent from the overall distribution. Just as repeated interactions between moons and ring particles can result in gaps in Saturn’s rings, resonant gravitational tugs from Jupiter leave “holes” not in space, but in the distribution of asteroid periods (or semimajor axes). ∞ (Sec. 12.4) These holes are called the **Kirkwood gaps**, after their discoverer, the 19th-century American astronomer Daniel Kirkwood.

### CONCEPT Check



- ✓ Why are astronomers so interested in Earth-crossing asteroids?

## 14.2 Comets

**Comets** are usually discovered as faint, fuzzy patches of light on the sky while they are still several astronomical units away from the Sun. Traveling in a highly elliptical orbit with the Sun at one focus (Figure 14.7), a comet brightens and develops an extended **tail** as it nears the Sun. (The name “comet” derives from the Greek word *kome*, meaning “hair.”) As the comet departs the Sun’s vicinity, its brightness and tail diminish until it once again becomes a faint point of light receding into the distance. Like the planets, comets emit no visible light of their own—they shine by reflected (or reemitted) sunlight. Each year, a few dozen are detected as they pass through the inner solar system. Many more must pass by unseen.



▲ **FIGURE 14.7 Distant Orbit** Comets move on highly eccentric paths that carry them far beyond the known planets. Their long orbital periods mean that most comets entering the inner solar system have never been seen before in human history—making their appearances impossible to predict and giving little warning of any (rare) close encounters with Earth.



◀ **FIGURE 14.8 Comet Structure** (a) Diagram of a typical comet, showing its nucleus, coma, envelope, and tail. The tail is not a sudden streak in time across the sky, as in the case of meteors or fireworks. Instead, it travels along with the rest of the comet, always pointing away from the Sun. Note how the invisible hydrogen envelope is usually larger than the visible extent of the comet; it is often even much larger than drawn here. (b) Halley's Comet in 1986, about 1 month before it rounded the Sun at perihelion, is shown here approximately to the same scale as in part (a). (NOAO)

## Comet Appearance and Structure


The various parts of a typical comet are shown in Figure 14.8. Even through a large telescope, the **nucleus**, or main solid body, of a comet is no more than a minute point of light. A typical cometary nucleus is extremely small—only a few kilometers in diameter. During most of the comet's orbit, far from the Sun, only this frozen nucleus exists. When a comet comes within a few astronomical units of the Sun, however, its icy surface becomes too warm to remain stable. Part of the comet becomes gaseous and expands into space, forming a diffuse **coma** (“halo”) of dust and evaporated gas around the nucleus. The process by which a solid (in this case ice) changes directly into a gas, without first becoming liquid, is called *sublimation*. Frozen carbon dioxide (dry ice) on Earth provides a familiar example of this process. In space, sublimation is the rule, rather than the exception, for the behavior of ice when it is exposed to heat.

The coma becomes larger and brighter as the comet nears the Sun. At maximum size, the coma can measure 100,000 km in diameter—almost as large as Saturn or Jupiter. Engulfing the coma, an invisible **hydrogen envelope**, usually distorted by the solar wind, stretches across millions of kilometers of space. The comet's tail, which is most pronounced when the comet is closest to the Sun and the rate of sublimation from the nucleus is greatest, is much larger still, sometimes spanning as much as 1 AU. From Earth, only the coma and tail of a comet are visible to the naked eye. Despite the size of the tail, most of the light comes from the coma. However, most of the comet's mass resides in the nucleus.

Two types of comet tails may be distinguished. **Ion tails** are approximately straight and are often made up of glowing, linear streamers like those seen in Figure 14.9(a). Their spectra show emission lines of numerous ionized molecules—molecules that have lost some of their normal complement of electrons—including carbon monoxide, nitrogen, and water, among many others. ∞ (Sec. 4.4) **Dust tails** are usually broad, diffuse, and gently curved (Figure 14.9b). They are rich in microscopic dust particles that reflect sunlight, making the tail visible from afar.

Comets' tails are in all cases directed *away* from the Sun by the solar wind (the invisible stream of matter and radiation escaping the Sun). Consequently, as depicted in Figure 14.10, the tail always lies outside the comet's orbit and actually leads the comet during the portion of the orbit that is outbound from the Sun.

Ion tails and dust tails differ in shape because of the different responses of gas and dust to the forces acting in interplanetary space. Every tiny particle in space in our solar system—including those in comets' tails—follows an orbit determined by gravity and the solar wind. If gravity alone were acting, the particle would follow the same curved path as its parent comet, in accordance with Newton's laws of motion. ∞ (Sec. 2.7) If the solar wind were the only influence, the tail would be swept up by it and would trail radially outward from the Sun. Ion tails are much more strongly influenced by the solar wind than by the Sun's gravity, so those tails always point directly away from the Sun. The heavier dust particles have more of a tendency to follow the comet's orbit, giving rise to the slightly curved dust tails.



(a)



(b)

◀ **FIGURE 14.9 Comet Tails** (a) Comet Giacobini-Zinner, seen here in 1959, showed mainly an ion tail; its coma measured 70,000 km across and its tail was well over 500,000 km long. (b) Comet Hale-Bopp in 1997, displayed both an ion tail (dark blue) and a dust tail (white blue), showing also the gentle curvature and inherent fuzziness of the dust. At the comet's closest approach to the Sun, its tail stretched nearly 40° across the sky. (U.S. Naval Observatory; W. Pacholka)

has been observed at every one of its passages since 240 B.C. A spectacular show, the tail of Halley's comet can reach almost a full astronomical unit in length, stretching many tens of degrees across the sky. Figure 14.11(a) shows Halley's comet as seen from Earth in 1910. Its most recent appearance, in 1986 (Figure 14.11b and also Figure 14.8b), was not ideal for terrestrial viewing, as the perihelion happened to occur on roughly

the opposite side of the Sun from Earth, but the comet was closely scrutinized by spacecraft (see below). The comet's orbit is shown in Figure 14.12. Its next scheduled visit to the inner solar system is in 2061.

## Physical Properties of Comets

The mass of a comet can occasionally be estimated by watching how the comet interacts with other solar system objects or by determining the size of the nucleus and assuming a density characteristic of icy composition. These methods yield typical cometary masses ranging from  $10^{12}$  to  $10^{16}$  kg, comparable to the masses of small asteroids.


A comet's mass decreases with time, because some material is lost each time the comet


rounds the Sun as material evaporates from its surface. For comets that travel within an astronomical unit of the Sun, the evaporation rate can reach as high as  $10^{30}$  molecules per second—about 30 tons of cometary material lost for every second the comet spends near the Sun (within Earth's orbit, say). Astronomers have estimated that this loss of material will destroy even a large comet, such as Halley or Hale-Bopp (Figure 14.9b), in just a few thousand orbits.

In seeking the physical makeup of a cometary body itself, astronomers are guided by the observation that comets have dust that reflects light and also certain gas that emits spectral lines of hydrogen, nitrogen, carbon, and oxygen. Even as the atoms, molecules, and dust particles boil off, creating the coma and tail, the nucleus itself

Arguably the most famous comet of all is Halley's comet. In 1705, the British astronomer Edmund Halley realized that the appearance of this comet in 1682 was not a one-time event. Basing his work on previous sightings of the comet, Halley calculated its path and found that the comet orbited the Sun with a period of 76 years. He predicted its reappearance in 1758. Halley's successful determination of the comet's trajectory and his prediction of its return was an early triumph of Newton's laws of motion and gravity. ∞ (Sec. 2.7) Although Halley did not live to see his calculations proved correct, the comet was named in his honor.

Once astronomers knew the comet's period, they traced the appearances of the comet backward in time. Historical records from many ancient cultures show that Halley's comet





◀ **FIGURE 14.10 Comet Trajectory** As it approaches the Sun, a comet develops an ion tail, which is always directed away from the Sun. Closer in, the dust tail displays marked curvature and tends to lag behind the ion tail. Compare this figure with a photo of a real comet, for example Figure 14.9.

∞ (Secs. 12.5, 13.5) Because of this composition, comets are often described as “dirty snowballs.”


## Space Missions to Comets

To date, three comets have received close-up visits from human spacecraft. When Halley’s comet rounded the Sun in 1986, a small armada of spacecraft launched by the (former) USSR, Japan, and a group of western European countries went to meet it. One of the Soviet craft, *Vega 2*, traveled through the comet’s coma and came within 8000 km of the nucleus. Using positional knowledge of the comet gained from the Soviet craft encounter, the European *Giotto* spacecraft (named after the Italian artist who painted an image of Halley’s comet not long after its appearance in the year 1301) was navigated to within 600 km of the nucleus. Figure 14.13 shows *Giotto*’s view of the comet’s nucleus, along with a sketch of its structure.

Halley’s nucleus is an irregular, potato-shaped object, 15 km long by as much as 10 km wide, and almost jet black—as dark as finely ground charcoal or soot. The solid nucleus was enveloped by a cloud of dust, which scattered light throughout the coma. Partly because of this scattering and partly because of dimming by the dust, none of the visiting spacecraft were able to discern much surface detail on the nucleus.

The spacecraft found direct evidence for several jets of matter streaming from the nucleus. Instead of evaporating uniformly from the whole surface to form the comet’s coma and tail, gas and dust vent from small areas on the sunlit side of Halley’s nucleus. The force of these jets may be largely responsible for the comet’s observed 53-hour rotation


remains a cold mixture of gas and dust, hardly more than a ball of loosely packed ice with a density of about  $100 \text{ kg/m}^3$  and a temperature of only a few tens of kelvins. Experts now consider cometary nuclei to be made up largely of dust particles trapped within a mixture of methane, ammonia, carbon dioxide, and ordinary water ice. (These constituents should be fairly familiar to you as the main components of most of the small moons in the outer solar system.)




(a)




(b)



▲ **FIGURE 14.11 Halley’s Comet** (a) Halley’s comet as it appeared in 1910. Top, on May 10, with a  $30^\circ$  tail; bottom, on May 12, with a  $40^\circ$  tail. (b) Halley on its return visit, here photographed with higher resolution on March 14, 1986. (Caltech; Mt. Stromlo and Siding Springs Observatories)



▲ **FIGURE 14.12 Halley's Orbit** Halley's comet has a smaller orbit and a shorter period than most comets, but its orbital orientation is not typical of a short-period comet. Sometime in the past, this comet must have encountered a jovian planet (probably Jupiter itself), which threw it into a tighter orbit that extends not to the Oort cloud, but merely a little beyond Neptune.



▲ **FIGURE 14.13 Halley, Up Close** (a) The European *Giotto* spacecraft resolved the nucleus of Halley's comet, showing it to be very dark, although heavy dust in the area obscured any surface features. Resolution here is about 50 m—half the length of a football field. At the time this image was made, in March 1986, the comet was within days of perihelion, and the Sun was toward the right. The brightest areas are jets of evaporated gas and dust spewing from the comet's nucleus. (b) A diagram of Halley's nucleus, showing its size, shape, jets, and other physical and chemical properties. (ESA/Max Planck Institut)

## DISCOVERY 14-1

### What Killed the Dinosaurs?

The name *dinosaur* derives from the Greek words *deinos* (“terrible”) and *sauros* (“lizard”). Dinosaurs were no ordinary reptiles: In their prime, roughly 100 million years ago, the dinosaurs were the all-powerful rulers of Earth. Their fossilized remains have been uncovered on all the world’s continents. Despite their dominance, according to the fossil record, these creatures vanished from Earth quite suddenly about 65 million years ago. What happened to them?

Until fairly recently, the prevailing view among paleontologists—scientists who study prehistoric life—was that dinosaurs were rather small-brained, cold-blooded creatures. In chilly climates, or even at night, the metabolisms of these huge reptiles would have become sluggish, making it difficult for them to move around and secure food. The suggestion was that they were poorly equipped to adapt to sudden changes in Earth’s climate, so they eventually died out. However, a competing, and still controversial, view of dinosaurs has emerged: Recent fossil evidence suggests that many of these monsters may in fact have been warm-blooded and relatively fast-moving creatures—not at all the dull-witted, slow-moving giants of earlier conception. In any case, no species able to dominate Earth for more than 100 million years could have been too poorly equipped for survival. For comparison, humans have thus far dominated for a little over 2 million years.

If the dinosaurs didn’t die out simply because of stupidity and inflexibility, then what happened to cause their sudden and complete disappearance? Many explanations have been offered for the extinction of the dinosaurs. Devastating plagues, magnetic field reversals, increased tectonic activity, severe climate changes, and supernova explosions have all been proposed. ∞ (Secs. 7.4, 7.5) In the 1980s, it was suggested that a huge extraterrestrial object collided with Earth 65 million years ago, and this is now (arguably) the leading explanation for the demise of the dinosaurs.

According to this idea, a 10- to 15-km-wide asteroid or comet struck Earth, releasing as much energy as 10 million or more of the largest hydrogen bombs humans have ever constructed and kicking huge quantities of dust (including the

pulverized remnants of the impactor itself) high into the atmosphere. (See the first figure.) The dust may have shrouded our planet for many years, virtually extinguishing the Sun’s rays during that time. On the darkened surface, plants could not survive. The entire food chain was disrupted, and the dinosaurs, at the top of that chain, eventually became extinct.

Although we have no direct astronomical evidence to confirm or refute this idea, we can estimate the chances that a large asteroid or comet will strike Earth today on the basis of observations



(D. Hardy)

period. Like maneuvering rockets on a spacecraft, such jets can cause a comet to change its rotation rate and even to veer away from a perfectly elliptical orbit. Astronomers had hypothesized the existence of these nongravitational forces based on slight deviations from Kepler’s laws observed in some cometary trajectories. The Halley encounter was the first time they actually saw the jets at work.

In 1999, NASA launched the *Stardust* mission, with the objective of collecting the first ever samples of cometary material and returning them to Earth. In 2001, the spacecraft used a gravity assist from Earth to boost it onto a path to intercept comet P/Wild 2 (“Wild” is German, pronounced “Vilt”). The comet was chosen because it is a relative


newcomer to the inner solar system, having been deflected onto its present orbit by an encounter with Jupiter in 1974. It therefore has not been subject to much solar heating or loss of mass by evaporation since it formed long ago.

In 2004, *Stardust* approached within 200 km of the comet’s nucleus (Figure 14.14a), collecting cometary particles in a specially designed foamlike “aerogel” detector (see Figure 14.14b). *Stardust* returned to Earth in 2006, returning the debris to mission scientists, who are now studying the detailed physical, chemical, and even biological properties of a body that most probably has not changed significantly since our solar system formed. As shown in Figure 14.14(c), the aerogel performed flawlessly, providing researchers

of the number of objects that are presently on Earth-crossing orbits. The second figure shows the likelihood of an impact as a function of the size of the colliding body. The horizontal scale indicates the energy released by the collision, measured in *megatons* of TNT. The megaton— $4.2 \times 10^{16}$  joules, the explosive yield of a large nuclear warhead—is the only common terrestrial measure of energy adequate to describe the violence of these occurrences.

We see that 100-million megaton events, like the planetwide catastrophe that supposedly wiped out the dinosaurs, are very rare, occurring only once every 10 million years or so. However, smaller impacts, equivalent to “only” a few tens of kilotons of TNT (roughly equivalent to the bomb that destroyed Hiroshima in 1945), could happen every few years—and we may be long overdue for one. The most recent large impact was the Tunguska explosion in Siberia, in 1908, which packed a roughly 1-megaton punch (see Figure 14.28).

The main geological evidence supporting the theory that the dinosaurs’ extinction was the result of an asteroid impact is a layer of clay enriched with the element iridium. The layer is found in 65-million-year-old rocky sediments all around our



with the first-ever samples of cometary material. Among other findings, detailed chemical analysis of the samples has revealed evidence for nitrogen-rich organic material apparently formed in deep space and the unexpected presence of silicate materials that should only have formed at high temperatures, possibly challenging astronomers’ current models of solar system formation. [\(Sec. 6.7\)](#)

A more recent NASA mission had a much more violent end. On July 4, 2005, a 400-kg projectile from NASA’s *Deep Impact* spacecraft crashed into comet Tempel 1 at more than 10 km/s (23,000 mph), blasting gas and debris from the comet’s surface into interplanetary space, while the spacecraft itself watched from a safe distance of 500 km. Figure 14.15 shows

planet. Iridium on Earth’s surface is rare, because most of it sank into our planet’s interior long ago. The abundance of iridium in this one layer of clay is about 10 times greater than in other terrestrial rocks, but it matches closely the abundance of iridium found in meteorites (and, we assume, in asteroids and comets, too). The site of the catastrophic impact has also been tentatively identified as being near Chicxulub, in the Yucatán Peninsula in Mexico, where evidence of a heavily eroded, but not completely obliterated, crater of just the right size and age has been found.

The theory is not without its detractors, however. Perhaps predictably, the idea of catastrophic change on Earth being precipitated by events in interplanetary space was rapidly accepted by most astronomers, but it remains controversial among some paleontologists and geologists. Opponents argue that the amount of iridium in the clay layer varies greatly from place to place across the globe, and there is no complete explanation of why that should be so. Perhaps, they suggest, the iridium was produced by volcanoes, and has nothing to do with an extraterrestrial impact at all.

Still, in the 30 years since the idea was first suggested, the focus of the debate seems to have shifted. The reality of a major impact 65 million years ago has become widely accepted, and much of the argument now revolves around the question of whether that event actually caused the extinction of the dinosaurs or merely accelerated a process that was already underway. Either way, the realization that such catastrophic events can and do occur marks an important milestone in our understanding of evolution on our planet. This realization was bolstered by the Shoemaker–Levy 9 impact on Jupiter in 1994. [\(Discovery 11-1\)](#) In addition, there is growing evidence for even larger impacts in the more distant past, with yet more sweeping evolutionary consequences. As is often the case in science, the debate has evolved, sometimes erratically, as new data have been obtained, but a real measure of consensus has already been achieved, and important new insights into our planetary environment have been gained. [\(Sec. 1.2\)](#)


As a general rule, we can expect that global catastrophes are bad for the dominant species on a planet. As the dominant species on Earth, we are the ones who now stand to lose the most.

an image of the resulting explosion about 1 minute after impact. Spectroscopic analysis of the ejected gas has provided scientists with their clearest view yet of the internal composition of a comet, and hence of the primordial matter of the early solar system, confirming the presence of water ice and many organic molecules. Observations of the crater suggest a low-density “fluffy” internal composition, consistent with the “snowball” picture of cometary structure presented earlier.

### CONCEPT Check

✓ In terms of composition, how do comets differ from asteroids?





**Interactive FIGURE 14.14 Comet Wild-2** (a) The *Stardust* spacecraft captured this image of comet Wild-2 in 2004, just before the craft passed through the comet's coma. (b) Onboard is a detector made of a foaml like jelly (called aerogel) that is 99.8% air, yet is strong enough to stop and store cometary dust particles as they hit the spacecraft. (c) Upon return of the craft to Earth in 2006, analysis began of the minute tracks in the aerogel, the ends of which contain captured comet dust fragments. (NASA)



**Interactive FIGURE 14.15 Deep Impact** The nucleus of comet Tempel 1 is shown at top before impact in 2005 and at bottom shortly after collision with a small projectile launched by the *Deep Impact* mother ship. (JPL)

## Comet Orbits

Comets that survive their close encounter with the Sun—some break up entirely—continue their outward journey to the edge of the solar system. Their highly elliptical orbits take many comets far beyond Pluto, perhaps even as far as 50,000 AU, where, in accord with Kepler's second law, they move more slowly and so spend most of their time. ∞ (Sec. 2.5)

Most comets take hundreds of thousands, and some even take millions, of years to complete a single orbit around the Sun. These comets are known as *long-period comets*. However, a few *short-period comets*, conventionally defined as those having orbital periods of less than 200 years, return for another encounter within a relatively short time. According to Kepler's third law, short-period comets do not venture far beyond the distance of Neptune at aphelion.

Unlike the orbits of the other solar system objects we have studied so far, the orbits of comets are not necessarily confined to within a few degrees of the ecliptic plane. Short-period comets do tend to have prograde orbits lying close to the ecliptic, but long-period comets exhibit all inclinations and all orientations, both prograde and retrograde, roughly uniformly distributed in all directions from the Sun.

The short-period comets originate beyond the orbit of Neptune, in the **Kuiper belt** (named after Gerard Kuiper, a pioneer in infrared and planetary astronomy). A little like the asteroids in the inner solar system, most Kuiper belt comets move in roughly circular orbits between about 30 and 50 AU from the Sun, never venturing inside the orbits of the jovian planets. Occasionally, however, a close encounter between two comets, or (more likely) the cumulative gravitational influence of one of the outer planets, “kicks” a Kuiper belt comet into an eccentric orbit that brings it into the inner solar system and into our view. The observed orbits of these comets reflect the flattened structure of the Kuiper belt.

What of the long-period comets? How do we account for their apparently random orbital orientations? Only a tiny portion of a typical long-period cometary orbit lies within the inner solar system, so it follows that, for every comet we see, there must be many more similar objects at great distances from the Sun. On these general grounds,


many astronomers reason that there must be a huge “cloud” of comets far beyond the orbit of Pluto, completely surrounding the Sun. This region, which may contain trillions of comets, with a total mass comparable to the mass of the inner planets, is named the **Oort cloud**, after the Dutch astronomer Jan Oort, who first wrote (in the 1950s) of the possibility of such a vast reservoir of inactive, frozen comets orbiting far from the Sun. The Kuiper belt and the orbits of some typical Oort cloud comets are sketched in Figure 14.16.

The observed orbital properties of long-period comets have led researchers to conclude that the Oort cloud may be up to 100,000 AU in diameter. Like those of the Kuiper belt, however, most of the comets of the Oort cloud never come anywhere near the Sun. Indeed, Oort cloud comets rarely approach even the orbit of Pluto, let alone that of Earth. Only when the gravitational field of a passing star happens to deflect a comet into an extremely eccentric orbit that passes through the inner solar system do we actually get to see the comet.

Because the Oort cloud surrounds the Sun in all directions, instead of being confined near the plane of the ecliptic like the Kuiper belt, the long-period comets we see can come from any direction in the sky. Despite their great distances and long orbital periods, Oort cloud comets are still gravitationally bound to the Sun. Their orbits are governed by precisely the same laws of motion that control the planets’ orbits.

### PROCESS OF SCIENCE Check

✓ In what sense are the comets we see *unrepresentative* of comets in general?



(a)

## 14.3 Beyond Neptune


No one has ever observed any comets in the faraway Oort cloud—they are just too small and dim for us to see from Earth. But in the 1990s such faint objects began to be inventoried in the relatively nearby Kuiper belt, just beyond Neptune’s orbit, some 30 to 50 AU from the Sun. They are collectively referred to as **Kuiper belt objects**. And as the search has broadened, other, even more distant bodies have been found. The generic term for any small body orbiting beyond Neptune—including members of the Kuiper belt—is **trans-Neptunian object**.

Ground-based telescopes have led the way in recent years in the painstaking effort to capture the meager amounts of sunlight reflected from these dark inhabitants of the outer solar system. However, one resident of the Kuiper belt has been known for decades—Pluto. Let’s begin our study of this distant region by reviewing what is known about its most prominent member.

### The Serendipitous Discovery of Pluto

Around the end of the 19th century, observations of the orbits of Uranus and Neptune suggested that Neptune’s influence was not sufficient to account for all the irregularities in Uranus’s motion. Furthermore, it seemed that Neptune itself might be affected by some other unknown body—perhaps even another planet. Following their success in the discovery of Neptune, astronomers hoped to pinpoint the location of this new object by using similar techniques. One of the most ardent searchers was Percival Lowell, a capable, persistent observer and one of the best-known astronomers of his day. (Recall that he was also the leading proponent of the theory that the “canals” on Mars were constructed by an intelligent race of Martians—see the Part 2 Opener on p. 130).

Basing his investigation primarily on the motion of Uranus (Neptune’s orbit was still relatively poorly determined at the time), and using techniques similar to those developed earlier in the search for Neptune by Adams and Leverrier,



(b)

◀ **FIGURE 14.16 Comet Reservoirs** (a) Diagram of the Oort cloud, showing a few cometary orbits. The solar system is much smaller than the overlaid box at the center of the figure. (b) The Kuiper belt, the source of short-period comets, whose orbits hug the plane of the ecliptic.

Lowell set about calculating where the supposed new body should be. He sought it, without success, during the decade preceding his death in 1916. Not until 14 years later did American astronomer Clyde Tombaugh, working with improved equipment and better photographic techniques at the Lowell Observatory, succeed in finding the new body, only  $6^\circ$  away from Lowell's predicted position. It was named Pluto, for the Roman god of the dead who presided over eternal darkness (and also because its first two letters and its astronomical symbol, ♇, are Lowell's initials). The discovery of Pluto was announced on March 13, 1930, Percival Lowell's birthday, and also the anniversary of Herschel's discovery of Uranus.


On the face of it, the discovery of Pluto looked like another spectacular success for celestial mechanics. However, it now appears that the supposed irregularities in the motions of Uranus and Neptune did not exist, and that the mass of Pluto, not measured accurately until the 1980s, is far too small to have caused them anyway. The discovery of Pluto owed much more to simple luck than to complex mathematics!

## Pluto's Orbital and Physical Properties

Pluto's orbit is quite elongated, with an eccentricity of 0.25. It is also inclined at  $17.2^\circ$  to the plane of the ecliptic. Because of its substantial orbital eccentricity, Pluto's distance from the Sun varies considerably. At perihelion, it lies 29.7 AU (4.4 billion km) from the Sun, *inside* the orbit of Neptune. At aphelion, the distance is 49.3 AU (7.4 billion km), well outside Neptune's orbit. Pluto last passed perihelion in 1989 and remained inside Neptune's orbit until February 1999. Given Pluto's 248-year orbital period, this will not occur again until the middle of the 23rd century. The orbits of Neptune and Pluto are sketched in Figure 14.17.

Pluto's orbital period is exactly 1.5 times that of Neptune—in other words, the two bodies are locked in a 3:2 resonance (two orbits of Pluto for every three of Neptune) as they orbit the Sun. As a result, even though their orbits appear to cross, Pluto and Neptune are in no danger of colliding with each other. Because of the resonance and Pluto's tilted orbital plane, the distance between the two bodies at closest approach is actually about 17 AU (compare with Pluto's closest approach to Uranus of just 11 AU). As with other solar system resonances, Pluto's 3:2 synchronization with Neptune is not a matter of chance. In fact, it is a direct consequence of the evolution of the outer solar system billions of years ago.

Recall from Chapter 6 that Neptune is thought to have migrated slowly outward as the planet interacted with planetesimals and helped form the Kuiper belt. ∞ (Sec. 6.7) As it did so, the radius corresponding to the 3:2 resonance also swept outward through the surviving planetesimals. Many planetesimals—Pluto included—on near-resonant orbits were captured and carried along with Neptune, locked in synchrony to it forever, as it drifted toward its present location. About 15 percent of all Kuiper belt objects—dubbed *plutinos*—share Pluto's orbital resonance.




**Interactive FIGURE 14.17 Neptune and Pluto** The orbits of Neptune and Pluto cross, although Pluto's orbital inclination and a 3:2 resonance prevent the planets from coming close to each other. Between 1979 and 1999, Pluto was actually inside Neptune's orbit.

Pluto is so far away that little is known of its physical nature. Until the late 1970s, studies of sunlight reflected from its surface suggested a rotation period of just under a week, but measurements of its mass and radius were uncertain. All this changed in 1978, when astronomers at the U.S. Naval Observatory discovered that Pluto has a companion. It was named Charon, after the mythical boatman who ferried the dead across the river Styx into Hades, Pluto's domain. The discovery photograph of Charon is shown in Figure 14.18(a). Charon is the small bump near the top of the image. Figure 14.18(b) shows a 2005 *Hubble Space Telescope* view that clearly resolves the two bodies and in addition shows two more small satellites orbiting the Pluto–Charon system. The new moons are perhaps 100 km in diameter and orbit Pluto at roughly twice the distance of Charon. They have since been named Nix (for the goddess of darkness and mother of Charon) and Hydra (a mythical nine-headed monster). Two more, even smaller (10 km), moons, known simply as S/2011 P1 and S/2012 P1, were spotted in 2011 and 2012, respectively.

The discovery of Charon permitted astronomers to measure the masses and radii of both bodies with great accuracy. Charon's orbit is inclined at an angle of  $118^\circ$  to the plane of Pluto's path around the Sun. By pure luck, over the 6-year period from 1985 to 1991 (less than 10 years after Charon's discovery), the two bodies happened to be oriented in such a way that viewers on Earth saw a series of *eclipses*, in which Pluto and Charon repeatedly passed in front of each other, as seen from our vantage point. Figure 14.19 sketches this orbital configuration. With more good fortune, these eclipses took place while Pluto was closest to the Sun, making for the best possible Earth-based observations.

Basing their calculations principally on the variations in reflected light as Pluto and Charon periodically hid each

▼ **FIGURE 14.18 Pluto and Charon** (a) The discovery photograph of Pluto's moon, Charon, shows the moon as a small blotch of light at the top right part of the image. (b) The Pluto–Charon system, shown to the same scale and better resolved than in part (a), as seen by the *Hubble Space Telescope*. (U.S. Naval Observatory; NASA)



(a)




(b)

other, astronomers calculated that Pluto and Charon move in a circular, tidally locked orbit with a period of 6.4 days and a separation of 19,700 km, implying a mass for Pluto of 0.0021 Earth mass ( $1.3 \times 10^{22}$  kg), far smaller than any previous estimate. ∞ (More Precisely 2-2) Pluto's diameter is 2270 km, about one-fifth the size of Earth. Charon is about 1300 km across. If both bodies have the same composition (probably a reasonable assumption), Charon's mass must be about one-sixth that of Pluto.



The masses and radii of Pluto and Charon imply average densities of  $2100 \text{ kg/m}^3$ —just what we would expect for bodies of that size made up mostly of water ice, like the large moons of the outer planets. ∞ (Sec. 6.2) In fact, Pluto is very

similar in both mass and radius to Neptune's large moon, Triton—which, as we have seen, is thought to be a captured Kuiper belt object. ∞ (Sec. 13.5) Spectroscopy reveals the presence of *frozen* methane as a major surface constituent of Pluto, implying a temperature of no more than 50 K. Pluto may also have a thin methane atmosphere, associated with the methane ice on its surface.


Recent computer-generated maps (Figure 14.20) have begun to hint at surface features on Pluto and suggest that




▲ **FIGURE 14.19 Pluto–Charon Eclipses** The orbital orientation of Charon produced a series of eclipses between 1985 and 1991. Observations of eclipses of Charon by Pluto and of Pluto by Charon have provided detailed information about the sizes and orbits of both bodies.



▲ **FIGURE 14.20 Pluto** A three-dimensional surface map of Pluto—not a photograph, but rather a “modeled” view created by carefully combining 24 *Hubble Space Telescope* images with a mathematical description of the surface. (NASA)



◀ **FIGURE 14.21** **Kuiper Belt Object** (a) These frames show some of the best available images of the Kuiper belt object, Pholus—the fuzzy blob (marked with an arrow) that changes position over the course of a few days. It’s almost 1000 km across and lies more than 40 AU from Earth. (b) The trans-Neptunian object Eris and its small moon Dysnomia (named after the Greek goddess of discord and her daughter, goddess of chaos and lawlessness) were imaged in the infrared at the Keck Observatory in Hawaii. (LPL/Keck)



Charon may have bright polar caps, although their composition and nature are unknown. Some of the dark regions may be craters or impact basins, as on Earth’s Moon. ∞ (Sec. 8.3) However, astronomers eager for a closer look will have to wait until 2015, when NASA’s *New Horizons* mission to Pluto and the Kuiper belt, launched in 2006, is scheduled to reach its destination.

## Properties of Trans-Neptunian Objects


Most Kuiper belt objects are not nearly as well observed as Pluto, which happens to be the largest known member of the class and orbits near the inner edge of the belt, making it appear relatively bright as seen from Earth. Most trans-Neptunian objects—even the large ones—are very faint: Figure 14.21 shows some of the best available images (apart from those of Pluto and Charon) of some of these distant bodies. Astronomers generally have only limited information on them—a *segment of an orbit*, from which

distance, semimajor axis, period, and eccentricity can be inferred; its *brightness*, which translates into an estimated diameter; and sometimes *brightness variations*, which may imply rotation or the presence of a companion.

Nevertheless, as astronomers have refined their observational techniques, the number of known objects beyond Neptune has risen rapidly. As of early 2013, the count stands at just over 1600; most are in the Kuiper belt. Because they are so small and distant, researchers reason that only a tiny fraction of the total have so far been observed, and the total number of Kuiper belt objects larger than 100 km is estimated to be more than 100,000. If that is so, then the combined mass of all the debris in the Kuiper belt could well be hundreds of times larger than the mass of the inner asteroid belt (although still less than the mass of Earth).

Unfortunately for Pluto’s planetary status, as the details were filled in and the numbers of known trans-Neptunian objects increased, it became more and more clear to astronomers that Pluto is not distinctly different from the other small bodies in the outer solar system, as had once been supposed. The Kuiper belt object Quaoar (pronounced “KWAH-o-wahr,” and named for a Native American creation god), discovered in 2002, is roughly 1200 km across—larger than the largest asteroid, Ceres, and more than half the size of Pluto. The Kuiper belt objects Haumea, discovered in 2003, and Makemake, discovered in 2005, are larger still—both some 1500–2000 km across (their names are drawn from Hawaiian mythology). But the final blow came with the confirmation in 2005 of the (non-Kuiper belt) object Eris (appropriately named for the Greek goddess of discord, and shown in Figure 14.21b), whose diameter, measured in 2006 by *HST*, is 2400 km—*larger* than Pluto.

The sizes of some of the biggest known trans-Neptunian objects are compared in Figure 14.22, and their orbits are sketched in Figure 14.23. These figures include yet another intriguing object called Sedna, discovered in 2003 and thought to have a diameter of about 1500 km. It is the




▲ **FIGURE 14.22 Trans-Neptunian Objects** Some large trans-Neptunian objects, including Pluto and the largest known, called Eris, with part of Earth and the Moon added for scale. Most diameters are approximate, as they are estimated from the object's observed brightness. (NASA; Caltech)

farthest known object in the solar system. Its highly elliptical orbit takes it out to almost 1000 AU from the Sun—almost to the (theoretical) inner edge of the Oort cloud. There may well be more Pluto-sized (or larger) objects still out there, waiting to be discovered—the possibility has not been conclusively ruled out. Systematic faint surveys of a broad swath of the sky that includes the entire plane of the ecliptic (as are planned within the next decade) will be needed before any definitive statement can be made.

## The King of the Kuiper Belt

Even before the discovery of Eris, many astronomers had already concluded that Pluto was not a different type of



▲ **FIGURE 14.23 Orbits in the Outer Solar System** Orbits of some prominent residents of the outer solar system: Jupiter, Neptune, the Kuiper belt, Pluto, and several more trans-Neptunian objects. Sedna's orbit extends for 10 times the outermost distance to Eris.

object at all, but simply the largest known member of the Kuiper belt, playing much the same role in the Kuiper belt as Ceres does among the asteroids. Once it became clear that Eris was most likely larger than Pluto, pressure mounted to find a classification that reflected astronomers' new understanding of the outer solar system. In 2006 the International Astronomical Union (IAU) adopted the first ever *definition* of a planet: A body is considered a planet if

1. it orbits the Sun,
2. it is massive enough that its own gravity has caused its shape to be approximately spherical, and
3. it has "cleared the neighborhood" around its orbit of other bodies.

"Clearing the neighborhood" means that the body has swept up (collided with) any debris whose orbit happens to intersect its path, or that its gravity has kicked most such debris to other parts of the solar system. ∞ (Discovery 6-1)

Pluto certainly satisfies the first two criteria. However, part 3 of this definition *excludes* Pluto from planetary status. The wording is chosen specifically to ensure that a planet is massive enough to dominate its immediate neighborhood, and Pluto, which orbits within the congested Kuiper belt and is resonantly tied to Neptune's orbital motion, clearly does not meet that condition. Indeed, as we saw in the chapter on the solar system (Chapter 6), the Kuiper belt was formed as Neptune and Uranus established themselves as planets by clearing their own orbital neighborhoods of interplanetary debris. ∞ (Sec. 6.7) As if for consolation, so as not to completely strip Pluto of its title, the International Astronomical Union invented a new term for bodies that satisfy criteria 1 and 2 but not 3—**dwarf planet**. Eris, Pluto, Makemake, Haumea, and Ceres all fall into this category. Some of the other bodies in Figure 14.22 may also be classified as dwarf planets once their properties are better determined. In 2008 the IAU decided that the icy dwarf planets beyond Neptune would henceforth be known collectively as **plutoids**.

The IAU decision has sparked controversy among astronomers. Some are unhappy at Pluto's demotion from the solar system "A list." They argue (probably correctly) that the new definition was concocted largely to exclude Pluto and Eris from planetary status and that criterion 3 in its current form is really too vague to be of much scientific value. Others applaud the redefinition as long-overdue recognition of Pluto's true identity as a large Kuiper belt object, but object to the new term "dwarf planet" as redundant and unnecessarily confusing.

The arguments over terminology are probably not over, but it seems unlikely that future changes will restore Pluto to its former status. Simply put, it is not sufficiently different from the other known Kuiper belt objects to warrant inclusion in a different category. Few astronomers doubt that if Pluto were discovered today, its classification as a member (the largest yet, the headlines would say!) of the Kuiper belt would be assured.

Pluto's reclassification illustrates the way in which science evolves. Our conception of the cosmos has undergone many changes—some radical, others more gradual—since the time of Copernicus. ∞ (Sec. 2.3) We have seen several examples already, and we will see many more later in this book. As our understanding grows, our terminology and classifications change. The situation with Pluto has a close parallel in the discovery of the first asteroids in the early 19th century. They, too, were initially classified as planets—indeed, in the 1840s, leading astronomy texts listed no fewer than 11 planets in our solar system, including as numbers 5 through 8 the asteroids Vesta, Juno, Ceres, and Pallas. Within a couple of decades, however, the discovery

of several dozen more asteroids had made it clear that these small bodies represented a whole new class of solar system objects, separate from the major planets, and the number of planets fell to 8 (including the then newly discovered Neptune).

Much of the observational work on the Kuiper belt began as a search for a 10th planet. It is ironic that the end result of these efforts has been a reduction in the number of "true" solar system planets back to eight!

### CONCEPT Check

✓ Why do astronomers no longer regard Pluto as a planet?

## 14.4 Meteoroids

On a clear night, it is possible to see a few *meteors*—"shooting stars"—every hour. A **meteor** is a sudden streak of light in the night sky caused by friction between air molecules in Earth's atmosphere and an incoming piece of interplanetary matter—an asteroid, a comet, or a **meteoroid**. The friction heats and excites the air molecules, which then emit light as they return to their ground states, producing the characteristic bright streak shown in Figure 14.24. Recall from Section 6.5 that the distinction between an asteroid and a meteoroid is simply a matter of size. Both are chunks of rocky interplanetary debris; meteoroids are conventionally taken to be less than 100 m in diameter.

Note that the brief flash that is a meteor is in no way similar to the broad, steady swath of light associated with a




(a)



(b)



▲ **FIGURE 14.24 Meteor Trails** A bright streak of light called a meteor is produced when a fragment of interplanetary debris plunges into the atmosphere, heating the air to incandescence. (a) A small meteor photographed against a backdrop of stars and the Northern Lights. (b) These meteors (one with a red smoke trail) streaked across the sky during the height of the Leonid meteor storm of November 2001. (*P. Parviainen; J. Lodriguss*)



**▲ FIGURE 14.25 Meteor Showers** A meteoroid swarm associated with a given comet intersects Earth's orbit at specific locations, giving rise to meteor showers at specific times of the year. If the comet's path happens to intersect Earth's, the result is a meteor shower each time Earth passes through the intersection (point 4).

comet's tail. A meteor is a fleeting event in Earth's atmosphere, whereas a comet tail exists in deep space and can be visible in the sky for weeks or even months. Before encountering the atmosphere, the piece of debris causing a meteor was almost certainly a meteoroid, because these small interplanetary fragments are far more common than either asteroids or comets. Any piece of interplanetary debris that survives its fiery passage through our atmosphere and finds its way to the ground is called a **meteorite**.

## Cometary Fragments

Smaller meteoroids are mainly the rocky remains of broken-up comets. Each time a comet passes near the Sun, some cometary fragments are dislodged from the main body. The fragments initially travel in a tightly knit group of dust or pebble-sized objects, called a **meteoroid swarm**, moving in nearly the same orbit as the parent comet. Over the course of time, the swarm gradually disperses along the orbit, and

eventually the **micrometeoroids**, as these small meteoroids are known, become more or less smoothly spread all the way around the parent comet's orbit.

If Earth's orbit happens to intersect the orbit of such a young cluster of meteoroids, a spectacular **meteor shower** can result. Earth's motion takes our planet across a given comet's orbit at most twice a year (depending on the precise orbit of each body). Intersection occurs at the same time each year (see Figure 14.25), so the appearance of certain meteor showers is a regular and (fairly) predictable event.

Table 14.1 lists some prominent meteor showers, the dates they are visible from Earth, and the comets from which they are thought to originate. Meteor showers are usually named for their *radiant*, the constellation from whose direction they appear to come (Figure 14.26). For example, the Perseid shower is seen to emanate from the constellation Perseus. It can last for several days, but reaches maximum every year on the morning of August 12, when upward of 50 meteors per hour can be observed. Astronomers can use the speed and direction of a meteor's flight to compute the meteor's interplanetary trajectory. This is how certain meteoroid swarms have come to be identified with well-known comet orbits. For example, the Perseid shower shares the same orbit as comet 1862III (also known as comet Swift-Tuttle), the third comet discovered in the year 1862.

**TABLE 14.1** Some Prominent Meteor Showers


Morning of Maximum Activity	Name of Shower	Rough Hourly Count	Parent Comet
Jan. 3	Quadrantid	40	—
Apr. 21	Lyrid	10	1861I (Thatcher)
May 4	Eta Aquarid	20	Halley
June 30	Beta Taurid	25 <sup>‡</sup>	Encke
July 30	Delta Aquarid	20	—
Aug. 11	Perseid	50	1862III (Swift-Tuttle)
Oct. 9	Draconid	up to 500	Giacobini-Zinner
Oct. 20	Orionid	30	Halley
Nov. 7	Taurid	10	Encke
Nov. 16	Leonid	12*	1866I (Tuttle)
Dec. 13	Geminid	50	3200 (Phaeton) <sup>†</sup>

\* Every 33 years, as Earth passes through the densest region of this meteoroid swarm, we see intense showers that can exceed 1000 meteors per minute for brief periods. This intense activity is next expected to occur in 2032.


<sup>†</sup> Phaeton is actually an asteroid and shows no signs of cometary activity, but its orbit matches the meteoroid paths very well.

<sup>‡</sup> Meteor count peaks after sunrise.





(a)



(b)



(c)

▲ **FIGURE 14.26 Radiant** (a) A group of meteoroids approaches an observer, all of them moving in the same direction at the same speed. (b) From the observer's viewpoint, the trajectories of the meteoroids (and the meteor shower they produce) appear to spread out from a central point, called the radiant, in much the same way as parallel railroad tracks seem to converge in the distance (c).

## Stray Asteroids


Larger meteoroids—more than a few centimeters in diameter—are usually *not* associated with swarms of cometary debris. Generally regarded as small bodies that have strayed from the asteroid belt, possibly as the result of collisions with or between asteroids, these objects have produced most of the cratering on the surfaces of the Moon, Mercury, Venus, Mars, and some of the moons of the jovian planets. When these large meteoroids enter Earth's atmosphere with a typical velocity of nearly 20 km/s, they produce energetic shock waves, or “sonic booms,” as well as bright streaks in the sky and dusty trails of discarded debris. Such large meteors are

sometimes known as *fireballs*. The greater the speed of the incoming object, the hotter its surface becomes and the faster it burns up. A few large meteoroids enter the atmosphere at such high speed (about 75 km/s) that they either fragment or disperse entirely at high altitudes.

The more massive meteoroids (at least a ton in mass and a meter across) do make it to Earth's surface, producing a crater such as the kilometer-wide Barringer Crater shown in Figure 8.18. From the size of this crater, we can estimate that the meteoroid responsible for its formation must have had a mass of about 200,000 tons. Since only 25 tons of iron meteorite fragments have been found at the crash site, the remaining mass must have been scattered by the explosion at impact, broken down by subsequent erosion, or buried in the ground.

Currently, Earth is scarred with nearly 100 craters larger than 0.1 km in diameter. Most of these craters are so heavily eroded by weather and distorted by crustal activity that they can be identified only in satellite photography, as shown in Figure 14.27. Fortunately, such major collisions between Earth and large meteoroids are thought to be rare events now. Researchers estimate that, on average, they occur only once every few hundred thousand years (see *Discovery 14-1*).

The orbits of large meteorites that survive their plunge through Earth's atmosphere can be reconstructed in a manner similar to that used to determine the orbits of meteor showers. In most cases, their computed orbits do indeed intersect the asteroid belt, providing the strongest evidence we have that they were once part of the belt before being



▲ **FIGURE 14.27 Manicouagan Reservoir** This infrared photograph, taken from orbit by the U.S. *Skylab* space station, shows the ancient impact basin that forms Quebec's Manicouagan Reservoir. A large meteorite landed there about 200 million years ago. (NASA)



**▲ FIGURE 14.28 Tunguska Debris** The Tunguska event of 1908 leveled trees over a vast area. Although the impact of the blast was tremendous and its sound audible for hundreds of kilometers, this Siberian site is so remote that little was known about the event until scientific expeditions arrived to study it many years later. (*Sovfoto/ Eastfoto*)

redirected, probably by a collision with another asteroid, into the Earth-crossing orbit that led to the impact with our planet. Not all meteoroids come from the asteroid belt, however: As we have already seen, some are known to have originated on the surface of Mars. [∞ \(Discovery 10-1\)](#) In addition, detailed composition studies reveal that others most

likely came from the Moon, blasted off the lunar surface by violent impacts long ago.

Not all encounters of meteoroids with Earth result in an impact. One of the most recent meteoritic events occurred in central Siberia on June 30, 1908 (Figure 14.28). The presence of only a shallow depression, as well as a complete absence of fragments, implies that this Siberian intruder exploded several kilometers above the ground, leaving a blasted depression at ground level, but no well-formed crater. Recent calculations suggest that the object in question was a rocky meteoroid about 30 m across. The explosion, estimated to have been equal in energy to a 10-megaton nuclear detonation, was heard hundreds of kilometers away and produced measurable increases in atmospheric dust levels all across the Northern Hemisphere.

## Meteorite Properties

One feature that distinguishes small micrometeoroids, which burn up in Earth's atmosphere, from larger meteoroids, which reach the ground, is their composition—as evidenced by their strikingly different densities. The average density of meteoritic fireballs that are too small to reach the ground (but that can be captured by high-flying aircraft) is about 500–1000 kg/m<sup>3</sup>. Such a low density is typical of comets, which are made of loosely packed ice and dust. In contrast, the meteorites that reach Earth's surface are often much denser—up to 5000 kg/m<sup>3</sup>—suggesting a composition more like that of the asteroids. Meteorites like those shown in Figure 14.29 have received close scrutiny from




(a)

**▲ FIGURE 14.29 Large Meteorites** (a) The world's second largest meteorite, the Ahnighito, on display at the American Museum of Natural History in New York, serves as a jungle gym for curious children. This 34-ton rock is so heavy that the Museum floor had to be specially reinforced to support its weight. (b) The Wabar meteorite, discovered in the Arabian desert. Although small fragments of the original meteorite had been collected more than a century before, the 2000-kg main body was not found until 1965. (*Corbis-Blair; Jon Mandaville/Aramco World*)



(b)



(a)



(b)

▲ **FIGURE 14.30 Meteorite Samples** (a) A stony (silicate) meteorite often has a dark fusion crust, created when its surface is melted during its passage through the atmosphere. The coin at the bottom is for scale. (b) Iron meteorites are much rarer than the stony variety and often contain some nickel as well. Most show characteristic crystalline patterns when their surfaces are cut, polished, and etched with acid. (*Science Graphics*)

planetary scientists—prior to the Space Age, they were the only type of extraterrestrial matter we could touch and examine in terrestrial laboratories.

Most meteorites are rocky in composition (Figure 14.30a), although a few percent are composed mainly of iron and nickel (Figure 14.30b). The basic composition of the rocky meteorites is much like that of the inner planets and the Moon, except that some of their lighter elements—such as hydrogen and oxygen—appear to have boiled away long ago when the bodies from which the meteorites originated were molten. Some meteorites show clear evidence of strong heating at some time in their past, most likely indicating that they originated on a larger body that either underwent some geological activity or was partially melted during the collision that liberated the fragments that eventually became the meteorites. Others show no such evidence and probably date from the formation of the solar system.

Most primitive of all are the *carbonaceous* meteorites, so called because of their relatively high carbon content.

These meteorites are black or dark gray and may well be related to the carbon-rich C-type asteroids that populate the outer asteroid belt. (Similarly, the silicate-rich stony meteorites are probably associated with the inner S-type asteroids.) Many carbonaceous meteorites contain significant amounts of ice and other volatile substances, and they are usually rich in organic molecules.

Finally, almost all meteorites are *old*. Direct radioactive dating shows most of them to be between 4.4 and 4.6 billion years old—roughly the age of the oldest lunar rocks. Meteorites, along with lunar rocks, comets, and Kuiper belt objects like Pluto, provide essential clues to the original state of matter in the solar neighborhood. ∞ (Sec. 6.7)

### CONCEPT Check


✓ What are meteoroids, and why are they important to planetary scientists?

**The Big Question** When will the next asteroid or comet hit Earth? That's the biggest question on the minds of many people, not just astronomers. Over the course of billions of years, life on Earth has been repeatedly disrupted by impacts with these stray objects. Smaller ones whiz by us often, bigger ones less frequently. More are surely on the way, and several agencies are now trying to take inventory of the trash in our cosmic neighborhood that might hurt us. Although today there is little we could do, many scientists are working hard to find ways to deflect future Earth-killers before they devastate our planet.

# Chapter Review

## SUMMARY

1 More than 100,000 **asteroids** (p. 340) have been cataloged. Most orbit in a broad band called the **asteroid belt** (p. 340) between the orbits of Mars and Jupiter. They are probably primal rocks that never clumped together to form a planet. The **Trojan asteroids** (p. 344) share Jupiter's orbit, remaining 60° ahead of or behind that planet as it moves around the Sun. A few **Earth-crossing asteroids** (p. 343) have orbits that intersect Earth's orbit and will probably collide with our planet one day.



2 The largest asteroids are a few hundred kilometers across. Most are much smaller. The total mass of all asteroids combined is less than the mass of Earth's Moon. Asteroids are classified according to the properties of their reflected light: Brighter, S-type (silicate) asteroids dominate the inner asteroid belt, whereas darker, C-type (carbonaceous) asteroids are more plentiful in the outer regions. The C-type asteroids are thought to have changed little since the solar system formed. Smaller asteroids tend to be irregular in shape and may have undergone violent collisions in the past.




3 **Comets** (p. 345) are fragments of icy material that normally orbit far from the Sun. As a comet approaches the Sun, its surface ice begins to vaporize. We see the comet by the sunlight reflected from the dust and vapor released. The **nucleus** (p. 346) of a comet may be only a few kilometers in diameter. It is surrounded by a **coma** (p. 346) of dust and gas and an extensive invisible **hydrogen envelope** (p. 346). Stretching behind the comet is a long **tail** (p. 345), formed by the interaction between the cometary material and the solar wind. The comet's **ion tail** (p. 346) consists of ionized gas particles and always points directly away from the Sun. The comet's **dust tail** (p. 346) is less affected by the solar wind and has a somewhat curved shape. Comets are icy, dusty bodies, sometimes called "dirty snowballs," that are thought to be leftover material unchanged since the formation of the solar system. Their masses are comparable to the masses of small asteroids.



4 Unlike the orbits of most other bodies in the solar system, comets' orbits are often highly elongated and not confined to the ecliptic plane. Most comets are thought to reside in the **Oort cloud** (p. 353), a vast "reservoir" of cometary material, tens of


thousands of astronomical units across, completely surrounding the Sun. A very small fraction of comets happen to have highly elliptical orbits that bring them into the inner solar system. Comets with orbital periods less than about 200 years are thought to originate not in the Oort cloud, but in the **Kuiper belt** (p. 352), a broad band lying roughly in the plane of the ecliptic, beyond the orbit of Neptune. More than 1100 **Kuiper belt objects** (p. 353) are now known. Pluto is the largest member of the class.



5 Pluto is the best-known body orbiting beyond Neptune. It was discovered in the 20th century after a laborious search for a planet that was supposedly affecting Uranus's orbital motion. However, we now know that Pluto is far too small to have any detectable influence on Uranus's path. Pluto has a moon, Charon, whose mass is about one-sixth that of the planet itself. Studies of Charon's orbit around Pluto have allowed the masses and radii of both bodies to be accurately determined. Several bodies comparable in size to Pluto and Charon orbit beyond Neptune. At least one—Eris—is larger than Pluto. Eris and Pluto are currently classified as **dwarf planets** (p. 357) because their masses are too low to have cleared their orbital neighborhoods of other bodies.



6 **Meteors** (p. 358), or "shooting stars," are bright streaks of light that flash across the sky as **meteoroids** (p. 358)—pieces of interplanetary debris—enter Earth's atmosphere. If a meteoroid reaches the ground, it is called a **meteorite** (p. 359). Each time a comet rounds the Sun, some cometary material becomes dislodged, forming a **meteoroid swarm** (p. 359)—a group of small **micrometeoroids** (p. 359) following the comet's original orbit. If Earth happens to pass through the comet's orbit, a **meteor shower** (p. 359) occurs. Larger meteoroids are probably pieces of material chipped off asteroids following collisions in the asteroid belt.



7 The major difference between meteoroids and asteroids is their size: The dividing line between them is conventionally taken to be 100 m. Meteorites are thought to be composed of the same material that makes up the asteroids, and the few orbits that have been determined are consistent with an origin in the asteroid belt. Some meteorites show evidence of heating, but the oldest ones do not. Most meteorites are between 4.4 and 4.6 billion years old. Comets and stray asteroids are responsible for most of the cratering on the various worlds in the solar system. The most recent large impact on Earth occurred in 1908, when an asteroid apparently exploded several miles above Siberia.



**MasteringAstronomy**® For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- LO1** What are the Trojan, Apollo, and Amor asteroids?
- How are asteroid masses measured?
- What do you think might happen if a near-Earth asteroid were found to be on a collision course with Earth?
- LO2** How do the C-type and S-type asteroids differ?
- Are all asteroids found in the asteroid belt?
- What are comets like when they are far from the Sun? What happens when they enter the inner solar system?
- Where in the solar system do most comets reside?
- LO3** Describe the various parts of a comet while it is near the Sun. What are the typical ingredients of a comet nucleus?
- POS** How do we know what comets are made of?
- LO4** Why can comets approach the Sun from any direction, but asteroids generally orbit close to the plane of the ecliptic?
- LO5** In what ways is the Kuiper belt similar to the asteroid belt? In what ways do they differ?
- POS** Why has the number of planets in the solar system recently decreased?
- LO6** Explain the difference between a meteor, a meteoroid, and a meteorite.
- What causes a meteor shower?
- LO7 POS** What do meteorites reveal about the age and formation of the solar system?

## Conceptual Self-Test: Multiple Choice

- VIS** According to Figure 14.1 (“Inner Solar System”), the asteroid groups with the smallest perihelion distances also tend to have orbits that (a) are slowest; (b) are nearly circular; (c) are most eccentric; (d) extend nearly to Jupiter.
- Most main-belt asteroids are about the size of (a) the Moon; (b) North America; (c) a U.S. state; (d) a small U.S. city.
- Spectroscopic studies indicate that the majority of asteroids contain large fractions of (a) carbon; (b) silicate rocks; (c) iron and nickel; (d) ice.
- Trojan asteroids orbiting at Jupiter’s Lagrangian points are located (a) far outside Jupiter’s orbit; (b) close to Jupiter; (c) behind and in front of Jupiter, sharing its orbit; (d) between Mars and Jupiter.
- The tails of a comet (a) point away from the Sun; (b) point opposite the direction of motion of the comet; (c) curve from right to left; (d) curve clockwise with the interplanetary magnetic field.
- Compared with the orbits of the short-period comets, the orbits of long-period comets (a) tend to lie in the plane of the ecliptic; (b) look like short-period orbits, but are simply much larger; (c) are much less eccentric; (d) can come from all directions.
- Kuiper belt objects are not regarded as planets because (a) they orbit too far from the Sun; (b) their masses are too low to clear other bodies from their orbital paths; (c) they are all irregular in shape; (d) they are predominantly icy in composition.
- According to the figure in *Discovery 14-1*, an impact resulting in global catastrophe is expected to occur roughly once per (a) year; (b) century; (c) millennium; (d) million years.
- A meteorite is a piece of interplanetary debris that (a) burns up in Earth’s atmosphere; (b) misses Earth’s surface; (c) glances off Earth’s atmosphere; (d) survives the trip to the surface.
- According to Table 14.1, the meteor shower that occurs closest to the autumnal equinox is the (a) Lyrids; (b) Beta Taurids; (c) Perseids; (d) Orionids.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

- (a) The asteroid Pallas has an average diameter of 520 km and a mass of  $3.2 \times 10^{20}$  kg. How much would a 100-kg astronaut weigh there? (b) What is the asteroid’s escape speed?
- You are standing on the surface of a spherical asteroid 10 km in diameter, of density  $3000 \text{ kg/m}^3$ . Could you throw a small rock fast enough that it escapes? Give the speed required in km/s and mph.
- Using the data given in the text, estimate Dactyl’s orbital period as it orbits Ida.
- (a) Calculate the orbital period of a comet with a perihelion distance of 0.5 AU and aphelion in the Oort cloud, at a distance of 50,000 AU from the Sun. (b) A short-period comet has a perihelion distance of 1 AU and an orbital period of 125 years. What is its maximum distance from the Sun?

5. ● Astronomers estimate that comet Hale-Bopp lost mass at an average rate of about 350,000 kg/s during the time it spent close to the Sun—a total of about 100 days. Estimate the total amount of mass lost and compare it with the comet's estimated mass of  $5 \times 10^{15}$  kg.
6. ●● It has been hypothesized that Earth is under continuous bombardment by house-sized “minicometes” with typical diameters of 10 m, at the rate of some 30,000 per day. Assuming spherical shapes and average densities of  $100 \text{ kg/m}^3$ , calculate the total mass of material reaching Earth each year. Compare the total mass received in the past 1 billion years (assuming that all rates were the same in the past) with the mass of Earth's oceans (see Chapter 7, problem 3).
7. ● A particular comet has a total mass of  $10^{13}$  kg, 95 percent of which is ice and dust. The remaining 5 percent is in the form of rocky fragments with an average mass of 100 g. How many meteoroids would you expect to find in the swarm formed by the breakup of this comet?
8. ●● It is observed that the number of asteroids or meteoroids of a given diameter is roughly inversely proportional to the square of the diameter. Approximating the actual distribution of asteroids first as a single 1000-km body (e.g., Ceres), then as one hundred 100-km bodies, then as ten thousand 10-km asteroids, and so on, and assuming constant densities of  $3000 \text{ kg/m}^3$ , calculate the total mass (in units of Ceres's mass) in the form of 1000-km bodies, 100-km bodies, 10-km bodies, 1-km bodies, and 100-m bodies.

## Activities

---

### Collaborative

1. The only way to tell an asteroid from a star is to watch it over several nights. The magazines *Sky & Telescope* and *Astronomy* often publish charts for especially prominent asteroids. Look for Ceres, Pallas, or Vesta, the brightest asteroids. Use the chart to locate the appropriate star field and aim binoculars at that location in the sky. You may be able to pick out the asteroid from the chart. If you can't, make a sketch of the entire field. Come back a few nights later and look again. The “star” that has moved is the asteroid.

### Individual

1. There are a number of major meteor showers every year, but if you plan to watch one, be sure to notice the phase of the Moon. Bright moonlight or city lights can obliterate a meteor shower. A common misconception about meteor watching is that most meteors are seen in the direction of the shower's radiant. In fact, they can appear in all parts of the sky! Just relax and let your eyes rove among the stars. You will generally see many more meteors in the hours before dawn than in the hours after sunset. Why do you suppose meteors have different brightnesses? Can you detect their variety of colors? Watch for meteors that appear to “explode” as they fall and vapor trails that linger after the meteor itself has disappeared.



# Exoplanets

## PLANETARY SYSTEMS BEYOND OUR OWN

The planets, moons, and small bodies of the solar system present us with a long list of interesting features and bizarre peculiarities. Every object has its idiosyncrasies, some of them due to particular circumstances and others the result of planetary evolution. Each time a new discovery is made, we learn a little more about the history of our planetary system and refine our theories of solar system formation and evolution.

Today, discoveries of planets orbiting other stars are flooding in at an unprecedented rate, giving us unexpected new examples of planetary systems in action and posing fundamental challenges to our understanding of planet formation. How will our solar system theories stand up to this onslaught of new data? Will they be overwhelmed and abandoned, perhaps leading to deeper understanding in the long run? Or are there underlying principles that unify our local knowledge with planetary systems beyond our own? The answer, as we will see, is “Stay tuned...”

**The Big Picture** Our own solar system formed some 4.5 billion years ago, a time so ancient that it’s virtually impossible to reconstruct the details of that remarkable event. Ironically, it is other such planetary systems far beyond our own that are now helping us decipher, like Rosetta stones, our own origins. Just as comparative planetology of our eight neighboring planets guides our knowledge of Earth’s history, extrasolar planets seemingly have much to teach us about how they all came to be in the first place.

**LEFT:** Astronomers routinely observe other young star systems, hoping to gain insight into the origins of our own solar system. This is actually a composite image, taken in the optical domain by two telescopes: The *Hubble Space Telescope* imaged the central parts and Japan’s Subaru Telescope extended the field of view around the edges. It shows the region called S106, a nebula about 3300 light-years away in the constellation Cygnus. Amid its chaotic gas and dust spanning a few light-years (hence thousands of times larger than a typical planetary system), many young stars—and probably planets—are now forming. (NASA; NAOJ)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 List some regular and irregular aspects of the solar system, and explain them in the context of the condensation theory.
- 2 Describe some techniques astronomers use to detect planets beyond the solar system.
- 3 Outline the properties of known extrasolar planets, and list some categories of exoplanet not found in the solar system.
- 4 Explain how extrasolar planets fit in with current theories of solar system formation.
- 5 Describe the current observational evidence for habitable Earth-like planets beyond our solar system.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.



## 15.1 Modeling Planet Formation

The origin of the planets and their moons is a complex and as yet incompletely solved puzzle, although the basic outlines of the processes involved are becoming understood. ∞ (Sec. 6.6) Most of our knowledge of the solar system's formative stages has come from studies of interstellar gas clouds, fallen meteorites, and Earth's Moon, as well as of the various planets observed with ground-based telescopes and planetary space probes. Ironically, studies of Earth itself do not help much because information about our planet's early stages eroded away long ago. Meteorites and comets provide perhaps the most useful information, for nearly all have preserved within them traces of solid and gaseous matter from the earliest times.

Until the mid-1990s, theories of the formation of planetary systems concentrated almost exclusively on our own solar system, for the very good reason that astronomers had no other examples of planetary systems against which to test their ideas. However, all that has now changed, and today astronomers have literally thousands of **extrasolar planets**—planets orbiting stars other than the Sun—to challenge their theories. And challenge them they do! As of mid-2013, we know of almost 900 “official” exoplanets, having confirmed observations and well-measured properties. In addition, some 2500 likely “candidates” await confirmation. As we will see, the exoplanetary systems discovered to date have a broad range of properties, many of them quite different from our own, and may well require us to rethink our conception of how stars and planets form. Figure 15.1 is one of the first images ever obtained of an extrasolar planet. Taken with the European Very Large Telescope in Chile, it shows a nearby system containing a planet a few times more massive than Jupiter. ∞ (Sec. 5.2) Astronomers hope that studies of the diversity of extrasolar planetary systems will shed further light on the formation of our own solar system and on Earth's place in the universe as an abode for life.

Still, realize that although we now know of many extrasolar planets, we have only limited information on each—estimates of orbits and masses, and fragmentary data on composition in a relatively few cases. Accordingly, we begin our study of planetary systems by reviewing the theory that accounts for most of the observed properties of the planetary system we know best: the solar system. Then we will be ready to confront the condensation theory with the fast-growing list of exoplanetary systems to assess how it holds up in the face of new observational data. ∞ (Sec. 1.2)

Figure 15.2 summarizes our earlier discussion in Chapter 6 of the formative stages of our planetary system. ∞ (Secs. 6.6, 6.7) Here, the condensation theory traces the formation of the solar system along the following broad lines, each part keyed to the figure:

(a) An interstellar cloud, actually very much larger than the resulting planetary system, began to contract. As the cloud



▲ **FIGURE 15.1 Extrasolar Planet** Most known extrasolar planets are too faint to be detectable against the glare of their parent stars. However, in this system, called 2M1207, the parent itself (centered) is very faint—a so-called *brown dwarf* (see Chapter 19)—allowing the planet (lower left) to be detected in the infrared. This planet has a mass about 5 times that of Jupiter and orbits 55 AU from the star, which is 230 light-years away. (ESO)

collapsed, it rotated faster, because of the law of conservation of angular momentum, and began to flatten. ∞ (More Precisely 6-1)

(b) By the time it had shrunk to a diameter of about 100 AU, the *solar nebula* had formed an extended, rotating disk. The temperature was greatest in the center, near the red proto-Sun, and coolest at the edges.


(c) Dust grains acted as condensation nuclei, forming clumps of matter that collided, stuck together, and grew into moon-sized (and larger) planetesimals. The composition of the grains and planetesimals depended on location in the nebula: rocky near the center, icy farther out.

(d) After a few million years, strong winds from the still-forming Sun expelled the nebular gas, but some massive planetesimals in the outer solar system had already accreted gas from the nebula.

(e) With the gas ejected, planetesimals in the inner solar system continued to collide and grow, forming the terrestrial planets. The outer jovian planets had already formed, and the Sun had become a star.

(f) Over the course of 100 million years or so, most planetesimals were accreted or ejected, leaving a few large planets moving in roughly circular orbits.

## 15.2 Solar System Regularities and Irregularities



**▲ FIGURE 15.2 Solar System Origin** The condensation theory of planet formation is artistically illustrated by these half-dozen changes, from infalling interstellar cloud at the top to newly emerged planetary system at the bottom. Consult the text on the opposite page for descriptions of each of the frames of this figure. The condensation theory was devised to explain the observed properties of our own solar system. Now astronomers have the opportunity to test it against observations of planetary systems elsewhere in the universe.

The condensation theory naturally accounts for the eight “characteristic” solar system properties listed and discussed in Chapter 6. ∞ (Sec. 6.6) Specifically, the planets’ orbits are nearly circular (2), in the same plane (3), and in the same direction as the Sun’s rotation on its axis (4) as a direct consequence of the nebula’s shape and rotation. The growth of planetesimals throughout the nebula, with each protoplanet ultimately sweeping up the material near it, accounts for the fact (1) that the planets are widely spaced—even if the theory does not quite explain the regularity of the spacing. The heating of the nebula and the Sun’s eventual ignition resulted in the observed differentiation (5), and the debris from the accretion–fragmentation stage naturally accounts for the asteroids (6), the Kuiper belt (7), and the Oort cloud comets (8).

It is important, in the wider context of extrasolar planetary systems, to realize that *all* aspects of the theory just described rest squarely on physical principles thought to apply to all stars in their early, planet-forming stages. In other words—in broad terms, at least—we should expect *all* exoplanetary systems to share these overall properties.

The condensation theory is an example of an *evolutionary* theory—one that describes the development of the solar system as a series of gradual and natural steps, understandable in terms of well-established physical principles. Evolutionary theories may be contrasted with *catastrophic* theories, which invoke accidental or unlikely celestial events to interpret observations. A good example of such a theory is the *collision hypothesis*, which imagines that the planets were torn from the Sun by a close encounter with a passing star. This hypothesis enjoyed some measure of popularity during the 19th century, due in part to the inability of the nebular theory to account for the observed properties of the solar system, but no scientist takes it seriously today. Aside from its extreme improbability,\* the collision hypothesis is completely unable to explain the orbits, the rotations, or the composition of the planets and their moons.

The nebular/condensation theory was constructed in large part to “predict” the regular solar system properties just discussed. But in addition to its many regularities, our solar system has many notable *irregularities*, as we have seen repeatedly in previous chapters. Far from threatening our theory, however, these irregularities are important facts for us to consider in shaping our explanations. For example, the explanation for the solar system must

\*A simple calculation indicates that a star like the Sun would have to wait more than 1 million trillion years—100 million times the age of the universe—for a single close encounter to occur.

not insist that *all* planets rotate in the same sense or have *only* prograde moons because that is not what we observe. Instead, the theory of the solar system should provide strong reasons for observed planetary characteristics, yet be flexible enough to allow for and explain the deviations, too. This aspect of the condensation theory becomes even more important when we try to understand the diverse properties of observed exoplanets.

In the condensation theory, the ability to accommodate the possibility of imperfections—deviations from the otherwise well-ordered scheme of things—is provided by the randomness inherent in the encounters that combined the planetesimals into protoplanets. Scientists generally try not to invoke chance to explain the universe, but there are many instances where random events really were critical in determining the present state of the solar system. As the numbers of large bodies decreased and their masses increased, individual collisions acquired greater and greater importance. The effects of these events can still be seen today in many parts of the solar system—for example, the large craters on many of the moons we have studied thus far. Some larger-scale planetary features thought to be attributable to random events include:

1. Two large bodies could have merged to form Venus, giving it its abnormally low rotation rate. ∞ (Sec. 9.2)
2. The Earth–Moon system may have formed from a collision between the proto-Earth and a Mars-sized object. ∞ (Sec. 8.8)
3. A late collision with a large planetesimal may have caused Mars’s curious north–south asymmetry and ejected much of the planet’s atmosphere. ∞ (Sec. 10.4)
4. The tilted rotation axis of Uranus may have been caused by grazing collisions with two or more large protoplanets late in the planet’s formation. ∞ (Sec. 13.2)
5. Uranus’s moon Miranda may have been almost destroyed by a planetesimal collision, accounting for its bizarre surface terrain. ∞ (Sec. 13.5)
6. Interactions between the jovian protoplanets and one or more planetesimals may account for the irregular moons of those planets and, in particular, Triton’s retrograde motion. ∞ (Sec. 13.5)
7. The Pluto–Charon system and the other known binary Kuiper belt objects may have formed by collisions or near misses between two icy planetesimals before most were ejected by interactions with the jovian planets. ∞ (Sec. 14.3)

Note that it is difficult to test any of these assertions directly, but it is reasonable to suppose that some (or even all) of these “odd” aspects of the solar system can be explained in terms of collisions late in the formative stages of the protoplanetary system. Not all astronomers agree with all of the explanations. However, most would accept at least some.

Finally, we must recognize that not all aspects of the present-day solar system were laid down at its formation long ago. The solar system has had ample time for significant evolutionary changes over its 4.6-billion-year history. Examples discussed in previous chapters include the atmospheres of the terrestrial planets, the synchronous orbits of many satellites (including Earth’s), the Trojan asteroids, the structure of the rings and large moons of the jovian planets, and Pluto’s orbital resonance with Neptune. ∞ (Secs. 7.6, 9.5, 10.6, 11.5, 12.4, 14.2, 14.3) It is reasonable to suppose that many exoplanetary systems have had similar opportunities to evolve—and diverge—since their formation.

### PROCESS OF SCIENCE Check


- ✓ Why is it important that a theory of solar system formation make clear statements about how planets arose, yet not be too rigid in its predictions?

## 15.3 Searching for Extrasolar Planets

The test of any scientific theory is how well it holds up in situations different from those in which it was originally conceived. ∞ (Sec. 1.2) With the discovery in recent years of numerous extrasolar planets orbiting other stars, astronomers now have the opportunity—indeed, the scientific obligation—to test their theories of solar system formation.

The detection of planets orbiting other stars has been a long-standing goal of generations of astronomers. Many claims of extrasolar planets have been made since the middle of the 20th century, but before 1994 none had been confirmed and most have been discredited. Since then, however, this fast-growing field of research has made enormous advances and is now one of the most vibrant areas in all of astronomy. These advances have been achieved through steady improvements in both telescope and detector technology and computerized data analysis.

Extrasolar planets are very faint and generally lie close to their parent stars, making them hard to resolve with current equipment. Figure 15.1 shows a nearby system containing a planet somewhat more massive than Jupiter. In this case, the parent “star” (actually, an object called a *brown dwarf*, to be discussed below) happens to be very faint, so the planet can be seen. In fact, only a few dozen extrasolar planets have so far been detected by direct imaging. In all other cases, the techniques used to find the planets have been *indirect*, based on analyses of light from the parent stars, not from the planets themselves.



**FIGURE 15.3 Detecting Extrasolar Planets** As a planet orbits its parent star, it causes the star to “wobble” back and forth. The greater the mass of the planet, the larger is the wobble. The center of mass of the planet–star system stays fixed. If the wobble occurs along our line of sight to the star, as shown by the yellow arrow, we can detect it by the Doppler effect.

## Radial Velocity Measurements

Figure 15.3 illustrates the basic detection method responsible for the discovery of many extrasolar planetary systems. As a planet orbits a star, gravitationally pulling first one way and then the other, the star “wobbles” slightly as the planet and star orbit their common center of mass.  $\infty$  (Sec. 2.8) The more massive the planet, or the less massive the star, the greater is the star’s movement. If the wobble happens to occur along our line of sight to the star, then we see small fluctuations in the star’s radial velocity, which can be measured using the Doppler effect.  $\infty$  (Sec. 3.5) Those fluctuations allow us to estimate the planet’s mass.

Figure 15.4 shows two sets of radial velocity data that betray the presence of planets orbiting other stars. Part (a) shows the line-of-sight velocity of the star 51 Pegasi, a near twin to our Sun lying some 40 light-years away. The data, acquired in 1994, were the first substantiated evidence for an extrasolar planet orbiting a Sun-like star.\*



The regular 50 m/s fluctuations in the star’s velocity have since been confirmed by several groups of astronomers and imply that a planet of at least half the mass of Jupiter orbits 51 Pegasi in a circular orbit with a period of just 4.2 days. (For comparison, the corresponding fluctuation in the Sun’s velocity due to Jupiter is roughly 12 m/s.) Note that

\*As we will see in Chapter 22, two other planets having masses comparable to Earth, and one planet with a mass comparable to that of Earth’s Moon, had previously been detected orbiting a particular kind of collapsed star called a pulsar. However, their formation was the result of a chain of events very different from those that formed Earth and the solar system.

we say “at least half” here because Doppler observations suffer from a fundamental limitation: They cannot distinguish between low-speed orbits seen edge-on and high-speed orbits seen almost face-on (so only a small component of the orbital motion contributes to the line-of-sight Doppler effect). As a result, only lower limits to planetary masses can be obtained by this method.

Figure 15.4(b) shows another set of Doppler data, this time revealing a more complex system: a triple-planet system orbiting another nearby Sun-like star named Upsilon Andromedae. The three planets have minimum masses of 0.7, 2.1, and 4.3 times the mass of Jupiter and orbital semimajor axes of 0.06, 0.83, and 2.6 AU, respectively. Figure 15.4(c) sketches their orbits, with the orbits of the solar terrestrial planets shown for scale. All told, as of mid-2013, more than 500 planets have been detected by means of radial velocity searches.

If the wobble produced in a star’s motion is predominantly *perpendicular* to our line of sight, then little or no Doppler effect will be observed; thus, the radial velocity technique cannot be used to detect a planet. However, in that case, the star’s *position* in the sky changes slightly from night to night, and, in principle at least, measuring this transverse motion can provide an alternative means of detecting extrasolar planets. Unfortunately, these side-to-side wobbles have proved difficult to measure accurately, as the angles involved are very small and the star in question has to be quite close to the Sun for useful observation to be possible. Based on observations of this type, a few candidate planetary systems have been proposed, but none has yet been placed on the “official” list of confirmed observations.



### Interactive **FIGURE 15.4** Planets Revealed

**MA** (a) Measurements of the Doppler shift of the star 51 Pegasi reveal a clear periodic signal indicating the presence of a planetary companion of mass at least half the mass of Jupiter. (b) Radial-velocity data for Upsilon Andromedae are much more complex, but are well fit (solid black, wobbly line) by a three-planet system orbiting the star. (c) A sketch of the inferred orbits of three planets from the Upsilon Andromedae system (in orange), with the orbits of the terrestrial planets superimposed for comparison (in white).


## Planetary Transits

As just noted, the radial velocity technique suffers from the limitation that the angle between the line of sight and the planet's orbital plane cannot be determined. However, in some systems, many of them originally discovered through Doppler measurements, that is not the case. Figure 15.5(a) shows how observations of a distant solar-type star (known only by its catalog name of HD 209458 and lying some 150 light-years from Earth) reveal a clear drop in brightness each time its 0.6-Jupiter-mass companion, orbiting at a distance of just 7 million km, passes between the star and Earth (Figure 15.5b). The drop in brightness is just 1.7 percent, but it occurs precisely on schedule every 3.5 days, the orbital period inferred from radial-velocity measurements.


Such **planetary transits**, similar to the transit of Mercury shown in Figure 2.17, are relatively rare, as they require us to see the orbit almost exactly edge-on. When they do occur, however, taken in conjunction with radial velocity measurements, they allow the unambiguous determination of the planet's mass and radius, and hence its mean density.  $\infty$  (Sec. 6.2) In the case shown in Figure 15.5, the planet's radius is found to be 100,000 km (1.4 times the radius of Jupiter), implying a density of just  $200 \text{ kg/m}^3$ , indicating a high-temperature gas-giant planet orbiting very close to its parent star.

Given that only a small fraction of planetary systems are expected to be oriented in just the right way to show transits, planet hunters adopt the strategy of repeatedly surveying thousands of stars in the hope of detecting a transit should one occur. Space-based telescopes are particularly well suited to this task, as they can stare continuously at a given region of the sky, making simultaneous, high-precision observations of the target stars. The stable observing conditions above the clouds, haze, and turbulence of Earth's atmosphere mean that orbiting instruments can measure tiny brightness changes—less than the 1 part in  $10^4$  needed to detect an Earth-like planet orbiting a Sun-like star.

The European *CoRoT* mission (short for **C**onvection, **R**otation, and planetary **T**ransits), launched in December 2006, was designed to monitor some 120,000 Sun-like stars for (among other things) brightness fluctuations due to planetary transits. So far, it has observed 31 confirmed planets, including one of the smallest exoplanets yet discovered, just 5 times the mass and 1.7 times the diameter of Earth. Roughly 200 additional candidate planets await confirmation. *CoRoT*'s originally planned 2.5-year lifetime had been extended into 2013, but a computer failure in late 2012 may have marked the end of this pioneering mission. NASA's *Kepler* spacecraft, launched in March 2009, monitored a similar number of stars (about 145,000), but with its larger mirror (0.95-m aperture) and darker location (in Earth-trailing orbit), it has found



(a)



(b)

many more confirmed planets (more than 130) and planetary candidates (more than 2700) than did *CoRoT*. Follow-up observations will undoubtedly show that some of the candidates are “false alarms,” but mission scientists are confident that the vast majority of them will be found to be real, greatly expanding the exoplanetary database. *Kepler*'s highly successful mission was expected to continue into 2016, but it appears to have ended prematurely in May 2013 with the failure of the second of four gyroscopes needed to accurately point the spacecraft.

### PROCESS OF SCIENCE Check

- ✓ Describe the two main ways in which astronomers search for extrasolar planets. Why don't we have many images of these bodies?

### Interactive FIGURE 15.5 An Extrasolar Transit

(a) If an extrasolar planet happens to pass between us and its parent star, the light from the star dims in a characteristic way. (b) Artist's conception of the planet orbiting a Sun-like star known as HD 209458. The planet is 200,000 km across and transits every 3.5 days, blocking about 2 percent of the star's light each time it does so.


## 15.4 Exoplanet Properties

To date, astronomers have discovered some 900 official extrasolar planets and 2700 candidates orbiting a total of 2500 stars. Most of the planets discovered by radial velocity measurements lie within about 500 light-years of the Sun; the planets and candidates discovered by *CoRoT*, *Kepler*, and some specialized techniques to be described in later chapters (see Sections 22.3 and 23.6) generally lie at much greater distances. *Discovery 15-1* describes the closest known exoplanet, orbiting one of the Sun's nearest neighbors in space. About 10 percent of the nearby stars surveyed to date have been found to host planets. In most cases, only a single, often quite massive, planet has been detected, but roughly 20 percent of stars with official planets, and about one-third of the *Kepler* candidate stars, have *multiple-planet systems* containing more than one planet. The three-planet system shown in Figure 15.4(c) is an example. Currently, the most extensive known exoplanetary system (called HD 10810) has seven planets orbiting the parent Sun-like star. These numbers are almost certainly limited by technological shortcomings, however, and most astronomers expect both the fraction of stars with planets and the number of planets per star to increase as detection capabilities and data analysis techniques continue to improve.


### Hot Jupiters and Super-Earths

Figure 15.6(a) presents the observed masses and semimajor axes of roughly 400 extrasolar planets with masses determined by radial velocity measurements and introduces some jargon used in the field. Each dot in the figure represents a planet, and we have added points corresponding to Earth, Neptune, and Jupiter in our own solar system.

Massive exoplanets are often referred to as *Jupiters*, while less massive, but (presumably) still “jovian” planets are called *Neptunes*. The dividing line between Jupiters and Neptunes is somewhat arbitrary, but is typically taken at about twice the mass of Neptune, or 0.1 Jupiter masses. The terminology is




(a)



(b)

**▲ FIGURE 15.6 Extrasolar Planetary Parameters** (a) Masses and orbital semimajor axes of hundreds of known extrasolar planets. Each point represents one planetary orbit. Planets are classified by familiar solar system names, depending on mass, and as hot or cold, depending on distance from their parent star. (b) Radii and semimajor axes of thousands of extrasolar *Kepler* candidates.

suggestive and is intended to distinguish between planets that are mostly gas, like Jupiter, and those that have substantial rocky cores, like Neptune, but realize that the practical division is based only on mass—we have little or no information on most of these planets' composition or internal structure.

Planets with masses between roughly 2 and 10 Earth masses (about half the mass of Neptune) are known as **super-Earths**. The upper limit in this case is significant, as theorists think that 10 Earth masses represents the minimum mass of a planetary core needed to accrete large amounts of nebular gas and begin to form a gas giant.  (Sec. 6.6) Below 2 Earth masses, exoplanets are simply called *Earths*.

Exoplanets are further subdivided depending on their distances from their parent stars. Planets with orbital semimajor axes less than 0.1 AU are said to be *hot*, while those on wider orbits are called *cold*. Again, the dividing line is somewhat arbitrary—the actual temperature of a planet depends not just on the size of its orbit, but also on the composition of the planet's atmosphere and temperature and brightness of the central star.

Only a small fraction (about 4 percent) of the more than 2700 *Kepler* candidates have so far been officially confirmed as planets. Still, the sheer size of the list, and the high probability that most planets on it are real, mean that astronomers take their properties very seriously. Because these candidates have been detected by transit, rather than by radial velocities, we generally don't know their masses, so we can't display them on Figure 15.6(a). However, *Kepler* has measured their radii quite accurately, and Figure 15.6(b) shows the radii and semimajor axes of the planets on the *Kepler* candidate list. The radii of Earth, Neptune, and Jupiter are also marked.

Converting between masses and radii is uncertain because in most cases the planet's composition is unknown. But, making reasonable assumptions, we can say, approximately, that on Figure 15.6(b) Jupiters (as just defined) have radii greater than about 5 Earth radii, Neptunes are between 2 and 5 Earth radii, super-Earths between 1.25 and 2 Earth radii, and Earths less than 1.25 Earth radii. Notice that the horizontal scales on the two figures are different—the *Kepler* candidates in part (b) tend to be smaller and/or lighter and lie much closer to their parent stars than the (mainly radial velocity) official exoplanets in part (a).

Most of the official planets observed so far fall into the “cold Jupiter” or “cold Neptune” categories, like the jovian planets in our own solar system, although their orbits are generally somewhat smaller than those of the jovian planets—less than a few astronomical units across—and considerably more eccentric. Fewer than 20 percent have eccentricities less than 0.1, whereas no jovian planet in our solar system has an eccentricity greater than 0.06. Figure 15.7 plots the actual orbits of some of these planets, with Earth's orbit superimposed for comparison.

A sizable minority of all observed exoplanets—about one-third of the official list and more than half of the *Kepler* candidates—move in “hot” orbits very close to their parent stars and have surface temperatures as high as 1000–2000 K. The most massive ones were the first to be discovered, and they were quickly dubbed **hot Jupiters**. They represent a new class of planet and have no counterparts in our own solar system. In most cases, they reside so close to their parent stars that their orbits have been circularized and synchronized by tidal effects similar to those controlling the orbits of Earth’s Moon and the Galilean moons of Jupiter. [∞ \(Secs. 8.4, 11.5\)](#)

About 70 super-Earths are currently known in the official planet list, with many hundreds more among the *Kepler* candidates. They are found in both hot and cold orbits. Some, especially the lower mass ones, might be large terrestrial planets. Others could be icy planetary cores that never managed to accrete significant amounts of nebular gas. Still others may have substantial atmospheres of light gases, but never grew to “Neptune” status—they are sometimes referred to as *gas dwarfs*. These latter two categories, if real, would represent two more classes of planet unknown in the solar system.


So far, only about a dozen official exo-Earths have been found. Most move on hot orbits close to their parent stars and are unlikely to resemble anything we’d want to call home, although a few may move in more comfortable orbits. In addition, though, the *Kepler* candidate list contains more than 300 possible Earths spanning a broad range of orbits—some potentially habitable, as we discuss in more detail below. One candidate “hot Mercury” was reported in early 2013.

The fact that we don’t see many low-mass planets or more massive planets on wide orbits (i.e., toward the right

or lower parts of Figures 15.6a and b) is not too surprising. Lightweight and/or distant planets simply don’t produce large enough velocity fluctuations for them to be easily detectable by the radial velocity method: Compare the current practical detection limit of about 1 m/s with the 12 m/s solar velocity variation produced by Jupiter’s motion and the 0.5 m/s and 0.1 m/s due to Neptune and Earth, respectively. And small planets or planets on wide orbits are least likely to produce a detectable transit. In other words, the methods employed so far are biased toward finding large and/or massive objects orbiting close to their parent stars. Those systems would be expected to give the strongest signal, and they are precisely what have been observed.

## Planetary Composition

By themselves, radial velocity measurements yield only a mass limit and some orbital parameters for an exoplanet. But as we have just seen, if the planet happens also to transit its star, then we can accurately determine both its mass and its radius and hence estimate its density and even its composition. More than 200 transiting hot Jupiters have been measured in this way, but when astronomers compute their densities they run into a problem—the numbers they obtain are much lower than the values predicted by theory. The computed densities range from 1300 kg/m<sup>3</sup> (the density of Jupiter) down to just 200 kg/m<sup>3</sup> (roughly the density of styrofoam) and are generally *inconsistent*, by a wide margin, with theoretical models, even those assuming the lightest possible composition of pure hydrogen and helium. The leading explanation for this discrepancy is that the heat of the nearby parent star



◀ **FIGURE 15.7 Extrasolar Orbits** The orbits of many extrasolar planets residing more than 0.15 AU from their parent star, superimposed on a single plot, with Earth’s orbit shown for comparison (in white). All these extrasolar planets are comparable in mass to Jupiter. A plot of all known extrasolar planets would be very cluttered, but the message would be much the same: These planetary systems don’t look much like ours!



## DISCOVERY 15-1

### The Closest Exoplanet

Everyone wants to know if “another Earth” exists and how far away it is. As of 2012, a few candidates for Earth-sized exoplanets have been identified, yet so far they all seem rare, distant, hot, and orbit stars unlike the Sun. Recently, though, the discovery of a near twin to Earth has caused quite a stir, since it is surprisingly close to us. European astronomers used a ground-based telescope in the Chilean Andes to examine in detail the nearest star system to the Sun—and what they found is tantalizingly similar to Earth.

The closest star to us in the night sky is not a single star but a group of three stars orbiting one another. This is the Alpha Centauri star system, whose A and B components (currently 4.4 light-years away) closely orbit each other and together appear as a bright source of light easily seen from Earth. Both are Sun-like stars. The third member, called C or Proxima Centauri (currently 4.2 light-years away), is a cool dwarf star invisible to the naked eye; it takes about a million years to orbit the A and B components.



The above map depicts most of the stars visible from Earth in the direction of Alpha Centauri. It shows what part of the southern sky looks like on a clear night without a telescope. The A and B members of the Alpha-Centauri system together form one of the brightest objects in the southern hemisphere; both are marked on the chart within the yellow circle. Proxima is too dim to see here.

The inset at the top right of the map is an actual photograph of just the region in and around the yellow circle on the map. Alpha Centauri (at center) appears so big mainly because this image contains both A and B members and because it is intentionally overexposed (which causes the small

artificial rings of light around all the other stars). Astronomers have searched many such images for dim planets orbiting all of the brightest stars but are always hampered by the intense glare of the star, whose emitted light tends to overwhelm the much dimmer, reflected light from any neighboring planets. It's a little like trying to photograph a minute speck of dust on the surface of a very bright searchlight—virtually impossible.

Instead, astronomers discovered an Earth-sized planet by analyzing the gravitationally induced Doppler motions of Alpha Centauri B, which it orbits. Observations were made several times a night, nearly every night, for more than 3 years. The new exoplanet, which as yet is nameless pending confirmation, has a mass similar to Earth's and a density of rock, but it orbits its parent star about 25 times closer (0.04 AU) than Earth orbits

the Sun. Alas, it's likely a roasted world incapable of hosting water-based life as we know it. So a true twin of Earth has not really been found—but there's a good chance that cooler, rocky planets reside farther out in the star's habitable zone.

The final illustration below is an artist's conception of the new exoplanet—drawn suggestively as a crescent of reflected light (at right) because little is currently known about this alien world. The art also gives an impression of what it would be like to live on this new world: Its central star, Alpha Centauri B (center) would be the brightest star in its sky, followed by Alpha Centauri A

(lower left) and then our own dim Sun at upper right.


Before mounting an expedition to check out our exoplanetary neighbor, humans will need to invent much better ways to travel in space. Four light-years, or about 42 trillion kilometers, is surprisingly far away. Rocket science tells us that it would take about 40,000 years to reach even this nearest star system. Of course, future technologies, such as a nuclear-propelled starship, might get us there quicker, but that future remains science fiction for now.



(D. DeMartin/ESO)



(L. Calcada, N. Risinger/ESO)



◀ **FIGURE 15.8 Earth-like Comparison** Two transiting super-Earths whose masses and radii are accurately known are depicted at the top alongside Earth and Neptune. Based on their average densities, these two new worlds seem to be very different from one another—one is rocky, somewhat like Earth or Neptune’s core, but the other may well be composed predominantly of water and ice. Depicted below are the nine official or candidate “habitable” exo-Earths shown in Figure 15.12. All objects in this figure are drawn to scale and have a color scheme of brown (rocky), blue (icy), yellow (gassy), gray (unknown).

has puffed these planets up far beyond their normal sizes. Possibly, the star’s tidal effect also contributes. However, no current model can account for the observed range in densities, and for now a complete explanation eludes astronomers.

Currently, some 25 transiting Earths and super-Earths have known masses and radii. Their mean densities range from 500 to 9000 kg/m<sup>3</sup>. ∞ (Sec. 6.1) The low end of this range suggests planets harboring large amounts of light gases, probably with rocky/icy cores and hydrogen/helium atmosphere—gas dwarfs. The high end suggests mainly rocky composition—compressed Earths. The intermediate densities suggest planets composed of water and/or other ices.

Figure 15.8 presents a visual comparison of Earth, Neptune, and two super-Earths whose physical properties are relatively well-known. CoRoT 7b is 4.8 times more massive and 1.7 times larger than Earth, implying a mean density of 5300 kg/m<sup>3</sup>—very similar to that of Earth, although because this planet orbits just 0.02 AU from its parent Sun-like star, surface conditions are much more extreme than those on Earth. Figure 15.9(a) is an artist’s impression of what this planet might be like. Second, the planet GJ 1214b has a mass 5.7 times that of Earth and a radius of 2.7 Earth radii, for a mean density of 1600 kg/m<sup>3</sup>. Definitely not rocky, this planet may be composed mainly of water and/or ice, possibly surrounding a small rocky core, with an atmosphere of hydrogen and helium.

Also shown in Figure 15.8 are the nine exo-Earths on the official and *Kepler* candidate lists (see also Figure 15.12) that are thought to reside at just the right distances from their parent stars that liquid water might exist on their surfaces, making them potential sites on which life might have evolved. Figure 15.9(b) is an artistic rendering of one of the more comfortable worlds recently discovered on the edges of habitable zones around their parent stars, such as those in Figure 15.8, where water might flood oceans, gases swirl in atmospheres, and even life potentially exist on the surface. Alas, it’s only art.

Planetary transits provide another important benefit to observers. Because the transit times, once measured, can be accurately predicted, astronomers can time their observations to obtain spectroscopic and other information on the starlit face of the planet during its “quarter” phase, allowing them to probe the planet’s atmospheric composition and dynamics. ∞ (Sec. 4.5) Since planets—even the hot ones—are cooler than their parent stars, their reflected light is most easily distinguished from the background starlight at infrared wavelengths, and the *Spitzer Space Telescope* has played a vital role in these studies. ∞ (Secs. 3.4, 5.7) So far, a variety of atoms and molecules, including hydrogen, sodium, methane (CH<sub>4</sub>), and carbon dioxide (CO<sub>2</sub>), have been detected in numerous planetary atmospheres; *Spitzer* has also observed water vapor in a few cases.

The infrared observations also allow scientists to probe the atmospheric temperatures of a few of these



(a)



(b)

**▲ FIGURE 15.9 Artist Impressions of Exoplanets** Shown here are artists' conceptions of two of the Earth-like planets noted in the previous figure. (a) The rocky exoplanet CoRoT 7b has properties comparable to Earth but must be intensely hot, as depicted here with a molten surface very close to its parent star. (b) The exo-Earth Kepler 54c may resemble this more hospitable world, but it's only an impression based on very limited data. (L. Calçada; M. Kommesser/ESO)

worlds. The tidal effects of their parent stars mean that these planets rotate synchronously, with one face permanently lit and the other permanently dark. ∞ (Sec. 8.4) Yet despite the 2000-K dayside temperatures, the data reveal remarkably small (200 K) temperature differences between the bright and dark sides, suggesting efficient transport of heat between the hemispheres by dynamic, windy atmospheres, probably via zonal flows much like those seen in the jovian planets of our own solar system. ∞ (Secs. 11.2, 12.2, 13.3)

Finally, the mass of the parent stars of the known planets are mostly comparable to or less than that of the Sun—that is, they are typical of stars in the solar neighborhood (see Chapter 17). Spectroscopic observations of

planet-hosting stars reveal what may well be a crucial piece in the puzzle of extrasolar planet formation. Stars having compositions similar to that of the Sun are statistically *much more likely* to have planets orbiting them than are stars containing smaller fractions of the key elements carbon, nitrogen, oxygen, silicon, and iron. Because the elements found in a star reflect the composition of the nebula from which it formed, and the elements just listed are the main ingredients of interstellar dust, this finding provides strong support for the condensation theory. ∞ (Sec. 6.7) Dusty disks really are more likely to form planets.

## Are They Really Planets?

Given that many exoplanetary systems seem to have properties quite different from our own for a time some astronomers questioned whether the mass measurements, and hence the identification of some of these objects as “planets,” could be trusted. Eccentric orbits are known to be common among double-star systems (pairs of stars in orbit around one another—see Section 17.7), and some researchers suggested that many of the newly found planets were really **brown dwarfs**—“failed stars” having insufficient mass to become true stars (Section 19.3).

The dividing line between genuine Jupiter-like planets and star-like brown dwarfs is uncertain, but it is thought to be around 12 Jupiter masses, a number comparable to the largest extrasolar planet masses measured (see Figure 15.6a). Planet proponents argue that this is not a coincidence, but rather indicates that planets up to the maximum possible mass have in fact been observed. Detractors have suggested that we might just be seeing some orbits almost face-on, greatly reducing the parent star's radial velocity and fooling us into thinking that we are observing low-mass planets instead of higher-mass brown dwarfs. However, the latter view has a serious problem, which worsens with every new low-mass extrasolar planet reported: Since the orientations of the actual orbits are presumably random, it is extremely unlikely that we would just happen to see *all* of them face-on, and there is no observational evidence for the many edge-on (and much easier to detect) systems we would also expect to see on statistical grounds.

However, the deciding factor against the brown-dwarf suggestion was the discovery of many *transiting* systems in which the orbital inclination is known and the planet masses well determined. The masses of those planets range from a few Earth masses to 10 Jupiter masses, completely consistent with the “planet” interpretation of the radial velocity data. Consequently, astronomers have concluded that, although there may be a few brown dwarfs lurking among the list of extrasolar planets, they probably do not constitute a significant fraction of the total.

**CONCEPT Check**

- ✓ Describe three distinct ways in which observed extrasolar planetary systems differ from the solar system.

## 15.5 Is Our Solar System Unusual?

Not so long ago, many astronomers argued that the condensation scenario described in Chapter 6 was in no way unique to our own system. ∞ (Secs. 6.6, 6.7) The same basic processes could have occurred, and perhaps *did* occur, they said, during the formative stages of many of the stars in our Galaxy, so planetary systems like our own should be common. Today we know that planetary systems *are* quite common, but in many cases the ones we see differ significantly from our own. We can thus legitimately ask whether our solar system really is unusual and whether the observations described in the previous sections undermine our current theory of solar system formation.

### Overall Planetary Properties

The condensation theory was constructed to account for the observed properties of our solar system. Having studied the corresponding properties of extrasolar planetary systems, let's compare some basic solar system features with those found in extrasolar systems.

1. *Planetary orbits in the solar system are relatively isolated and coplanar.* The 80 or so multiple-planet fits to radial velocity observational data (like that shown in Figure 15.4) often *assume* coplanar motion in order to determine the planetary orbits, so those systems do little to support or challenge the condensation theory. However, as noted earlier, roughly 1 in 3 of the *Kepler* candidates—more than 800 in all—are found in multiple systems, and these orbits must be coplanar (since we necessarily see these systems edge on), strongly supporting this key prediction of the condensation theory. Of course, the *Kepler* data can't rule out the possibility that these systems contain additional planets on very inclined orbits, but we currently have no indication that this might be the case. In addition, the known multiple systems generally appear to have fairly widely spaced orbits, broadly consistent with the orbits in our solar system.
2. *The planets in the solar system all orbit in the same direction as the Sun's rotation.* This generally seems to be the case in extrasolar planetary systems, to the extent that the orbit planes can be determined, although in at least one transiting hot Jupiter system the planet orbit is roughly perpendicular to the star's rotation, or may even be retrograde. The most likely explanation for this

unusual state of affairs seems to be an early encounter with another body that kicked the aberrant planet out of the disk and into its current inclined orbit.


3. *The solar system contains interplanetary debris left over from its formation.* Individual small bodies, such as asteroids and Kuiper belt objects, cannot be seen in extrasolar planetary systems. However, the debris disks observed around many stars (see Figures 6.10 and 6.12) have convinced most astronomers that the processes of accretion and fragmentation predicted by the condensation theory really are occurring in extrasolar systems.

Interestingly, numerous exoplanetary orbital *resonances*—configurations where the orbital periods of different planets are related to one another in some simple way—are known, something not found in the solar system. ∞ (Sec. 8.4) The origin of these resonances is still a matter of debate, but as a practical matter, in systems detected via transits, they afford astronomers an important alternate means of estimating planetary masses, which would otherwise be unknown.


### Cold and Hot Jupiters

As noted in the previous section, most cold extrasolar Jupiters and Neptunes move in orbits that are considerably more eccentric than the orbits of the jovian planets in the solar system. Does this make our system fundamentally different from the others? Probably not. To a large extent, this discrepancy can be explained by the selection effect discussed in Section 15.4—eccentric orbits tend to produce larger velocities and hence are more readily discovered. As search techniques improve, astronomers are finding more and more Jupiter-mass (and lower-mass) planets on wider and less eccentric orbits. Figure 15.10(a) shows evidence for one of the most “Jupiter-like” planets yet detected and 15.10(b) is an artist's conception of what it might look like: a 0.95-Jupiter-mass object moving on a roughly circular orbit around a near-twin of our own Sun. The planet's period is 9.1 years. It is too early to say whether cold Jupiters on nearly circular orbits will turn out to be unusual or common among exoplanets, but they clearly exist among the systems already observed.

Are the eccentric extrasolar orbits we see consistent with the condensation theory? The answer seems to be yes. The theory actually allows many ways in which massive planets can end up in eccentric orbits. Indeed, an important aspect of solar system formation not mentioned in our earlier discussion is the fact that many theorists have worried about how Jupiter could have remained in a stable orbit after it formed in the protosolar disk! Jupiter-sized planets may be knocked into eccentric orbits by random interactions with other Jupiter-sized planets or by the tidal effects of nearby stars. Some researchers have even suggested that violent interactions in the early solar system may have ejected some jovian planets completely. And, alternatively, if the jovian planets formed by



(a)



(b)

### Interactive FIGURE 15.10 Jupiter-like Planet?

**MA** (a) Velocity “wobbles” in the star HD 154345 reveal the presence of an extrasolar planet with one of the most “Jupiter-like” orbits yet discovered. The parent star is almost identical to the Sun, and the 0.95-Jupiter-mass planet orbits at a distance of 4.2 AU with an orbital eccentricity of 0.04. (b) An artist’s impression of a Jupiter-like planet orbiting a Sun-like star. (T. Pyle/NASA)


gravitational instability, they could have had eccentric orbits right from the start (and then we must explain how those orbits circularized in the case of the solar system).

Note that although the observed eccentric Jupiters probably do not represent a serious challenge to the condensation theory, the most eccentric orbits may pose problems for any terrestrial planets those systems may contain. In our solar system, the presence of a massive Jupiter on a nearly circular orbit is known to have a stabilizing influence on the other planetary orbits, tending to preserve the relative tranquility of our planetary environment. In extrasolar systems with very eccentric giant orbits, not only is this stabilization absent, but also having a Jupiter-sized planet repeatedly


plow through the inner parts of the system means that any terrestrial planets have almost certainly been ejected from the system. Fortunately, only a relatively small fraction of the wide (few astronomical units) Jupiter orbits seen so far have eccentricities high enough for this to be the case.

What of the hot Jupiters, which have no counterparts among the planets of the solar system? Here too there is an explanation that fits within the condensation theory! In Section 15.2 we described how the gravitational interaction between massive planets and the gas disk in which they formed probably led to the inward migration of the jovian planets early in the history of the solar system. However, even before the first hot Jupiter was observed, theorists had realized that, depending on how long the disk survived before being dispersed by the newborn Sun, the process could easily have deposited the planet in an orbit very close to the parent star, as illustrated in Figure 15.11. The theorists were right, and the observed hot Jupiters may provide a much-needed connection between extrasolar planetary systems and our own. Interestingly, it appears that the presence of Saturn may have helped stabilize Jupiter’s orbit against this last effect. Isolated or particularly massive Jupiters are precisely the planets one would expect to find on hot orbits.

Finally, strange as it might seem, having a Jupiter-sized planet sink inward through the disk in which planetesimals are still forming and merging is not necessarily detrimental



**Interactive FIGURE 15.11 Sinking Planet** Friction between a giant planet and the nebular disk in which it formed tends to make the planet spiral inward. The process continues until the disk is dispersed by the wind from the central star, possibly leaving the planet in a “hot-Jupiter” orbit.



◀ **FIGURE 15.12 Habitable Zones** Every star is surrounded by a habitable zone, within which an Earth-like planet could have liquid water on its surface. Marked in or near their habitable zones are the eight planets of our solar system, 12 extrasolar super-Earths, and three exo-Earths. About 30 more Kepler candidates, as yet unconfirmed, are marked with smaller dots.

to the formation of terrestrial planets. Note from the timeline in Figure 6.18 that the hot Jupiters must have reached their scorched orbits before the gas disk dispersed—that is, within a few million years of the formation of the solar nebula and long *before* the formation of the terrestrial planets was complete. Computer simulations indicate that the inward migration was a fairly rapid process that stirred up the planetesimals as the giant planet moved through, but for the most part did not disrupt or eject them. The main consequence may have been to mix in more icy material from further out, possibly resulting in planets more massive and more water-rich than would otherwise have been the case.

## Searching for Earths


Giant planets are interesting objects, and hot Jupiters present a fascinating variety of novel problems to observers and theorists alike. And it is still too early to say if the growing number of observed super-Earths—and their absence in our own solar system—will come to pose a problem for the condensation theory. But to many astronomers, the real goal of extrasolar planet research is the detection of terrestrial planets with conditions similar to those found on Earth—and, by extension, the possible

discovery of life elsewhere in the universe. As we have seen in Chapters 10–12, this latter goal is increasingly becoming the motivation behind solar system exploration, too. The numerous Earths and low-mass, terrestrial super-Earths discussed in Section 15.4 represent our starting point in the search for potentially habitable worlds.

If Earth-like environments are our ultimate goal, what orbital parameters are of greatest interest to astronomers? As we saw earlier, many astronomers think that

a key requirement for the development of life as we know it is the existence of liquid water on (or under) a planet's surface, implying a surface temperature roughly in the 0–100° C range. ∞ (Sec. 10.5) The planet's temperature depends both on its distance from its parent star and on the star's intrinsic brightness. Figure 15.12 illustrates how a **habitable zone** (really, a three-dimensional shell) surrounds any given star. Within that zone, liquid water can exist on the planet, making it a possible abode for life.\* Notice how, for low-mass, faint stars, the habitable zone is small and lies close to the star, while for more massive, brighter stars it lies much farther out and can be 1 AU or more wide. Three terrestrial planets—Venus, Earth, and Mars—lie in or near the Sun's habitable zone. As we saw in Chapter 10, any or all of them might have seen the development of life, given the right circumstances. ∞ (Sec. 10.6)

\*Of course, planetary atmospheres can greatly affect surface conditions. The greenhouse effect can warm planets that would otherwise be frozen, whereas the runaway greenhouse effect may transform an apparently habitable planet into an inferno. ∞ (Secs. 7.2, 9.5) Our choice of planets in Figure 15.12 tries to take these effects into account. In addition, there are ways in which liquid water might exist at other locations—under the surface of a jovian moon, for example—but the habitable zone still provides a handy rule of thumb to indicate where planets like Earth might reside. ∞ (Secs. 11.5, 12.5)



◀ **FIGURE 15.13 Whole New Worlds** This artist's illustration depicts the variety of planets detected by the *Kepler* spacecraft—some mostly gaseous, others perhaps with rocky or wet surfaces. Additional observations will be needed to pin down the nature of these alien worlds. (C. Pulliam and D. Aguilar, CFA)

At the time of writing, nine “official” super-Earths and three exo-Earths are known to orbit in or near the habitable zones of their parent stars. They are indicated as labeled points on Figure 15.11. In addition, about 30 more Earths or super-Earths from the *Kepler* candidate list move in habitable orbits. They are also marked in the figure. A few of the official planets with measured radii also have estimated masses, implying densities consistent with rocky/metallic terrestrial composition. The other planets marked on Figure 15.11 are of unknown density and hence composition. Most of the planets in the figure orbit near the “hot” edge of the habitable zone, but this is just another aspect of the observational bias described earlier—planets in close orbits are the most likely to be detected. Figure 15.13 places some of the new findings into perspective, although with a considerable amount of artistic license.

Many planet hunters are confident that within the next decade (or sooner), observational techniques will reach the level of sophistication at which jovian and even terrestrial planets similar to those in our solar system should be readily detectable—if they exist. Advances during the next decade will either bring numerous detections of extrasolar planets in “solar system” orbits or allow astronomers to conclude that systems like our own really are a small minority. Either way, the consequences are profound.

#### PROCESS OF SCIENCE Check


- ✓ How does the condensation theory accommodate the dissimilarities between the properties of planets in our solar system and those of the known extrasolar planets?

**The Big Question** People from all walks of life—not just astronomers—eagerly await the discovery of a true Earth-like planet around another star. When will Earth’s twin be found, and will it have blue skies, deep oceans, and livable lands? Most intriguing of all, will it be inhabited? We live at a remarkable time when we are actually addressing—and often answering—some of the most profound questions that human beings have pondered for thousands of years.

# Chapter Review

## SUMMARY

1 Many large-scale, regular properties of the solar system (such as its coplanar, prograde planetary orbits) are well explained by the condensation theory and are expected to be found in exoplanetary systems, too. Other, irregular properties (such as anomalous rotation or evidence of major impacts) are the result of random events that occurred after the solar system formed. We can't predict the outcomes of these random processes, but they are a necessary part of the theory, and we expect that similar processes may have shaped the detailed evolution of most, if not all, exoplanetary systems.




2 Many of the currently known extrasolar planets were discovered by observing their parent star wobble back and forth as the planet orbits. Most of the rest have been observed as they **transit** (p. 000) their parent star, passing directly in front of it and slightly reducing its brightness. Other detection methods include direct imaging and studies of circumstellar disks.




3 Astronomers have identified some 900 official **extrasolar planets** (p. 368) and 2500 planetary candidates. About 500 of the official planets have been identified by radial velocity measurements. The remainder (including all the candidates) have been detected via transits. About 20 percent of all official planetary systems, and one-third of the candidates, contain more than one planet. Known exoplanetary masses range from that of Earth to many times that of Jupiter. Some planets move on “hot” orbits close to their parent star, while others move on wide, “cold” orbits similar to those of the jovian planets in

the solar system. **Hot Jupiters** (p. 375) and **super-Earths** (p. 374) are new classes of planet not known in the solar system. It is not yet known whether our solar system is unusual among planetary systems. The observed exoplanetary classes are compatible with the condensation theory, and the fact that we see the extrasolar planets we do is due at least in part to the fact that these are the planets we can best see using current techniques. Stars containing larger fractions of “dusty” elements, such as carbon and silicon, are more likely to have planets, again consistent with the condensation theory.



4 It is not known whether systems like our solar system are common or rare among extrasolar planets. A growing number of observed exoplanetary systems contain Jupiter-sized planets on wide, roughly circular orbits, the eccentric Jupiters, hot Jupiters, and super-Earths seen elsewhere can be accommodated within the condensation theory, even though they have no counterparts in the solar system.



5 About 20 Earths and super-Earths are known to orbit within the **habitable zone** (p. 381) of their parent star—the region surrounding the star where liquid water can exist on a planet's surface. Their structure and composition are unknown, but in the few cases where both masses and radii have been measured, the inferred densities are consistent with those of the terrestrial planets.



**MasteringAstronomy**<sup>®</sup> For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

1. **LO1 POS** Give three examples of present-day properties that our solar system model does not have to explain, and say why no explanation is necessary.
2. **POS** Explain the difference between evolutionary theories and catastrophic theories of the solar system's origin.
3. Describe some ways in which random processes played a role in the determination of planetary properties.
4. **LO2 POS** Describe three ways in which astronomers set about looking for extrasolar planets.
5. Why are current detection techniques biased toward finding large or massive exoplanets orbiting close to their parent stars?
6. If transits are so rare, why do astronomers think they are the best way to search for extrasolar Earths?



7. **LO3** In what ways do extrasolar planetary systems differ from our own solar system?
8. Describe some ways in which observed exoplanetary systems are similar to our solar system.
9. What is a hot Jupiter?
10. What is a super-Earth?
11. **POS** Do the observed extrasolar planets imply that Earth-like planets are rare?
12. **LO4 POS** Is our solar system unusual among planetary systems?
13. What is the habitable zone of a star?
14. **LO5** What evidence do we have for habitable Earth-like planets orbiting other stars?
15. Do you think humans could survive on a 10-Earth mass terrestrial super-Earth?

## Conceptual Self-Test: Multiple Choice

1. A successful scientific model of the origin of planetary systems must be able to account for all of the following solar system features, except for (a) intelligent life; (b) the roughly circular planetary orbits; (c) the roughly coplanar planetary orbits; (d) the extremely distant orbits of the comets.
2. Using the standard model of planetary system formation, scientists invoke catastrophic events to explain why (a) Mercury has no moon; (b) Pluto is not a gas giant; (c) Uranus has an extremely tilted rotation axis; (d) there is no planet between Mars and Jupiter.
3. Astronomers have confirmed the existence of at least (a) one; (b) ten; (c) several hundred; (d) several thousand planets beyond our own solar system.
4. The distance to the nearest exoplanetary system is (a) a few light-years; (b) about 100 light-years; (c) about 1000 light-years; (d) thousands of light-years.
5. **VIS** From Figure 15.6(a) (“Extrasolar Planetary Parameters”), most official extrasolar planets are (a) hot and massive (comparable in mass to Jupiter); (b) cool and light (similar in mass to Earth); (c) hot and light; (d) cool and massive.
6. **VIS** From Figure 15.6(b) (“Extrasolar Planetary Parameters”), most planets detected by *Kepler* are (a) very massive; (b) much farther from their star than Earth is from the Sun; (c) rocky; (d) larger than Earth.
7. Super-Earths are (a) made of nickel and iron; (b) comparable in size to Neptune; (c) a few times more massive than Earth; (d) usually found in “hot” orbits.
8. **VIS** In Figure 15.17 (“Habitable Zones”), the habitable zone of a star twice as massive as the Sun (a) is centered at roughly 3 AU from the star; (b) is more than 10 AU wide; (c) lies entirely within 1 AU of the star; (d) is the same size as the Sun’s habitable zone.
9. A planet in the habitable zone (a) has living creatures on it; (b) may have liquid water on its surface; (c) is rocky, like Earth; (d) has an oxygen atmosphere.
10. The total number of habitable, Earth-like exoplanets is approximately (a) 10; (b) 100; (c) 1000; (d) unknown.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

1. ●● A hot Jupiter orbiting an 0.5 solar mass star has an orbital period of 4 days. What is its distance from the star?
2. ● By what fraction would a Neptune-size planet dim the light from a star half the size of the Sun during a transit?
3. ● Estimate how long *Kepler* would have to observe a planet orbiting 10 AU from a Sun-like star in order to report a detection.
4. ● A Jupiter-sized planet is observed to lie 5 arc seconds from its parent star, which lies 50 light-years from Earth. What is the distance from the planet to the star?
5. ●● The two planets orbiting the nearby star Gliese 876 are observed to be in a 2:1 resonance (i.e., the period of one is twice that of the other). The inner planet has an orbital period of 30 days. If the star’s mass is the mass of the Sun, calculate the semimajor axis of the outer planet’s orbit.
6. ●● The planet orbiting star HD187123 has a semimajor axis of 0.042 AU. If the star’s mass is 1.06 times the mass of the Sun, calculate how many times the planet has orbited its star since the paper announcing its discovery was published on December 1, 1998.
7. ●● Stellar radial velocity variations as small as 1 m/s can be detected with current technology. For a Jupiter-mass planet orbiting a Sun-like star, this corresponds to a planetary orbital velocity of approximately 1 km/s. Based on this information, what is the radius of the widest circular orbit on which Jupiter could currently be detected orbiting the Sun?
8. ● Given the data provided in the text, calculate the gravitational acceleration at the surfaces of the two transiting super-Earths (CoRoT 7b and GJ 1214B) discussed in Section 15.4.

## Activities

---

### Collaborative

1. Which of the “irregular” characteristics of the solar system listed in Section 15.2 do you think are most and least likely to have occurred by pure chance? Debate your choice and support your arguments with observational data. Can you think of other irregular features not on the list?

### Individual

1. Do you think our solar system is unusual? Justify your opinion using data obtained from the Extrasolar Planets Encyclopedia, which you can find online at <http://exoplanet.eu>, and the *Kepler* database at <http://exoplanetarchive.ipac.caltech.edu>. Use these resources to find examples of (1) a hot Jupiter, (2) a cold Jupiter on an orbit like Jupiter’s, (3) a hot super-Earth, (4) a super-Earth in the habitable zone, and (5) an Earth in the habitable zone.

## PART THREE

# Stars and Stellar Evolution



Portrait of Cecilia Payne-Gaposchkin  
(Harvard)

**Life can be hard** for graduate students in astronomy. They take some tough courses, many in physics, and they assist in teaching undergraduate courses, but mostly they strive to do original research. Ideally, on the (typically 5- or 6-year) road to their Ph.D. degree, they make a discovery or gain some unique insight that they then write up as part of their doctoral dissertation. The process is exhausting, and some leave the field after the grad school grind, never again to publish in a scientific journal. Others, though, find it exhilarating, and go on to highly productive careers in astronomy.


Arguably one of the most brilliant doctoral theses in astronomy was written in 1925 by a student at Harvard—and she did it in 2 years. Cecilia Payne-Gaposchkin (1900–1979) was an English student who crossed the ocean to pursue graduate studies at Radcliffe College, and she quickly gravitated to the nearby Harvard Observatory, then perhaps the leading center for research on stars. It was also a place where women, though they often did not get the credit at the time, were making some of the most fundamental advances in stellar astronomy. It was Nirvana for her, and she never left.

Cecilia Payne knew far more physics than most astronomers of the time. She was one of the first to apply the then revolutionary quantum theory of atoms to the spectra of stars, thereby ascertaining stellar temperatures and chemical abundances. Of fundamental importance, her work proved that hydrogen and helium—not the heavy elements, as was then supposed—are the most abundant elements in stars and, therefore, in the universe. Her findings were so revolutionary that the leading theorist of the time, Henry Norris Russell of Princeton, declared her work to be “clearly impossible.” It took years to convince the astronomical community that hydrogen is about a million times more abundant in stars than are most of the common elements found on Earth. Yet we take these findings for granted today.


In collaboration with her husband, the exiled Russian astronomer Sergei Gaposchkin, Cecilia Payne-Gaposchkin spent decades making literally millions of observations of thousands of star clusters, variable stars, and galactic novae. Her analysis provided a firm theoretical basis for many properties of stars and their use as distance indicators in the universe. Much of her work has stood the test of time. Despite a flood of new data and new theoretical ideas, it remains the bedrock of modern astronomy.



Women “computers” at work; Payne at the  
inclined desk (Harvard)



Variable Star WR 124 (STScI)




Pistol Star (STScI)

**Today**, the landscape of stellar research is even richer than Cecilia Payne-Gaposchkin knew. We now see stars much more clearly, their spectra in much finer detail, and all of it to much greater distances. Not only do we know what stars are made of, in most cases we also know *why* their composition is as it is and how many of those stars are born, live, and die. Yet the picture is still very much unfinished, as 21st-century astronomy continues to uncover new and exciting features of stars and stellar systems.


In 1976, well past her retirement, Payne-Gaposchkin was chosen (perhaps ironically, given Russell's initial reaction to her work half a century earlier) as the Henry Norris Russell Lecturer, the highest accolade of the American Astronomical Society. Her talk was an enthusiastic summary of a lifetime of astronomical research—an encyclopedic talk with no notes and no prompts, given in perfectly punctuated English. It was very clear that she knew some individual giant stars as well as she knew her best friends. Her talk ended with the following advice to astronomers young and old:

"The reward of the young scientist is the emotional thrill of being the first person in the history of the world to see something or to understand something. Nothing can compare with that experience. The reward of the old scientist is the sense of having seen a vague sketch grow into a masterly landscape. Not a finished picture, of course; a picture that is still growing in scope and detail with the application of new techniques and new skills. The old scientist cannot claim that the masterpiece is his own work. He may have roughed out part of the design, laid on a few strokes, but he has learned to accept the discoveries of others with the same delight that he experienced on his own when he was young."


Illustrated on this page are some recent findings in stellar research—work that undoubtedly would have caused Cecilia Payne-Gaposchkin to express more of her trademark enthusiasm. Today's research also would have made her justly proud, for so much of it relies on the insights gained by her and her colleagues during the first half of the 20th century.




Rosebud Nebula (JPL)



Henize Nebula (JPL)





# The Sun

## OUR PARENT STAR

Living in the solar system, we have the chance to study at close range perhaps the most common type of cosmic object—a star. Our Sun is a star, and a fairly average one at that, but with a unique feature: It is very close to us—some 300,000 times closer than our next nearest neighbor, Alpha Centauri. Whereas Alpha Centauri is 4.3 light-years distant, the Sun is only 8 light-minutes away from us. Consequently, astronomers know far more about the properties of the Sun than about any of the other distant points of light in the universe.

A good fraction of all our astronomical knowledge is based on modern studies of the Sun—from the production of seemingly boundless energy in its core to the surprisingly complex activity in its atmosphere. Just as we studied our parent planet, Earth, to set the stage for our exploration of the solar system, we now examine our parent star, the Sun, as the next step in our exploration of the universe.

**The Big Picture** The Sun is our star—the main source of energy that powers weather, climate, and life on Earth. Imagine our planet without the Sun—no light, no heat, no comforting “parent” in the sky. Although we take it for granted each and every day, the Sun is vitally important to us in the cosmic scheme of things. Simply put, without the Sun, we would not exist.

**LEFT:** The Sun, much like the planets, experiences a kind of weather, including storms. This spectacular image, taken with a filter in the ultraviolet part of the spectrum to diminish the glare and enhance contrast, shows moderate surface activity (in white). The image was observed in 2011 with the *Solar Dynamics Observatory*—a robot that orbits Earth but stares at the Sun unblinkingly 24 hours a day, eavesdropping on its atmosphere, surface, and interior. (NASA)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Summarize the overall properties and internal structure of the Sun.
- 2 Describe the concept of luminosity, and explain how it is measured.
- 3 Explain how studies of the solar surface tell us about the Sun’s interior.
- 4 List and describe the outer layers of the Sun.
- 5 Describe the nature and variability of the Sun’s magnetic field.
- 6 List the various types of solar activity, and explain their relation to solar magnetism.
- 7 Outline the process by which energy is produced in the Sun’s interior.
- 8 Explain how observations of the Sun’s core changed our understanding of fundamental physics.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.

## 16.1 Physical Properties of the Sun

The Sun is the sole source of light and heat for the maintenance of life on Earth. The Sun is a **star**—a glowing ball of gas held together by its own gravity and powered by nuclear fusion at its center. In its physical and chemical properties, the Sun is similar to most other stars, regardless of when and where they formed. Indeed, our Sun appears to be a rather typical star, lying right in the middle of the observed ranges of stellar mass, radius, brightness, and composition. Far from detracting from our interest in the Sun, this very mediocrity is one of the main reasons that astronomers study it—they can apply knowledge of solar phenomena to many other stars in the universe.

### Overall Properties

The Sun's radius, roughly 700,000 km, is determined most directly by measuring the angular size ( $0.5^\circ$ ) of the Sun and then employing elementary geometry. [∞ \(Sec. 1.6\)](#) The Sun's mass,  $2.0 \times 10^{30}$  kg, follows from Newton's laws of motion and gravity, applied to the observed orbits of the planets. [∞ \(More Precisely 2-2\)](#) The average solar density derived from its mass and volume, approximately  $1400 \text{ kg/m}^3$ , is quite similar to that of the jovian planets and about one-quarter the average density of Earth.

Solar rotation can be measured by timing sunspots and other surface features as they traverse the solar disk. [∞ \(Sec. 2.4\)](#) These observations indicate that the Sun rotates in about a month, but it does not do so as a solid body. Instead, it spins *differentially*, like Jupiter and Saturn—faster at the equator and slower at the poles. [∞ \(Sec. 11.1\)](#) The equatorial rotation period at the equator is about 25 days. Sunspots are never seen above latitude  $60^\circ$  (north or south), but at that latitude they indicate a 31-day period. Other measurement techniques, such as those discussed in Section 16.2, reveal that the Sun's rotation period continues to increase as we approach the poles. The polar rotation period is not known with certainty, but it may be as long as 36 days.

The Sun's surface temperature is measured by applying the radiation laws to the observed solar spectrum. [∞ \(Sec. 3.4\)](#) The distribution of solar radiation has the approximate shape of a blackbody curve for an object at about 5800 K. The average solar temperature obtained in this way is known as the Sun's *effective temperature*.

Having a radius of more than 100 Earth radii, a mass of more than 300,000 Earth masses, and a surface temperature well above the melting point of any known material, the Sun is clearly a body that is very different from any other we have encountered so far.

### Solar Structure

The Sun has a surface of sorts—not a solid surface (the Sun contains no solid material), but rather that part of the


brilliant gas ball we perceive with our eyes or view through a heavily filtered telescope. This “surface”—the part of the Sun that emits the radiation we see—is called the **photosphere**. Its radius is about 700,000 km. However, the thickness of the photosphere is probably no more than 500 km, less than 0.1 percent of the radius, which is why we perceive the Sun as having a well-defined, sharp edge (Figure 16.1).

The main regions of the Sun are illustrated in Figure 16.2 and summarized in Table 16.1. We will discuss them all in more detail later in the chapter. Just above the photosphere is the Sun's lower atmosphere, called the **chromosphere**, about 1500 km thick. From 1500 km to 10,000 km above the top of the photosphere lies a region called the **transition zone**, in which the temperature rises dramatically. Above 10,000 km, and stretching far beyond, is a tenuous (thin), hot upper atmosphere: the solar **corona**. At still greater distances, the corona turns into the **solar wind**, which flows away from the Sun and permeates the entire solar system. [∞ \(Sec. 6.5\)](#) Extending down some 200,000 km below the photosphere is the **convection zone**, a region where the material of the Sun is in constant convective motion. Below the convection zone lies the **radiation zone**, in which solar energy is transported toward the surface by radiation rather than by convection. The term *solar interior* is often used to mean both the radiation and convection zones. The central **core**, roughly 200,000 km in radius, is the site of powerful nuclear reactions that generate the Sun's enormous energy output.

### Luminosity


The properties of size, mass, density, rotation rate, and temperature are familiar from our study of the planets. But the Sun has an additional property, perhaps the most important of all from the point of view of life on Earth: The Sun *radiates* a great deal of energy into space, uniformly (we assume) in all directions. By holding a light-sensitive device—a photoelectric cell, perhaps—perpendicular to the Sun's rays, we can measure how much solar energy is received per square meter of surface area every second. Imagine our detector as having a surface area of 1 square meter ( $1 \text{ m}^2$ ) and as being placed at the top of Earth's atmosphere. The amount of solar energy reaching this surface each second is a quantity known as the **solar constant**, whose value is approximately 1400 watts per square meter ( $\text{W/m}^2$ ).

About 50 to 70 percent of the incoming energy from the Sun reaches Earth's surface; the rest is intercepted by the atmosphere (30 percent) or reflected away by clouds (0 to 20 percent). Thus, on a clear day, a sunbather's body having a total surface area of about  $0.5 \text{ m}^2$  receives solar energy at a rate of roughly  $1400 \text{ W/m}^2 \times 0.70$  (70 percent)  $\times 0.5 \text{ m}^2 \approx 500 \text{ W}$ , equivalent to the output of a small electric room heater or five 100-watt lightbulbs.



▲ **FIGURE 16.1 The Sun** The inner part of this composite, filtered image of the Sun shows a sharp solar edge, although our star, like all stars, is made of a gradually thinning gas. The edge appears sharp because the solar photosphere is so thin. The outer portion of the image is the solar corona, normally too faint to be seen, but visible during an eclipse, when the light from the solar disk is blotted out. Note the blemishes; they are sunspots. 🔗 (Sec. 2.4) (NOAO)

Let us now ask about the *total* amount of energy radiated in all directions from the Sun, not just the small fraction intercepted by our detector or by Earth. Imagine a three-dimensional sphere is centered on the Sun and just large enough that its surface intersects Earth's center (Figure 16.3). The sphere's radius is 1 AU, and its surface area is therefore  $4\pi \times (1 \text{ AU})^2$ , or approximately  $2.8 \times 10^{23} \text{ m}^2$ . Multiplying the rate at which solar



▲ **FIGURE 16.2 Solar Structure** The main regions of the Sun, not drawn to scale, with some physical dimensions labeled. The photosphere is the visible “surface” of the Sun. Below it lie the convection zone, the radiation zone, and the core. Above the photosphere, the solar atmosphere consists of the chromosphere, the transition zone, and the corona.

energy falls on each square meter of the sphere (i.e., the solar constant) by the total surface area of our imaginary sphere, we can determine the total rate at which energy leaves the Sun's surface. This quantity is known as the **luminosity** of the Sun. It turns out to be just under  $4 \times 10^{26} \text{ W}$ .


The Sun is an enormously powerful source of energy. *Every second*, it produces an amount of energy equivalent to the detonation of about 10 billion 1-megaton nuclear bombs. Six seconds worth of solar energy output, suitably

**TABLE 16.1 The Standard Solar Model**

Region	Inner Radius (km)	Temperature (K)	Density ( $\text{kg}/\text{m}^3$ )	Defining Properties
Core	0	15,000,000	150,000	Energy generated by nuclear fusion
Radiation zone	200,000	7,000,000	15,000	Energy transported by electromagnetic radiation
Convection zone	496,000*	2,000,000	150	Energy carried by convection
Photosphere	696,000*	5800	$2 \times 10^{-4}$	Electromagnetic radiation can escape—the part of the Sun we see
Chromosphere	696,500*	4500	$5 \times 10^{-6}$	Cool lower atmosphere
Transition zone	698,000*	8000	$2 \times 10^{-10}$	Rapid increase in temperature
Corona	706,000*	3,000,000	$10^{-12}$	Hot, low-density upper atmosphere
Solar wind	10,000,000	> 1,000,000	$10^{-23}$	Solar material escapes into space and flows outward through the solar system

\* These radii are based on the accurately determined radius of the photosphere. The other radii quoted are approximate, round numbers.





▲ **FIGURE 16.3 Solar Luminosity** If we draw an imaginary sphere around the Sun so that the sphere's surface passes through Earth's center, then the radius of this imaginary sphere equals 1 AU. The "solar constant" equals the power striking a 1-m<sup>2</sup> detector at Earth's distance, as implied in the inset. The Sun's luminosity is then determined by multiplying the sphere's surface area by the solar constant. (NASA)

focused, would evaporate all of Earth's oceans. Three minutes would melt our planet's crust. The scale on which the Sun operates simply defies earthly comparison. Let's begin our more detailed study with a look at where all this energy comes from.

### PROCESS OF SCIENCE Check

- ✓ Why must we assume that the Sun radiates equally in all directions when we compute the solar luminosity from the solar constant?

## 16.2 The Solar Interior


How do astronomers know about conditions in the interior of the Sun? As we have just seen, the fact that the Sun shines tells us that its center must be very hot, but our direct knowledge of the solar interior is actually quite limited. (See Section 16.7 for a discussion of one important "window" we do have into the solar core.) Lacking direct measurements, researchers must use other means to probe the inner workings of our parent star. To this end, they construct *mathematical models* of the Sun, combining all available data with theoretical insight into solar physics

to find the model that agrees most closely with observations. ∞ (Sec. 1.2) Recall from Chapter 11 how similar techniques are used to infer the structures of the jovian planets. ∞ (Sec. 11.3) The result in the case of the Sun is the **standard solar model**, which has gained widespread acceptance among astronomers.

## Modeling the Structure of the Sun

The Sun's bulk properties—its mass, radius, temperature, and luminosity—do not vary much from day to day or from year to year. Although we will see in Chapter 20 that stars like the Sun do change significantly over periods of *billions* of years, for our purposes here this slow evolution may be ignored. On "human" time scales, the Sun may reasonably be thought of as unchanging.

Based on this simple observation, as illustrated in Figure 16.4, theoretical models generally begin by assuming that the Sun is in a state of **hydrostatic equilibrium**, in which pressure's outward push exactly counteracts gravity's inward pull. This stable balance between opposing forces is the basic reason that the Sun neither collapses under its own weight nor explodes into interstellar space. ∞ (More Precisely 8-1) The assumption of hydrostatic equilibrium, coupled with our knowledge of some basic physics, then lets us predict the density and temperature in the solar interior. This information, in turn, allows the model to make predictions about other observable solar properties—luminosity, radius, spectrum, and so on—and the internal details of the model are fine-tuned until the



▲ **Narrated FIGURE 16.4 Stellar Balance** In the interior of a star such as the Sun, the outward pressure of hot gas balances the inward pull of gravity. This is true at every point within the star, guaranteeing its stability.

predictions agree with observations. This is the scientific method at work; the standard solar model is the result. ∞ (Sec. 1.2)


Hydrostatic equilibrium has an important consequence for the solar interior. Because the Sun is very massive, its gravitational pull is very strong, so very high internal pressure is needed to maintain the balance. This high pressure in turn requires a very high central temperature, a fact crucial to our understanding of solar energy generation (Section 16.6). Indeed, calculations of this sort carried out by British astrophysicist Sir Arthur Eddington around 1920 provided astronomers with the first inkling that fusion might be the process that powers the Sun.

To test and refine the standard solar model, astronomers are eager to obtain information about the solar interior. However, with so little direct information about conditions below the photosphere, we must rely on more indirect techniques. In the 1960s, measurements of the Doppler shifts of solar spectral lines revealed that the surface of the Sun oscillates, or vibrates, like a complex set of bells. ∞ (Secs. 3.5, 4.5) These vibrations, illustrated in Figure 16.5(a), are the result of internal pressure waves (somewhat like sound waves in air) that reflect off the photosphere and repeatedly cross the solar interior (Figure 16.5b). Because the waves can penetrate deep inside the Sun, analysis of their surface patterns allows scientists to study conditions far below the Sun's surface. The process is similar to the way in which seismologists learn about the interior of Earth by observing the P- and S-waves produced by earthquakes. ∞ (Sec. 7.3) For this reason, the study of solar surface patterns is usually called

**helioseismology**, even though solar pressure waves have nothing whatever to do with solar seismic activity—there is no such thing.

The most extensive study of solar vibrations is the ongoing GONG (short for Global Oscillations Network Group) project. By making continuous observations of the Sun from many clear sites around Earth, solar astronomers can obtain uninterrupted high-quality solar data spanning weeks at a time. The space-based *Solar and Heliospheric Observatory (SOHO)*, launched by the European Space Agency in 1995 and now permanently stationed between Earth and the Sun some 1.5 million kilometers from our planet (see Discovery 16-1), has provided continuous monitoring of the Sun's surface and atmosphere since 1995. Analysis of these data sets provides detailed information about the temperature, density, rotation, and convective state of the solar interior, permitting direct comparison with theory over a large portion of the Sun's volume. The agreement between the standard solar model and observations is spectacular—the frequencies and wavelengths of observed solar oscillations are within 0.1 percent of the model predictions.

The data also allow scientists to monitor global circulation patterns—large-scale gas flows in the solar interior, including two gigantic “conveyor belts” that transport subsurface material from the equator to the poles, then return it to the equator at a depth of some 300,000 km, far below the convection zone. These circulation patterns, which move at 10–15 m/s and take roughly 40 years to complete a single loop, are thought to play crucial roles in regulating the sunspot cycle (see Section 16.4).




▲ **FIGURE 16.5 Solar Oscillations** (a) By observing the motion of the solar surface, scientists can determine the wavelengths and the frequencies of the individual waves and deduce information about the Sun's complex vibrations. (b) Waves contributing to the observed oscillations can travel deep inside the Sun, providing vital information about the solar interior. (*National Solar Observatory*)


Figure 16.6 shows the solar density and temperature, plotted as functions of distance from the Sun's center, according to the standard solar model. Notice how the density drops rather sharply at first and then decreases more slowly near the solar photosphere, some 700,000 km from the center. The variation in density is large, ranging from a core value of about  $150,000 \text{ kg/m}^3$ , 20 times the density of iron, to an intermediate value (at 350,000 km) of about  $1000 \text{ kg/m}^3$ , the density of water, to an extremely small photospheric value of  $2 \times 10^{-4} \text{ kg/m}^3$ , 10,000 times less dense than air at the surface of Earth. Because the density is so high in the core, roughly 90 percent of the Sun's mass is contained within the inner half of its radius. The solar density continues to decrease out beyond the photosphere, reaching values as low as  $10^{-23} \text{ kg/m}^3$  in the far corona—about as thin as the best vacuum physicists can create in laboratories on Earth.

The solar temperature also decreases with increasing radius in the solar interior, but not as rapidly as the density. Computer models indicate a temperature of about 15 million K at the core, consistent with the minimum 10 million K needed to initiate the nuclear reactions known to power most stars, decreasing to the observed value of about 5800 K at the photosphere.

As the data improve and old mysteries are resolved, new ones often emerge. For example, helioseismology indicates that the Sun's rotation speed varies with depth—perhaps not too surprising, given the surface differential rotation mentioned earlier and the fact that similar behavior has been noted in the outer planets. What is puzzling, though, is the *complexity* of the differential motion. The surface layers show a “zonal flow” of sorts, with alternating bands of higher- and lower-than-average rotation rates. Just below the surface are wide “rivers” of lower speed (at the equator) and higher speed (polar) rotation. The material at the base of the convection zone appears to oscillate in rotation speed, sometimes moving faster (by about 10 percent) than the surface layers, sometimes slower, with a period of about 1.3 years. Deeper still, the radiative interior rotates more or less as a solid body, once every 26.9 days. A full explanation of the Sun's rotation currently eludes theorists.

## Energy Transport

The very hot solar interior ensures violent and frequent collisions among gas particles. Particles move in all directions at high speeds, bumping into one another unceasingly. In and near the core, the extremely high temperatures guarantee that the gas is completely ionized. Recall from Chapter 4 that, under less extreme conditions, atoms absorb photons that can boost their electrons to more excited states.  (Sec. 4.2) With no electrons left on atoms to capture the photons, however, the deep solar interior is relatively



▲ **FIGURE 16.6 Solar Interior** Density and temperature vary greatly inside the Sun. Parts (b) and (c) show the changes in solar density and temperature, relative to the cutaway diagram of the Sun's interior (a).

transparent to radiation. Only occasionally does a photon encounter and scatter off of a free electron or proton. The energy produced by nuclear reactions in the core travels outward toward the surface in the form of radiation with relative ease.

As we move outward from the core, the temperature falls, atoms collide less frequently and less violently, and more and more electrons manage to remain bound to their


## DISCOVERY 16-1

## Eavesdropping on the Sun


Throughout the few decades of the Space Age, various nations, led by the United States, have sent spacecraft to most of the major bodies in the solar system. One of the as-yet-unexplored bodies is Pluto, the most notable member of the Kuiper belt, which has never been visited by a robot orbiter or even a flyby craft—although this may change soon. ∞ (Sec. 14.3) The other unexplored body is the Sun. Currently, the next best things to a dedicated, close-up reconnoitering spacecraft are the *Solar and Heliospheric Observatory* (*SOHO*) and the *Solar Dynamics Observatory* (*SDO*). Both spacecraft have radioed back to Earth volumes of new data—and more than a few new puzzles—about our parent star.

*SOHO* is a billion-dollar mission operated primarily by the European Space Agency. Launched in 1995 and (as of early 2013) still operating, 18 years into its planned 3-year mission, this 2-ton robot is now on-station about 1.5 million kilometers sunward of Earth—about 1 percent of the distance from Earth to the Sun. This is the so-called  $L_1$  Lagrangian point, where the gravitational pull of the Sun and Earth are precisely equal—a good place to park a monitoring platform. In contrast, the U.S. *SDO* spacecraft, launched in 2010, travels around Earth in an inclined, geosynchronous orbit. Both automated vehicles stare at the Sun 24 hours a day and carry instruments capable of measuring almost everything from the Sun's corona and magnetic field to its solar wind and internal vibrations. The accompanying figure shows a false-color ultraviolet image of the Sun's lower corona, recently obtained by *SDO*.

Both spacecraft are positioned just beyond Earth's magnetosphere, so their instruments can study cleanly the high-speed charged particles of the solar wind. Coordinating these on-site measurements with *SOHO* and *SDO* images of the Sun itself, astronomers can study solar weather in great detail and in real time. The accumulated data are sufficiently good that mission scientists now think they can follow solar magnetic field loops expanding and breaking as the Sun prepares itself for mass ejections several days



(NASA)




before they actually occur (see Section 16.5). Given that such coronal storms can endanger pilots and astronauts and play havoc with communications, power grids, satellite electronics, and other human activity, the prospect of having accurate forecasts of disruptive solar events is a very welcome development.

By monitoring all aspects of the Sun, from surface oscillations to the details of its magnetic field structure, these spacecraft are steadily refining astronomers' models of solar structure, solar magnetism, and solar activity. These remarkable spacecraft have radioed back to Earth a wealth of new scientific information about our parent star. Since our detailed understanding of stars in general rests squarely on our knowledge of the Sun, *SOHO* and *SDO* are continually expanding the foundations of our study of the universe on every scale.

parent nuclei. With more and more atoms retaining electrons that can absorb the outgoing radiation, the gas in the interior changes from being relatively transparent to being almost totally opaque. By the outer edge of the radiation zone, roughly 500,000 km from the center (actually, 496,000 km, according to the best available *SOHO* data), *all* the photons produced in the Sun's core have been absorbed. Not one of them reaches the surface. But what happens to the energy they carry?

The photons' energy must travel beyond the Sun's interior: That we see sunlight—visible energy—proves that energy escapes. The escaping energy reaches the

surface by *convection*—the same basic physical process we saw in our study of Earth's atmosphere, although it operates in a very different environment in the Sun. ∞ (Sec. 7.2) Hot solar gas moves outward while cooler gas above it sinks, creating a characteristic pattern of convection cells. All through the convection zone, energy is transported to the surface by physical motion of the solar gas. (Note that this actually represents a departure from hydrostatic equilibrium, as defined above, but it can still be handled within the standard solar model.) Remember that there is no physical movement of material when radiation is the energy-transport mechanism; convection and radiation



◀ **FIGURE 16.7 Solar Convection** Energy is physically transported in the Sun's convection zone, which here is visualized as a boiling, seething sea of gas. As drawn, the convective cell sizes become progressively larger at greater depths. This is a highly simplified diagram; there are many different cell sizes, and they are not so neatly arranged.

are *fundamentally different* ways in which energy can be transported from one place to another.

Figure 16.7 is a schematic diagram of the solar convection zone. There is a hierarchy of convection cells, organized in tiers of many different sizes at different depths. The deepest tier, lying approximately 200,000 km below the photosphere, is thought to contain large cells some tens of thousands of kilometers in diameter. Heat is then successively carried upward through a series of progressively smaller cells, stacked one on another, until, at a depth of about 1000 km, the individual cells are about 1000 km across. The top of this uppermost tier of convection is the visible surface of the Sun, where astronomers can directly observe the cell sizes. Information about convection below that level is inferred mostly from computer models of the solar interior.


At some distance from the core, the solar gas becomes too thin to sustain further upwelling by convection. Theory suggests that this distance roughly coincides with the photospheric surface we see. Convection does not proceed into the solar atmosphere; there is simply not enough gas there—the density is so low that there are too few atoms or ions to intercept much sunlight, so the gas becomes transparent again and radiation once more becomes the mechanism of energy transport. Photons reaching the photosphere escape more or less freely into space, and the photosphere emits thermal radiation, like any other hot object. The photosphere is narrow, and the “edge” of the Sun sharp, because this transition from opacity to complete transparency is very rapid. Just below the bottom of the photosphere the gas is still convective, and radiation does not reach us directly. A few hundred kilometers higher, the gas is too thin to emit or absorb any significant amount of radiation.

## Granulation

Figure 16.8 is a high-resolution photograph of the solar surface. The visible surface is highly mottled, or **granulated**, with regions of bright and dark gas known as *granules*. Each

bright granule measures about 1000 km across—comparable in size to a continent on Earth—and has a lifetime of between 5 and 10 minutes. Together, several million granules constitute the top layer of the convection zone, immediately below the photosphere.

Each granule forms the topmost part of a solar convection cell. Spectroscopic observation within and around the bright regions shows direct evidence for the upward motion



▲ **FIGURE 16.8 Solar Granulation** This photograph of the granulated solar photosphere, taken with the 1-m Swedish Solar Telescope looking directly down on the Sun's surface, shows typical solar granules comparable in size to Earth's continents. The bright portions of the image are regions where hot material upwells from below, as illustrated in Figure 16.7. The darker (redder) regions correspond to cooler gas that is sinking back down into the interior. (SST/Royal Swedish Academy of Sciences)

of gas as it “boils” up from within—evidence that convection really does occur just below the photosphere. Spectral lines detected from the bright granules appear slightly bluer than normal, indicating Doppler-shifted matter approaching us at about 1 km/s. ∞ (Sec. 3.5) Spectroscopes focused on the darker portions of the granulated photosphere show the same spectral lines to be redshifted, indicating matter moving away from us.

The variations in brightness of the granules result strictly from differences in temperature. The upwelling gas is hotter and therefore emits more radiation than the cooler, downward-moving gas. The adjacent bright and dark gases appear to contrast considerably, but in reality their temperature difference is less than about 500 K. Careful measurements also reveal a much larger-scale flow on the solar surface. **Supergranulation** is a flow pattern quite similar to granulation, except that supergranulation cells measure some 30,000 km across. As with granulation, material upwells at the center of the cells, flows across the surface, then sinks down again at the edges. Scientists suspect that supergranules are the imprint on the photosphere of a deeper tier of large convective cells, like those depicted in Figure 16.7.

### CONCEPT Check

✓ What are the two distinct ways in which energy moves outward from the solar core to the photosphere?


## 16.3 The Sun's Atmosphere

Astronomers can glean an enormous amount of information about the Sun from an analysis of the absorption lines that arise in the photosphere and lower atmosphere. ∞ (Sec. 4.4) Figure 16.9 (see also Figure 4.4) is a detailed spectrum of the Sun spanning a range of wavelengths from 360 to 690 nm. Notice the intricate dark Fraunhofer absorption spectral lines superposed on the background continuous spectrum.

Tens of thousands of spectral lines have been observed and cataloged in the solar spectrum. In all, some 67 elements have been identified in the Sun in various states of ionization and excitation. ∞ (Sec. 4.2) More elements probably exist there, but they are present in such small quantities that our instruments are simply not sensitive enough to detect them. Table 16.2 lists the 10 most common elements in the Sun. Notice that hydrogen is by far the most abundant element, followed by helium. This distribution is just what we saw on the jovian planets, and it is what we will find for the universe as a whole.

### Solar Spectral Lines

As discussed in Chapter 4, spectral lines arise when electrons in atoms or ions make transitions between states of well-defined energies, emitting or absorbing photons of specific energies (i.e., wavelengths or colors) in the process. ∞ (Sec. 4.2) However, to explain the spectrum of the Sun (and, indeed, the spectra of all stars), we must slightly modify



◀ **FIGURE 16.9** Solar Spectrum

A detailed visible spectrum of our Sun shows thousands of dark Fraunhofer (absorption) spectral lines, indicating the presence of 67 different elements in various stages of excitation and ionization in the lower solar atmosphere. The numbers give wavelengths, in nanometers. (Palomar Observatory/ Caltech)

**TABLE 16.2** The Composition of the Sun

Element	Percentage of Total Number of Atoms	Percentage of Total Mass
Hydrogen	91.2	71.0
Helium	8.7	27.1
Oxygen	0.078	0.97
Carbon	0.043	0.40
Nitrogen	0.0088	0.096
Silicon	0.0045	0.099
Magnesium	0.0038	0.076
Neon	0.0035	0.058
Iron	0.0030	0.14
Sulfur	0.0015	0.040

our earlier description of the formation of absorption lines. We explained these lines in terms of cool foreground gas intercepting light from a hot background source. In actuality, both the bright background and the dark absorption lines in Figure 16.9 form at roughly the same locations in the Sun—the solar photosphere and lower chromosphere. To understand how these lines are formed, consider again the solar energy emission process in a little more detail.

Below the photosphere, the solar gas is sufficiently dense, and interactions among photons, electrons, and ions sufficiently common, that radiation cannot escape directly into space. In the solar atmosphere, however, the probability that a photon will escape without further interaction with matter depends on the photon's energy. Recall from Chapter 4 that an atom or ion can absorb a photon only if that photon's energy has just the right value to cause an electron to jump from one energy level to another.  $\infty$  (Sec. 4.3) Hence, if the photon energy happens to correspond to some electronic transition in an atom or ion in the gas, then the photon may be absorbed again before it can travel very far—the more elements present of the type suitable for absorption, the lower the escape probability. Conversely, if the photon's energy does not coincide with any such transition, then the photon cannot interact further with the gas, and it leaves the Sun headed for interstellar spaces, or perhaps the detector of an astronomer on Earth.


Thus, when we look at the Sun, we are actually peering down into the solar atmosphere to a depth that depends on the wavelength of the light under study. Photons with wavelengths far from any absorption feature (i.e., having energies far from any atomic transition) are less likely to interact with matter as they travel through the solar gas and so come from deeper in the photosphere. However, photons with wavelengths near the centers of absorption lines are much more likely to be captured by an atom or ion and therefore escape mainly from higher (and cooler) levels. The lines are darker than their surroundings because the temperature where they form is lower than the 5800-K temperature at the base of the photosphere, where most of the continuous emission originates. (Recall that, by

Stefan's law, the brightness of a radiating object depends on its temperature—the cooler the gas, the less energy it radiates.)  $\infty$  (Sec. 3.4) Thus, the existence of Fraunhofer lines is direct evidence that the temperature in the Sun's atmosphere decreases with height above the photosphere.



Strictly speaking, spectral analysis allows us to draw conclusions only about the part of the Sun where the lines form—the photosphere and chromosphere. However, most astronomers think that, with the exception of the solar core (where nuclear reactions are steadily changing the composition—see Sec. 16.6), the data in Table 16.2 are representative of the entire Sun. That assumption is strongly supported by the excellent agreement between the standard solar model, which makes the same assumption, and helioseismological observations of the solar interior.

## The Chromosphere

Above the photosphere lies the cooler chromosphere, the inner part of the solar atmosphere. This region emits very little light of its own and cannot be observed visually under normal conditions. The photosphere is just too bright, dominating the chromosphere's radiation. The relative dimness of the chromosphere results from its low density—large numbers of photons simply cannot be emitted by a tenuous gas containing very few atoms per unit volume. Still, although it is not normally seen, astronomers have long been aware of the chromosphere's existence. Figure 16.10 shows the Sun during an eclipse in which the photosphere—but not the chromosphere—is obscured by the Moon. The chromosphere's characteristic reddish hue is



**▲ FIGURE 16.10 Solar Chromosphere** This photograph of a total solar eclipse shows the solar chromosphere a few thousand kilometers above the Sun's surface. Note the prominence at left. (G. Schneider)



▲ **FIGURE 16.11 Solar Spicules** Short-lived, narrow jets of gas that typically last mere minutes can be seen sprouting up from the solar chromosphere in this ultraviolet image of the Sun. These so-called spicules are the thin spikelike regions whose gas escapes from the Sun at speeds of about 100 km/s. (NASA)

plainly visible. This coloration is due to the red H $\alpha$  (hydrogen alpha) emission line of hydrogen, which dominates the chromospheric spectrum. [∞ \(More Precisely 4-1\)](#)

The chromosphere is far from tranquil. Every few minutes, small solar storms erupt, expelling jets of hot matter known as *spicules* into the Sun's upper atmosphere (Figure 16.11). These long, thin spikes of matter leave the Sun's surface at typical speeds of about 100 km/s and reach several thousand kilometers above the photosphere. Spicules are not spread evenly across the solar surface. Instead, they cover only about 1 percent of the total area, tending to accumulate around the edges of supergranules. The Sun's magnetic field is also known to be somewhat stronger than average in those regions. Scientists speculate that the downward-moving material there tends to strengthen the solar magnetic field, and spicules are the result of magnetic disturbances in the Sun's churning outer layers.

## The Transition Zone and the Corona

During the brief moments of an eclipse, if the Moon's angular size is large enough that both the photosphere and the chromosphere are blocked, the ghostly solar corona can be seen (Figure 16.12). With the photospheric light removed, the pattern of spectral lines changes dramatically. The intensities of the usual lines alter (suggesting changes in composition or temperature, or both), the spectrum shifts from absorption to emission, and an entirely new set of spectral lines suddenly appears. The shift from absorption to emission is entirely in accordance with Kirchhoff's laws, because we see the corona against the blackness of space, not against the bright continuous spectrum from the photosphere below. [∞ \(Sec. 4.1\)](#)

These new coronal (and in some cases chromospheric) lines were first observed during eclipses in the 1920s. For years afterward, some researchers (for want of any better

explanation) attributed them to a new nonterrestrial element, which they dubbed “coronium.” We now recognize that these new spectral lines do not indicate any new kind of atom. Coronium does not exist. Rather, the new lines arise because atoms in the corona have lost several more electrons than atoms in the photosphere—that is, the coronal atoms are much more highly ionized. Therefore, their internal electronic structures, and hence their spectra, are quite different from the structure and spectra of atoms and ions in the photosphere. For example, astronomers have identified coronal lines corresponding to iron ions with as many as 13 of their normal 26 electrons missing. In the photosphere, most iron atoms have lost only 1 or 2 of their electrons.


The cause of this extensive electron stripping is the high coronal *temperature*. The degree of ionization inferred from spectra observed during solar eclipses tells us that the temperature of the upper chromosphere exceeds that of the photosphere. Furthermore, the temperature of the solar corona, where even more ionization is seen, is higher still. Figure 16.13 shows how the temperature of the Sun's atmosphere varies with altitude. The temperature decreases to a minimum of about 4500 K some 500 km above the photosphere, after which it rises steadily. About 1500 km above the photosphere, in the transition zone, the temperature begins to rise rapidly, reaching more than 1 million K at an altitude of 10,000 km. Thereafter, in the corona, the temperature remains roughly constant at around 3 million K, although *SOHO* and other orbiting instruments have detected coronal “hot spots” having temperatures many times higher than this average value.

The cause of the rapid temperature rise is not fully understood. The temperature profile runs contrary to intuition: Moving away from a heat source, we would normally expect the heat to diminish, but this is not the case in the lower atmosphere of the Sun. The corona must have another energy source. Astronomers now think that magnetic





▲ **FIGURE 16.12 Solar Corona** When both the photosphere and the chromosphere are obscured by the Moon during a solar eclipse, the faint and extended corona becomes visible. (National Solar Observatory)



▲ **FIGURE 16.13 Solar Atmospheric Temperature** The change of gas temperature in the lower solar atmosphere is dramatic. The temperature, indicated by the blue line, reaches a minimum of 4500 K in the chromosphere and then rises sharply in the transition zone, finally leveling off at around 3 million K in the corona.

disturbances in the solar photosphere are ultimately responsible for heating the corona (Section 16.5).

## The Solar Wind

Electromagnetic radiation and fast-moving particles—mostly protons and electrons—escape from the Sun all the time. The radiation moves away from the photosphere at the speed of light, taking 8 minutes to reach Earth. The particles travel more slowly, although at the still considerable speed of about 500 km/s, reaching Earth in a few days. This constant stream of escaping solar particles is the solar wind.

The solar wind results from the high temperature of the corona. About 10 million km above the photosphere, the coronal gas is hot enough to escape the Sun's gravity, and it begins to flow outward into space. At the same time, the solar atmosphere is continuously replenished from below. If that were not the case, the corona would disappear in about a day. The Sun is, in effect, “evaporating”—constantly shedding mass through the solar wind. The wind is an extremely thin medium, however. Even though it carries away roughly 2 million tons of solar matter each


second, less than 0.1 percent of the Sun's mass has been lost this way since the solar system formed 4.6 billion years ago.

### CONCEPT Check


- ✓ Describe two ways in which the spectrum of the solar corona differs from that of the photosphere.

## 16.4 Solar Magnetism


The Sun has a powerful and complex magnetic field. Discovered in 1908 by American astronomer George Ellery Hale, the solar field still presents puzzles to scientists today. The structure of the Sun's magnetic field lines is crucial to understanding many aspects of the Sun's appearance and surface activity, yet the details of the field-line geometry, and even the mechanism responsible for generating and sustaining the entire solar field, remain subjects of intense research. Curiously, the keys to understanding many aspects of solar magnetism lie in a phenomenon first observed nearly three centuries before Hale's groundbreaking discovery.



▲ **FIGURE 16.14 Sunspots** This photograph of the entire Sun, taken during a period of maximum solar activity, shows several groups of sunspots. The largest spots in the image are more than 20,000 km across, nearly twice the diameter of Earth. Typical sunspots are only about half that size. (Palomar Observatory/Caltech)



Sunspots appear dark because they are slightly cooler than the surrounding gas.



This spot is about the size of Earth.

## Sunspots

Figure 16.14 is an optical photograph of the entire Sun, showing numerous dark blemishes on its surface. First studied in detail by Galileo around 1613, these “spots” provided one of the first clues that the Sun was not a perfect, unvarying creation, but rather a place of constant change. ∞ (Sec. 2.4) The dark areas are called **sunspots** and typically measure about 10,000 km across, approximately the size of Earth. As shown in the figure, they often occur in groups. At any given time, the Sun may have hundreds of sunspots, or it may have none at all.

Studies of sunspots show an **umbra**, or dark center, surrounded by a grayish **penumbra**. The close-up views in Figure 16.15 show each of these dark areas and the brighter undisturbed photosphere nearby. This gradation in darkness is really a gradual change in photospheric temperature—sunspots are simply *cooler* regions of the photospheric gas. The temperature of the umbra is about 4500 K, compared with the penumbra’s 5500 K. The spots, then, are certainly composed of hot gases. They seem dark only because they appear against an even brighter background (the 5800 K photosphere). If we could magically remove a sunspot from the Sun (or just block out the rest of the Sun’s emission), the spot would glow brightly, just like any other hot object having a temperature of roughly 5000 K.



▲ **FIGURE 16.15 Sunspots, Up Close** (a) An enlarged photo of the largest pair of sunspots in Figure 16.14 shows how each spot consists of a cool, dark umbra surrounded by a warmer, brighter penumbra. (b) A high-resolution image of a single typical sunspot shows details of its structure as well as the surface granules surrounding it. (Palomar Observatory/Caltech; SST/Royal Swedish Academy of Sciences)

## The Sun’s Magnetic Field

What causes a sunspot? Why is it cooler than the surrounding photosphere? The answers to these questions are closely tied to the structure of the Sun’s magnetic field. We saw in Chapter 4 that analysis of spectral lines can yield detailed information about the magnetic field at the location where the lines originate. ∞ (Sec. 4.5) Indeed, Hale’s discovery of solar magnetism was made through observations of the Zeeman effect (broadening or splitting of spectral lines by a magnetic field) in H $\alpha$  lines observed in sunspots. Most importantly, both the *strength* of the magnetic field and the *orientation* of a field line along the line of sight (toward or away from the observer) can be determined.


The magnetic field in a typical sunspot is about 1000 times greater than the field in neighboring, undisturbed photospheric regions (which is itself several times stronger than Earth’s magnetic field). Furthermore, the field lines are not

► **FIGURE 16.16 Solar Magnetism** (a) The Sun’s magnetic field lines emerge from the surface through one member of a sunspot pair and reenter the Sun through the other member. If the magnetic field lines are directed into the Sun in one leading spot, they are inwardly directed in all other leading spots in that hemisphere. The opposite is the case in the southern hemisphere, where the polarities are always opposite those in the north. (b) An ultraviolet image taken by the *Transition Region and Coronal Explorer (TRACE)* satellite, showing magnetic field lines arching between two sunspot groups. (NASA)




Sunspots almost always come in pairs whose members lie at roughly the same latitude and have opposite magnetic polarities. Figure 16.16(a) illustrates how magnetic field lines emerge from the solar interior through one member (S) of a sunspot pair, loop through the solar atmosphere, and then reenter the photosphere through the other member (N). As in Earth’s magnetosphere, charged particles tend to follow the solar magnetic field lines. ∞ (Sec. 7.5) Figure 16.16(b) shows an actual image of solar magnetic loops, revealing high-temperature gas flowing along a complex network of magnetic field lines connecting two sunspot groups.

Despite the irregular appearance of the sunspots themselves, there is a great deal of order in the underlying solar field. *All* the sunspot pairs in the same solar hemisphere (north or south) at any instant have the *same* magnetic configuration. That is, if the leading spot (measured in the direction of the Sun’s rotation) of one pair has N polarity, as shown in the figure, then all leading spots in that hemisphere have the same polarity. What’s more, in the other hemisphere at the same time, all sunspot pairs have the *opposite* magnetic configuration (S polarity leading). To understand these regularities in sunspot polarities, we must look at the Sun’s magnetic field in a little more detail.



randomly oriented, but instead are directed roughly perpendicular to (out of or into) the Sun’s surface. Scientists think that sunspots are cooler than their surroundings because these abnormally strong fields tend to block (or redirect) the convective flow of hot gas, which is normally toward the surface of the Sun.

The **polarity** of a sunspot simply indicates which way its magnetic field is directed relative to the solar surface. We conventionally label spots where field lines emerge from the interior as “S” and those where the lines dive below the photosphere as “N” (so field lines above the surface always run from S to N, as on Earth).




**▲ FIGURE 16.17 Solar Rotation** (a, b) The Sun's differential rotation wraps and distorts the solar magnetic field. (c) Occasionally, the field lines burst out of the surface and loop through the lower atmosphere, thereby creating a sunspot pair. The underlying pattern of the solar field lines explains the observed pattern of sunspot polarities. (See Figure 16.21.)

The combination of differential rotation and convection radically affects the character of the Sun's magnetic field, which in turn plays a major role in determining the numbers and location of sunspots. As illustrated in Figure 16.17, the Sun's differential rotation distorts the solar magnetic field, “wrapping” it around the solar equator and eventually causing any originally north–south magnetic field to reorient itself in an east–west direction. At the same time, convection causes the magnetized gas to well up toward the surface, twisting and tangling the magnetic field pattern. In some places, the field lines become kinked like a twisted garden hose, causing the field strength to increase. Occasionally, the field becomes so strong that it overwhelms the Sun's gravity, and a “tube” of field lines bursts out of the surface and loops through the lower atmosphere, forming a sunspot pair. The general east–west orientation of the underlying solar field accounts for the observed polarities of the resulting sunspot pairs in each hemisphere.


## The Solar Cycle

Sunspots are not steady. Most change their size and shape, and all come and go. Figure 16.18 shows a time sequence in which a number of spots varied—sometimes growing, sometimes dissipating—over a period of several days. Individual spots may last anywhere from 1 to 100 days; a large group typically lasts 50 days. Not only do sunspots come and go with time, but their numbers and distribution across the face of the Sun also change fairly regularly. Centuries of observations have established a clear

**▼ FIGURE 16.18 Sunspot Rotation** This sequence, running left to right, shows the evolution of some sunspots and lower chromospheric activity over a period of 12 days. An  $H\alpha$  filter was used to make these photographs, taken from the *SkyLab* space station. An arrow follows one set of sunspots over the course of a week as they are carried around the Sun by its rotation. (NASA)



► **FIGURE 16.19 Sunspot Cycle** (a) Monthly number of sunspots during the 20th century clearly displays the (roughly) 11-year solar cycle. At the time of minimum solar activity, hardly any sunspots are seen. About 4 years later, at maximum solar activity, about 100 spots are observed per month. (b) Sunspots cluster at high latitudes when solar activity is at a minimum. They appear at lower and lower latitudes as the number of sunspots peaks. They are again prominent near the Sun's equator as solar minimum is approached once more.



**sunspot cycle.** Figure 16.19(a) shows the number of sunspots observed each year during the 20th century. The average number of spots reaches a maximum every 11 or so years and then falls off almost to zero before the cycle begins afresh.

The latitudes at which sunspots appear vary as the sunspot cycle progresses. Individual sunspots do not move up or down in latitude, but new spots appear closer to the equator as older ones at higher latitudes fade away. Figure 16.19(b) is a plot of observed sunspot latitude as a function of time. At the start of each cycle, at *solar minimum*, only a few spots are seen, and these are generally confined to two narrow zones about 25° to 30° north and south of the solar equator. Approximately four years into the cycle, around *solar maximum*, the number of spots has increased markedly, and they are found within about 15° to 20° of the equator. Finally, by the end of the cycle, at solar minimum, the number has fallen again, and most sunspots lie within about 10° of the solar equator. The beginning of each new cycle appears to overlap the end of the last.


Complicating this picture further, the 11-year sunspot cycle is actually only half of a longer 22-year **solar cycle**. During the first 11 years of the cycle, the leading spots of all the pairs in the northern hemisphere have the same

Blue lines indicate how the “average” sunspot latitude varies over the course of a cycle.

polarity, while spots in the southern hemisphere have the opposite polarity (Figure 16.16). These polarities then reverse their signs for the next 11 years, so the full solar cycle takes 22 years.

Astronomers think that the Sun’s magnetic field is both generated and amplified by the constant stretching, twisting, and folding of magnetic field lines that results from the combined effects of differential rotation and convection, although the details are still not well understood. The theory is similar to the “dynamo” theory that accounts for the magnetic fields of Earth and the jovian planets, except that the solar dynamo operates much faster and on a much larger scale. ∞ (Sec. 7.5) One prediction of this theory is that the Sun’s magnetic field should rise to a maximum, then fall to zero, and reverse itself, more or less periodically, just as is observed. Solar surface activity, such as the sunspot cycle, simply follows the variations in the magnetic field. The changing numbers of sunspots and their migration to lower latitudes are both consequences of the strengthening and eventual decay of the field lines as they become more and more tightly wrapped around the solar equator.

Figure 16.20 plots sunspot data extending back to the invention of the telescope. As can be seen, the 11-year “periodicity” of the solar sunspot cycle is far from regular.



▲ **FIGURE 16.20 Maunder Minimum** Average number of sunspots occurring each month over the past four centuries. Note the absence of spots during the late 17th century.

Not only does the period range from 7 to 15 years, but the sunspot cycle disappeared entirely over a number of years in the relatively recent past. The lengthy period of solar inactivity that extended from 1645 to 1715 is called the **Maunder minimum**, after the British astronomer who drew attention to these historical records. The corona was apparently also less prominent during total solar eclipses around that time, and Earth aurorae were sparse throughout the late 17th century. Lacking a complete understanding of the solar cycle, we cannot easily explain how it could shut down entirely. Most astronomers suspect changes in the Sun's convection zone or rotation pattern, but the specific causes of the Sun's century-long variations, as well as the details of the connection between solar activity and Earth's climate, remain a mystery (see *Discovery 16-2*).

In fact, the most recent sunspot minimum (in 2008–2009) has resulted in the least active Sun in almost a century; no sunspots at all were seen on almost 80 percent of the days during those years, and the solar wind was anomalously weak. The new cycle, due to peak in mid-2013, is also predicted to be much less active than normal. Scientists attribute this reduced activity to changes in the subsurface “conveyor belt” flow described in Section 16.2, which is thought to directly affect sunspot behavior. For unknown reasons, the flow sped up during the 1990s by a few meters per second, greatly suppressing the numbers of spots during the next cycle (and quite possibly the one after that, now underway). It has since slowed, but for now no one is sure how long the effects will last.

### CONCEPT Check

- ✓ **What do observations of sunspot polarities tell us about the solar magnetic field?**

## 16.5 The Active Sun


Most of the Sun's luminosity results from continuous emission from the photosphere. However, superimposed on this steady, predictable aspect of our star's energy output is a much more irregular component, characterized by explosive and unpredictable surface activity. Solar activity contributes little to the Sun's total luminosity and probably has no significant bearing on the evolution of the Sun, but it does affect us here on Earth. The size and duration of coronal holes are strongly influenced by the level of solar activity. Hence, so is the strength of the solar wind, and that in turn directly affects Earth's magnetosphere.

### Active Regions


The photosphere surrounding a pair or group of sunspots can be a violent place, sometimes erupting explosively, spewing forth large quantities of energetic particles into the corona. The sites of these energetic events are known as **active regions**. Most groups of sunspots have active regions associated with them. Like all other aspects of solar activity, these phenomena tend to follow the solar cycle and are most frequent and violent around the time of solar maximum.

Figure 16.21 shows two large solar **prominences**—loops or sheets of glowing gas ejected from active regions on the solar surface, moving through the inner parts of the corona under the influence of the Sun's magnetic field. Magnetic instabilities in the strong fields found in and near sunspot groups may cause the prominences, although the details are not fully understood. The arching magnetic field lines in and around the active region are also easily seen (see also Figure 16.16b). The rapidly changing structure of the field lines and the fact that they can quickly transport mass and energy from one part of the solar surface to another, possibly tens of thousands of kilometers away, make the theoretical study of active regions an extraordinarily difficult task.

*Quiescent prominences* persist for days or even weeks, hovering high above the photosphere, suspended by the Sun's magnetic field. *Active prominences* come and go much more erratically, changing their appearance in a matter of hours or surging up from the solar photosphere and then immediately falling back on themselves. A typical solar prominence measures some 100,000 km in extent, nearly 10 times the diameter of planet Earth. Prominences as large as the one shown in Figure 16.21(a) (which traversed almost half a million kilometers of the solar surface) are less common and usually appear only at times of greatest solar activity. The largest prominences can release up to  $10^{25}$  joules of energy, counting both particles and radiation—not much compared with the total solar luminosity of  $4 \times 10^{26}$  W, but still



(a)



(b)

**▲ FIGURE 16.21 Solar Prominences** (a) This particularly large solar prominence was observed by ultraviolet detectors aboard the *SOHO* spacecraft in 2002. (b) Like a phoenix rising from the solar surface, this filament of hot gas measures more than 100,000 km in length. Earth could easily fit between its outstretched “arms.” Dark regions in this *TRACE* image have temperatures less than 20,000 K; the brightest regions are about 1 million K. Most of the gas will subsequently cool and fall back into the photosphere. (NASA)


enormous by terrestrial standards. (All the power plants on Earth would take a billion years to produce that much energy.)

**Flares** are another type of solar activity observed low in the Sun’s atmosphere near active regions. Also the result of magnetic instabilities, flares, like that shown in Figure 16.22, are even more violent (and even less well understood) than prominences. They often flash across a region of the Sun in minutes, releasing enormous amounts of energy as they go. Space-based observations indicate that X-ray and ultraviolet emissions are especially intense in the extremely compact hearts of flares, where temperatures can reach 100 million K.

So energetic are these cataclysmic explosions that some researchers have likened flares to bombs exploding in the lower regions of the Sun’s atmosphere. A major flare can release as much energy as the largest prominences, but in a matter of minutes or hours rather than days or weeks. Unlike the gas that makes up the characteristic loop of a prominence, the particles produced by a flare are so energetic that the Sun’s magnetic field is unable to hold them and shepherd them back to the surface. Instead, the particles are simply blasted into space by the violence of the explosion. Flares are thought to be responsible for most of the internal pressure waves that give rise to solar surface oscillations.

Figure 16.23 shows a **coronal mass ejection** from the Sun. Sometimes (but not always) associated with flares and prominences, these phenomena are giant magnetic “bubbles” of ionized gas that separate from the rest of the solar atmosphere and escape into interplanetary space. Such ejections occur about once per week at times of sunspot minimum, but up to two or three times per day at solar maximum. Carrying an enormous amount of energy, they can—if their fields are properly oriented—merge with Earth’s magnetic field via a process known as *reconnection*, dumping some of their energy into the magnetosphere and potentially causing widespread communications and power disruptions on our planet (Figure 16.23b; see also *Discovery 16-2*).

▼ **FIGURE 16.22 Solar Flare** Much more violent than a prominence, a solar flare is an explosion on the Sun's surface that sweeps across an active region in a matter of minutes, accelerating solar material to high speeds and blasting it into space. (USAF)




► **FIGURE 16.23 Coronal Mass Ejection** (a) A few times per week, on average, a giant magnetized “bubble” of solar material detaches itself from the Sun and rapidly escapes into space, as shown in this *SOHO* image taken in 2002. The circles are artifacts of an imaging system designed to block out the light from the Sun itself and exaggerate faint features at larger radii. (b) Should a coronal mass ejection encounter Earth with its magnetic field oriented opposite to our own, as illustrated, the field lines can join together as in part (c), allowing high-energy particles to enter and possibly severely disrupt our planet's magnetosphere. By contrast, if the fields are oriented in the same direction, the coronal mass ejection can slide by Earth with little effect. (NASA/ESA)


## The Changing Solar Corona

Unlike the 5800 K photosphere, which emits most strongly in the visible part of the electromagnetic spectrum, the hot coronal gas radiates at much higher frequencies—primarily in the X-ray range. ∞ (Sec. 3.4) For this reason, X-ray telescopes have become important tools in the study of the solar corona. Figure 16.24(a) shows several X-ray images of the Sun. The full corona extends well beyond the regions shown, but the density of coronal particles emitting the radiation diminishes rapidly with distance from the Sun. The intensity of X-ray radiation farther out is too dim to be seen here.


In the mid-1970s, instruments aboard NASA's *Skylab* space station revealed that the solar wind escapes mostly through solar “windows” called **coronal holes**. The dark area moving from left to right in Figure 16.24(a), which shows more recent data from the Japanese *Yohkoh* X-ray solar observatory, represents a coronal hole. Not really



(a)




(b)



(c)





**▲ FIGURE 16.24 Coronal Hole** (a) Images of X-ray emission from the Sun observed by the *Yohkoh* satellite. Note the dark, V-shaped coronal hole traveling from left to right, where the X-ray observations outline in dramatic detail the abnormally thin regions through which the high-speed solar wind streams forth. (b) Charged particles follow magnetic field lines (blue curves) that compete with gravity. When the field is trapped and loops back toward the photosphere, the particles are also trapped; otherwise, they can escape as part of the solar wind. (ISAS/Lockheed Martin)

holes, such structures are simply deficient in matter—vast regions of the Sun’s atmosphere where the density is about 10 times lower than the already tenuous, normal corona. Note that the underlying solar photosphere looks black in these images because it is far too cool to emit X-rays in any significant quantity.

Coronal holes are lacking in matter because the gas there is able to stream freely into space at high speeds, driven by disturbances the Sun’s atmosphere and magnetic field. Figure 16.24(b) illustrates how, in coronal holes, the solar magnetic field lines extend from the surface far out into interplanetary space. Because charged particles tend to

follow the field lines, they can escape, particularly from the Sun’s polar regions, according to findings from *SOHO* and NASA’s *Ulysses* spacecraft, which flew high above the ecliptic plane to explore the Sun’s polar regions. The speed of the solar wind can reach 800 km/s over some coronal holes. In other regions of the corona, the solar magnetic field lines stay close to the Sun, keeping charged particles near the surface and inhibiting the outward flow of the solar wind (just as Earth’s magnetic field tends to prevent the incoming solar wind from striking Earth), so the density remains relatively high. Because of the “open” field structure in coronal holes, flares and other magnetic activity (which, as we have seen, are related to magnetic loops near the solar photosphere) tend to be suppressed there.

The largest coronal holes, like that shown in Figure 16.24(a), can be hundreds of thousands of kilometers across and may survive for many months. Structures of this size are seen only a few times per decade. Smaller holes—perhaps only a few tens of thousand kilometers in size—are much more common, appearing every few hours.

Coronal holes appear to be an integral part of the process by which the Sun’s large-scale field reverses and replenishes itself over the course of the solar cycle. Long-lived holes persist at the Sun’s polar regions over much of the magnetic cycle, and the numbers and locations of other holes appear to change in step with solar activity. However, like many aspects of the solar magnetic field, the structure and evolution of coronal holes are not fully understood; they are currently the subject of intense research.

Finally, the solar corona varies with the sunspot cycle. The photograph of the corona in Figure 16.12 shows the quiet Sun, at sunspot minimum. At such times, the corona is fairly regular in appearance and seems to surround the Sun more or less uniformly. Compare that image with Figure 16.25, which was taken in 1994 near a peak in the sunspot cycle. The active corona is much more irregular in

## DISCOVERY 16-2

### Solar–Terrestrial Relations

Our Sun has often been worshipped as a god with power over human destinies. Obviously, the steady stream of solar energy arriving at our planet every day is essential to our lives, but over the past century there have also been repeated claims of a correlation between the Sun’s activity and Earth’s weather. Only recently, however, has the subject become scientifically respectable—that is, more natural than supernatural.

In fact, there do seem to be some correlations between the 22-year solar cycle (two sunspot cycles with oppositely directed magnetic fields) and periods of climatic dryness here on Earth. For example, near the start of the past eight cycles, there have been droughts in North America—at least within the middle and western plains from South Dakota to New Mexico. Another possible Sun–Earth connection is a link between solar activity and increased atmospheric circulation on our planet. As circulation increases, terrestrial storm systems deepen, extend over wider ranges of latitude, and carry more moisture. The relationship is complex and the subject controversial, because no one has yet shown any physical mechanism (other than the Sun’s heat, which does not vary much during the solar cycle) that would allow solar activity to stir our terrestrial atmosphere. Without a better understanding of the physical mechanism involved, none of these effects can be incorporated into our weather-forecasting models.

Solar activity may also influence long-term climate on Earth. For example, the Maunder minimum (see Section 16.6) seems to correspond fairly well with the coldest years of the so-called Little Ice Age that chilled northern Europe and North America during the late 1600s. The accompanying “winter” scene actually captured one summer season in 17th-century Holland. How the active Sun and its abundance of sunspots may affect Earth’s climate is a frontier problem in terrestrial climatology.

Measurements of the solar constant made over the past two

decades indicate that the Sun’s energy output varies with the solar cycle. Paradoxically, the Sun’s luminosity is greatest when many dark sunspots cover its surface! Thus, the Maunder minimum does correspond to an extended period of lower-than-average solar emission. However, recent observed changes in the Sun’s luminosity have been small—no more than 0.2 or 0.3 percent. It is not known by how much, if at all, the Sun’s output declined during the Maunder minimum, nor how large a change would be needed to account for the alterations in climate that occurred.

One correlation that is definitely established, and also better understood, is that between solar activity and geomagnetic disturbances at Earth. The extra radiation and particles thrown off by flares or coronal mass ejections impinge on Earth’s environment, overloading the Van Allen belts, thereby causing brilliant auroras in our atmosphere and degrading our communication networks. We are only beginning to understand how the radiation and particles emitted by solar phenomena also interfere with terrestrial radars, power networks, and other technological equipment. Some power outages on Earth are actually caused, not by increased customer demand or malfunctioning equipment, but by weather on the Sun!

We cannot yet predict just when and where solar flares or coronal mass ejections will occur. However, it would certainly be to our advantage to be able to do so, as that aspect of the active Sun affects our lives. This is a highly fertile area of astronomical research and one with clear terrestrial applications.



(Rijksmuseum, Amsterdam, Holland/The Bridgman Art Library)


appearance and extends farther from the solar surface. The “streamers” of coronal material pointing away from the Sun are characteristic of this phase.

Astronomers think that the corona is heated primarily by activity in the photosphere, which can inject large amounts of energy into the lower solar atmosphere. The myriad spicules and small-scale magnetic disturbances in the photosphere probably provide most of the energy needed to heat the corona. More extensive disturbances

often move through the corona above an active site in the photosphere, distributing the energy throughout the coronal gas. Given this connection, it is hardly surprising that both the appearance of the corona and the strength of the solar wind are closely correlated with the solar cycle.

### CONCEPT Check

✓ Why is solar activity important to life on Earth?



▲ **FIGURE 16.25 Active Corona** Photograph of the solar corona during the July 1994 eclipse, near the peak of the sunspot cycle. At these times, the corona is much less regular and much more extended than at sunspot minimum (compare to Figure 16.12). The changing shape and size of the corona is the direct result of variations in prominence and flare activity over the course of the solar cycle. (NCAR High Altitude Observatory)

## 16.6 The Heart of the Sun

What powers the Sun? What forces are at work in the Sun's core to produce its enormous luminosity? By what process does the Sun shine, day after day, year after year, eon after eon? Answers to these questions are central to all astronomy. Without them, we can understand neither the physical existence of stars and galaxies in the universe nor the biological existence of life on Earth.

### Solar Energy Production

In round numbers, the Sun's luminosity is  $4 \times 10^{26}$  W and its mass is  $2 \times 10^{30}$  kg. We can quantify how efficiently the Sun generates energy by dividing the solar luminosity by the solar mass:

$$\frac{\text{solar luminosity}}{\text{solar mass}} = 2 \times 10^{-4} \text{ W/kg.}$$

This simply means that, on average, every kilogram of solar material yields about 0.2 milliwatt of energy—0.0002 joule (J) of energy every second.


This is not much energy—a piece of burning wood generates about a million times more energy per unit mass per unit time than does our Sun, so the equivalent solar luminosity could (in principle) be created by a pile of burning logs comparable in mass to planet Earth. But there is one very important difference: The logs cannot continue to burn at this rate for billions of years.

To appreciate the magnitude of the energy generated by our Sun, we must consider not the ratio of the solar luminosity to the solar mass, but instead the total amount of energy generated by each gram of solar matter *over the entire lifetime of the Sun as a star*. This is easy to do. We simply multiply the rate at which the Sun generates energy by the age of the Sun, about 5 billion years. We obtain a value of  $3 \times 10^{13}$  J/kg. This is the average amount of energy radiated by every kilogram of solar material since the Sun formed. It represents a *minimum* value for the total energy radiated by the Sun, for more energy will be needed for every additional day the Sun shines. Should the Sun endure for another 5 billion years (as is predicted by theory), we would have to double this value.

Either way, this energy-to-mass ratio is very large. At least 60 trillion joules (on average) of energy must arise from every kilogram of solar matter to power the Sun throughout its lifetime. But the Sun's generation of energy is not explosive, releasing large amounts of energy in a short period. Instead, it is slow and steady, providing a uniform and long-lived rate of energy production. Only one known energy-generation mechanism can conceivably power the Sun in this way: **nuclear fusion**—the combining of light nuclei into heavier ones.

### Nuclear Fusion

We can represent a typical fusion reaction symbolically as



For powering the Sun and other stars, the most important piece of this equation is the energy produced.

The essential point here is that, during a fusion reaction, the total mass *decreases*—the mass of nucleus 3 is less than the combined masses of nuclei 1 and 2. Where does this mass go? It is converted into energy in accordance with Einstein's famous equation of **mass-energy equivalence**

$$E = mc^2,$$

or


$$\text{energy} = \text{mass} \times (\text{speed of light})^2.$$

This equation expresses the discovery, made by Albert Einstein at the beginning of the 20th century, that matter and energy are interchangeable—one can be converted into the other. To determine the amount of energy corresponding

to a given mass, simply multiply the mass by the square of the speed of light ( $c$  in the equation). For example, the energy equivalent of 1 kg of matter is  $1 \times (3 \times 10^8)^2$ , or  $9 \times 10^{16}$  J. The speed of light is so large that even small amounts of mass translate into enormous amounts of energy.

The production of energy by a nuclear fusion reaction is an example of the **law of conservation of mass and energy**, which states that the *sum* of mass and energy (suitably converted into the same units, using Einstein's equation) must always remain *constant* during any physical process. *There are no known exceptions.* According to this law, an object can literally disappear, provided that some energy appears in its place. If magicians really made rabbits disappear, the result would be a flash of energy equaling the product of the rabbit's mass and the square of the speed of light—enough to destroy the magician, everyone in the audience, and probably all of the surrounding state as well! In the case of fusion reactions in the solar core, the energy is produced primarily in the form of electromagnetic radiation. The light we see coming from the Sun means that the Sun's mass must be slowly, but steadily, decreasing with time.

## Charged Particle Interactions

All atomic nuclei are positively charged, so they repel one another. Furthermore, by the inverse-square law, the closer two nuclei come to one another, the greater is the repulsive force between them (Figure 16.26a).  (Sec. 3.2) How, then, do nuclei—two protons, say—ever manage to fuse into anything heavier? The answer is that if they collide at high enough speeds, one proton can momentarily plow deep into the other, eventually coming within the exceedingly short range of the **strong nuclear force**, which binds nuclei together (see *More Precisely 16-1*). At distances less than about  $10^{-15}$  m, the attraction of the nuclear force overwhelms the electromagnetic repulsion, and fusion occurs. Speeds in excess of a few hundred kilometers per second, corresponding to a gas temperature of  $10^7$  K or more, are needed to slam protons together fast enough to initiate fusion. Such conditions are found in the core of the Sun and at the centers of all stars.


The fusion of two protons is illustrated schematically in Figure 16.26(b). In effect, one of the protons turns into a neutron, creating new particles in the process, and combines with the other proton to form a **deuteron**, the nucleus of a special form of hydrogen called *deuterium*. Deuterium (also referred to as “heavy hydrogen”) differs from ordinary hydrogen by virtue of an extra neutron in its nucleus. We can represent the reaction as follows:



The **positron** in this reaction is a positively charged electron. Its properties are identical to those of a normal,

negatively charged electron, except for the positive charge. Scientists call the electron and the positron a “matter–anti-matter pair”—the positron is said to be the *antiparticle* of the electron. The newly created positrons find themselves in the midst of a sea of electrons with which they interact immediately and violently. The particles and antiparticles annihilate (destroy) one another, producing pure energy in the form of gamma-ray photons.

The final product of the reaction is a particle known as a **neutrino**, a word derived from the Italian for “little neutral one.” Neutrinos carry no electrical charge and are of very low



**▲ FIGURE 16.26 Proton Interactions** (a) Since like charges repel, two low-speed protons veer away from one another, never coming close enough for fusion to occur. (b) Higher-speed protons can overcome their mutual repulsion, approaching close enough for the strong force to bind them together—in which case they collide violently, triggering nuclear fusion that ultimately powers the Sun.

mass—at most 1/100,000 the mass of an electron, which itself has only 1/2000 the mass of a proton. (The exact mass of the neutrino remains uncertain, although experimental evidence strongly suggests that it is not zero.) Neutrinos move at almost the speed of light and interact with hardly anything. They can penetrate, without stopping, several light-years of lead (a very dense material, widely used in terrestrial laboratories as an effective shield against radiation). Their interactions with matter are governed by the **weak nuclear force**, described in *More Precisely 16-1*. Despite their elusiveness, neutrinos can be detected with carefully constructed instruments. In the final section of this chapter we discuss some rudimentary neutrino “telescopes” and the important contribution they have made to solar astronomy.


Nuclei such as normal hydrogen and deuterium, containing the same number of protons, but different numbers of neutrons, represent different forms of the same element—they are known as **isotopes** of that element. Usually, there are about as many neutrons in a nucleus as protons, but the exact number of neutrons can vary, and most elements exist in a number of isotopic forms. To

avoid confusion when talking about isotopes of the same element, nuclear physicists attach a number to the symbol representing the element. This number indicates the total number of particles (protons plus neutrons) in the nucleus of an atom of the element. Thus, ordinary hydrogen is denoted by  ${}^1\text{H}$ , deuterium by  ${}^2\text{H}$ . Normal helium (two protons plus two neutrons) is  ${}^4\text{He}$  (also referred to as helium-4), and so on. We will adopt this convention for the rest of the book.

## The Proton–Proton Chain

The basic set of nuclear reactions powering the Sun (and the vast majority of all stars) is sketched in Figure 16.27. It is not a single reaction, but rather a sequence called the **proton–proton chain**. The following stages are shown in the figure.

- I. First, two protons combine to form deuterium, as in Figure 16.26.
- II. The resulting positrons annihilate with electrons, releasing energy in the form of gamma rays. The



**MA** **Interactive FIGURE 16.27 Solar Fusion** In the proton–proton chain, a total of six protons (and two electrons) are converted into two protons, one helium-4 nucleus, and two neutrinos. The two leftover protons are available as fuel for new proton–proton reactions, so the net effect is that four protons are fused to form one helium-4 nucleus. Energy, in the form of gamma rays, is produced at each stage. (Most of the photons are omitted for clarity.) The three stages indicated here correspond to reactions (I), (II), and (III) described in the text.

## MORE PRECISELY 16-1

### Fundamental Forces

Our studies of nuclear reactions have uncovered new ways in which matter can interact with matter at the subatomic level. Let's pause to consider in a slightly more systematic fashion the relationships among the various forces of nature.

As best we can tell, the behavior of all matter in the universe—from elementary particles to clusters of galaxies—is ruled by just four (or fewer) basic forces, which are *fundamental* to everything in the universe. In a sense, the search to understand the nature of the universe is the quest to understand the nature of these forces.

The **gravitational force** is probably the best known of the four. Gravity binds galaxies, stars, and planets together and holds humans on the surface of Earth. As we saw in Chapter 2, its magnitude decreases with distance according to an inverse-square law. [☞ \(Sec. 2.7\)](#) Its strength is also proportional to the masses of each of the two objects involved. Thus, the gravitational field of an atom is extremely weak, but that of a galaxy, consisting of huge numbers of atoms, is very powerful. Gravity is by far the weakest of the forces of nature, but its effect accumulates as we move to larger and larger volumes of space, and nothing can cancel its attractive pull. As a result, gravity is the dominant force in the universe on all scales larger than that of Earth.

The **electromagnetic force** is another of nature's basic agents. Any particle having a net electric charge, such as an electron or a proton in an atom, exerts an electromagnetic force on any other charged particle. The everyday things we see around us are held together by this force. As with gravity, the strength of the electromagnetic force decreases with distance according to an inverse-square law. [☞ \(Sec. 3.2\)](#) However, for subatomic particles, electromagnetism is much stronger than gravity. For example, the electromagnetic force between two protons exceeds their gravitational attraction by a factor of about  $10^{36}$ . Unlike gravity, electromagnetic forces can repel (like charges) as well as attract (opposite charges). Positive and negative charges tend to neutralize each other, greatly diminishing their net electromagnetic influence. Above the microscopic level, most objects are in fact very close to being electrically neutral. Thus, except in unusual circumstances, the electromagnetic force is relatively unimportant on macroscopic scales.

A third fundamental force of nature is simply termed the **weak nuclear force**. This force is much weaker than electromagnetism, and its influence is somewhat more subtle. The

weak nuclear force governs the emission of radiation from some radioactive atoms; the emission of a neutrino during the first stage of the proton–proton reaction (Figure 16.26) is also the result of a weak interaction. The weak nuclear force does not obey the inverse-square law. Its effective range is much less than the size of an atomic nucleus—about  $10^{-18}$  m.

It is now known that electromagnetism and the weak force are not really separate forces at all, but rather two different aspects of a more basic **electroweak force**. At “low” temperatures, such as those found on Earth or even in stars, the electromagnetic and weak forces have quite distinct properties. However, as we will see in Chapter 27, at very high temperatures, such as those that prevailed in the universe when it was much less than a second old, the two are indistinguishable. Under those conditions, electromagnetism and the weak force are said to be “unified” into the electroweak force, and the universe has only three fundamental forces, rather than four.

Strongest of all the forces is the **strong nuclear force**. This force binds atomic nuclei and subnuclear particles (e.g., protons and neutrons) together and governs the generation of energy in the Sun and all other stars. Like the weak force, and unlike the forces of gravity and electromagnetism, the strong force operates only at very close range. It is unimportant outside a distance of a hundredth of a millionth of a millionth ( $10^{-15}$ ) of a meter. However, within that range (e.g., in atomic nuclei), it binds particles with enormous strength. In fact, it is the range of the strong force that determines the typical sizes of atomic nuclei. Only when two protons are brought within about  $10^{-15}$  m of one another can the attractive strong force overcome their electromagnetic repulsion. High-energy accelerator experiments suggest that, at very close quarters (less than  $10^{-16}$  m), the strong force has a “hard” core where the attraction turns into repulsion. (This scale is too small to affect atomic nuclei, but it may be crucial in determining the physics of supernovae explosions—see Section 21.2.)

Very loosely speaking, we can say that the strong force is about 100 times stronger than electromagnetism, 1 million times stronger than the weak force, and  $10^{38}$  times stronger than gravity. But not all particles are subject to all types of force. All particles interact through gravity because all have mass. However, only charged particles interact electromagnetically. Protons and neutrons are affected by the strong nuclear force, but electrons are not. Finally, under the right circumstances, the weak force can affect any type of subatomic particle, regardless of its charge.


deuterons combine with protons to create an isotope of helium called helium-3 (containing only one neutron), releasing additional energy, again in the form of gamma-ray photons. Two of each of these sets of reactions are shown in Figure 16.27.

III. Finally, two helium-3 nuclei combine to produce helium-4, two protons, and still more gamma-ray energy.


Gargantuan quantities of protons are fused into helium by the proton–proton chain in the core of the Sun each and every second. The energy released ultimately becomes the sunlight that warms our planet.

Setting aside the temporary intermediate nuclei produced, we see that the *net* effect of the proton–proton chain is that four hydrogen nuclei (six protons consumed, two

returned) combine to create one nucleus of the next lightest element, helium-4 (containing two protons and two neutrons, for a total mass of four), creating two neutrinos, and releasing energy in the form of gamma rays:  $\infty$  (Sec. 4.2)



or



As discussed in more detail in *More Precisely 16-2*, to fuel the Sun's present energy output, hydrogen must be fused into helium in the core at a rate of 600 million tons per second—a lot of mass, but only a tiny fraction of the total amount available. The Sun will be able to sustain this rate of core burning for about another 5 billion years (see Chapter 20).

The Sun's nuclear energy is produced in the core in the form of gamma rays. However, as it passes through the cooler layers of the solar interior and photons are absorbed and reemitted, the radiation's blackbody spectrum steadily shifts toward lower and lower temperatures and, by Wien's law, the characteristic wavelength of the radiation increases.  $\infty$  (Sec. 2.4) The energy eventually leaves the photosphere mainly in the form of visible and infrared radiation. A comparable amount of energy is carried off by the neutrinos, which escape unhindered into space at almost the speed of light.

Other reaction sequences can produce the same end result as the proton–proton chain (see *More Precisely 20-1*). However, the sequence shown in Figure 16.27 is the simplest and produces almost 90 percent of the Sun's luminosity. Note how, at each stage of the chain, more massive, complex nuclei are created from simpler, lighter ones. We will see in Chapters 20 and 21 that not all stars are powered by hydrogen fusion. Nevertheless, *nuclear fusion*—the slow, but steady, transformation of light elements into heavier ones, creating energy in the process—is responsible for virtually *all* of the starlight we see.

### CONCEPT Check

✓ Why does the fact that we see sunlight imply that the Sun's mass is slowly decreasing?

## 16.7 Observations of Solar Neutrinos


Theorists are quite sure that the proton–proton chain operates in the core of the Sun. However, because the gamma-ray energy created in the proton–proton chain is transformed into visible and infrared radiation by the time it emerges from the Sun, astronomers have no direct electromagnetic evidence of the core nuclear reactions.

Instead, the *neutrinos* created in the proton–proton chain are our best bet for learning about conditions in the solar core. They travel cleanly out of the Sun, interacting with virtually nothing, and escape into space a few seconds after being created. Of course, the fact that they can pass through the entire Sun without interacting also makes neutrinos difficult to detect on Earth! Nevertheless, with knowledge of neutrino physics it is possible to construct neutrino detectors.

Over the past four decades, several experiments have been designed to detect solar neutrinos reaching Earth. Some detectors use large quantities of the elements chlorine or gallium, which happen to be slightly more likely than most to interact with neutrinos. The interactions turn chlorine nuclei into argon or gallium into germanium. The new nuclei are radioactive, and the detection of the radiation from their decay signals the neutrino capture. Other detectors (two of which are shown in Figure 16.28) look for light produced when a high-energy neutrino occasionally collides with an electron in a water molecule, accelerating the electron to almost the speed of light. As the high-speed electron moves through the water, it emits electromagnetic radiation, mainly in the ultraviolet part of the spectrum. At visible wavelengths, the water appears blue. Large photomultiplier tubes (light-amplification devices) detect the resultant faint glow that betrays the neutrino's passage. In all cases, the probability of a given neutrino interacting with matter in the detector is extraordinarily small—only 1 in  $10^{15}$  of the neutrinos passing through the apparatus is actually detected. Large amounts (tons) of target material and long-duration experiments (months or years) are needed to obtain accurate measurements.

The detectors' designs differ widely, they are sensitive to neutrinos of very different energies, and they disagree somewhat in the details of their results, but they all agree on one very important point: Although solar neutrinos are observed (and in fact have measured energies in the range predicted by the standard solar model), there is a real difference between the Sun's theoretical neutrino output and the number of neutrinos actually detected on Earth. The number of solar neutrinos reaching Earth is substantially less (by 50 to 70 percent) than the prediction of the standard solar model. This discrepancy is known as the **solar neutrino problem**.

How can we explain this clear disagreement between theory and observation? The broad agreement among several independent, well-designed, and thoroughly tested experiments implies to most scientists that experimental error is not the cause, and researchers are confident that the experimental results can be trusted. Indeed, the lead investigators of the two experiments shown in Figure 16.28 received the strongest possible scientific endorsement of their work in the form of the 2002 Nobel Prize.



(a)



(b)

In that case, there are really only two possibilities: Either *solar neutrinos are not produced as frequently as we think, or not all of them make it to Earth.*

It is very unlikely that the resolution of the solar neutrino problem lies in the physics of the Sun's interior. For example, we might think of reducing the theoretical number of neutrinos by postulating a lower temperature in the solar core, but the nuclear reactions described in

◀ **FIGURE 16.28 Neutrino Telescopes** (a) This swimming pool-sized “neutrino telescope” is buried beneath a mountain near Tokyo, Japan. Called Super Kamiokande, it is filled with 50,000 tons of purified water, and contains 13,000 individual light detectors (some shown here being inspected by technicians in a rubber raft) to sense the telltale signature—a brief burst of light—of a neutrino passing through the apparatus. (b) The Sudbury Neutrino Observatory (SNO), situated 2 km underground in Ontario, Canada. The SNO detector is similar in design to the Kamiokande device, but by using “heavy” water (with hydrogen replaced by deuterium) instead of ordinary water, and adding 2 tons of salt, it also becomes sensitive to other neutrino types. The device contains 10,000 light-sensitive detectors arranged on the inside of the large sphere shown here. (ICRR, SNO; LBNL)

the previous section are just too well known, and the agreement between the standard solar model and helioseismological observations (Section 16.2) is far too close for conditions in the core to deviate much from the predictions of the model.

Instead, the answer involves the properties of the neutrinos themselves and has caused scientists to rethink some very fundamental concepts in particle physics. If neutrinos have even a minute amount of mass, theory indicates that it should be possible for them to change their properties—even to transform into other particles—during their 8-minute flight from the solar core to Earth through a process known as **neutrino oscillations**. In this picture, neutrinos are produced in the Sun at the rate required by the standard solar model, but some turn into something else—actually, other types of neutrinos—on their way to Earth and hence go undetected in the experiments just described. (In the jargon of the field, the neutrinos are said to “oscillate” into other particles.)

In 1998 the Japanese group operating the Super Kamiokande detector shown in Figure 16.28(a) reported the first experimental evidence of neutrino oscillations (and hence of nonzero neutrino masses), although the observed oscillations did not involve neutrinos of the type produced in the Sun. Then, in 2001, measurements made at the Sudbury Neutrino Observatory (SNO) in Ontario, Canada (Figure 16.28b), revealed strong evidence for the “other” neutrinos into which the Sun's neutrinos have been transformed. Subsequent SNO observations with a modified detector confirmed the result. The total numbers of neutrinos observed were completely consistent with the standard solar model. The solar neutrino problem was solved, the scientific method proved itself again—and neutrino astronomy claimed its first major triumph!

### PROCESS OF SCIENCE Check




- ✓ Using the solar neutrino problem as an example, discuss how scientific theory and observation evolve and adapt when they come into conflict.



## MORE PRECISELY 16-2


### Energy Generation in the Proton-Proton Chain

Let's look in a little more detail at the energy produced by fusion in the solar core and compare it with the energy needed to account for the Sun's luminosity. Using the notation presented in the text, the proton-proton chain may be compactly described by the following reactions:



As discussed in the text and illustrated below, the net effect of the fusion process is that four protons combine to produce a nucleus of helium-4, in the process creating two neutrinos and two positrons (which are quickly converted into energy by annihilation with electrons).

We can calculate the total amount of energy released by accounting carefully for the total masses of the nuclei involved and applying Einstein's famous formula  $E = mc^2$ .



This in turn allows us to relate the Sun's total luminosity to the consumption of hydrogen fuel in the core. Careful laboratory experiments have determined the masses of all the particles involved in the above reaction: The total mass of the protons is  $6.6943 \times 10^{-27}$  kg, the mass of the helium-4 nucleus is  $6.6466 \times 10^{-27}$  kg, and the neutrinos are virtually massless. We omit the positrons here—their masses will end up being counted as part of the total energy released. The difference between the total mass of the four protons and that of the final helium-4 nucleus,  $0.0477 \times 10^{-27}$  kg, is not great, but it is easily measurable.

Multiplying the vanished mass by the square of the speed of light yields  $0.0477 \times 10^{-27} \text{ kg} \times (3.00 \times 10^8 \text{ m/s})^2 = 4.28 \times 10^{-12} \text{ J}$ . This is the energy produced in the form of radiation when  $6.69 \times 10^{-27}$  kg (the rounded-off mass of the four protons) of hydrogen fuses to helium. It follows that fusion of 1 kg of hydrogen generates  $4.28 \times 10^{-12} / 6.69 \times 10^{-27} = 6.40 \times 10^{14} \text{ J}$ . Put another way, the process converts about 0.71 percent of the original mass into energy. A negligible fraction (actually, about 2 percent) of this energy is carried away by the neutrinos. The rest appears in the form of gamma rays and ultimately is radiated away from the solar photosphere—that is, it becomes the solar luminosity.


Thus we have established a direct connection between the Sun's energy output and the consumption of hydrogen in the core. The Sun's luminosity of  $3.84 \times 10^{26} \text{ W}$  (see Table 16.1), or  $3.84 \times 10^{26} \text{ J/s}$  (joules per second), implies a mass consumption rate of  $3.84 \times 10^{26} \text{ J/s} / 6.40 \times 10^{14} \text{ J/kg} = 6.00 \times 10^{11} \text{ kg/s}$ —600 million tons of hydrogen every second (1 ton = 1000 kg). A mass of 600 million tons sounds like a lot—the mass of a small mountain—but it represents only a few million millionths of the total mass of the Sun. Put another way, of this 600 million tons, roughly  $600 \text{ million tons/s} \times 0.0071 = 4.3 \text{ million tons per second}$  of solar matter is converted into radiation—comparable to the mass carried away by the solar wind. Our parent star will be able to sustain this loss rate for a very long time.

**The Big Question** How does the Sun's corona get so hot? What causes the 11-year solar cycle of activity? Why do sunspots exist and why do they look so messy? There is not just one big question regarding the Sun, but many smaller, nagging ones that have perplexed scientists for decades. Although we understand the basic physics of how our star shines, we still have much to learn about its unpredictable behavior that can sometimes affect life here on Earth.


# Chapter Review

## SUMMARY


1 Our Sun is a **star** (p. 390), a glowing ball of gas held together by its own gravity and powered by nuclear fusion at its center. The **photosphere** (p. 390) is the region at the Sun's surface from which virtually all the visible light is emitted. The main interior regions of the Sun are the **core** (p. 390), where nuclear reactions generate energy; the **radiation zone** (p. 390), where the energy travels outward in the form of electromagnetic radiation; and the **convection zone** (p. 390), where the Sun's matter is in constant convective motion.



2 The amount of solar energy reaching a  $1 \text{ m}^2$  at the top of Earth's atmosphere each second is a quantity known as the **solar constant** (p. 390). The Sun's **luminosity** (p. 391) is the total amount of energy radiated from the solar surface per second. It is determined by multiplying the solar constant by the area of an imaginary sphere of radius 1 AU.



3 Much of our knowledge of the solar interior comes from mathematical models. The model that best fits the observed properties of the Sun is the **standard solar model** (p. 392). **Helioseismology** (p. 393)—the study of vibrations of the solar surface caused by pressure waves in the interior—provides further insight into the Sun's structure. The effect of the solar convection zone can be seen on the surface in the form of **granulation** (p. 396) of the photosphere. Lower levels in the convection zone also leave their mark on the photosphere in the form of larger transient patterns called **supergranulation** (p. 397).




4 Above the photosphere lies the **chromosphere** (p. 390), the Sun's lower atmosphere. Most of the absorption lines seen in the solar spectrum are produced in the upper photosphere and the chromosphere. In the **transition zone** (p. 390) above the chromosphere, the temperature increases from a few thousand to around a million kelvins. Above the transition zone is the Sun's thin, hot upper atmosphere, the




solar **corona** (p. 390). At a distance of about 15 solar radii, the gas in the corona is hot enough to escape the Sun's gravity, and the corona begins to flow outward as the **solar wind** (p. 390).


5 **Sunspots** (p. 401) are Earth-sized regions on the solar surface that are a little cooler than the surrounding photosphere. They are regions of intense magnetism. Both the numbers and locations of sun spots vary in a roughly 11-year **sunspot cycle** (p. 404) as the Sun's magnetic field rises and falls. The overall direction of the field reverses from one sunspot cycle to the next. The 22-year cycle that results when the direction of the field is taken into account is called the **solar cycle** (p. 404).



6 Solar activity tends to be concentrated in **active regions** (p. 405) associated with groups of sunspots. **Prominences** (p. 405) are looplike or sheetlike structures produced when hot gas ejected by activity on the solar surface interacts with the Sun's magnetic field. The more intense **flares** (p. 406) are violent surface explosions that blast particles and radiation into interplanetary space. **Coronal mass ejections** (p. 406) are huge blobs of magnetized gas escaping into interplanetary space. Most of the solar wind flows outward from low-density regions of the corona called **coronal holes** (p. 407).



7 The Sun generates energy by converting hydrogen to helium in its core by the process of **nuclear fusion** (p. 410). Nuclei are held together by the **strong nuclear force** (p. 411). When four protons are converted to a helium nucleus in the **proton-proton chain** (p. 412), some mass is lost. The **law of conservation of mass and energy** (p. 411) requires that this mass appear as energy, eventually resulting in the light we see. Very high temperatures are needed for fusion to occur.



8 **Neutrinos** (p. 411) are nearly massless particles that are produced in the proton-proton chain and escape from the Sun. They interact via the **weak nuclear force** (p. 412). Despite their elusiveness, it is possible to detect a small fraction of the neutrinos streaming from the Sun. Observations over several decades led to the **solar neutrino problem** (p. 414)—substantially fewer neutrinos are observed than are predicted by theory. The accepted explanation, supported by recent observational evidence, is that **neutrino oscillations** (p. 415) convert some neutrinos to other (undetected) particles en route from the Sun to Earth.



**MasteringAstronomy**<sup>®</sup> For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- LO1** Name and briefly describe the main regions of the Sun. How hot is the solar surface? The solar core?
- LO2** What is luminosity, and how is it measured in the case of the Sun?
- POS** How do scientists construct models of the Sun?
- What is helioseismology, and what does it tell us about the Sun?
- LO3 POS** How do observations of the Sun's surface tell us about conditions in the solar interior?
- Describe how energy generated in the solar core eventually reaches Earth.
- LO4** Why does the Sun appear to have a sharp edge?
- LOS** What is the solar wind?
- Why do we say that the solar cycle is 22 years long?
- LO6** What is the cause of sunspots, flares, and prominences?
- Describe how coronal mass ejections may influence life on Earth.
- What fuels the Sun's enormous energy output?
- LO7** What are the ingredients and the end result of the proton-proton chain in the Sun? Why is energy released in the proton-proton chain?
- LO8 POS** Why are scientists so interested in solar neutrinos? What is the most likely solution to the solar neutrino problem?
- What would we observe on Earth if the Sun's internal energy source suddenly shut off? How long do you think it might take—minutes, days, years, or millions of years—for the Sun's light to begin to fade? Repeat the question for solar neutrinos.

## Conceptual Self-Test: Multiple Choice

- Compared with Earth's diameter, the Sun's diameter is about (a) the same; (b) ten times larger; (c) one hundred times larger; (d) one million times larger.
- Overall, the Sun's average density is roughly the same as that of (a) rain clouds; (b) water; (c) silicate rocks; (d) iron-nickel meteorites.
- The Sun spins on its axis roughly once each (a) hour; (b) day; (c) month; (d) year.
- If astronomers lived on Venus instead of on Earth, the solar constant they measure would be (a) larger; (b) smaller; (c) the same.
- The primary source of the Sun's energy is (a) fusion of light nuclei to make heavier ones; (b) fission of heavy nuclei into lighter ones; (c) the slow release of heat left over from the Sun's formation; (d) the solar magnetic field.
- VIS** According to the standard model of the Sun (Figure 16.6), as the distance from the center increases, the density decreases (a) at about the same rate as the temperature decreases; (b) faster than the temperature decreases; (c) more slowly than the temperature decreases; (d) but the temperature increases.
- A typical solar granule is about the size of (a) a U.S. city; (b) a large U.S. state; (c) the Moon; (d) Earth.
- As we move to greater and greater distances above the solar photosphere, the temperature in the Sun's atmosphere (a) steadily increases; (b) steadily decreases; (c) first decreases and then increases; (d) stays the same.
- The time between successive sunspot maxima is about (a) a month; (b) a year; (c) a decade; (d) a century.
- The solar neutrino problem is that (a) we detect more solar neutrinos than we expect; (b) we detect fewer solar neutrinos than we expect; (c) we detect the wrong type of neutrinos; (d) we can't detect solar neutrinos.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

- Use the reasoning presented in Section 16.1 to calculate the value of the "solar constant" (a) on Mercury at perihelion, (b) on Jupiter.
- Use Wien's law to determine the wavelength corresponding to the peak of the blackbody curve (a) in the core of the Sun, where the temperature is  $10^7$  K, (b) in the solar

convection zone ( $10^5$  K), and (c) just below the solar photosphere ( $10^4$  K). [∞ \(Sec. 3.4\)](#). What form (visible, infrared, X-ray, etc.) does the radiation take in each case?

3. ●● The largest-amplitude solar pressure waves have periods of about 5 minutes and move at the speed of sound in the outer layers of the Sun, roughly 10 km/s. (a) How far does such a wave move during one wave period? (b) Approximately how many wavelengths are needed to completely encircle the Sun's equator? (c) Compare the wave period with the orbital period of an object moving just above the solar photosphere. [∞ \(More Precisely 2-2\)](#)
4. ● If convected solar material moves at 1 km/s, how long does it take to flow across the 1000-km expanse of a typical granule? Compare your answer with the roughly 10-minute lifetimes observed for most solar granules.
5. ●● Use Stefan's law ( $\text{flux} \propto T^4$ , where  $T$  is the temperature in kelvins) to calculate how much less energy (as a fraction) is emitted per unit area of a 4500-K sunspot than from the surrounding 5800-K photosphere. [∞ \(Sec. 3.4\)](#)
6. ● The solar wind carries mass away from the Sun at a rate of about 2 million tons/s (1 ton = 1000 kg). At this rate, how long would it take for all of the Sun's mass to escape?
7. ● Use the information presented in this chapter and in *More Precisely 8-1* to estimate the radius at which the speed of protons in the corona first exceeds the solar escape speed.
8. ●● (a) Assuming constant luminosity, calculate how much equivalent mass (relative to the current mass of the Sun) the Sun has radiated into space in the 4.6 billion years since it formed. How much hydrogen has been consumed? (b) How long would it take the Sun to radiate its entire mass into space?

## Activities

*Never look directly at the Sun without a filter!*

### Collaborative

1. The safest way to observe the Sun—and one that lets several people share the experience—is to *project* the Sun's image onto a screen. Here are two ways to do it.
  - (a) To build a “pinhole camera,” you'll need two sheets of stiff white paper and a pin. Use the pin to punch a hole in the center of one of your pieces of paper. Go outside, hold the paper up, and aim the hole at the Sun. **Don't look directly at the Sun—through the hole or in any other way!** Find the image of the Sun coming through the hole. This is not just a circle of light, but a true image of the Sun! Move the other piece of paper back and forth until the image looks best. What happens to the image as you vary the size of the pinhole?
  - (b) Alternatively, you can project an image of the Sun using a pair of binoculars or a small telescope. You'll need a sheet of stiff white paper and a tripod. Mount the binoculars or telescope firmly on the tripod and point it at the Sun. **Don't attempt to view the Sun directly through the binoculars or telescope!** Hold the paper about 10 inches behind the eyepiece to act as a screen. You should see a bright circle of light, probably blurred. Focus the instrument until the circle is sharp. This is the disc of the Sun. Experiment with moving the card closer and farther away from the telescope. What effect does this have on the image?

Build one of these projectors and use it to study some sunspots, as in individual project 2. If you are fortunate enough to be in the right place at the right time, use it to view a partial or total solar eclipse. For comparison, have half your group construct a pinhole camera and the other half build a telescopic projector, and compare the results. What are the pros and cons of each?

### Individual

1. A filtered telescope will easily show you sunspots. Count the number of sunspots you see on the Sun's surface. Notice that sunspots often come in pairs or groups. Come back and look again a few days later and you'll see that the Sun's rotation has caused the spots to move, and the spots themselves have changed. If a sufficiently large sunspot (or sunspot group) is seen, continue to watch it as the Sun rotates. It will be out of sight for about 2 weeks. Can you determine the solar rotation period from these observations?
2. Solar granulation is not too hard to see. Earth's atmosphere is most stable in the morning hours. Observe the Sun on a cool morning, 1 or 2 hours after it has risen. Use high magnification and look initially at the middle of the Sun's disk. Can you see changes in the granulation pattern? They are there, but they are not always obvious or easy to see.



# The Stars

## GIANTS, DWARFS, AND THE MAIN SEQUENCE

We have now studied Earth, the Moon, the solar system, and the Sun. To continue our inventory of the contents of the universe, we must move away from our local environment into the depths of space. In this chapter, we take a great leap in distance and consider stars in general. Our primary goal is to comprehend the nature of the stars that make up the constellations, as well as the myriad more distant stars we cannot perceive with our unaided eyes. Rather than studying their individual peculiarities, however, we will concentrate on determining the physical and chemical properties they share.

There is order in the legions of stars scattered across the sky. By cataloging and comparing their basic properties—luminosities, temperatures, composition, masses, and radii—astronomers gain new insights into how stars form and evolve. Like comparative planetology in the solar system, the study of the stars plays a vital role in furthering our understanding of the Galaxy and the universe we inhabit.

**The Big Picture** Stars are everywhere in the nighttime sky. The naked eye can spot about 6000 of them, spread across 88 constellations. Millions more are visible even with binoculars or a small telescope. The total number of stars is impossible to count, and relatively few have been studied in detail. Yet, it is stars that tell us more about the fundamentals of astronomy than any other objects in the universe.

**LEFT:** The *Hubble Space Telescope* recently imaged the magnificent globular star cluster M9, a rich group of about 300,000 stars spread across some 25 light-years. Resembling a swirling beehive, this region is about 25,000 light-years away in the constellation Ophiuchus. Its reddish stars are about twice as old as the Sun. (ESA/NASA)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Explain how stellar distances are determined.
- 2 Describe the motions of the stars through space, and say how those motions are measured from Earth.
- 3 Distinguish between luminosity and apparent brightness, and explain how stellar luminosity is determined.
- 4 Explain the usefulness of classifying stars according to their colors, surface temperatures, and spectral characteristics.
- 5 Say how physical laws are used to estimate stellar sizes.
- 6 Describe how a Hertzsprung–Russell diagram is used to identify stellar properties.
- 7 Outline how knowledge of a star's spectroscopic properties can lead to an estimate of its distance.
- 8 Explain how the masses of stars are measured and how mass is related to other stellar properties.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.

## 17.1 The Solar Neighborhood

The Galaxy in which we live—the Milky Way—is an enormous collection of stars and interstellar matter held together by gravity. It contains more than 100 billion stars, spread throughout a volume of space nearly 100,000 light-years across, all orbiting the *galactic center* some 25,000 light-years from Earth (see Chapter 23). The Sun—and, in fact, every other star you see while gazing at the night sky—is part of this vast system. In this chapter we describe some of the distance-measurement techniques by which astronomers have extended their studies to larger and larger volumes of space, mapping out the distribution of stars on these vast scales.


As with the planets, knowing the distances to the stars is essential to determining many of their other properties. Looking deep into space within our Galaxy, astronomers can observe and study literally millions of individual stars. By observing other distant galaxies, we can statistically infer the properties of trillions more. All told, the observable universe probably contains several tens of sextillions (1 sextillion =  $10^{21}$ ) stars. Yet, remarkably, despite their incredible numbers, the essential properties of stars—their appearance in the sky, their births, lives, deaths, and even their interactions with their environment—can be understood in terms of just a few basic physical stellar quantities: luminosity (brightness), temperature (color), chemical composition, size, and mass. ∞ (Sec. 16.1)

The twin goals of measuring stellar distances and categorizing stellar properties have advanced hand in hand. As we will see, as more stellar distances become known, new insights into stellar properties are obtained, and these in turn present new techniques for distance measurement, applicable to even greater distances. In many ways, the story of how these techniques have evolved in tandem is the history of modern astronomy.


### Stellar Parallax

Recall from Chapter 1 how surveyors and astronomers use *parallax* to measure distances to terrestrial and solar system objects. Parallax is the apparent shift of a foreground object relative to some distant background as the observer's point of view changes. ∞ (Sec. 1.6) To determine an object's parallax, we observe it from either end of some baseline and measure the angle through which the line of sight to the object shifts. In astronomical contexts, the angle is usually obtained by comparing photographs made from the two ends of the baseline.

As the distance to the object increases, the parallax becomes smaller and therefore harder to measure. Even the closest stars are so far away that no baseline on Earth is sufficient to allow an accurate determination of their distances. Their parallactic shifts, as seen from different points on Earth, are just too small. However, by observing a star *at different times of the year*, as shown in Figure 17.1 (see also Figure 1.30)



(a)




This is the view as seen in January . . . . . and in July, when the star shifts.

(b)

▲ **FIGURE 17.1 Stellar Parallax** (a) For observations of stars made 6 months apart, the baseline is twice the Earth–Sun distance, or 2 AU. Compare with Figure 1.30, which shows the same geometry, but on a much smaller scale. (b) The parallactic shift (exaggerated here, and indicated by the white arrow) is usually measured photographically, as illustrated by the red star seen here. Images of the same region of the sky made at different times of the year determine a star's apparent movement relative to background stars.


and then comparing our observations, we can extend our baseline to the diameter of Earth’s orbit around the Sun, namely, 2 AU. Only with this enormously longer baseline do some stellar parallaxes become measurable. As indicated in the figure, a star’s parallactic angle—or, more commonly, just its “parallax”—is conventionally defined to be *half* its apparent shift relative to the background as we move from one side of Earth’s orbit to the other.

Because stellar parallaxes are so small, astronomers generally find it convenient to measure parallax in arc seconds rather than in degrees. If we ask at what distance a star must lie in order for its observed parallax to be exactly 1", we get an answer of 206,265 AU, or  $3.1 \times 10^{16}$  m  (More Precisely 1-2) Astronomers call this distance 1 **parsec** (1 pc), from “*parallax* in arc *seconds*.” Because parallax decreases as distance increases, we can relate a star’s parallax to its distance by the following simple formula:

$$\text{distance (in parsecs)} = \frac{1}{\text{parallax (in arc seconds)}}.$$

Thus, a star with a measured parallax of 1" lies at a distance of 1 pc from the Sun. The parsec is defined so as to make the conversion between distance and parallactic angle easy: An object with a parallax of 0.5" lies at a distance of  $1/0.5 = 2$  pc, an object with a parallax of 0.1" lies at  $1/0.1 = 10$  pc, and so on. One parsec is approximately equal to 3.3 light-years.



## Our Nearest Neighbors


The closest star to Earth (after the Sun) is called Proxima Centauri. This star is a member of a triple-star system (three separate stars orbiting one another, bound together by gravity) known as the Alpha Centauri complex.  (Discovery 15-1) Proxima Centauri displays the largest known stellar parallax, 0.77", which means that it is about  $1/0.77 = 1.3$  pc away—approximately 270,000 AU, or 4.3 light-years. That’s the distance of the *nearest* star to Earth—almost 300,000 times the distance from Earth to the Sun! This is a fairly typical interstellar distance in the Milky Way Galaxy.

Vast distances can sometimes be grasped by means of analogies. Imagine Earth as a grain of sand orbiting a marble-sized Sun at a distance of 1 m. The nearest star,

also a marble-sized object, is then more than 270 *kilometers* away. Except for the other planets in our solar system, themselves ranging in size from grains of sand to millimeter-sized pellets and all lying within 50 m of the “Sun,” nothing else of consequence exists in the 270 km separating the two stars. Such is the void of interstellar space.

The next nearest neighbor to the Sun beyond the Alpha Centauri system is called Barnard’s star. Its parallax is 0.55", so it lies at a distance of 1.8 pc, or 6.0 light-years—370 km in our model—from Earth. Figure 17.2 is a map of our nearest Galactic neighbors—the 30 or so stars lying within 4 pc of Earth.

Ground-based images of stars are generally smeared out into a disk of radius 1" or so by turbulence in Earth’s atmosphere.  (Sec. 5.4) However, astronomers have special equipment that can routinely measure stellar parallaxes of 0.03" or less, corresponding to stars within about 30 pc (100 light-years) of Earth. Several thousand stars lie within this range, most of them of much lower luminosity than the Sun and invisible to the naked eye. High-resolution adaptive optics systems allow even more accurate measurements of stellar positions, extending the parallax range to over 100 pc in a few cases, although such measurements are not yet “routine.”  (Sec. 5.4)



▲ **FIGURE 17.2 The Solar Neighborhood** A plot of the 30 closest stars to the Sun, projected so as to reveal their three-dimensional relationships. All lie within 4 pc (about 13 light-years) of Earth. The circular gridlines represent distances from the Sun in the galactic plane; the vertical lines denote distances perpendicular to that plane.



Even greater precision can be achieved by placing instruments in space, above Earth's atmosphere. In the 1990s, data from the European *Hipparcos* satellite extended the range of accurately measured parallaxes to well over 200 pc, encompassing nearly a million stars. Even so, the vast majority of stars in our Galaxy are far more distant. Following *Hipparcos*, the European Space Agency (ESA) has ambitious plans to greatly expand the scope of its stellar measurements. ESA's GAIA project, due to launch in 2013, will have the astonishing range of 10,000 pc spanning much of the Milky Way Galaxy and encompassing roughly 1 billion stars! In addition to mapping out the structure of the Milky Way Galaxy to unprecedented precision, this mission will allow astronomers to study in detail the properties of nearby stars of all masses and will also greatly expand our knowledge of extrasolar planetary systems. [∞ \(Sec. 15.6\)](#) With these new data, in a time span of just three decades, the fundamental stellar database upon which almost all of astronomy depends will have increased in size by a factor of a million. The results may be nothing short of revolutionary.

## Stellar Motion

In addition to the apparent motion caused by parallax, stars have real spatial motion through the Galaxy. In Chapter 23, we will see how astronomers have measured the Sun's actual motion around the galactic center. However, *relative* to the Sun—that is, as seen by astronomers on Earth as we travel through space along with our parent star—stellar motion has two components. A star's *radial* velocity—along the line of sight—can be measured using the Doppler effect. [∞ \(Sec. 3.5\)](#) For many nearby stars, their *transverse* velocity—perpendicular to our line of sight—can also be determined by careful monitoring of the star's position in the sky.


Figure 17.3 compares two photographs of the sky around Barnard's star. The photographs were made on the same day of the year, but 22 years apart. Note that the star, marked by the arrow, has moved during the 22-year interval shown: If the two images were superimposed, the images of the other stars in the field of view would coincide, but those

of Barnard's star would not. Because Earth was at the same point in its orbit when these photographs were taken, the observed displacement is *not* the result of parallax caused by Earth's motion around the Sun. Instead, it indicates *real* spatial motion of Barnard's star relative to the Sun.


The annual movement of a star across the sky, as seen from Earth and corrected for parallax, is called **proper motion**. It describes the transverse component of a star's velocity relative to the Sun. (Both the star and the Sun are moving through space as they travel through the Galaxy; however, only their *relative* motion causes the star's position in the sky to change, as seen from Earth.) Like parallax, proper motion is measured in terms of angular displacement. Since the angles involved are typically very small, proper motion is usually expressed in arc seconds per year. Barnard's star moved 228" in 22 years, so its proper motion is  $228''/22$  years, or  $10.4''/\text{yr}$ .

A star's transverse velocity is easily calculated once its proper motion and its distance are known. At the distance of Barnard's star (1.8 pc), an angle of  $10.4''$  corresponds to a physical displacement of 0.000091 pc, or about 2.8 billion km. Barnard's star takes a year ( $3.2 \times 10^7$  s) to travel this distance, so its transverse velocity is 2.8 billion km/ $3.2 \times 10^7$  s, or 89 km/s. [∞ \(More Precisely 1-2\)](#) Even though stars' transverse velocities are often quite large—tens or even hundreds of kilometers per second—their great distances from the Sun mean that their proper motion is small, and it usually takes many years for us to discern their movement across the sky. In all probability, every star in Figure 17.3 has some transverse motion relative to the Sun. However, only Barnard's star has proper motion large enough to be visible in these frames. In fact, Barnard's star has the largest known proper motion of any star. Only a few hundred stars have proper motions greater than  $1''/\text{yr}$ .

Now consider the three-dimensional motion of our nearest neighbor, the Alpha Centauri system, sketched in Figure 17.4 in relation to our own solar system. Alpha Centauri's proper motion has been measured to be  $3.7''/\text{yr}$ . At Alpha Centauri's distance of 1.35 pc, that measurement implies a transverse velocity of 24 km/s. We can determine



▲ **FIGURE 17.3 Proper Motion** A comparison of two photographic plates taken 22 years apart shows evidence of real spatial motion for Barnard's star (denoted by an arrow). (*Harvard College Observatory*)



▲ **FIGURE 17.4 Real Spatial Motion** This sketch shows the motion of the Alpha Centauri star system relative to our solar system. The transverse component of its velocity is derived by observing the system's proper motion. The radial component is measured by using the Doppler shift of lines in Alpha Centauri's spectrum. The true spatial velocity, indicated by the red arrow, results from the combination of the two.

the other component of motion—the radial velocity—by means of the Doppler effect. Spectral lines from Alpha Centauri are blueshifted by a tiny amount—about 0.0067 percent—allowing astronomers to measure the star system's radial velocity (relative to the Sun) as  $300,000 \text{ km/s} \times 6.7 \times 10^{-5} = 20 \text{ km/s}$  toward us. ∞ (Sec. 3.5)

What is the true spatial motion of Alpha Centauri? Will this alien system collide with our own sometime in the future? The answer is no: Alpha Centauri's transverse velocity will steer it well clear of the Sun. We can combine the transverse (24 km/s) and radial (20 km/s) velocities according to the Pythagorean theorem, as indicated in Figure 17.4. The total velocity is  $\sqrt{24^2 + 20^2}$ , or about 31 km/s, in the direction shown by the horizontal red arrow. As the figure indicates, Alpha Centauri will get no closer to us than about 1 pc, and that won't happen until 280 centuries from now.

### CONCEPT Check

- ✓ Why can't astronomers use simultaneous observations from different parts of Earth's surface to determine stellar distances?
- ✓ Why are the spatial velocities of distant stars generally poorly known?


## 17.2 Luminosity and Apparent Brightness

Luminosity is an *intrinsic* property of a star—it does not depend in any way on the location or motion of the observer. Luminosity is sometimes referred to as the star's *absolute brightness*. However, when we look at a star, we see, not its luminosity, but rather its **apparent brightness**—the amount of energy striking a unit area of some light-sensitive surface or device (such as a charge-coupled device [CCD] chip or a human eye) per unit time. Apparent brightness is a measure, not of a star's luminosity, but of the *energy flux* (energy per unit area per unit time) produced by the star, as seen from Earth. A star's apparent brightness depends on our *distance* from the star. In this section, we discuss in more detail how these important quantities are related to one another.


### Another Inverse-Square Law

Figure 17.5 shows light leaving a star and traveling through space. Moving outward, the radiation passes through imaginary spheres of increasing radius surrounding the source. The amount of radiation leaving the star per unit time—the star's luminosity—is constant, so the farther the light travels from the source, the less energy passes through each unit of area. Think of the energy as being spread out over an ever-larger area and therefore spread more thinly, or “diluted,” as it expands into space.

Because the area of a sphere grows as the square of the radius, the energy per unit area—the star's apparent brightness, as seen by our eye or our telescope—is inversely proportional to the square of the distance from the star.



▲ **Narrated FIGURE 17.5 Inverse-Square Law** As light moves away from a source such as a star, it steadily dilutes while spreading over progressively larger surface areas (depicted here as sections of spherical shells). Thus, the amount of radiation received by a detector (the source's apparent brightness) varies inversely as the square of its distance from the source.



▲ **FIGURE 17.6 Luminosity** Two stars A and B of different luminosities can appear equally bright to an observer on Earth if the brighter star B is more distant than the fainter star A.

Doubling the distance from a star makes it appear  $2^2$ , or 4, times dimmer. Tripling the distance reduces the apparent brightness by a factor of  $3^2$ , or 9, and so on.

Of course, the star's luminosity also affects its apparent brightness. Doubling the luminosity doubles the energy crossing any spherical shell surrounding the star and hence doubles the apparent brightness. We can therefore say that the apparent brightness of a star is directly proportional to the star's luminosity and inversely proportional to the square of its distance:

$$\text{apparent brightness (energy flux)} \propto \frac{\text{luminosity}}{\text{distance}^2}.$$

Thus, two identical stars can have the same apparent brightness if (and only if) they lie at the same distance from Earth.

However, as illustrated in Figure 17.6, two nonidentical stars can also have the same apparent brightness if the more luminous one lies farther away. A bright star (i.e., a star with large apparent brightness) is a powerful emitter of radiation (high luminosity), is near Earth, or both. Without additional information, we cannot distinguish between the effects of increasing luminosity and decreasing distance. Similarly, a faint star (a star with small apparent brightness) is a weak emitter (low luminosity), is far from Earth, or both.

Determining a star's luminosity is a twofold task. First, the astronomer must determine the star's apparent brightness by measuring the amount of energy detected through a telescope in a given amount of time. Second,


the star's distance must be measured—parallax for nearby stars and by other means (to be discussed later) for more distant stars. The luminosity can then be found from the inverse-square law. This is basically the same reasoning we used earlier in Chapter 16, in our discussion of how astronomers measure solar luminosity. (In our new terminology, the solar constant is just the apparent brightness of the Sun.) ∞ (Sec. 16.1)

## The Magnitude Scale

Instead of measuring apparent brightness in SI units (e.g., watts per square meter, the unit used for the solar constant in Chapter 16), astronomers often find it more convenient to work in terms of a construct called the **magnitude scale**. ∞ (Sec. 16.1) The scale dates from the second century B.C., when the Greek astronomer Hipparchus classified the naked-eye stars into six groups. The brightest stars were categorized as first magnitude. The next brightest stars were labeled second magnitude, and so on, down to the faintest stars visible to the naked eye, which were classified as sixth magnitude. The range 1 (brightest) through 6 (faintest) spanned all the stars known to the ancients. Notice that magnitudes are really *rankings* in terms of apparent brightness (energy flux)—a *large* magnitude means a *faint* star. Just as “first rate” means “good” in everyday speech, “first magnitude” in astronomy means “bright.”

When astronomers began using telescopes with sophisticated detectors to measure the light received from stars, they quickly discovered two important facts about the magnitude scale. First, the 1–6 magnitude range defined by Hipparchus spans about a factor of 100 in apparent brightness—a first-magnitude star is approximately 100 times brighter than a sixth-magnitude star. Second, the physiological characteristics of the human eye are such that each change in magnitude of 1 corresponds to a factor of about 2.5 in apparent brightness. In other words, to the human eye, a first-magnitude star is roughly 2.5 times brighter than a second-magnitude star, which is roughly 2.5 times brighter than a third-magnitude star, and so on. (By combining factors of 2.5, you can confirm that a first-magnitude star is indeed  $(2.5)^5 \approx 100$  times brighter than a sixth-magnitude star.)

Modern astronomers have modified and extended the magnitude scale in a number of ways. First, we now *define* a change of 5 in the magnitude of an object (going from magnitude 1 to magnitude 6, say, or from magnitude 7 to magnitude 2) to correspond to *exactly* a factor of 100 in apparent brightness. Second, because we are really talking about apparent (rather than absolute) brightnesses, the numbers in Hipparchus's ranking system are called **apparent magnitudes**. Third, the scale is no longer limited to whole numbers: A star of apparent magnitude



### Interactive FIGURE 17.7 Apparent Magnitude

This graph illustrates the apparent magnitudes of some astronomical objects and the limiting magnitudes (that is, the faintest magnitudes attainable) of some telescopes used to observe them.

Apparent magnitude measures a star's apparent brightness when the star is seen at its actual distance from the Sun. To compare intrinsic, or absolute, properties of stars, however, astronomers imagine looking at all stars from a standard distance of 10 pc. There is no particular reason to use 10 pc—it is simply convenient. A star's **absolute magnitude** is its apparent magnitude when it is placed at a distance of 10 pc from the observer. Because the distance to the star is fixed in this definition, *absolute magnitude is a measure of a star's absolute brightness, or luminosity.*

We can use the earlier discussion of the inverse-square law to relate absolute and apparent magnitudes if the distance to the star is known. When a star farther than 10 pc away from us is moved to a point 10 pc away, its apparent brightness increases and hence its apparent magnitude decreases. Stars more than 10 pc from Earth therefore have apparent magnitudes that are greater than their absolute magnitudes. For example, if a star at a distance of 100 pc were moved to the standard 10-pc distance, its distance would decrease by a factor of 10, so (by the inverse-square law) its apparent brightness would *increase* by a factor of  $10^2 = 100$ . Its apparent magnitude (by definition) would therefore decrease by 5. In other words, at 100 pc distance, the star's apparent magnitude exceeds its absolute magnitude by 5.

For stars closer than 10 pc, the reverse is true. An extreme example is our Sun. Because of its proximity to Earth, it appears very bright and thus has a large negative apparent magnitude (Figure 17.7). However, the Sun's absolute magnitude is 4.83. If the Sun were moved to a distance of 10 pc from Earth, it would be only slightly brighter than the faintest stars visible to the naked eye in the night sky.

Knowledge of a star's apparent magnitude and distance allows us to compute its absolute magnitude (luminosity). Conversely, the numerical difference between a star's absolute and apparent magnitudes is a direct measure of the distance to the star. *More Precisely 17-1* presents more detail and some examples for the connection between absolute magnitude and luminosity and on the use of the magnitude scale in computing stellar luminosities and distances.

### CONCEPT Check

- ✓ Two stars are observed to have the same apparent magnitude. On the basis of this information, what, if anything, can be said about their luminosities?

4.5 is intermediate in apparent brightness between a star of apparent magnitude 4 and one of apparent magnitude 5. Finally, magnitudes outside the range from 1 to 6 are allowed: Very bright objects can have apparent magnitudes much less than 1, and very faint objects can have apparent magnitudes far greater than 6.

Figure 17.7 illustrates the apparent magnitudes of some astronomical objects, ranging from the Sun, at  $-26.7$ , to the faintest object detectable by the *Hubble* or *Keck* telescopes, an object having an apparent magnitude of 30—about as faint as a firefly seen from a distance equal to Earth's diameter. Note that this range in magnitudes corresponds to a very large factor (actually, of  $10^{56.7/2.5} = 10^{22.7} \approx 5 \times 10^{22}$ ) in apparent brightness. Indeed, one of the main reasons that astronomers use this (admittedly rather intimidating) scale is that it allows them to compress a large spread in observed stellar properties into more “manageable” form.\*

\*Putting in the numbers, we can calculate that magnitude 1 corresponds to a flux of  $1.1 \times 10^{-8} \text{ W/m}^2$ , magnitude 20 to  $2.9 \times 10^{-16} \text{ W/m}^2$ , and so on, but astronomers find the “magnitude” versions more intuitive and much easier to remember.

## 17.3 Stellar Temperatures

Looking at the night sky, you can tell at a glance which stars are hot and which are cool. In Figure 17.8, which shows the constellation Orion as it appears through a small telescope, the colors of the cool red star Betelgeuse ( $\alpha$ ) and the hot blue star Rigel ( $\beta$ ) are clearly evident. Note that these colors are intrinsic properties of the stars and have *nothing* to do with Doppler redshifts or blueshifts. However, to obtain these stars' temperatures (3200 K for Betelgeuse and 11,000 K for Rigel), more precise observations are required. To make such measurements, astronomers turn to the radiation laws and the detailed properties of stellar spectra.  $\infty$  (Secs. 3.4, 4.3)


### Color and the Blackbody Curve

Astronomers can determine a star's surface temperature by measuring the star's apparent brightness (energy flux) at several frequencies and then matching the observations to the appropriate blackbody curve.  $\infty$  (Sec. 3.4) In the case of the Sun, the theoretical curve that best fits the emission describes a 5800-K emitter.  $\infty$  (Sec. 16.1) The same technique works for any other star, regardless of its distance from Earth.

Because the basic shape of the blackbody curve is so well understood, astronomers can estimate a star's temperature using as few as *two* measurements at selected wavelengths (which is fortunate, as detailed spectra of faint stars are often difficult and time consuming to obtain). This is accomplished through the use of telescope filters that block out all radiation except that within specific wavelength ranges. For example, a B (blue) filter rejects all radiation except for a certain range from violet to blue light. Defined by international agreement to extend from 380 to 480 nm, this range corresponds to wavelengths to which photographic film happens to be most sensitive. Similarly, a V (visual) filter passes only radiation within the 490- to 590-nm range (green to yellow), corresponding to that part of the spectrum to which human eyes are particularly sensitive. Many other filters are also in routine use—a U (ultraviolet) filter covers the near ultraviolet, and infrared filters span longer wavelength parts of the spectrum.

Figure 17.9 shows how the B and V filters admit different amounts of light for objects radiating at different


temperatures. In curve (a), corresponding to a very hot 30,000-K emitter, considerably more radiation (about 30 percent more) is received through the B filter than through the V filter, so this object looks brighter in B than in V. In curve (b), the temperature is 10,000 K, and the B and V fluxes are about the same. In the cool 3000-K curve (c), about five times more energy is received in the V range than in the B range, so the B image now is much fainter than the V image. In each case, it is possible to reconstruct the entire blackbody curve on the basis of only those two measurements, because no other blackbody curve can be




(a)



(b)



**► FIGURE 17.8 Star Colors** (a) The different colors of the stars composing the constellation Orion are easily distinguished in this photograph taken by a wide-field camera attached to a small telescope. The bright red star at the upper left is Betelgeuse ( $\alpha$ ); the bright blue-white star at the lower right is Rigel ( $\beta$ ). (Compare with Figure 1.8.) The field of view of this photograph is wide, about  $20^\circ$  across. (b) An incredibly rich field of colorful stars, this time in the direction of the center of the Milky Way. Here, the field of view is just 2 arc minutes across—much smaller than in (a). (P. Sanz/Alamy; NASA)



▲ **FIGURE 17.9 Blackbody Curves** Star (a) is very hot—30,000 K—so its B (blue) intensity is greater than its V (visual) intensity (as is actually the case for Rigel in Figure 17.8a). Star (b) has roughly equal B and V readings and so appears white, and its temperature is about 10,000 K. Star (c) is red; its V intensity greatly exceeds the B value, and its temperature is 3000 K (much as for Betelgeuse in Figure 17.8a).

drawn through both measured points. To the extent that a star's spectrum is well approximated as a blackbody, measurements of the B and V fluxes are enough to specify the star's blackbody curve and thus yield its surface temperature.

Thus, astronomers can estimate a star's temperature simply by measuring and comparing the amount of light received through different colored filters. As discussed in Chapter 5, this type of non-spectral-line analysis using a standard set of filters is known as **photometry**. ∞ (Sec. 5.3) Table 17.1 lists, for several prominent stars, the surface temperatures derived by photometric means, along with the color that would be perceived in the absence of filters.


**TABLE 17.1 Stellar Colors and Temperatures**

Surface Temperature (K)	Color	Familiar Examples
30,000	Blue-violet	Mintaka ( $\delta$ Orionis)
20,000	Blue	Rigel
10,000	White	Vega, Sirius
7000	Yellow-white	Canopus
6000	Yellow	Sun, Alpha Centauri
4000	Orange	Arcturus, Aldebaran
3000	Red	Betelgeuse, Barnard's star

## Stellar Spectra

Color is a useful way to describe a star, but astronomers often use a more detailed scheme to classify stellar properties, incorporating additional knowledge of stellar physics obtained through spectroscopy. Figure 17.10 compares the spectra of several different stars, arranged in order of decreasing surface temperature (as determined from measurements of their colors). All the spectra extend from 400 to 650 nm, and each shows a series of dark absorption lines superimposed on a background of continuous color, like the spectrum of the Sun. ∞ (Sec. 16.3) However, the precise patterns of lines reveal many differences. Some stars display strong lines in the long-wavelength part of the spectrum (to the left in the figure). Other stars have their strongest lines at short wavelengths (to the right). Still others show strong absorption lines spread across the whole visible spectrum. What do these differences tell us?

Although spectral lines of many elements are present with widely varying strengths, the differences among the spectra in Figure 17.10 are not due to differences in composition.




▲ **FIGURE 17.10 Stellar Spectra** Comparison of spectra observed for seven different stars having a range of surface temperatures. These are not actual spectra, which are messy and complex, but simplified artists' renderings illustrating notable spectral features. The spectra of the hottest stars, at the top, show lines of helium and multiply ionized heavy elements. In the coolest stars, at the bottom, helium lines are absent, but lines of neutral atoms and molecules are plentiful. At intermediate temperatures, hydrogen lines are strongest.

## MORE PRECISELY 17-1

### More on the Magnitude Scale

Let's recast our discussion of two important topics—stellar luminosity and the inverse-square law—in terms of magnitudes.

*Absolute magnitude* is equivalent to luminosity—an intrinsic property of a star. Given that the Sun's absolute magnitude is 4.83 (see Appendix 3, Table 6), we can construct a conversion chart (shown below) relating these two quantities. Since an increase in brightness by a factor of 100 corresponds to a decrease in magnitude by 5 units, it follows that a star with luminosity 100 times that of the Sun has absolute magnitude  $4.83 - 5 = -0.17$ , while a 0.01-solar luminosity star has absolute magnitude  $4.83 + 5 = 9.83$ . We can fill in the gaps by noting that 1 magnitude corresponds to a factor of  $100^{1/5} \approx 2.512$ ; 2 magnitudes to  $100^{2/5} \approx 6.310$ , and so on. A factor of 10 in brightness corresponds to 2.5 magnitudes. You can use this chart to convert between solar luminosities and absolute magnitudes in many of the figures in this and later chapters.



**EXAMPLE 1** Let's calculate the luminosity (in solar units) of a star having absolute magnitude  $M$  (the conventional symbol for absolute magnitude, not to be confused with mass!). The star's absolute magnitude differs from that of the Sun by  $(M - 4.83)$  magnitudes, so, in accordance with the reasoning just presented,

the luminosity  $L$  differs from the solar luminosity by a factor of  $10^{-(M-4.83)/5}$ , or  $10^{-(M-4.83)/2.5}$ . We can therefore write

$$L \text{ (solar units)} = 10^{-(M-4.83)/2.5}.$$

Plugging in some numbers (taken from Appendix 3, Tables 5 and 6), we find that the Sun, with  $M = 4.83$ , of course has  $L = 10^0 = 1$ . Sirius A, with  $M = 1.45$ , has luminosity  $10^{1.35} = 22$  solar units, Barnard's star, with  $M = 13.24$ , has luminosity  $10^{-3.35} = 4.3 \times 10^{-4}$  solar units, Betelgeuse has  $M = -5.14$  and luminosity 9700 Suns, and so on.

To cast the *inverse-square law* in these terms, recall that increasing the distance to a star by a factor of 10 decreases its apparent brightness by a factor of 100 (by the inverse-square law) and hence increases its apparent magnitude by 5 units. Increasing the distance by a factor of 100 increases the apparent magnitude by 10, and so on. Every increase in distance by a factor of 10 increases the apparent magnitude by 5. Since absolute magnitude is simply apparent magnitude at a distance of 10 pc, we can write

apparent magnitude – absolute magnitude

$$= 5 \log_{10} \left( \frac{\text{distance}}{10 \text{ pc}} \right).$$

(The *logarithm* function—the LOG key on your calculator—is defined by the property that if  $a = \log_{10}(b)$ , then  $b = 10^a$ .) Even though it doesn't look much like it, this equation is precisely equivalent to the inverse-square law presented in the text! Note that for stars more than 10 pc from Earth, the apparent magnitude is greater than the absolute magnitude, while the reverse is true for stars closer than 10 pc.

**EXAMPLE 2** The Sun, with an absolute magnitude of 4.83, seen from a distance of 100 pc, would have an apparent magnitude of  $4.83 + 5 \log_{10}(100) = 14.83$ , since  $\log_{10}(100) = 2$ . This is well below the threshold of visibility for binoculars or even a large amateur telescope (see Figure 17.7). We can also turn this around to illustrate how knowledge of a star's absolute and apparent magnitudes tells us its distance. The star Rigel Kentaurus (also known as Alpha Centauri) has absolute magnitude +4.34 and is observed to have apparent magnitude –0.01. The magnitude difference is –4.35, so its distance must therefore be  $10 \text{ pc} \times 10^{-4.34/5} = 1.35 \text{ pc}$ , in agreement with the result (obtained by parallax) presented in the text.

Detailed spectral analysis indicates that the seven stars shown have similar elemental abundances—all are more or less solar in makeup.  $\infty$  (Sec. 16.3) Rather, as discussed in Chapter 4, the differences are due almost entirely to the stars' *temperatures*.  $\infty$  (Sec. 4.5) The spectrum at the top of the figure is exactly what we would expect from a star with solar composition and a surface temperature of about 30,000 K, the second

spectrum is what we would anticipate from a 20,000-K star, and so on, down to the 3000-K star at the bottom.

The main differences among the spectra in Figure 17.10 are as follows:

- Spectra of stars having surface temperatures exceeding 25,000 K usually show *strong* absorption lines of singly

ionized helium (i.e., helium atoms that have lost one orbiting electron) and multiply ionized heavier elements, such as oxygen, nitrogen, and silicon (the latter lines are not shown in the figure). These strong lines are not seen in the spectra of cooler stars because only very hot stars can excite and ionize such tightly bound atoms.

- In contrast, the hydrogen absorption lines in the spectra of very hot stars are relatively *weak*. The reason is not a lack of hydrogen, which is by far the most abundant element in all stars. At these high temperatures, however, much of the hydrogen is ionized, so there are few intact hydrogen atoms to produce strong spectral lines.
- Hydrogen lines are strongest in stars having intermediate surface temperatures of around 10,000 K. This temperature is just right for electrons to move frequently between hydrogen's second and higher orbitals, producing the characteristic visible hydrogen spectrum. ∞ (*More Precisely 4-1*) Lines of tightly bound atoms—for example, of helium and nitrogen—which need lots of energy for excitation, are rarely observed in the spectra of these stars, whereas lines from more loosely bound atoms—such as those of calcium and titanium—are relatively common.
- Hydrogen lines are again weak in stars with surface temperatures below about 4000 K, but now because the temperature is too low to boost many electrons out of the ground state. ∞ (*Sec. 4.2*) The most intense spectral lines in these stars are due to weakly excited heavy atoms; no lines from ionized elements are seen. Temperatures in the coolest stars are low enough for molecules to survive, and many of the observed absorption lines are produced by molecules rather than by atoms. ∞ (*Sec. 4.4*)

Stellar spectra are the source of all the detailed information we have on stellar composition, and they do in fact reveal significant differences in composition among stars, particularly in the abundances of carbon, nitrogen, oxygen, and heavier elements. However, as we have just seen, these differences are *not* the primary reason for the different spectra that are observed. Instead, the main determinant of a star's spectral appearance is its temperature, and stellar spectroscopy is a powerful and precise tool for measuring this important stellar property.

## Spectral Classification

Stellar spectra like those shown in Figure 17.10 were obtained for many stars well before the start of the 20th century as observatories around the world amassed spectra from stars in both hemispheres of the sky. Between 1880 and 1920, researchers correctly identified some of the observed spectral lines on the basis of comparisons between those

lines and lines obtained in the laboratory. The researchers, though, had no firm understanding of how the lines were produced. Modern atomic theory had not yet been developed, so the correct interpretation of the line strengths, as just described, was impossible at the time.

Lacking a full understanding of how atoms produce spectra, early workers classified stars primarily according to their hydrogen-line intensities. They adopted an alphabetic A, B, C, D, E... scheme in which A stars, with the strongest hydrogen lines, were thought to have more hydrogen than did B stars, and so on. The classification extended as far as the letter P.

In the 1920s, scientists began to understand the intricacies of atomic structure and the causes of spectral lines. Astronomers quickly realized that stars could be more meaningfully classified according to their surface temperature. Instead of adopting an entirely new scheme; however, they chose to shuffle the existing alphabetical categories—those based on the strengths of the hydrogen lines—into a new sequence based on temperature. In the modern scheme, the hottest stars are designated O, because they have very weak absorption lines of hydrogen and were classified toward the end of the original scheme. In order of decreasing temperature, the surviving letters now run O, B, A, F, G, K, M. (The other letter classes have been dropped.) These stellar designations are called **spectral classes** (or *spectral types*). Use the time-honored (and politically incorrect) mnemonic “**Oh, Be A Fine Girl, Kiss Me**” to remember them in the correct order.\*

Astronomers further subdivide each lettered spectral classification into 10 subdivisions, denoted by the numbers 0–9. By convention, the lower the number, the hotter is the star. For example, our Sun is classified as a G2 star (a little cooler than G1 and a little hotter than G3), Vega is a type A0, Barnard's star is M5, Betelgeuse is M2, and so on. Table 17.2 lists the main properties of each stellar spectral class for the stars presented in Table 17.1.

We should not underestimate the importance of the early work in classifying stellar spectra. Even though the original classification was based on erroneous assumptions, the painstaking accumulation of large quantities of accurate data paved the way for rapid improvements in understanding once a theory came along that explained the observations.

### CONCEPT Check

- ✓ Why does a star's spectral classification depend on its temperature?

\*Astronomers have since added two new spectral classes—L and T—for low-mass, low-temperature stars whose odd spectra distinguish them from the M-class stars in the current scheme. These objects are not “true” stars, fusing hydrogen into helium in their cores; rather, they are “brown dwarfs” (see Chapter 20) that never achieved high enough central temperature for fusion to begin.



TABLE 17.2 Stellar Spectral Classes

Spectral Class	Surface Temperature (K)	Noteworthy Absorption Lines	Familiar Examples
O	30,000	Ionized helium strong; multiply ionized heavy elements; hydrogen faint	Mintaka (O9)
B	20,000	Neutral helium moderate; singly ionized heavy elements; hydrogen moderate	Rigel (B8)
A	10,000	Neutral helium very faint; singly ionized heavy elements; hydrogen strong	Vega (A0), Sirius (A1)
F	7000	Singly ionized heavy elements; neutral metals; hydrogen moderate	Canopus (F0)
G	6000	Singly ionized heavy elements; neutral metals; hydrogen relatively faint	Sun (G2), Alpha Centauri (G2)
K	4000	Singly ionized heavy elements; neutral metals strong; hydrogen faint	Arcturus (K2), Aldebaran (K5)
M	3000	Neutral atoms strong; molecules moderate; hydrogen very faint	Betelgeuse (M2), Barnard's star (M5)

## 17.4 Stellar Sizes

Most stars are unresolved points of light in the sky, even when viewed through the largest telescopes. Even so, astronomers can often make quite accurate determinations of their sizes.


### Direct and Indirect Measurements

Some stars are big enough, bright enough, and close enough to allow us to measure their sizes *directly*. One well-known example is the bright red star Betelgeuse, a prominent member of the constellation Orion (Figure 17.8). As shown in Figure 17.11(a), Betelgeuse is barely large enough to be resolvable by the *Hubble Space Telescope*

at short wavelengths. Steadily improving interferometric and adaptive-optics techniques have allowed astronomers to construct very-high-resolution stellar images in a small number of cases. Some results show enough detail to show a few surface features, as noted in Figure 17.11(b) for the same star Betelgeuse. [∞ \(Sec. 5.4 and 5.6\)](#)

Once a star's angular size has been measured, if its distance is also known, we can determine its radius by simple geometry. [∞ \(Sec. 1.6\)](#) For example, with a distance of 130 pc and an angular diameter of up to 0.045", Betelgeuse's maximum radius is 630 times that of the Sun. (We say "maximum radius" here because, as it happens, Betelgeuse is a *variable star*—its radius and luminosity vary somewhat irregularly, with a period of roughly 6 years.) All told, the sizes of perhaps a few dozen stars have been measured in this way.

Most stars are too distant or too small for such direct measurements to be made. Instead, their sizes must be inferred by indirect means, using



◀ **FIGURE 17.11 Betelgeuse** The swollen star Betelgeuse (shown here in false color) is close enough for us to resolve its size directly, along with some surface features thought to be storms similar to those that occur on the Sun. (a) An ultraviolet view of Betelgeuse, as seen by a European camera onboard the *Hubble Space Telescope*, nearly resolves this huge star. (b) An infrared image, acquired by a three-telescope interferometer in Arizona does better, showing two spots on Betelgeuse's surface. (ESA/NASA; SAO)

## MORE PRECISELY 17-2

### Estimating Stellar Radii

We can combine the Stefan–Boltzmann law  $F = \sigma T^4$  with the formula for the area of a sphere,  $A = 4\pi R^2$ , to obtain the relationship between a star’s radius ( $R$ ), luminosity ( $L$ ), and temperature ( $T$ ) described in the text:

$$L = 4\pi\sigma R^2 T^4,$$


or

$$\text{luminosity} \propto \text{radius}^2 \times \text{temperature}^4.$$

If we adopt convenient “solar” units, in which  $L$  is measured in solar luminosities ( $3.9 \times 10^{26} \text{ W}$ ),  $R$  in solar radii (696,000 km), and  $T$  in units of the solar temperature (5800 K), we can eliminate the constant  $4\pi\sigma$  and write this equation as

$$R^2 (\text{in solar radii}) \times T^4 (\text{in units of } 5800 \text{ K}).$$

As illustrated in the accompanying figure, both the radius and the temperature are important in determining the star’s luminosity.



To compute the radius of a star from its luminosity and temperature, we rearrange terms so that the equation reads (in the same units)

$$R = \sqrt{L} / T^2$$

This simple application of the radiation laws is the basis for almost every estimate of stellar size made in this text. Let’s illustrate the process by computing the radii of two stars discussed in the text.

**Examples** In the solar units defined above, the star Aldebaran has luminosity  $L = 1.3 \times 10^{29} \text{ W} / 3.9 \times 10^{26} \text{ W} = 330$  units and temperature  $T = 4000 \text{ K} / 5800 \text{ K} = 0.69$  unit. Thus, according to the equation, its radius is  $R = \sqrt{330 / 0.69^2} = 18 / 0.48$  units = 39 solar radii—Aldebaran is a giant star. At the opposite extreme, Procyon B has  $L = 2.3 \times 10^{23} \text{ W} / 3.9 \times 10^{26} \text{ W} = 0.0006$  unit and  $T = 8500 \text{ K} / 5800 = 1.5$  units, so its radius is  $R = \sqrt{0.0006 / 1.5^2} = 0.01$  times the radius of the Sun—Procyon B is a dwarf.

The following table lists luminosities, temperatures, and calculated radii (all in solar units) for some of the other stars mentioned in this chapter.

Star	Luminosity (L)	Temperature (T)	Radius (R)
Sirius B	0.025	4.7	0.007
Barnard’s star	0.0045	0.56	0.2
Sun	1	1	1
Sirius A	23	2.1	1.9
Vega	55	1.6	2.8
Arcturus	160	0.78	21
Rigel	63,000	1.9	70
Betelgeuse	36,000	0.55	630

Note that some of the luminosities shown in the table differ significantly from those in Appendix 6, Tables 5 and 6, and elsewhere in the text. This is because the values used elsewhere refer to visible light only, whereas the radiation laws (and the values in the table) refer to *total* luminosities. As we saw in Chapter 3, a star radiates its energy over a broad range of wavelengths that often extends well beyond the visible domain. [∞ \(Sec. 3.4\)](#) A star like the Sun, whose emission happens to peak near the middle of the visible spectrum, emits most (roughly 80 percent) of its energy in the form of visible light. However, the cooler Aldebaran emits the bulk (about 75 percent) of its energy at infrared wavelengths, while the hot white dwarf Sirius B shines mainly in the ultraviolet—only about 10 percent of its energy is visible.

the radiation laws. [∞ \(Sec. 3.4\)](#) The radiation emitted by a star is governed by the Stefan–Boltzmann law, which states that the energy emitted *per unit area* per unit of time increases as the fourth power of the star’s surface temperature. [∞ \(More Precisely 3-2\)](#) To determine the star’s luminosity, we must multiply by its surface area—large bodies radiate more energy than do small bodies at the same temperature. Because the surface area is proportional to the square of the radius, we have


$$\text{luminosity} \propto \text{radius}^2 \times \text{temperature}^4.$$

This **radius–luminosity–temperature relationship** is important because it demonstrates that knowledge of a star’s

luminosity and temperature can yield an estimate of the star’s radius—an *indirect* determination of stellar size.

### Giants and Dwarfs

Let’s consider some examples to clarify these ideas. The star known as Aldebaran (the orange-red “eye of the bull” in the constellation Taurus) has a surface temperature of about 4000 K and a luminosity of  $1.3 \times 10^{29} \text{ W}$ . Thus, its surface temperature is 0.7 times and its luminosity about 330 times, the corresponding quantities for our Sun. The radius–luminosity–temperature relationship (see *More Precisely 17-2*) then implies



▲ **FIGURE 17.12 Stellar Sizes** Illustrated here are the different sizes of several well-known stars. Only part of the red-giant star Antares can be shown on this scale, and the supergiant Betelgeuse would fill the entire page. (Here and in other figures, the symbol “ $\odot$ ” stands for the Sun, so the symbol “ $R_{\odot}$ ” means “solar radius.”)

that the star’s radius is almost 40 times the solar value. If our Sun were that large, its photosphere would extend halfway to the orbit of Mercury and, seen from Earth, would cover more than 20 degrees on the sky. A star as large as Aldebaran is known as a *giant*. More precisely, **giants** are stars having radii between 10 and 100 times that of the Sun. Since any 4000-K object is reddish in color, Aldebaran is known as a **red giant**. Even larger stars, ranging up to 1000 solar radii in size, are known as **supergiants**. Betelgeuse is a prime example of a **red supergiant**.

Now consider Procyon B, a faint companion to Procyon A, one of the brightest stars in the night sky (see Appendix 3, Table 5). Procyon B’s surface temperature is roughly 8500 K, about one and a half times that of the Sun. The star’s total luminosity is  $2.3 \times 10^{25}$  W, about 0.0006 times the solar value. Again using the radius–luminosity–temperature relationship, we obtain a radius of 0.01 solar radii—slightly

larger than that of Earth. Procyon B is hotter, but smaller and much less luminous, than our Sun. Such a star is known as a *dwarf*. In astronomy, the term **dwarf** refers to any star of radius comparable to or smaller than the Sun (including the Sun itself). Because any 8500 K object glows white-hot, Procyon B is an example of a **white dwarf**.

The radii of the vast majority of stars (measured mostly with the radius–luminosity–temperature relationship) range from less than 0.01 to over 100 times the radius of the Sun. Figure 17.12 illustrates the estimated sizes of a few well-known stars.

### CONCEPT Check



- ✓ Can we measure the radius of a star without knowing the star’s distance from Earth?

## 17.5 The Hertzsprung–Russell Diagram

Astronomers use luminosity and surface temperature to classify stars in much the same way that height and weight serve to classify the bulk properties of human beings. We know that people’s height and weight are well correlated: Tall people tend to weigh more than short ones. We might naturally wonder if the two basic stellar properties are also related in some way.

Figure 17.13 plots luminosity versus temperature for several well-known stars. Figures of this sort are called *Hertzsprung–Russell diagrams*, or **H–R diagrams**, after Danish astronomer Ejnar Hertzsprung and U.S. astronomer Henry Norris Russell, who independently pioneered the use of such plots in the second decade of the 20th century. The vertical scale, expressed in units of the solar luminosity ( $3.9 \times 10^{26}$  W), extends over a large range, from  $10^{-4}$  to  $10^4$ ; the Sun appears right in the middle of the luminosity range, at a luminosity of 1. Surface temperature is plotted on the horizontal axis, although in the unconventional sense of temperature increasing to the *left* (so that the spectral sequence O, B, A... reads from left to right). To change the horizontal scale so that temperature would increase conventionally to the right would play havoc with historical precedent.

As we have just seen, astronomers often use a star’s *color* to measure its temperature. Indeed, the spectral classes plotted along the horizontal axis of Figure 17.13 are equivalent to the B/V color index. Also, because astronomers commonly express a star’s luminosity as an absolute magnitude, stellar *magnitude* instead of stellar luminosity could be plotted on the vertical axis (see *More Precisely 17-1*). Many astronomers prefer to present their data in these more “observational” terms, and the diagrams corresponding to plots like Figure 17.13 are called **color-magnitude diagrams**. In this book, we will cast our



**MA** **Interactive FIGURE 17.13 H–R Diagram of Well-Known Stars** A plot of luminosity against surface temperature (or spectral class) is a useful way to compare stars. Plotted here are the data for some stars mentioned earlier in the text. The Sun, which has a luminosity of 1 solar unit and a temperature of 5800 K, is a G-type star. The B-type star Rigel, at top left, has a temperature of about 11,000 K and a luminosity more than 10,000 times that of the Sun. The M-type star Proxima Centauri, at bottom right, has a temperature of about 3000 K and a luminosity less than  $\frac{1}{10,000}$  that of the Sun. (See also Overlay 1 of the acetate insert.)

**MA** **Interactive FIGURE 17.14 H–R Diagram of Nearby Stars** Most stars have properties within the long, thin, shaded region of the H–R diagram known as the main sequence. The points plotted here are for stars lying within about 5 pc of the Sun. Each dashed diagonal line corresponds to a constant stellar radius. (Recall that the symbol “ $R_{\odot}$ ” means “solar radius.”)

discussion mainly in terms of the more “theoretical” quantities, temperature and luminosity, but realize that, for many purposes, color-magnitude and H–R diagrams represent pretty much the same thing.


## The Main Sequence

The handful of stars plotted in Figure 17.13 gives little indication of any particular connection between stellar properties. However, as Hertzsprung and Russell plotted more and more stellar temperatures and luminosities, they found that a relationship does in fact exist: Stars are *not* uniformly scattered across the H–R diagram; instead, most are confined to a fairly well-defined band stretching diagonally from the top left (high temperature, high luminosity) to the bottom right (low temperature, low luminosity). In other words, cool stars tend to be faint (less luminous) and hot stars tend to be bright (more luminous). This band of stars spanning the H–R diagram is known as the **main sequence**.

Figure 17.14 shows a more systematic study of stellar properties, covering the 80 or so stars that lie within 5 pc of the Sun. As more points are included in the diagram, the main sequence “fills up,” and the pattern becomes more evident. The vast majority of stars in the immediate vicinity of the Sun lie on the main sequence.

The surface temperatures of main-sequence stars range from about 3000 K (spectral class M) to over 30,000 K (spectral class O). This relatively small temperature range—a difference of only a factor of 10—is determined mainly by the rates at which nuclear reactions occur in stellar cores.  $\infty$  (Sec. 16.6) In contrast, the observed range in luminosities is very large, covering some eight orders of magnitude (i.e., a factor of 100 million), ranging from  $10^{-4}$  to  $10^4$  times the luminosity of the Sun.

Using the radius–luminosity–temperature relationship (Section 17.4), astronomers find that stellar radii also vary along the main sequence. The faint, red M-type stars at the bottom right of the H–R diagram are only about one-tenth the size of the Sun, whereas the bright, blue O-type stars in the upper left are about 10 times larger than the Sun. The diagonal dashed lines in Figure 17.14 represent constant stellar radii, meaning that any star lying on a given line has the same radius, regardless of its luminosity or temperature.



▲ **FIGURE 17.15 H-R Diagram of Brightest Stars** An H-R diagram for the 100 brightest stars in the sky is biased in favor of the most luminous stars—which appear toward the upper left—because we can see them more easily than we can the faintest stars. (Compare with Figure 17.14, which shows only the closest stars.)

Along a constant-radius line, the radius–luminosity–temperature relationship implies that

$$\text{luminosity} \propto \text{temperature}^4.$$

By including such lines on our H-R diagrams, we can indicate stellar temperatures, luminosities, and radii on a single plot.

We see a very clear trend as we traverse the main sequence from top to bottom. At one end, the stars are large, hot, and bright. Because of their size and color, they are referred to as **blue giants**. The very largest are called **blue supergiants**. At the other end, stars are small, cool, and faint. They are known as **red dwarfs**. Our Sun lies right in the middle.

Figure 17.15 shows an H-R diagram for a different group of stars—the 100 stars of known distance having the greatest apparent brightness, as seen from Earth. Notice the much larger number of very luminous stars at the upper end of the main sequence than at the lower end. The reason for this excess of blue giants is simple: We can see very luminous stars a long way off. The stars shown in this figure are scattered through a much greater volume of space than those depicted in Figure 17.14, but the sample is heavily biased toward the brightest objects. In fact, of the 20 brightest stars in the sky, only 6 lie within 10 pc of us; the rest are visible, despite their great distances, because of their high luminosities.

If very luminous blue giants are overrepresented in Figure 17.15, low-luminosity red dwarfs are surely underrepresented. In fact, no dwarfs appear on the diagram. This absence is not surprising because low-luminosity stars are difficult to observe from Earth. In the 1970s, astronomers began to realize that they had greatly underestimated the number of red dwarfs in our galaxy. As hinted at by the H-R diagram in Figure 17.14, which shows an unbiased sample of stars in the solar neighborhood, red dwarfs are actually the most common type of star in the sky. In fact, they probably account for upward of 80 percent of all stars in the universe. In contrast, O- and B-type supergiants are extremely rare, with only about 1 star in 10,000 falling into these categories.


## White Dwarfs and Red Giants

Most stars lie on the main sequence. However, some of the points plotted in Figures 17.13 through 17.15 clearly do not. One such point in Figure 17.13 represents Procyon B, the white dwarf discussed earlier (Section 17.4), with surface temperature 8500 K and luminosity about 0.0006 times the solar value. A few more such faint, hot stars can be seen in Figure 17.14 in the bottom left-hand corner of the H-R diagram. This region, known as the **white-dwarf region**, is marked on Figure 17.14.

Also shown in Figure 17.13 is Aldebaran (discussed in Section 17.4), whose surface temperature is 4000 K and whose luminosity is some 300 times greater than the Sun's. Another point represents Betelgeuse (Alpha Orionis), the ninth-brightest star in the sky, a little cooler than Aldebaran, but more than 100 times brighter. The upper right-hand corner of the H-R diagram, where these stars lie (marked on Figure 17.15), is called the **red-giant region**. No red giants are found within 5 pc of the Sun (Figure 17.14), but many of the brightest stars seen in the sky are in fact red giants (Figure 17.15). Though relatively rare, red giants are so bright that they are visible to great distances. They form a third distinct class of stars on the H-R diagram, very different in their properties from both main-sequence stars and white dwarfs.

The *Hipparcos* mission (Section 17.1), in addition to determining hundreds of thousands of stellar parallaxes with unprecedented accuracy, also measured the colors and luminosities of more than 2 million stars. Figure 17.16 shows an H-R diagram based on a tiny portion of the enormous *Hipparcos* dataset. The main-sequence and red-giant regions are clearly evident. Few white dwarfs appear, however, simply because the telescope was limited to observations of relatively bright objects—brighter than apparent magnitude 12. Almost no white dwarfs lie close enough to Earth that their magnitudes fall below this limit.

About 90 percent of all stars in our solar neighborhood, and probably a similar percentage elsewhere in the universe, are main-sequence stars. About 9 percent of stars are white dwarfs, and 1 percent are red giants.



▲ **FIGURE 17.16 Hipparcos H–R Diagram** This simplified version of one of the most complete H–R diagrams ever compiled represents more than 20,000 data points, as measured by the European *Hipparcos* spacecraft for stars within a few hundred parsecs of the Sun.

### CONCEPT Check

- ✓ Only a tiny fraction of all stars are giants. Why, then, do giants account for so many of the brightest stars in the night sky?

## 17.6 Extending the Cosmic Distance Scale

We have already discussed the connections between luminosity, apparent brightness, and distance. Knowledge of a star’s apparent brightness and its distance allows us to determine its luminosity using the inverse-square law. But we can also turn the problem around. If we somehow knew a star’s luminosity and then measured its apparent brightness, the inverse-square law would tell us its distance from the Sun.

### Spectroscopic Parallax

Most of us have a rough idea of the approximate intrinsic brightness (that is, the luminosity) of a typical traffic signal. Suppose you are driving down an unfamiliar street and see a red traffic light in the distance. Your knowledge of the light’s luminosity enables you immediately to make

a mental estimate of its distance. A light that appears relatively dim (low apparent brightness) must be quite distant (assuming it’s not just dirty). A bright one must be relatively close. Thus, *measurement of the apparent brightness of a light source, combined with some knowledge of its luminosity, can yield an estimate of its distance.*

For stars, the trick is to find an independent measure of the luminosity without knowing the distance. The H–R diagram can provide just that. For example, suppose we observe a star and determine its apparent magnitude to be 10. By itself, that doesn’t tell us much—the star could equally well be faint and close, or bright and distant (Figure 17.6). But suppose we have some additional information: The star lies on the main sequence and has spectral type A0. Then we can read the star’s luminosity off Figure 17.15. A main-sequence A0 star has a luminosity of approximately 100 solar units. According to *More Precisely 17-1*, this corresponds to an absolute magnitude of 0 and hence to a distance of 1000 pc.


This process of using stellar spectra to infer distances is called **spectroscopic parallax**.<sup>\*</sup> The key steps are as follows:

1. We measure the star’s apparent brightness and spectral type *without* knowing how far away it is.
2. Then we use the spectral type to estimate the star’s luminosity.
3. Finally, we apply the inverse-square law to determine the distance to the star.

The existence of the main sequence allows us to make a connection between an easily measured quantity (spectral type) and the star’s luminosity, which would otherwise be unknown. The term *spectroscopic parallax* refers to the specific process of using stellar spectra to infer luminosities and hence distances. However, as we will see in upcoming chapters, this essential logic (with a variety of different techniques replacing step 2) is used again and again as a means of distance measurement in astronomy. In practice, the “fuzziness” of the main sequence translates into a small (10–20 percent) uncertainty in the distance, but the basic idea remains valid.

In Chapter 2 we introduced the first “rung” on a ladder of distance-measurement techniques that will ultimately carry us to the edge of the observable universe. That rung is radar ranging on the inner planets. ∞ (Sec. 2.6) It establishes the scale of the solar system and defines the astronomical unit. In Section 17.1, we discussed a second rung in the cosmic distance ladder—stellar parallax—which is based on the first, since Earth’s orbit is the baseline. Now, having used the first two rungs to determine the distances and other physical properties of

<sup>\*</sup>This unfortunate name is very misleading, as the method has nothing in common with stellar (geometric) parallax other than its use as a means of determining stellar distances.



▲ **FIGURE 17.17 Stellar Distances** Knowledge of a star's luminosity and apparent brightness can yield an estimate of its distance. Astronomers use this third rung on the distance ladder, called spectroscopic parallax, to measure distances as far out as individual stars can be clearly discerned—several thousand parsecs.

many nearby stars, we can employ that knowledge in turn to construct a third rung in the ladder—spectroscopic parallax. As illustrated schematically in Figure 17.17, this new rung expands our cosmic field of view still deeper into space.

Spectroscopic parallax can be used to determine stellar distances out to several thousand parsecs. Beyond that, spectra and colors of individual stars are difficult to obtain. The “standard” main sequence is obtained from H–R diagrams of stars whose distances can be measured by (geometric) parallax, so the method of spectroscopic parallax is calibrated by using nearby stars. Note that, in employing this method, we are assuming (without proof) that distant stars are basically similar to nearby stars and that *they fall on the same main sequence as nearby stars*. Only by making this assumption can we expand the boundaries of our distance-measurement techniques.

Of course, the main sequence is not really a line in the H–R diagram: It has some thickness. For example, the luminosities of main-sequence stars categorized as type A0 (Vega, for example) can actually range from about 30 to 100 times the luminosity of the Sun. The main reason for this range is the variation in stellar composition and age from place to place in our Galaxy. As a result, there is considerable uncertainty in the luminosity obtained by this method and hence a corresponding uncertainty in the distance of the star. Distances obtained by spectroscopic parallax are generally accurate to no better than about 25 percent.

Although this may not seem very accurate—a cross-country traveler in the United States would hardly be impressed to be told that the best estimate of the distance between Los Angeles and New York is somewhere between 3000 and 5000 km—it illustrates the point that, in astronomy, even something as simple as the distance

to another star can be very difficult to measure. Still, an estimate with an uncertainty of  $\pm 25$  percent is far better than no estimate at all. The deployment of the next generation of astrometry satellites (Section 17.1) promises to remedy this situation, combining radical improvements in both the range and accuracy of stellar parallax measurements.

Finally, realize that, because each rung in the distance ladder is calibrated using data from the lower rungs, changes made at *any* level will affect measurements made on *all* larger scales. As a result, the impact of new high-quality observations, such as those made by the *Hipparcos* mission (Section 17.1), extends far beyond the volume of space actually surveyed. By recalibrating the local foundations of the cosmic distance scale, *Hipparcos* caused astronomers to revise their estimates of distances on *all* scales—up to and including the scale of the universe itself. All distances quoted throughout this text reflect updated values based on *Hipparcos* data.


## Luminosity Class

What if the star in question happens to be a red giant or a white dwarf and does not lie on the main sequence? Recall from Chapter 4 that detailed analysis of spectral *line widths* can provide information on the *density* of the gas where the line formed. ∞ (Sec. 4.5) The atmosphere of a red giant is much less dense than that of a main-sequence star, and this in turn is much less dense than the atmosphere of a white dwarf. Figure 17.18(b) and (c) illustrate the difference between the spectra of a main-sequence star and a red giant of the same spectral type.


Over the years, astronomers have developed a system for classifying stars according to the widths of their spectral lines. Because line width depends on density in the stellar photosphere, and because this density in turn is well correlated with luminosity, this stellar property is known as **luminosity class**. The standard luminosity classes are listed in Table 17.3 and shown on the H–R diagram in Figure 17.18(a). By determining a star's luminosity class,

**TABLE 17.3** Stellar Luminosity Classes

Class	Description
Ia	Bright supergiants
Ib	Supergiants
II	Bright giants
III	Giants
IV	Subgiants
V	Main-sequence stars and dwarfs



◀ **FIGURE 17.18 Luminosity Classes** (a) Approximate locations of the standard stellar luminosity classes in the H–R diagram. The widths of absorption lines also provide information on the density of a star’s atmosphere. The denser atmosphere of a main-sequence K-type star has broader lines (c) than a giant star of the same spectral class (b).



astronomers can usually tell with a high degree of confidence what sort of object it is. Now we have a way of specifying a star’s location in the diagram in terms of properties that are measurable by purely spectroscopic means: *Spectral type and luminosity class define a star on the H–R diagram just as surely as do temperature and luminosity.* The full specification of a star’s spectral properties includes its luminosity class. For example, the Sun, a G2 main-sequence star, is of class G2V, the B8 blue supergiant Rigel is B8Ia, the red dwarf Barnard’s star is M5V, the red supergiant Betelgeuse is M2Ia, and so on.

Consider, for example, a K2-type star (Table 17.4) with a surface temperature of approximately 4500 K. If the widths of the star’s spectral lines tell us that it lies on the main sequence (i.e., it is a K2V star), then its luminosity is about 0.3 times the solar value. If the star’s spectral lines are observed to be narrower than lines

normally found in main-sequence stars, the star may be recognized as a K2III giant, with a luminosity 100 times that of the Sun (Figure 17.18a). If the lines are very narrow, the star might instead be classified as a K2Ib supergiant, brighter by a further factor of 40, at 4000 solar luminosities. In each case, knowledge of luminosity classes allows astronomers to identify the object and make an appropriate estimate of its luminosity and hence its distance.

### CONCEPT Check

- ✓ Suppose astronomers discover that, due to a calibration error, *all* distances measured by geometric parallax are 10 percent larger than currently thought. What effect would this finding have on the “standard” main sequence used in spectroscopic parallax?

**TABLE 17.4** Variation in Stellar Properties within a Spectral Class

Surface Temperature (K)	Luminosity (solar luminosities)	Radius (solar radii)	Object	Example
4900	0.3	0.8	K2V main-sequence star	ε Eridani
4500	110	21	K2III red giant	Arcturus
4300	4000	140	K2Ib red supergiant	ε Pegasi



## 17.7 Stellar Masses


What ultimately determines a star's position on the main sequence? The answer is its *mass* and its *composition*. Mass and composition are fundamental properties of any star. Together, they uniquely determine the star's internal structure, its external appearance, and even (as we will see in Chapter 20) its future evolution. The ability to measure these two key stellar properties is of the utmost importance if we are to understand how stars work. We have already seen how spectroscopy is used to determine composition. ∞ (Sec. 16.3) Now let's turn to the problem of finding a star's mass.

As with all other objects, we measure a star's mass by observing its gravitational influence on some nearby body—another star, perhaps, or a planet. If we know the distance between the two bodies, then we can use Newton's laws to calculate their masses. The extrasolar planetary systems that have recently been detected have not been studied well enough to provide independent stellar mass measurements, and we are a long way from placing our own spacecraft in orbit around other stars. ∞ (Sec. 15.5) Nevertheless, there are ways of determining stellar masses.

### Binary Stars


Most stars are members of *multiple-star systems*—groups of two or more stars in orbit around one another. The majority of stars are found in **binary-star systems**, which consist of two stars in orbit about a common center of mass, held together by their mutual gravitational attraction. ∞ (Sec. 2.7) Other stars are members of triple, quadruple, or even more complex systems. The Sun is not part of a multiple-star system—if it has anything at all uncommon about it, it may be its lack of stellar companions.

Astronomers classify binary-star systems (or simply *binaries*) according to their appearance from Earth and the ease with which they can be observed. **Visual binaries** have widely separated members that are bright enough to be observed and monitored separately, as shown in




### Interactive FIGURE 17.20 Spectroscopic Binary

MA Properties of binary stars can be determined by measuring the periodic Doppler shift of one star relative to the other while moving in their orbits. This is a single-line system, in which only one spectrum (from the brighter component) is visible.



▲ **FIGURE 17.19 Visual Binary** The period and separation of a binary-star system can be observed directly if each star is clearly seen. At the left is an orbital diagram for the double star Kruger 60; at the right are actual photographs taken in some of the years indicated. (Harvard College Observatory)

Figure 17.19. The more common **spectroscopic binaries** are too distant to be resolved into separate stars, but they can be indirectly perceived by monitoring the back-and-forth Doppler shifts of their spectral lines as the stars orbit each other. Recall that motion toward an observer shifts the lines toward the blue end of the electromagnetic spectrum and motion away from the observer shifts them toward the red end. ∞ (Sec. 3.5) In a *double-line* spectroscopic binary, two distinct sets of spectral lines—one for each star—shift back and forth as the stars move. Because we see particular lines alternately approaching and receding, we know that the objects emitting the lines are in orbit. In the more common *single-line* systems, such as that shown in Figure 17.20, one star is too faint for its



**Interactive FIGURE 17.21 Eclipsing Binary** If the two stars in a binary-star system eclipse one another, additional information on their radii and masses can be obtained by observing the periodic decrease in starlight as one star passes in front of the other.

spectrum to be distinguished, so only one set of lines is observed to shift back and forth. The shifting means that the star that is observed must be in orbit around another star, even though the companion cannot be observed directly. (If this idea sounds familiar, it should—all of the extrasolar planetary systems discovered to date are extreme examples of single-line spectroscopic binaries.)

∞ (Sec. 15.5)

In the much rarer **eclipsing binaries**, the orbital plane of the pair of stars is almost edge-on to our line of sight. In this situation, depicted in Figure 17.21, we observe a periodic decrease in starlight as one star passes in front of (transits) the other. By studying the variation in the light from the binary system—the binary’s **light curve**—astronomers can derive detailed information not only about the stars’ orbits and masses, but also about their radii. Thus, eclipsing binaries provide an alternative means of measuring stellar radii that is *independent* of either the direct or the indirect methods described in Section 17.4.

For example, in the sequence shown in Figure 17.21, the maximum brightness (frames 1, 3, and 5) represents the combined brightnesses of the two stars, whereas the shallower minimum (frame 4) represents the brighter (larger) component only. These two pieces of information allow us to infer the individual brightnesses of the two stars. The deeper minima (frames 2 and 6) occur because the fainter red star partially blocks the light of the much brighter yellow star. The change in brightness tells us what fraction of the brighter star is obscured, and that in turn tells us the ratio of the *areas* of the two stars and hence (since area is proportional to radius squared) the ratio of their radii. If we also knew the components’ orbital *speeds*—from Doppler measurements, say—then the widths of the minima and the time taken to go from minimum to maximum light would tell us the actual radii of the stars.

The preceding categories of binary-star systems are not mutually exclusive. For example, a single-line spectroscopic binary may also happen to be an eclipsing system. In that case, astronomers can use the eclipses to gain extra information about the fainter member of the pair. Occasionally, two unrelated stars just happen to lie close together in the sky, even though they are actually widely separated. These *optical doubles* are just chance superpositions and carry no useful information about stellar properties.

## Mass Determination

By observing the actual orbits of the stars, the back-and-forth motion of the spectral lines, or the dips in the light curve—whatever information is available—astronomers can measure the binary’s orbital period. Observed periods span a broad range—from hours to centuries. How much additional information can be extracted depends on the type of binary involved.

If the distance to a *visual* binary is known, the semimajor axis of its orbit can be determined directly by simple geometry. Knowledge of the binary period and orbital semimajor axis is all we need to determine the combined mass of the component stars using the modified form of Kepler’s third law. ∞ (Sec. 2.7) Since the orbits of both stars can be separately tracked, it is also possible to determine each of the individual stars’ masses. Recall from Section 2.8 that, in any system of orbiting objects, each object orbits the common center of mass. Measuring the distance from each star to the center of mass of a visual binary yields the ratio of the stellar masses. Knowing both the sum of the masses and their ratio, we can then find the mass of each star.


For *spectroscopic* binaries, it is not possible to determine the semimajor axis directly. Doppler shift measurements give us information on the orbital velocities of the two stars, but only with regard to their radial components—that is, along the line of sight. As a result, we cannot determine the *inclination* of the orbit to our line of sight, and this imposes a limitation on how much information we can obtain—simply put, we cannot distinguish between a slow-moving binary seen edge-on and a fast-moving binary seen almost face-on (so that only a small component of the orbital motion is along the line of sight). We have already encountered this limitation in our study of extrasolar planets. ∞ (Sec. 15.6)

For a double-line spectroscopic system, individual radial velocities, and hence the ratio of the component masses, can be determined, but the uncertainty in the orbital inclination means that only lower limits on the individual masses can be obtained. For single-line

systems, even less information is available, and only a fairly complicated relation between the component masses (known as the *mass function*) can be derived. However, if, as is often the case, the mass of the brighter star can be determined by other means (e.g., if the brighter star is recognized as a main-sequence star of a certain spectral class—see Figure 17.22), a lower limit can then be placed on the mass of the fainter, unseen star.

Finally, if a spectroscopic binary happens also to be an *eclipsing* system, then the uncertainty in the inclination is removed, as the binary is known to be edge-on or very nearly so. In that case, both masses can be determined for a double-line binary. For a single-line system, the mass function is simplified to the point where the mass of the unseen star is known if the mass of the brighter star can be found by other means (e.g., by recognizing it as a main-sequence star of known spectral type).

Despite all these qualifications and difficulties, the masses of individual component stars have been obtained for many nearby binary systems. *More Precisely 17-3* presents a simple example of how this is accomplished in practice.




▲ **FIGURE 17.22 Stellar Masses** More than any other stellar property, mass determines a star's position on the main sequence. Low-mass stars are cool and faint; they lie at the bottom of the main sequence. Very massive stars are hot and bright; they lie at the top of the main sequence. ("M<sub>☉</sub>" means "solar mass.")

## 17.8 Mass and Other Stellar Properties

We end our introduction to the stars with a brief look at how mass is correlated with the other stellar properties discussed in this chapter. Figure 17.22 is a schematic H–R diagram showing how stellar mass varies along the main sequence. There is a clear progression from low-mass red dwarfs to high-mass blue giants. With few exceptions, main-sequence stars range in mass from about 0.1 to 20 times the mass of the Sun. The hot O- and B-type stars are generally about 10 to 20 times more massive than our Sun. The coolest K- and M-type stars contain only a few tenths of a solar mass. The mass of a star at the time of its formation determines the star's location on the main sequence. Based on observations of stars within a few hundred light-years of the Sun, Figure 17.23 illustrates how the masses of main-sequence stars are distributed. Notice the huge fraction of low-mass stars, as well as the tiny fraction contributed by stars of more than a few solar masses.

Figure 17.24 illustrates how a main-sequence star's radius and luminosity depend on its mass. The two plots shown, of the *mass–radius* and *mass–luminosity* relations, are based on observations of binary-star systems. Along the main sequence, both radius and luminosity increase with mass. As an approximate rule of thumb, we can say that radius increases proportionally to stellar mass, whereas luminosity increases much faster—almost as the *fourth*

*Small stars are less common than big ones, in much the same way that grains of sand on a beach greatly outnumber bigger rocks.*




▲ **FIGURE 17.23 Stellar Mass Distribution** The range of masses of main-sequence stars, as determined from careful measurement of stars in the solar neighborhood.

## MORE PRECISELY 17-3

### Measuring Stellar Masses in Binary Stars



As discussed in the text, most stars are members of binary systems—where two stars orbit one another, bound together by gravity. Here we describe—in an idealized case where the relevant orbital parameters are known—how we can use the observed orbital data, together with our knowledge of basic physics, to determine the masses of the component stars.

Consider the nearby visual binary system made up of the bright star Sirius A and its faint companion Sirius B, sketched in the accompanying figure. The binary's orbital period can be measured simply by watching the stars orbit one another, or alternatively by following the back-and-forth velocity wobbles of Sirius A due to its faint companion. It is almost exactly 50 years. The orbital semimajor axis can also be obtained by direct observation of the orbit, although in this case we must use some additional knowledge of Kepler's laws to correct for the binary's  $46^\circ$  inclination to the line of sight. [☞ \(Sec. 2.5\)](#) It is 20 AU—a measured angular size of  $7.5''$  at a distance of 2.7 pc. [☞ \(More Precisely 1-2\)](#) Once we know these two key orbital parameters, we can use the modified version of Kepler's third law to calculate the sum of the masses of the two stars. The result is  $20^3/50^2 = 3.2$  times the mass of the Sun. [☞ \(Sec. 2.8\)](#)



Further study of the orbit allows us to determine the individual stellar masses. Doppler observations show that Sirius A moves at approximately half the speed of its companion relative to their center of mass. [☞ \(Secs. 2.8, 3.5\)](#) This implies that Sirius A must have twice the mass of Sirius B. It then follows that the masses of Sirius A and Sirius B are 2.1 and 1.1 solar masses, respectively.

Often the calculation of the masses of binary components is complicated by the fact that only partial information is available—we might only be able to see one star, or perhaps only spectroscopic velocity information is available (see Section 17.7). Nevertheless, this technique of combining elementary physical principles with detailed observations is how virtually every stellar mass quoted in this text has been determined.



**▲ FIGURE 17.24 Stellar Radii and Luminosities** (a) Dependence of stellar radius on mass for main-sequence stars. Actual measurements show that the radius increases nearly in proportion to the mass over much of the range (as indicated by the straight line drawn through the data points). (b) Dependence of main-sequence luminosity on mass. The luminosity increases roughly as the fourth power of the mass (indicated again by the straight line).

**TABLE 17.5** Key Properties of Some Well-Known Main-Sequence Stars

Star	Spectral Type	Mass ( $M$ ) (solar masses)	Central Temperature ( $10^6$ K)	Luminosity ( $L$ ) (solar luminosities)	Estimated Lifetime ( $M/L$ ) ( $10^6$ years)
Spica B*	B2V	6.8	25	800	90
Vega	A0V	2.6	21	50	500
Sirius A	A1V	2.1	20	22	1000
Alpha Centauri	G2V	1.1	17	1.6	7000
Sun	G2V	1.0	15	1.0	10,000
Proxima Centauri	M5V	0.1	0.6	0.00006	16,000,000

\* The “star” Spica is, in fact, a binary system comprising a B1III giant primary (Spica A) and a B2V main-sequence secondary (Spica B).

power of the mass (as indicated by the line in Figure 17.24b). Thus, a 2-solar-mass main-sequence star has a radius about twice that of the Sun and a luminosity of 16 ( $2^4$ ) solar luminosities; a 0.2-solar-mass main-sequence star has a radius of roughly 0.2 solar radii and a luminosity of around 0.0016 ( $0.2^4$ ) solar luminosity.

Table 17.5 compares some key properties of several well-known main-sequence stars, arranged in order of decreasing mass. Notice that the central temperature (obtained from mathematical models similar to those discussed in Chapter 16) differs relatively little from one star to another, compared with the large spread in stellar luminosities. [∞ \(Sec. 16.2\)](#) The rapid rate of nuclear burning deep inside a star releases vast amounts of energy per unit time. How long can the fire continue to burn? We can estimate a main-sequence star’s *lifetime* simply by dividing the amount of fuel available (the mass of the star) by the rate at which the fuel is being consumed (the star’s luminosity):

$$\text{stellar lifetime} \propto \frac{\text{stellar mass}}{\text{stellar luminosity}}.$$

The mass–luminosity relation tells us that a star’s luminosity is roughly proportional to the fourth power of its mass, so we can rewrite this expression to obtain, approximately,

$$\text{stellar lifetime} \propto \frac{1}{(\text{stellar mass})^3}.$$

The final column in Table 17.5 lists estimated lifetimes, based on the above proportionality and noting that the lifetime of the Sun (see Chapter 20) is about 10 billion years.

For example, the lifetime of a 10-solar-mass main-sequence O-type star is roughly  $10/10^4 = 1/1000$  of the lifetime of the Sun, or about 10 million years. The nuclear reactions in such a massive star proceed so rapidly that its fuel is quickly depleted, despite its large mass. We can be sure that all the O- and B-type stars we now observe are quite young—less than a few tens of millions of years old. Massive stars older than that have already exhausted their fuel and no longer emit large amounts of energy. They have, in effect, died.

At the opposite end of the main sequence, the cooler K- and M-type stars have less mass than our Sun has. With their low core densities and temperatures, their proton–reactions churn away rather sluggishly, much more slowly than those in the Sun’s core. The small energy release per unit time leads to low luminosities for these stars, so they have very long lifetimes. Many of the K- and M-type stars we now see in the night sky will shine on for at least another trillion years. The evolution of stars—large and small—is the subject of Chapters 20 and 21.

### PROCESS OF SCIENCE Check

- ✓ How do we know the masses of stars that aren’t components of binaries?

**The Big Question** Our Sun will expand as it ages, and it is destined to balloon rapidly into a red giant as it begins running out of fuel in about 5 billion years. A burning question, often asked and then quickly dismissed as being too remote in time is, will the red-giant Sun expand enough to engulf Earth? No one is certain. We do know that the Sun is losing lots of matter, thereby lessening its gravitational pull. Perhaps that will allow Earth to recede eventually to a relatively safe orbit.

# Chapter Review

## SUMMARY





1 The distances to the nearest stars can be measured by trigonometric parallax. A star with a parallax of 1 arc second (10) is 1 **parsec** (p. 423)—about 3.3 light-years—away from Earth.

2 Stars have real motion through space as well as apparent motion as Earth orbits the Sun. A star's **proper motion** (p. 424)—its true motion across the sky—is a measure of the star's velocity perpendicular to our line of sight. The star's radial velocity—along the line of sight—is measured by the Doppler shift of the spectral lines emitted by the star.

3 The **apparent brightness** (p. 425) of a star is the rate at which energy from the star reaches a detector. Apparent brightness falls off as the inverse square of the distance. Optical astronomers use the **magnitude scale** (p. 426) to express and compare stellar brightnesses. The greater the magnitude, the fainter the star; a difference of five magnitudes corresponds to a factor of 100 in brightness. **Apparent magnitude** (p. 426) is a measure of apparent brightness. The **absolute magnitude** (p. 427) of a star is the apparent magnitude it would have if placed at a standard distance of 10 pc from the viewer. It is a measure of the star's luminosity.


4 Astronomers often measure the temperatures of stars by measuring their brightnesses through two or more optical filters and then fitting a black-body curve to the results. The measurement of the amount of starlight received through each member of a set of filters is called **photometry** (p. 429). Spectroscopic observations of stars provide an accurate means of determining both stellar temperatures and stellar composition. Astronomers classify stars according to the absorption lines in their spectra. The standard stellar **spectral classes** (p. 431), in order of decreasing temperature, are O, B, A, F, G, K, and M.

5 Only a few stars are large enough and close enough that their radii can be measured directly. The sizes of most stars are estimated indirectly through the **radius–luminosity–temperature relationship** (p. 433). Stars comparable in size to,




or smaller than, the Sun are categorized as **dwarfs** (p. 434), stars up to 100 times larger than the Sun are called **giants** (p. 434), and stars more than 100 times larger than the Sun are known as **supergiants** (p. 434). In addition to “normal” stars such as the Sun, two other important classes of star are **red giants** (p. 434) (and **red supergiants**) (p. 434), which are large, cool, and luminous, and **white dwarfs** (p. 434), which are small, hot, and faint.


6 A plot of stellar luminosities versus stellar spectral classes (or temperatures) is called an **H–R diagram** (p. 434), or a **color–magnitude diagram** (p. 434). About 90 percent of all stars plotted on an H–R diagram lie on the **main sequence** (p. 435), which stretches from hot, bright **blue supergiants** (p. 436) and **blue giants** (p. 436), through intermediate stars such as the Sun, to cool, faint **red dwarfs** (p. 436). Most main-sequence stars are red dwarfs; blue giants are quite rare. About 9 percent of stars are in the **white-dwarf region** (p. 436), and the remaining 1 percent are in the **red-giant region** (p. 436).



7 If a star is known to be on the main sequence, measurement of its spectral type allows its luminosity to be estimated and its distance to be measured. This method of determining distance, which is valid for stars up to several thousand parsecs from Earth, is called **spectroscopic parallax** (p. 437). A star's **luminosity class** (p. 438) allows astronomers to distinguish main-sequence stars from giants and supergiants of the same spectral type.



8 Most stars are not isolated in space, but instead orbit other stars in **binary-star systems** (p. 440). In a **visual binary** (p. 440), both stars can be seen and their orbit charted. In a **spectroscopic binary** (p. 440), the stars cannot be resolved, but their orbital motion can be detected spectroscopically. In an **eclipsing binary** (p. 441), the orbit is oriented in such a way that one star periodically passes in front of the other as seen from Earth and dims the light we receive. Studies of binary-star systems often allow stellar masses to be measured. The mass of a star determines the star's size, temperature, and brightness. Hot blue giants are much more massive than the Sun; cool red dwarfs are much less massive. High-mass stars burn their fuel rapidly and have much shorter lifetimes than the Sun. Low-mass stars consume their fuel slowly and may remain on the main sequence for trillions of years.



**MasteringAstronomy**<sup>®</sup> For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- LO1** How is parallax used to measure the distances to stars? What is a parsec?
- LO2** Explain two ways in which a star's real motion through space translates into motion that is observable from Earth.
- LO3** How do astronomers go about measuring stellar luminosities? What is the difference between luminosity and apparent brightness?
- How do astronomers measure stellar temperatures?
- LO4 POS** Briefly describe how stars are classified according to their spectral characteristics.
- LO5** Describe how astronomers measure stellar radii. List some characteristics of red-giant and white-dwarf stars.
- Why do some stars have very few hydrogen lines in their spectra?
- What information is needed to plot a star on the H–R diagram?
- LO6** What is the main sequence? What basic property of a star determines where it lies on the main sequence?
- Why does the H–R diagram constructed from data on the brightest stars differ so much from the diagram constructed from data on the nearest stars?
- LO7** How are distances determined by spectroscopic parallax?
- Which stars are most common in our Galaxy? Why don't we see many of them in H–R diagrams? Which stars are least common in our Galaxy?
- LO8 POS** How can stellar masses be determined by observing binary-star systems?
- High-mass stars start off with much more fuel than low-mass stars. Why don't high-mass stars live longer?
- POS** In general, is it possible to determine the age of an individual star simply by noting its position on an H–R diagram? Explain.

## Conceptual Self-Test: Multiple Choice

- VIS** If Earth's orbit around the Sun were smaller, the parallactic angle to the star shown in Figure 17.1 ("Stellar Parallax") would be (a) smaller; (b) larger; (c) the same.
- From a distance of 1 parsec, the angular size of Earth's orbit would be (a) 1 degree; (b) 2 degrees; (c) 1 arc minute; (d) 2 arc seconds.
- According to the inverse-square law, if the distance to a lightbulb increases by a factor of 5, the bulb's apparent brightness (a) stays the same; (b) becomes 5 times less; (c) becomes 10 times less; (d) becomes 25 times less.
- Compared with a star of absolute magnitude  $-2$  at a distance of 100 pc, a star of absolute magnitude 5 at a distance of 10 pc will appear (a) brighter; (b) fainter; (c) to have the same brightness; (d) bluer.
- VIS** Pluto's apparent magnitude is approximately 14. According to Figure 17.7 ("Apparent Magnitude"), Pluto can be seen (a) with the naked eye on a dark night; (b) using binoculars; (c) using a 1-m telescope; (d) only with the *Hubble Space Telescope*.
- Stars of spectral class M do not show strong lines of hydrogen in their spectra because (a) they contain very little hydrogen; (b) their surfaces are so cool that most hydrogen is in the ground state; (c) their surfaces are so hot that most hydrogen is ionized; (d) the hydrogen lines are swamped by even stronger lines of other elements.
- Cool stars can be very luminous if they are very (a) small; (b) hot; (c) large; (d) close to our solar system.
- VIS** According to Figure 17.13 ("H–R Diagram of Prominent Stars"), Barnard's star must be (a) hotter; (b) larger; (c) closer to us; (d) bluer than Proxima Centauri.
- VIS** Figure 17.15 ("H–R Diagram of Brightest Stars") shows Vega and Arcturus at approximately the same level on the vertical axis. This means that Arcturus must be (a) hotter than; (b) fainter than; (c) larger than; (d) of the same spectral class as Vega.
- The mass of a star may be determined (a) by measuring its luminosity; (b) by determining its composition; (c) by measuring its Doppler shift; (d) by studying its orbit around a binary companion.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

1. ● How far away is the star Spica, whose parallax is 0.0120? What would Spica's parallax be if it were measured from an observatory on Neptune's moon Triton as Neptune orbited the Sun?
2. ●● A star lying 20 pc from the Sun has proper motion of 0.50/yr. What is the star's transverse velocity? If the star's spectral lines are observed to be redshifted by 0.01 percent, calculate the magnitude of its three-dimensional velocity relative to the Sun.
3. ● What is the luminosity of a star having three times the radius of the Sun and a surface temperature of 10,000 K?
4. ●● Calculate the solar energy flux (energy received per unit area per unit time), as seen from a distance of 10 pc from the Sun. Compare your answer with the solar constant at Earth.
5. ●● Two stars—A and B, with luminosities 0.5 and 4.5 times the luminosity of the Sun, respectively—are observed to have the same apparent brightness. Which star is more distant, and how much farther away is it than the other?
6. ● A star has apparent magnitude 10.0 and absolute magnitude 2.5. How far away is it?
7. ●● Using the data shown in Figure 17.7, calculate the greatest distance at which a star like the Sun could be seen with (a) binoculars, (b) a typical 1-m telescope, (c) a 4-m telescope, and (d) the *Hubble Space Telescope*.
8. ●● Given that the Sun's lifetime is about 10 billion years, estimate the life expectancy of (a) a 0.2-solar mass, 0.01-solar luminosity red dwarf, (b) a 3-solar mass, 30-solar luminosity star, (c) a 10-solar mass, 1000-solar luminosity blue giant.

## Activities

### Collaborative

1. Estimate the total number of stars visible in the night sky. Each member of your group should be equipped with identical cardboard tubes—the tube at the center of a roll of paper towels or toilet paper is prosaic but perfect for the task. On a clear, moonless night, hold your tube up to your eye and count the total number of stars you can see. Do this several times, randomly choosing different areas of the sky and avoiding obvious obstacles like clouds and trees. Try to sample all directions roughly equally. Hold the tube steady during each star count. Be sure to allow time for your eyes to adapt to the dark—10 to 15 minutes at least—before taking any measurements. Add up all your measurements and divide by the total number of observations to calculate the average number of stars observed—call it  $n$ , say. You can convert this number into an estimate of the total number  $N$  of visible stars by multiplying by the square of the ratio

of the length  $L$  of the tube to its diameter  $D$ , that is:  $N = (L/D)^2 \times n$ . (Can you figure out where this formula came from?) Repeat your measurement at a variety of sites—a city, the suburbs, and a dark rural location at least. Can you understand why astronomers are so concerned about how light pollution affects their work?

### Individual

1. Every winter you can find an astronomy lesson in the evening sky. The Winter Circle is an asterism—or pattern of stars—made up of six bright stars in five different constellations: Sirius, Rigel, Betelgeuse, Aldebaran, Capella, and Procyon. These stars span nearly the entire range of colors (and therefore temperatures) possible for normal stars. Rigel is a B-type star, Sirius, A; Procyon, F; Capella, G; Aldebaran, K; and Betelgeuse, M. The color differences of these stars are easy to see. Why do you suppose there is no O-type star in the Winter Circle?





# The Interstellar Medium

# 18

## GAS AND DUST AMONG THE STARS

Stars and planets are not the only inhabitants of our Galaxy. The space around us harbors invisible matter throughout the dark voids between the stars. The density of this matter is extremely low—approximately a trillion trillion times less dense than matter in either stars or planets, far more tenuous than the best vacuum attainable on Earth. Only because the volume of interstellar space is so vast does its mass amount to anything at all.

So why bother to study this near-perfect vacuum? We do so for three important reasons. First, there is nearly as much mass in the “voids” among the stars as there is in the stars themselves. Second, interstellar space is the region out of which new stars are born. Third, interstellar space is also the region into which old stars expel their matter when they die. It is one of the most significant crossroads through which matter passes anywhere in our universe.

**The Big Picture** Interstellar space comprises a much bigger domain of real estate than anything yet studied in this book. Extending into deeper space for hundreds and even thousands of light-years, on scales much larger than stars and planets, the interstellar medium is the place where nature conducts many of its changes. Rich in gas and dust, yet spread extraordinarily thinly throughout the vast, dark regions among the stars, interstellar matter occasionally reveals itself in silhouette, glows as nebulae, and contracts to form new stars.

**LEFT:** This remarkable image—a visual, true-color photo taken by the *Hubble Space Telescope*—shows pillars of gas and dust within the Carina Nebula. These flimsy structures, about 7500 light-years away and extending a few light-years across (thus much bigger than our solar system), will not survive long; radiation from hidden stars is slowly destroying them. In about 100,000 years, a cluster of stars will form here. (*STScI*)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Summarize the composition and physical properties of the interstellar medium.
- 2 Describe the characteristics of emission nebulae, and explain their significance in the life cycle of stars.
- 3 List the basic properties of dark interstellar clouds.
- 4 Specify the radio techniques used to probe the nature of interstellar matter.
- 5 Explain the nature and significance of interstellar molecules.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.

## 18.1 Interstellar Matter

Figure 18.1 is a mosaic of photographs covering a much greater expanse of universal “real estate” than anything we have studied thus far. From our vantage point on Earth, the panoramic view shown here stretches all the way across the sky. On a clear night, it is visible to the naked eye as the Milky Way. In Chapter 23, we will come to recognize this band as the flattened disk, or *plane*, of our Galaxy.

The bright regions in this image are congregations of innumerable unresolved stars, merging together into a continuous blur at the resolution of the telescope. However, the dark areas are not simply “holes” in the stellar distribution. They are regions of space where *interstellar matter* obscures (blocks) the light from stars beyond, blocking from our view what would otherwise be a rather smooth distribution of bright starlight. Their very darkness means that they cannot easily be studied by the optical methods used to examine stellar matter. There is, quite simply, nothing to see!

### Gas and Dust

From Figure 18.1 (see also Figure 18.4), it is evident that interstellar matter is distributed very unevenly throughout space. In some directions, the obscuring matter is largely absent, allowing astronomers to study objects literally billions of parsecs from the Sun. In other directions, there are small amounts of interstellar matter, so the obscuration is moderate, preventing us from seeing objects more than a few thousand parsecs away, but still allowing us to study nearby stars. Still other regions are so heavily obscured that starlight from even relatively nearby stars is completely absorbed before reaching Earth.

The matter among the stars is collectively termed the **interstellar medium**. It is made up of two components—*gas* and *dust*—intermixed throughout all space. The gas is made up mainly of individual atoms, of average size  $10^{-10}$  m (0.1 nm) or so, and small molecules, no larger than about  $10^{-9}$  m across. Interstellar dust is more complex, consisting of clumps of atoms and molecules—not unlike chalk dust or the microscopic particles that make up smoke, soot, or fog.

Apart from numerous narrow atomic and molecular absorption lines, the gas alone does not block radiation to any great extent. The obscuration that is evident in Figure 18.1 is caused by the dust. Light from distant stars cannot penetrate the densest accumulations of interstellar dust any more than a car’s headlights can illuminate the road ahead in a thick fog.

### Extinction and Reddening

We can use its effect on starlight to measure both the amount and the size of interstellar dust. As a rule of thumb, a beam of light can be absorbed or scattered only by particles having diameters comparable to or larger than the wavelength of the radiation involved. Thus, a range of dust particle sizes will tend to block shorter wavelengths most effectively. Furthermore, even for particles of a given size, the amount of obscuration (that is, absorption or scattering) produced by particles of a given size *increases* with *decreasing* wavelength. The size of a typical interstellar dust particle—or **dust grain**—is about  $10^{-7}$  m (0.1  $\mu\text{m}$ ), comparable in size to the wavelength of visible light. Consequently, dusty regions of interstellar space are transparent to long-wavelength radio and infrared radiation, but opaque to shorter wavelength optical and ultraviolet radiation. The overall dimming of starlight by interstellar matter is called **extinction**.






▲ **FIGURE 18.1 Milky Way Mosaic** The Milky Way Galaxy photographed panoramically, across  $360^\circ$  of the entire southern and northern celestial sphere. This band, which constitutes the central plane of our Galaxy, contains high concentrations of stars, as well as interstellar gas and dust. The white box shows the field of view of Figure 18.4. (ESO/S. Brunier)

**Narrated FIGURE 18.2 Reddening** (a) Starlight passing through a dusty region of space is both dimmed and reddened, but spectral lines are still recognizable in the light that reaches Earth. (b) This dusty interstellar cloud, called Barnard 68, is opaque to visible light, except near its edges, where some light from background stars can be seen. The cloud spans about 0.5 light-year and lies about 520 light-years away. Frame (c) illustrates (in false color) how infrared radiation can penetrate Barnard 68. (ESO)

Because the interstellar medium is more opaque to short-wavelength radiation than to radiation of longer wavelengths, light from distant stars is preferentially robbed of its higher frequency (“blue”) components. Hence, in addition to being generally diminished in brightness, stars also tend to appear redder than they really are. This effect, known as **reddening**, is conceptually similar to the process that produces spectacular red sunsets here on Earth. ∞ (More Precisely 7-1)

As illustrated in Figure 18.2(a), extinction and reddening change a star’s apparent brightness and color. However, the patterns of absorption lines in the original stellar spectrum are still recognizable in the radiation reaching Earth, so the star’s spectral class can be determined. Astronomers can use this fact to study the interstellar medium. From a main-sequence star’s spectral and luminosity classes, astronomers learn the star’s true luminosity and color. ∞ (Secs. 17.5, 17.6) They then measure the degree to which the starlight has been affected by extinction and reddening en route to Earth, and this,



in turn, allows them to estimate both the numbers and the sizes of interstellar dust particles along the line of sight to the star. By repeating these measurements for stars in many different directions and at many different distances from Earth, astronomers have built up a picture of the distribution and overall properties of the interstellar medium in the solar neighborhood.

Reddening can be seen very clearly in Figure 18.2(b), which shows a type of compact, dusty interstellar cloud called a *globule*. (We will discuss such clouds in more detail in Section 18.3.) The center of this cloud, called Barnard 68, is opaque to all optical wavelengths, so starlight cannot pass through it. However, near the edges, where there is less intervening cloud matter, some light does make it through. Notice how stars seen through the cloud are both dimmed and reddened relative to those seen directly. Figure 18.2(c) shows the same cloud in the infrared part of the spectrum. Much more of the radiation gets through, but even here reddening (or its infrared equivalent) can be seen.

## Overall Density

Gas and dust are found everywhere in interstellar space—no part of our Galaxy is truly devoid of matter. However, the density of the interstellar medium is extremely low. Overall, the gas averages roughly  $10^6$  atoms per cubic meter—just 1 atom per cubic centimeter—although there are large variations from place to place: Densities ranging from  $10^4$  to  $10^9$  atoms/m<sup>3</sup> have been found. Matter this diffuse is far less dense than the best vacuum—about  $10^{10}$  molecules/m<sup>3</sup>—ever attained in laboratories on Earth.

Interstellar dust is even rarer. On average, there is only one dust particle for every trillion or so atoms—just  $10^{-6}$  dust particles per cubic meter, or 1000 per cubic kilometer. Some parts of interstellar space are so thinly populated that harvesting all the gas and dust in a region the size of Earth would yield barely enough matter to make a pair of dice.

How can such fantastically sparse matter diminish light radiation so effectively? The key is size—interstellar space is vast. The typical distance between stars (1 pc or so in the vicinity of the Sun) is much, much greater than the typical size of the stars themselves (around  $10^{-7}$  pc). Stellar and planetary sizes pale in comparison to the vastness of interstellar space. Thus, matter can accumulate, regardless of how thinly it is spread. For example, an imaginary cylinder 1 m<sup>2</sup> in cross section and extending from Earth to Alpha Centauri would contain more than 10 billion billion dust particles. (Sec. 17.1) Over huge distances, dust particles accumulate slowly, but surely, to the point at which they can effectively block visible light and other short-wavelength radiation. Even though the density of matter is very low, interstellar space in the vicinity of the Sun contains about as much mass as exists in the form of stars.

Despite their rarity, dust particles make interstellar space a *relatively* dirty place. Earth's atmosphere, by comparison, is about a million times cleaner. Our air is tainted by only one dust particle for about every billion billion ( $10^{18}$ ) atoms of atmospheric gas. If we could compress a typical parcel of interstellar space to equal the density of air on Earth, this parcel would contain enough dust to make a fog so thick that we would be unable to see our hand held at arm's length in front of us.

## Composition


The composition of interstellar gas is reasonably well understood from spectroscopic studies of absorption lines formed when light from a distant star interacts with gas along the observer's line of sight (see Section 18.3). In most cases, the elemental abundances detected in interstellar gas mirror those found in other astronomical objects, such as the Sun, the stars, and the jovian planets. Most of the gas—about 90 percent by number—is

atomic or molecular hydrogen; some 9 percent is helium, and the remaining 1 percent consists of heavier elements. The abundances of some heavy elements, such as carbon, oxygen, silicon, magnesium, and iron, are much lower in interstellar gas than in our solar system or in stars. The most likely explanation for this finding is that substantial quantities of these elements have been used to form the interstellar dust, taking them out of the gas and locking them up in a form that is much harder to observe.

In contrast to interstellar gas, the composition of interstellar dust is currently not very well known. We have some infrared evidence for silicates, graphite, and iron—the same elements that are underabundant in the gas—lending support to the theory that interstellar dust forms out of interstellar gas. The dust probably also contains some “dirty ice,” a frozen mixture of ordinary water ice contaminated with trace amounts of ammonia, methane, and other chemical compounds. This composition is quite similar to that of cometary nuclei in our own solar system. (Sec. 14.2)

## Dust Shape

Curiously, astronomers know the *shapes* of interstellar dust particles better than their composition. Although the minute atoms in the interstellar gas are basically spherical, the dust particles are not. Individual dust grains are apparently elongated or rodlike, as shown in Figure 18.3(a), although recent theoretical studies of how dust particles collide, stick, and break up suggest that their larger scale structure may be considerably more complex (Figure 18.3b).



▲ **FIGURE 18.3 Interstellar Dust** (a) Typical interstellar dust particles, as inferred from polarization studies, have sizes of only about one ten-thousandth of a millimeter, yet space contains enough of them to obscure our view in certain directions. (b) This result of a computer simulation shows how grains may grow as dust particles collide, stick, and fragment in interstellar space.

Astronomers infer this elongated structure from the fact that the light emitted by stars is dimmed and partially **polarized**, or aligned, by the intervening dust. Recall from Chapter 3 that light consists of electromagnetic waves composed of vibrating electric and magnetic fields.  $\infty$  (Sec. 3.2, Fig. 3.7) Normally, these waves are randomly oriented, and the radiation is said to be unpolarized. Stars emit unpolarized radiation from their photospheres. However, under the right conditions, the radiation can become polarized en route to Earth, with the electric fields all vibrating in roughly the same plane. One way in which this can happen is if the radiation interacts with an elongated dust grain, which tends to absorb electric waves vibrating parallel to its length.

Thus, if the light detected by our telescope is polarized, it is because some interstellar dust lies between the emitting object and Earth. Based on this reasoning, astronomers have determined not only that interstellar dust particles must be elongated in shape, but also that they tend to be *aligned* over large regions of space.

The alignment of the interstellar dust is the subject of ongoing research among astronomers. The current view, accepted by most, is that the dust particles are affected by a weak interstellar magnetic field, perhaps a million times weaker than Earth's field. Each dust particle responds to the field in much the same way that small iron filings are aligned by an ordinary bar magnet. Measurements of the blockage and polarization of starlight thus yield information about the size and shape of interstellar dust particles, as well as about magnetic fields in interstellar space.

### CONCEPT Check

✓ If space is a near-perfect vacuum, how can there be enough dust in it to block starlight?

## 18.2 Emission Nebulae


Figure 18.4 shows a magnified view of the central part of Figure 18.1 (the region indicated by the rectangle in the earlier figure), in the general direction of the constellation Sagittarius. The field of view is mottled with stars and interstellar matter. The patchiness of the obscuration is evident. In addition, several large fuzzy patches of light are clearly visible. These fuzzy objects, labeled M8, M16, M17, and M20, correspond to the 8th, 16th, 17th, and 20th objects in a catalog compiled by Charles Messier, an 18th-century French astronomer.\* Today they are known as **emission nebulae**—glowing clouds of hot interstellar matter. Figure 18.5 enlarges the left side of Figure 18.4, showing the nebulae more clearly.

### Observations of Emission Nebulae

Historically, astronomers have used the term **nebula** to refer to any “fuzzy” patch (bright or dark) on the sky—any region of space that was clearly distinguishable through a telescope, but not sharply defined, unlike a star or a planet. We now know that many (although not all) nebulae are clouds of interstellar dust and gas.



If a cloud happens to obscure stars lying behind it, we see it as a dark patch on a bright background, as in Figures 18.1, 18.2(b), and 18.4—a dark nebula. But if something within the cloud—a group of hot young stars, for example—causes it to glow, then we see a bright emission nebula

*\*Messier was actually more concerned with making a list of celestial objects that might be confused with comets, his main astronomical interest. However, the catalog of 109 “Messier objects” is now regarded as a much more important contribution to astronomy than any comets Messier discovered.*



► **FIGURE 18.4 Milky Way in Sagittarius** Enlargement of the central part of Figure 18.1, showing regions of brightness (vast fields of stars) as well as regions of darkness (where interstellar matter obscures the light from more distant stars). The field of view is about 35° across. Two of the emission nebulae discussed in the text are labeled. (ESO/S. Guisard)

▼ **FIGURE 18.5 Galactic Plane** A black-and-white photograph of part (about  $12^\circ$  across) of the region of the sky shown in Figure 18.4, showing stars, gas, and dust, as well as several distinct fuzzy patches of light known as emission nebulae. The plane of the Milky Way is marked with a white diagonal line. (*Harvard College Observatory*)




▲ **FIGURE 18.6 M20–M8 Region** A true-color enlargement of the bottom of Figure 18.5, showing M20 (top) and M8 (bottom) more clearly. The two nebulae are only a few degrees apart on the sky. (*R. Gendler*)

instead. The method of spectroscopic parallax applied to stars that are visible within the emission nebulae shown in Figure 18.5 indicates that their distances from Earth range from 1200 pc (M8) to 1800 pc (M16). ∞ (Sec. 17.6) Thus, all four nebulae are near the limit of visibility for any object embedded in the dusty galactic plane. M16, at the top left, is approximately 1000 pc from M20, near the bottom.

We can gain a better appreciation of these nebulae by examining progressively smaller fields of view. Figure 18.6 is an enlargement of the region near the bottom of Figure 18.5, showing M20 at the top and M8 at the bottom, only a few degrees away. Figure 18.7 is an enlargement of the top of Figure 18.6, presenting a close-up of M20 and its immediate environment. The total area of the close-up view displayed measures some 10 pc across. Emission nebulae are among the most spectacular objects in the entire universe, yet they appear only as small, undistinguished patches of light when viewed in the larger context of the Milky Way, as in Figure 18.4. Perspective is crucial in astronomy.

The emission nebulae shown in Figures 18.5–18.7 are regions of glowing, ionized gas. At or near the center of each is at least one newly formed hot O- or B-type star



▲ **Interactive FIGURE 18.7 Trifid Nebula** (a) Further enlargement of the top of Figure 18.6, showing only M20 and its interstellar environment. ∞ Called the Trifid Nebula because of the dust lanes (in black) that trisect its midsection, the nebula itself (in red) is about 20 light-years across. The blue reflection nebula is unrelated to the red emission nebula; it is caused by starlight reflected from intervening dust particles. (b) A false-color infrared image taken by the *Spitzer Space Telescope* reveals bright regions of star-forming activity mostly in those lanes of dust. (*AURA; NASA*)


producing huge amounts of ultraviolet light. As ultraviolet photons travel outward from the star, they ionize the surrounding gas. As electrons recombine with nuclei, they emit visible radiation, causing the gas to fluoresce, or glow. ∞ (Sec. 4.2) The reddish hue of these nebulae—and, in fact, of all emission nebulae—results when hydrogen atoms emit light in the red part of the visible spectrum. Specifically, it is caused by the emission of radiation at 656.3 nm—the H $\alpha$  line discussed in Chapter 4. ∞ (More Precisely 4-1) Other elements in the nebula also emit radiation as their electrons recombine, but because hydrogen is so plentiful, its emission usually dominates.

Woven through the glowing nebular gas, and plainly visible in Figures 18.5–18.7, are lanes of dark, obscuring dust. These **dust lanes** are part of the nebulae and are not just unrelated dust clouds that happen to lie along our line of sight. The bluish region visible in Figures 18.6 and 18.7 immediately above M20 is another type of nebula unrelated to the red emission nebula itself. Called a **reflection nebula**, it is caused by starlight scattered from dust particles in interstellar clouds located just off the line of sight between Earth and the bright stars within M20. Reflection nebulae appear blue for much the same reason that Earth’s daytime sky is blue: short-wavelength blue light is more easily scattered by interstellar matter back toward Earth and into our detectors. ∞ (More Precisely 7-1) Figure 18.8 sketches some of the key features of emission nebulae, illustrating the connection between the central stars, the nebula itself, and the surrounding interstellar medium.

Figure 18.9 shows enlargements of two of the nebulae visible in Figure 18.5. Notice again the hot, bright stars embedded within the glowing nebular gas and the predominant red coloration of the emitted radiation in parts (a) and (c). The relationship between the nebulae and their dust lanes is again evident in Figures 18.9(b) and (d), where regions of gas and dust are simultaneously silhouetted against background nebular emission and illuminated by foreground nebular stars.


The interaction between stars and gas is particularly striking in Figure 18.9(b). The three dark “pillars” visible in this spectacular *Hubble Space Telescope* image are part of the interstellar cloud from which the stars formed. The rest of the cloud in the vicinity of the new stars has already been heated and dispersed by their radiation in a process known as *photoevaporation*. The fuzz around the edges of the pillars, especially at the top right and center, is the result of this ongoing process. (See also an up-close view of another such pillar in M16 in the chapter-opening photo.) As photoevaporation continues, it eats away the less dense material first, leaving behind delicate sculptures composed of the denser parts of the original cloud, just as wind and water create spectacular structures in Earth’s deserts and shores by eroding away the softest rock. The process is a dynamic one: The pillars will eventually be destroyed, but probably not for another hundred thousand or so years.

Spectroscopists often refer to the *ionization state* of an atom by attaching a roman numeral to the chemical symbol for the atom—I for the neutral (that is, not ionized) atom, II for a singly ionized atom (an atom missing one electron), III



▲ **FIGURE 18.8 Nebular Structure** An emission nebula results when ultraviolet radiation from one or more hot stars ionizes part of an interstellar cloud. If starlight happens to encounter another dusty cloud, some of the radiation, particularly at the shorter wavelength blue end of the spectrum, may be scattered back toward Earth, forming a reflection nebula.





▲ **FIGURE 18.9 Emission Nebulae** Enlargements of selected portions of Figure 18.5. (a) M16, the Eagle Nebula and (b) a close-up of its huge pillars of cold gas and dust within, showing delicate sculptures created by the action of stellar ultraviolet radiation on the original cloud. (c) M8, the Lagoon Nebula and (d) a high-resolution view of its core, a region known as the Hourglass. The varied colors of the insets result from observations at different wavelengths: Green represents emission from hydrogen atoms, red emission from singly ionized sulfur, and blue emission from doubly ionized oxygen. (ESA; AURA; NASA)

for a doubly ionized atom (one missing two electrons), and so on. Because emission nebulae are composed mainly of ionized hydrogen, they are often referred to as **HII regions**. Regions of space containing primarily neutral (atomic) hydrogen are known as **HI regions**.

## Nebular Spectra

Most of the photons emitted by the recombination of electrons with atomic nuclei escape from the emission nebulae. Unlike the ultraviolet photons originally emitted by

the embedded stars, these reemitted photons do not have enough energy to ionize the nebular gas, so they pass through the nebula relatively unhindered. Some eventually reach Earth. By studying these lower-energy photons, we can learn much about the detailed properties of emission nebulae.

Because at least one hot star resides near the center of every emission nebula, we might think that the combined spectrum of the star and the nebula would be hopelessly confused. In fact, they are not: We can easily distinguish nebular spectra from stellar spectra because the physical conditions in stars and emission nebulae differ so greatly. In particular, emission nebulae are made of hot, thin gas that, as we saw in Chapter 4, yields detectable *emission* lines. [∞ \(Sec. 4.1\)](#) When our spectroscope is trained on a star, we see a familiar stellar spectrum, consisting of a blackbody-like continuous spectrum and absorption lines, together with superimposed emission lines from the nebular gas. When no star appears in the field of view, only the emission lines are seen. Analyses of nebular spectra show compositions close to those derived from observations of the Sun and other stars and elsewhere in the interstellar medium: Hydrogen is about 90 percent abundant by number, followed by helium at about 9 percent; the heavier elements together make up the remaining 1 percent.

Unlike stars, nebulae are large enough for their actual sizes to be measurable by simple geometry. Coupling this information on size with estimates of the amount of matter along our line of sight (as revealed by the nebula's total emission of light), we can find the nebula's density. [∞ \(Sec. 6.2\)](#) Generally, emission nebulae have only a few hundred particles, mostly protons and electrons, in each cubic centimeter—a density some  $10^{22}$  times lower than that of a typical planet. Spectral-line widths imply that the gas atoms and ions have temperatures around 8000 K. [∞ \(Sec. 4.5\)](#) Table 18.1 lists some vital statistics for each of the nebulae shown in Figure 18.5.

## “Forbidden” Lines

When astronomers first studied the spectra of emission nebulae, they found many lines that did not correspond to anything observed in terrestrial laboratories. For example,

in addition to the dominant red coloration just discussed, many nebulae emit light with a characteristic green color (see Figure 18.10). The greenish tint of portions of these nebulae puzzled astronomers in the early 20th century and defied explanation in terms of the properties of spectral lines known at the time, prompting speculation that the nebulae contained elements that were unknown on Earth. Some scientists even went so far as to invent the term “nebulium” for a supposed new element, much as the name helium came about when that element was first discovered in the Sun (recall also the fictitious element “coronium” from Chapter 16). [∞ \(Sec. 16.3\)](#)


Later, with a fuller understanding of the workings of the atom, astronomers realized that these lines did in fact result from electron transitions within the atoms of familiar elements, but under unfamiliar conditions that were not reproducible in laboratories. Astronomers now understand that the greenish tint in Figure 18.10(b) and (c) is caused by a particular electron transition in doubly ionized oxygen. However, the structure of oxygen is such that an ion in the higher energy state for this transition tends to remain there for a very long time—many hours, in fact—before dropping back to the lower state and emitting a photon. Only if the ion is left undisturbed during this time, and not kicked into another energy state by a random interaction with another atom or molecule in the gas, will the transition actually occur and the photon be emitted.

In a terrestrial experiment, no atom or ion is left undisturbed for long. Even in a “low-density” laboratory gas, there are many trillions of particles per cubic meter, and each particle undergoes millions of collisions with other gas particles every second. The result is that an ion in the particular energy state that produces the peculiar green line in the nebular spectrum never has time to emit its photon in the lab—collisions kick it into some other state long before that occurs. For this reason, the line is usually called *forbidden*, even though it violates no law of physics; it simply occurs on Earth with such low probability that it is never seen.

In a typical emission nebula, the density is so low that collisions between particles are extremely rare. There is plenty of time for the excited ion to emit its photon, so the forbidden line is produced. Numerous forbidden lines are known in nebular spectra. These lines remind us once again that the environment in the interstellar medium is

**TABLE 18.1** Some Nebular Properties

Object	Approximate Distance (pc)	Average Diameter (pc)	Density ( $10^6$ particles/ $m^3$ )	Mass (solar masses)	Temperature (K)
M8	1200	14	80	2600	7500
M16	1800	8	90	600	8000
M17	1500	7	120	500	8700
M20	1600	6	100	250	8200




**▲ FIGURE 18.10 Orion Nebula** (a) Lying some 1400 light-years from Earth, the Orion Nebula (M42) is visible to the naked eye as the fuzzy middle “star” of Orion’s sword. (b) Like all emission nebulae, the Orion Nebula consists of hot, glowing gas powered by a group of bright stars in the center. In addition to exhibiting red  $H\alpha$  emission, parts of the nebula show a slight greenish tint, caused by a so-called forbidden transition in ionized oxygen. (c) A high-resolution image shows rich detail in a region about 0.5 light-year across. Structural details are visible down to a level of  $0.1''$ , or 6 light-hours—a scale comparable to the dimensions of our solar system. (NASA; ESO)

very different from conditions on Earth and warn us of the potential difficulties involved in extending our terrestrial experience from our laboratories to the study of interstellar space.

Some regions of interstellar space contain extremely dilute, even hotter gas than is found within emission nebulae. Ultraviolet observations by space-based instruments have found that these superheated interstellar “bubbles,” making up the *intercloud medium*, may extend far into interstellar space beyond our local neighborhood and, conceivably, into the even vaster spaces among the galaxies. This high-temperature gas is probably the result of the violent expansion of debris from stars that exploded long ago. Somewhat like the Sun’s faint corona, these regions are dark despite their high temperatures because the density of matter there is very low. [∞ \(Sec. 16.3\)](#)

The Sun seems to reside in one such low-density region—a huge cavity called the “Local Bubble,” sketched in Figure 18.11. The Local Bubble contains about 200,000 stars and extends for nearly 100 pc. It was probably carved out by multiple supernova explosions (see Chapters 20 and 21) that occurred several hundred thousand years ago in the Scorpius–Centaurus association, a rich cluster of bright young stars. Perhaps our hominid ancestors may have seen these ancient events—stellar catastrophes as bright as the full Moon—that now aid modern astronomers.



**▲ FIGURE 18.11 Local Bubble** The Sun resides in a vast low-density region of space that engulfs us nearly spherically. This cavity was likely caused by stellar explosions long ago, which then heated the nearby interstellar gas and expelled it well out of the solar neighborhood. Several prominent stars in our nighttime sky are plotted in this artist’s conception, which depicts what the “bubble” might look like from afar.

**CONCEPT Check**

- ✓ If emission nebulae are powered by ultraviolet radiation from very hot (blue-white) stars, why do they appear red?

## 18.3 Dark Dust Clouds

Emission nebulae and even the much larger interstellar bubbles are only small components of the interstellar medium. Most of space—in fact, more than 99 percent of it—is devoid of such regions and contains no stars. It is simply cold and dark. Look again at Figure 18.4, or just ponder the evening sky. The dark regions are by far the most representative of interstellar space. The average temperature of a typical dark region of interstellar matter is about 100 K. Compare this with 273 K, at which water freezes, and 0 K, at which atomic and molecular motions cease. [∞ \(More Precisely 3-1\)](#)

Within these vast, dark interstellar voids lurks another type of astronomical object: the **dark dust cloud**. Dark dust clouds are even colder than their surroundings (with temperatures as low as a few tens of kelvins) and thousands or even millions of times denser. Along any given line of sight, their densities can range from  $10^7$  atoms/m<sup>3</sup> to more than  $10^{12}$  atoms/m<sup>3</sup> ( $10^6$  atoms/cm<sup>3</sup>). Dark dust clouds are often called *dense* interstellar clouds by researchers, but we must recognize that even these densest interstellar regions are barely denser than the best vacuum achievable in terrestrial laboratories. Still, it is because their density is much larger than the average



value of  $10^6$  atoms/m<sup>3</sup> in interstellar space that we can distinguish these clouds from the surrounding expanse of the interstellar medium.

### Obscuration of Visible Light



Interstellar clouds bear little resemblance to terrestrial clouds. Most are much bigger than our solar system, and some are many parsecs across. (Yet even so, they make up no more than a few percent of the entire volume of interstellar space.) Despite their name, these clouds are made up primarily of gas, just like the rest of the interstellar medium. However, their absorption of starlight is due almost entirely to the dust they contain.

Figure 18.12(a) shows a region called L977, in the constellation Cygnus. It is a classic example of a dark dust cloud. The dense globule Barnard 68, shown in Figure 18.2(b), is another. Some early (18th-century) observers thought that these dark patches on the sky were simply empty regions of space that happened to contain no bright stars. However, by the late 19th century, astronomers had discounted this idea. They realized that seeing clear spaces among the stars would be like seeing clear tunnels between the trees in a huge forest, and it was statistically impossible that so many tunnels would lead directly away from Earth.

Despite this realization, before the advent of radio astronomy astronomers had no direct means of studying clouds such as L977. Emitting no visible light, they are generally undetectable to the eye, except by the degree to which they dim starlight. However, as shown in Figure 18.12(b), the cloud's radio emission—in this case from carbon monoxide (CO) molecules contained within its volume—outlines the



◀ **FIGURE 18.12 Obscuration and Emission** (a) At optical wavelengths, this dark dust cloud (known as L977) can be seen only by its obscuration of background stars. (b) At radio wavelengths, it emits strongly in the CO molecular line, with the most intense radiation coming from the densest part of the cloud. (C. and E. Lada)



◀ **FIGURE 18.13** Dark Dust Cloud

The Ophiuchus dark dust cloud resides only 550 light-years away, surrounded by colorful stars and nebulae that are actually small illuminated parts of a much bigger, and invisible, molecular cloud engulfing much of the 6-degree-wide region shown. Many stages of star formation can be seen in this spectacular four-image mosaic. The dark cloud itself is “visible” only because it blocks light coming from stars behind it. Notice the cloud’s irregular shape, and especially its long “streamers” at upper left. The bright, giant star Antares, the (much more distant) star cluster M4, and a nearby blue reflection nebula are also noted. (*R. Gendler/J. Misti/S. Mazlin*)

cloud clearly at radio wavelengths, providing an indispensable tool for the study of such objects. We will return to the subject of molecular emission from interstellar clouds in Section 18.5.

Figure 18.13 is a spectacular wide-field image of another dark dust cloud. Taking its name from a neighboring star system, Rho Ophiuchus, this dust cloud resides relatively nearby—about 170 pc from the Sun—making it one of the most intensely studied regions of star formation in the Milky Way. Pockets of heavy blackness mark regions where the dust and gas are especially concentrated and the light from the background stars is completely obscured. Measuring several parsecs across, the Ophiuchus cloud is only a tiny part of the grand mosaic shown in Figure 18.1. Note that this cloud, like most interstellar clouds, is very irregularly shaped. Note especially the long “streamers” of (relatively) dense dust and gas at upper left. By contrast, the bright patches within the dark regions are foreground objects—emission nebulae and groups of bright stars. Some of them are part of the cloud itself, where newly formed stars near the edge of the cloud have created “hot spots” in the cold, dark gas. Others have no connection to the cloud and just happen to lie along the line of sight.


Dark and dusty interstellar clouds are sprinkled throughout our Galaxy. We can study them at optical wavelengths only if they happen to block the light emitted by more distant stars or nebulae. The dark outline of the L997 cloud in Figure 18.12(a) and the dust lanes visible in Figures

18.7 and 18.9 are good examples of this obscuration. Figure 18.14 shows another well-known, and particularly striking, example of such a cloud—the Horsehead Nebula in Orion. This curiously shaped finger of gas and dust projects out from the much larger dark cloud (called L1630) that fills the bottom half of the image and stands out clearly against the red glow of a background emission nebula. For reference, the stars and bright emission nebulae lie in front of the dark cloud; the red glow that silhouettes the Horsehead lies behind and above it.

## Absorption Spectra

Astronomers first became aware of the true extent of dark interstellar clouds in the 1930s, as they studied the optical spectra of distant stars. The gas in such a cloud absorbs some of the stellar radiation in a manner that depends on the cloud’s own temperature, density, and elemental abundance. The absorption lines thus produced contain information about dark interstellar matter, just as stellar absorption lines reveal the properties of stars. ∞ (Sec. 4.1)

Because the interstellar absorption lines are produced by cold, low-density gas, astronomers can easily distinguish them from the much broader absorption lines formed in stars’ hot lower atmospheres. ∞ (Sec. 4.5) Figure 18.15(a) illustrates how light from a star may pass through several interstellar clouds on its way to Earth. These clouds need not be close to the star, and, indeed, they usually are not.



**▲ FIGURE 18.14 Horsehead Nebula** (a) Located in the constellation Orion, not far from the Orion Nebula, this Horsehead Nebula is a striking example of a dark dust cloud, silhouetted against the bright background of an emission nebula. (b) A stunning image of the Horsehead, taken at highest resolution by the Very Large Telescope (VLT) in Chile. [🔗 \(Sec. 5.2\)](#) This nebular region is roughly 5000 light-years from Earth, in the constellation Orion. (Royal Observatory of Belgium; ESO)



Each absorbs some of the stellar radiation in a manner that depends on its own temperature, density, velocity, and elemental abundance. Figure 18.15(b) depicts part of a typical spectrum produced in this way.

The narrow absorption lines contain information about dark interstellar clouds, just as stellar absorption lines reveal the properties of stars and nebular emission lines tell us about conditions in hot nebulae. By studying these lines, astronomers can probe the cold depths of interstellar space. In most

cases, the elemental abundances detected in interstellar clouds mirror those found in other astronomical objects—perhaps not surprising, since (as we will see in Chapter 19) interstellar clouds are the regions that spawn emission nebulae and stars.

### PROCESS OF SCIENCE Check

- ✓ How do astronomers use optical observations to probe the properties of dark dust clouds?



**◀ FIGURE 18.15 Absorption by Interstellar Clouds**

(a) Simplified diagram of some interstellar clouds between a hot star and Earth. Optical observations might show an absorption spectrum like that traced in (b). The wide, intense lines are formed in the star's hot atmosphere; narrower, weaker lines arise from the cold interstellar clouds. The smaller the cloud, the weaker are the lines. The redshifts or blueshifts of the narrow absorption lines provide information on cloud velocities. The widths of all the spectral lines depicted here are greatly exaggerated for the sake of clarity.

## 18.4 21-Centimeter Radiation

A basic difficulty with the optical technique just described is that we can examine interstellar clouds only along the line of sight to a distant star. To form an absorption line, a background source must provide radiation to absorb. The need to see stars through clouds also restricts this approach to relatively local regions, within a few thousand parsecs of Earth. Beyond that distance, stars are completely obscured, and optical observations are impossible. As we have seen, infrared observations provide a means of viewing the emission from some clouds, but they do not completely solve the problem because only the denser, dustier clouds emit enough infrared radiation for astronomers to study them in that part of the spectrum.

To probe interstellar space more thoroughly, we need a more general, more versatile observational method—one that does not rely on conveniently located stars and nebulae. In short, we need a way to detect cold, neutral interstellar matter anywhere in space through its *own* radiation. This may sound impossible, but such an observational technique does in fact exist. The method relies on low-energy *radio* emissions produced by the interstellar gas itself.


### Electron Spin

Recall that a hydrogen atom has one electron orbiting a single-proton nucleus. Besides its orbital motion around the central proton, the electron also has some rotational motion—that is, *spin*—about its own axis. The proton also spins. This model is analogous to a planetary system in which, in addition to the orbital motion of a planet about a central star, both the planet (electron) and the star (proton) rotate about their own axes. But bear in mind the crucial difference between planetary and atomic systems: A planet orbiting the Sun is free to move in any orbit and spin at any rate, but within an atom, all physical quantities, such as energy, momentum, and angular momentum (spin), are *quantized*—they are permitted to take on only specific, distinct values. [☞ \(Sec. 4.2\)](#)

The laws of physics dictate that there are exactly two possible spin configurations for a hydrogen atom in its ground state. The electron and proton can rotate in the same direction, with their spin axes parallel, or they can rotate with their axes antiparallel (i.e., parallel, but oppositely oriented). Figure 18.16 shows these two configurations. The antiparallel configuration has slightly less energy than the parallel state.

### Radio Emission

All matter in the universe tends to achieve its lowest possible energy state, and interstellar gas is no exception. A slightly excited hydrogen atom with the electron and proton spinning in the same direction eventually drops down to the less energetic, opposite-spin state as the electron suddenly and spontaneously reverses its spin. As with any other such change, the transition from a high-energy state to a




▲ **FIGURE 18.16 Hydrogen 21-cm Emission** A ground-level hydrogen atom changing from a higher-energy state (top) to a lower-energy state (bottom).

low-energy state releases a photon with energy equal to the energy difference between the two levels.

Because that energy difference is very small, the energy of the emitted photon is very low. [☞ \(Sec. 4.2\)](#) Consequently, the wavelength of the radiation is rather long—in fact, it is 21.1 cm, roughly the width of this book. That wavelength lies in the radio portion of the electromagnetic spectrum. Researchers refer to the spectral line that results from this hydrogen spin-flip process as **21-centimeter radiation**. This spectral line provides a vital probe into any region of the universe containing atomic hydrogen gas. Figure 18.17 shows typical spectral profiles of 21-cm radio signals observed from several different regions of space. These tracings are the characteristic signatures of cold, atomic hydrogen in our Galaxy. Needing no visible starlight to help calibrate their signals, radio astronomers can observe *any* interstellar region that contains enough hydrogen gas to produce a detectable signal. Even the low-density regions between the dark clouds can be studied.

As can be seen in the figure, actual 21-cm lines are quite jagged and irregular, somewhat like nebular emission lines in appearance. The irregularities arise because there are usually numerous clumps of interstellar gas along any given line of sight, each with its own density, temperature, radial velocity, and internal motion. Thus, the intensity, width, and Doppler shift of the resultant 21-cm line vary from place to place. [☞ \(Sec. 4.5\)](#) All these different lines are superimposed in the signal we eventually receive at Earth, and sophisticated computer analysis is generally required to disentangle them. The “average” figures quoted earlier for the temperatures (100 K) and densities ( $10^6$  atoms/m<sup>3</sup>) of the regions between the dark dust clouds are based on 21-cm measurements. Observations of the dark clouds themselves using 21-cm radiation yield densities and temperatures in good agreement with those obtained by optical spectroscopy.

All interstellar atomic hydrogen emits 21-cm radiation. But if all atoms eventually fall into their lowest-energy



▲ **FIGURE 18.17 21-cm Lines** Typical 21-cm radio spectral lines observed from several different regions of interstellar space. The peaks do not all occur at a wavelength of exactly 21.1 cm, corresponding to a frequency of 1420 MHz, because the gas in our Galaxy is moving with respect to Earth.

configuration, then why isn't all the hydrogen in the Galaxy in the lower energy state by now? Why do we see 21-cm radiation today? The answer is that the energy difference between the two states is comparable to the energy of a typical atom at a temperature of 100 K or so. As a result, atomic collisions in the interstellar medium are energetic enough to boost the electron into the higher energy configuration and so maintain comparable numbers of hydrogen atoms in either state. At any instant, any sample of interstellar hydrogen will contain many atoms in the upper level, so conditions will always be favorable for 21-cm radiation to be emitted.

Of great importance, the wavelength of this characteristic radiation is much larger than the typical size of interstellar dust particles. Accordingly, 21-cm radiation reaches Earth completely unscattered by interstellar debris. The opportunity to observe interstellar space well beyond a few thousand parsecs, and in directions lacking background stars, makes 21-cm observations among the most important and useful in all astronomy. We will see in Chapters 23 through 25 how such observations are indispensable in allowing astronomers to map out the large-scale structure of our Galaxy and many others.

## CONCEPT Check


- ✓ Why is 21-cm radiation so useful as a probe of galactic structure?

## 18.5 Interstellar Molecules

In some particularly cold (typically, 10–20 K) interstellar regions, densities can reach as high as  $10^{12}$  particles/m<sup>3</sup>. Until the late 1970s, astronomers regarded these regions simply as abnormally dense interstellar clouds, but it is now recognized that they belong to an entirely new class of interstellar matter. The gas particles in these regions are not in atomic form at all; they are molecules. Because of the predominance of molecules in these dense interstellar regions, they are known as **molecular clouds**. They literally dwarf even the largest emission nebulae, which were previously thought to be the most massive residents of interstellar space.

### Molecular Spectral Lines

As noted in Chapter 4, much like atoms, molecules can become excited through collisions or by absorbing radiation. ∞ (Sec. 4.4) Furthermore, again like atoms, molecules eventually return to their ground states, emitting radiation in the process. The energy states of molecules are much more complex than those of atoms, however. Once more like atoms, molecules can undergo internal electron transitions, but unlike atoms, they can also rotate and vibrate. They do so in specific ways, obeying the laws of quantum physics. Figure 18.18 depicts a simple molecule rotating rapidly—that is, a molecule in an excited rotational state. After a length of time that depends on its internal makeup, the molecule relaxes back to a slower rotational rate (a state of lower energy). This change causes a photon to be emitted, carrying an energy equal to the energy difference between the two rotational states involved. The energy differences between these states are generally very small, so the emitted radiation is usually in the radio range.



▲ **FIGURE 18.18 Molecular Emission** As a molecule changes from a rapid rotation (left) to a slower rotation (right), a photon is emitted that can be detected with a radio telescope. The lengths of the curved arrows are proportional to the spin rate of the molecule.



We are fortunate that molecules emit radio radiation, because they are invariably found in the densest and dustiest parts of interstellar space. These are regions where the absorption of shorter wavelength radiation is enough to prohibit the use of ultraviolet, optical, and most infrared techniques that might ordinarily detect changes in the energy states of the molecules. Only low-frequency radio radiation can escape.

Why are molecules found only in the densest and darkest of the interstellar clouds? One possible reason is that the dust serves to protect the fragile molecules from the normally harsh interstellar environment—the same absorption that prevents high-frequency radiation from getting out to our detectors also prevents it from getting in to destroy the molecules. Another possibility is that the dust acts as a catalyst that helps form the molecules. The grains provide both a place where atoms can stick together and react and a means of dissipating any heat associated with the reaction, which might otherwise destroy the newly formed molecules. Probably the dust plays both roles; the close association between dust grains and molecules in dense interstellar clouds argues strongly in favor of this view, although the details are still being debated.

## Molecular Tracers

In mapping molecular clouds, radio astronomers are faced with a problem. Molecular hydrogen ( $H_2$ ) is by far the most common constituent of these clouds, but unfortunately, despite its abundance, this molecule does not emit or absorb radio radiation. Rather, it emits only short-wavelength ultraviolet radiation, so it cannot easily be used as a probe of cloud structure. Nor are 21-cm observations helpful—they are sensitive only to *atomic* hydrogen, not to the *molecular* form of the gas. Theorists had expected  $H_2$  to abound in these dense, cold pockets of interstellar space, but proof of its existence was hard to obtain. Only when spacecraft measured the ultraviolet spectra of a few stars located near the edges of some dense clouds was the presence of molecular hydrogen confirmed.


With hydrogen effectively ruled out as a probe of molecular clouds, astronomers must use observations of other molecules to study the dark interiors of these dusty regions. Molecules such as carbon monoxide (CO), hydrogen cyanide (HCN), ammonia ( $NH_3$ ), water ( $H_2O$ ), methyl alcohol ( $CH_3OH$ ), formaldehyde ( $H_2CO$ ), and about 150 others, some quite complex, are now known to exist in interstellar space.\* These molecules are found only in very small quantities—they are generally 1 million to 1 billion times less abundant than  $H_2$ —but they are important as *tracers* of a cloud's structure and physical

\*Some remarkably complex organic molecules, including formaldehyde ( $H_2CO$ ), ethyl alcohol ( $CH_3CH_2OH$ ), methylamine ( $CH_3NH_2$ ), and formic acid ( $H_2CO_2$ ), have been found in the densest of the dark interstellar clouds. Their presence has fueled speculation about the origins of life, both on Earth and in the interstellar medium—especially since the report (still unconfirmed) by radio astronomers in the mid-1990s of evidence that glycine ( $NH_2CH_2COOH$ ), one of the key amino acids that form the large protein molecules in living cells, may also be present in interstellar space.


properties. They are produced by chemical reactions within molecular clouds. When we observe them, we know that the regions under study must also contain high densities of molecular hydrogen, dust, and other important constituents.

The rotational properties of different molecules often make them suitable as probes of regions with different physical properties. Formaldehyde may provide the most useful information on one region, carbon monoxide on another, and water on yet another, depending on the densities and temperatures of the regions involved. The data obtained thereby equip astronomers with a sophisticated spectroscopic “toolbox” for studying the interstellar medium.

For example, Figure 18.19 shows some of the sites where formaldehyde molecules have been detected near M20. At practically every dark area sampled between M16 and M8, the formaldehyde molecule is present in surprisingly large abundance (although it is still far less common than  $H_2$ ). Analyses of spectral lines at many locations along the  $12^\circ$ -wide swath shown in Figure 18.5 indicate that the temperature and density are much the same in all the molecular clouds studied (50 K and  $10^{11}$  molecules/ $m^3$ , on average). Figure 18.20 shows a contour map of the distribution



▲ **FIGURE 18.19 M20's Environment** Spectra indicate that formaldehyde molecules exist in the extended environment (arrows) around M20. Formed by the absorption of background radiation, the spectral lines are most intense both in the dark dust lanes trisecting the nebula and in the dark regions beyond the nebula. (Background image: AURA)



▲ **FIGURE 18.20 Molecules Near M20** This contour map of formaldehyde near the M20 nebula shows how that molecule is more abundant in the darkest interstellar regions. The contour values increase from the outside to the inside, so the maximum density of formaldehyde lies just to the bottom right of the visible nebula. The different colored contours outline the intensity of formaldehyde spectral lines at different frequencies. (*Background image: AURA*)

of formaldehyde molecules in the immediate vicinity of the M20 nebula. After radio spectral lines of formaldehyde were observed at various locations, contours connecting regions of similar abundance were drawn. Notice that the amount of formaldehyde (and, we assume, the amount of hydrogen) peaks in a dark region well away from the visible nebula.

Radio maps of interstellar gas and infrared maps of interstellar dust reveal that molecular clouds do not exist as distinct objects in space. Rather, they make up huge **molecular cloud complexes**, typically up to 50 pc across and containing enough gas to make a million stars like our Sun. About a thousand such giant complexes are currently known in our Galaxy.

In recent years, astronomers have come to realize that the interstellar medium is a dynamic, ever-changing environment, in which energy released by newborn stars (Chapter 19) and supernovae (Chapter 21) drives large-scale, turbulent motion in the interstellar gas. In this view, the cold molecular clouds we see are simply regions of dense gas temporarily compressed by the overall flow—transient islands in a sea of surrounding chaos.

The discovery of many interstellar molecules in the 1970s forced astronomers to rethink and reobserve interstellar space. In doing so, they realized that this active and vital domain is far from the void suspected by theorists not long before. As we will see in Chapter 19, regions of space once thought to contain nothing more than galactic “garbage”—the cool, tenuous darkness among the stars—now play a critical role in our understanding of stars and the interstellar medium from which they are born.

### PROCESS OF SCIENCE Check

- ✓ In mapping molecular clouds, why do astronomers use observations of “minority” molecules such as carbon monoxide and formaldehyde when these molecules constitute only a tiny fraction of the total number of molecules in interstellar space?

**The Big Question** Might life have originated in space and then been delivered to Earth billions of years ago? That’s a question often asked by astronomers, now that so many organic (carbon-rich) molecules have been detected in the dark depths of interstellar space. Yet the formation of complex molecules in extremely cold and tenuous environments still puzzles chemists, and geologists are unsure whether asteroids or comets could really have transported organic molecules to Earth’s surface.

# Chapter Review

## SUMMARY

1 The **interstellar medium** (p. 450) occupies the space among the stars. It is made up of cold (less than 100 K) gas, mostly atomic or molecular hydrogen and helium, and **dust grains** (p. 450). Interstellar dust is highly effective at blocking our view of distant stars, even though the density of the interstellar medium is very low. The spatial distribution of interstellar




matter is patchy. The general diminution of starlight by dust is called **extinction** (p. 450). In addition, the dust preferentially absorbs short-wavelength radiation, leading to a distinct **reddening** (p. 451) of light passing through interstellar clouds. Interstellar dust is thought to be composed of silicates, graphite, iron, and “dirty ice.” Interstellar dust particles are apparently elongated or rodlike. The **polarization** (p. 453) of starlight provides a means of studying these particles.

**2** A **nebula** (p. 453) is a fuzzy bright or dark patch on the sky. **Emission nebulae** (p. 453) are extended clouds of hot, glowing interstellar gas. Associated with star formation, they result when hot O- and B-type stars heat and ionize their surroundings. Studies of the emission lines produced by excited nebular atoms allow astronomers to measure the properties of nebulae. Nebulae are often crossed by dark **dust lanes** (p. 455), part of the larger cloud from which they formed.




**3** **Dark dust clouds** (p. 459) are cold, irregularly shaped regions in the interstellar medium whose constituent dust diminishes or completely obscures the light from background stars. The interstellar medium also contains many cold, dark **molecular clouds** (p. 463), which are cool and dense enough that much of the gas exists in molecular form. Dust within these clouds probably both protects the molecules and acts as a catalyst to




help them form. Often, several molecular clouds are found close to one another, forming an enormous **molecular cloud complex** (p. 465) millions of times more massive than the Sun.

**4** Cold, dark regions of interstellar space containing atomic hydrogen can be observed in the radio spectrum via the **21-centimeter radiation** (p. 462) produced when the electron in an atom of hydrogen reverses its spin, changing its energy slightly in the process. Molecular clouds are observed mainly through the radio radiation emitted by the molecules they contain. Radio waves are not appreciably absorbed by the interstellar medium, so astronomers making observations at these wavelengths can often “see” to great distances.



**5** Hydrogen is by far the most common constituent of molecular clouds, but molecular hydrogen is very hard to observe. Astronomers usually study these clouds via observations of other “tracer” molecules that are less common, but much easier to detect. Many complex molecules have been identified in these regions.



**MasteringAstronomy**® For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction’s numbered Learning Outcomes.

## Review and Discussion

- LO1** Give a brief description of the interstellar medium. How dense is it, and how is it distributed throughout space?
- What is the composition of interstellar gas? What about interstellar dust?
- Why is interstellar dust so much more effective than interstellar gas at absorbing starlight?
- Compare the reddening of stars by interstellar dust with the reddening of the setting Sun.
- What does the polarization of starlight tell us about the interstellar medium?
- What are some methods that astronomers use to study interstellar dust?
- LO2** What is an emission nebula?
- What is photoevaporation, and how does it change the structure and appearance of an emission nebula?
- Why are some spectral lines observed in emission nebulae not normally seen in laboratories on Earth?
- What is the Local Bubble? How did it form?
- LO3 POS** Describe some ways in which we can “see” a dark interstellar cloud.
- Give a brief description of a dark dust cloud.
- LO4** What is 21-cm radiation? With what element is it associated? Why is useful to astronomers?
- LOS POS** How do astronomers explore the structure of molecular cloud complexes?
- If our Sun were surrounded by a cloud of gas, would this cloud be an emission nebula? Why or why not?



## Conceptual Self-Test: Multiple Choice

- The chemical composition of the interstellar medium is basically similar to that of (a) the Sun; (b) Earth; (c) Venus; (d) Mars.
- The density of atoms in the interstellar medium is most similar to (a) wildfire smoke; (b) dark rain clouds; (c) deep ocean water; (d) the interior of a TV tube.

3. Of the following objects, the one that shines most like an emission nebula shines is (a) a regular incandescent light-bulb with a filament; (b) a red hot ember from a campfire; (c) a glowing fluorescent light tube; (d) a star like the Sun.
4. Stars interact with emission nebulae by (a) exciting their atoms enough to emit light; (b) illuminating them like an advertising billboard; (c) causing them to contract; (d) heating them so they explode.
5. A dark interstellar globule is about the same size as (a) a cloud in Earth's atmosphere; (b) the entire planet Earth; (c) a star like the Sun; (d) the Oort cloud.
6. **VIS** The Ophiuchi cloud, shown in Figure 18.13 ("Dark Dust Cloud"), is dark because (a) there are no stars in this region; (b) the stars in this region are young and faint; (c) starlight from behind the cloud does not penetrate the cloud; (d) the region is too cold to sustain stellar fusion.
7. If a proton and an electron within a hydrogen atom initially have parallel spins, then change to have antiparallel spins, the atom must (a) absorb energy; (b) emit energy; (c) become hotter; (d) become larger.
8. The telescope best suited to observing dark dust clouds is (a) an X-ray telescope; (b) a large visible-light telescope; (c) an orbiting ultraviolet telescope; (d) a radio telescope.
9. The largest interstellar clouds are (a) molecular clouds; (b) dark dust clouds; (c) emission nebulae; (d) globules.
10. Molecular clouds are routinely studied using spectral lines from all but which of the following? (a) Molecular hydrogen; (b) Carbon monoxide; (c) Formaldehyde; (d) Water.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

1. ● The average density of interstellar gas within the Local Bubble is much lower than the value mentioned in the text—in fact, it is roughly  $10^3$  hydrogen atoms/m<sup>3</sup>. Given that the mass of a hydrogen atom is  $1.7 \times 10^{-27}$  kg, calculate the total mass of interstellar matter contained within a bubble volume equal in size to planet Earth.
2. ● Assume the same average density as in the previous question, and calculate the total mass of interstellar hydrogen contained within a cylinder of cross-sectional area 1 m<sup>2</sup>, extending from Earth to Alpha Centauri.
3. ●● Given the average density of interstellar matter stated in Section 18.1, calculate how large a volume of space would have to be compressed to make a cubic meter of gas equal in density to air on Earth (1.2 kg/m<sup>3</sup>).
4. ● Interstellar extinction is sometimes measured in magnitudes per kiloparsec (1 kpc = 1000 pc). Light from a star 1500 pc away is observed to be diminished in intensity by a factor of 20 over and above the effect of the inverse-square law. What is the average interstellar extinction, in mag/kpc, along the line of sight?
5. ●● A beam of light shining through a dense molecular cloud is diminished in intensity by a factor of 2 for every 5 pc it travels. By how many magnitudes is the light from a background star dimmed if the total thickness of the cloud is 60 pc?
6. ●● A star of apparent magnitude 10 lies 500 pc from Earth. If interstellar absorption results in an average extinction of 2 mag/kpc, calculate the star's absolute magnitude and luminosity.
7. ●● Estimate the escape speeds near the edges of the four emission nebulae listed in Table 18.1, and compare them with the average speeds of hydrogen nuclei in those nebulae.  (More Precisely 8-1) Do you think it is possible that the nebulae are held together by their own gravity?
8. ● To carry enough energy to ionize a hydrogen atom, a photon must have a wavelength of less than  $9.12 \times 10^{-8}$  m (91.2 nm). Using Wien's law, calculate the temperature a star must have for the peak wavelength of its blackbody curve to equal this value.  (Sec. 3.4)

## Activities

### Collaborative

1. Observe the Messier objects M8, M16, M17, M20, and M42—star-forming regions described in this chapter. Not all are easily observable on any given night, so do some advance research online and make a list of which are visible. A small telescope will give the best results in most cases, and you may want to observe in shifts over the course of the night. For each object, carefully follow the instructions for locating it, and sketch it (or photograph it, if you have the equipment). Compare your sketch to the images in this chapter.

### Individual

1. Observe the Milky Way on a dark, clear night away from city lights. Is it a continuous band of light across the sky, or is it mottled? The parts of the Milky Way that appear missing are actually dark dust clouds lying relatively near the Sun. Identify the constellations in which you see these clouds. Make a sketch and compare with a star atlas. Find other small clouds in the atlas and try to find them with your eye or with binoculars.



Hydrogen sulphide

Méthanol

Formyl cation

Dimethyl ether

Sulphur dioxide

Water

Sulphur dioxide

Formaldehyde

Deuterium cyanide

Carbon monoxide

Water

Methanol

Acrylonitrile

Methyl formate

Methanol

Hydrogen cyanide

Sulphur dioxide

Hydrogen cyanide

# Star Formation

## A TRAUMATIC BIRTH

We now move from the interstellar medium—the gas and dust among the stars—back to the stars themselves. The next four chapters discuss the formation and evolution of stars. We have already seen that stars change as they consume their fuel supply, and we have extensive observational evidence of stars at many different evolutionary stages. With the help of these observations, astronomers have developed an understanding of stellar evolution—the complex changes undergone by stars as they form, mature, grow old, and die.

We begin by studying how interstellar clouds of gas and dust are transformed into the myriad stars throughout the night sky. As we will see, the process is far from gentle—stellar nurseries are scenes of violent outbursts, interstellar shock waves, even actual collisions, as prestellar fragments grow in mass and compete for resources in a newborn cluster. The Sun and planet Earth are survivors of a similarly violent environment, some 4.5 billion years ago.

**The Big Picture** Few issues in astronomy are more basic than knowing how stars originate. Stars are the most numerous and obvious residents of the nighttime sky—just look up on any clear night. Astronomers are eager to understand the details of how stars emerge from the dark messiness of interstellar space to become bright round balls of intense energy. The process is a remarkable one and we have learned much about it in the past few decades.

**LEFT:** This remarkable image—actually a large mosaic of a billion bits of data stitched together from hundreds of smaller images—shows a classic star-forming region. The *Hubble Space Telescope* captured this optical view of the Orion Nebula, a stellar nursery lying roughly 1400 light-years from Earth, populated with thousands of young stars that have recently emerged from the loose matter comprising the surrounding nebulosity. The infrared spectrum at bottom was acquired by Europe's *Herschel Space Observatory*, another outstanding telescope orbiting Earth. This chapter-opener represents state-of-the-art photography and spectroscopy in astronomy today. (STScI; ESA)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Summarize the sequence of events leading to the formation of a star like our Sun.
- 2 Explain how the formation of a star depends on its mass.
- 3 Describe some of the observational evidence supporting the modern theory of star formation.
- 4 Describe the nature of interstellar shock waves, and discuss their possible role in the formation of stars.
- 5 Explain why stars form in clusters, and distinguish between open and globular star clusters.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.

## 19.1 Star-Forming Regions

Our universe is constantly renewing itself. Literally billions of stars have been born, lived out their lives, and died since our Galaxy formed. We do not see this activity when we gaze at the nighttime sky, because the time scales on which stars play out this cosmic drama are enormously long by human standards. Even the shortest-lived O-type stars survive for millions of years. [∞ \(Sec. 17.8\)](#) Nevertheless, we have plenty of evidence for ongoing stellar evolution throughout the cosmos.

### Young Stars in the Universe

Our Sun, and probably most of the stars in our immediate cosmic neighborhood, formed billions of years ago. [∞ \(Sec. 6.7\)](#)

However, we know that many relatively nearby stars are much younger than this. The magnificent emission nebulae discussed in Chapter 18 and the ultraluminous, short-lived stars that power them are direct proof that star formation is a continuing process. [∞ \(Sec. 18.2\)](#) The hottest stars in these regions must have formed less than a few million years ago—the blink of an eye, in cosmic terms—and there is no reason to suppose that Galactic star formation has recently and abruptly ceased! Stars are forming all across the Milky Way, even as you read this.

In fact, star-forming regions are observed in many regions of the universe far beyond our own Galaxy. Figure 19.1 shows one of the most spectacular regions discovered to date. It lies in a small, companion galaxy to our own—one of the so-called Magellanic Clouds to be studied later in Chapter 23. Some 170,000 light-years away, this stunningly rich region of young blue

► **FIGURE 19.1 Stellar Nursery** This combined visible-infrared image captured by the new wide-field camera on the *Hubble Space Telescope* shows a highly detailed view of the star cluster R136, a huge group of bright young blue stars still embedded in the glowing reddish nebula, called Tarantula, in which they formed a few million years ago. The whole region shown is about 100 light-years across. (NASA/ESA)

stars is the largest stellar nursery in our local cosmic neighborhood. The Milky Way may well contain many similarly large star clusters, but if they exist, they must be obscured by much intervening interstellar material.

Simply put, a star forms when part of the interstellar medium—one of the cold dark clouds we studied in Chapter 18—begins to collapse under its own weight. [∞ \(Sec. 18.3\)](#) The cloud fragment heats up as it shrinks, and eventually its center becomes hot enough for nuclear fusion to begin. At that point, the contraction stops and a star is born. But what starts the collapse? How and why does it end? And what determines the mass of the star (or stars) that results? [∞ \(Sec. 17.8\)](#) As we will see, both the environment in which stars form and the effects of interactions among neighbors during the star formation process are critical in determining stellar properties.



## MORE PRECISELY 19-1

### Competition in Star Formation

In the text we present star formation as a competition between gravity, which tends to make interstellar clouds collapse, and heat, which opposes it. In fact, the interstellar medium is a lot more complicated, and heat isn't the only factor opposing gravitational contraction. Two other important factors affecting star formation are *rotation* and *magnetism*.


*Rotation*—spin—competes with gravity's inward pull. As we saw in Chapter 6, a contracting cloud having even a small spin will develop a bulge around its midsection.

∞ (Sec. 6.6) As the cloud contracts, it spins faster (to conserve its angular momentum), the bulge grows, and material on the edge tends to fly off into space. (Think of mud flung from a rapidly rotating bicycle wheel.) Eventually, as in Figure 6.9, the cloud forms a flattened, rotating disk.

For material to remain part of the cloud and not be spun off into space, a force must be applied—in this case, the force of gravity. The more rapid the rotation, the greater is the tendency for the gas to escape, and the greater is the gravitational force needed to retain it. Hence, we can regard rotation as opposing the inward pull of gravity. Should rotation overpower gravity, the cloud would disperse. Thus, more mass is needed for a rapidly rotating interstellar cloud to contract to form a star than is needed for a nonrotating cloud.

*Magnetism* can also hinder a cloud's contraction. Magnetic fields permeate most interstellar clouds. As a cloud contracts, it heats up, and atomic encounters become violent enough to (partly) ionize the gas. As noted in Chapter 7 in discussing Earth's Van Allen belts, and in Chapter 16 in discussing activity on the Sun, magnetic fields can exert electromagnetic control over charged particles. ∞ (Secs. 7.5, 16.5) In effect, the particles tend to become "tied" to the magnetic field—they are free to move *along* the field lines, but are inhibited from moving *perpendicular* to them.

Magnetism can hinder the contraction of an interstellar gas cloud, causing it to contract in a distorted way. Because the ions are tied to the magnetic field, the field lines (red) follow the contraction of the cloud, which itself shrinks more rapidly along the field lines than perpendicular to them. The three frames in the accompanying figure trace the evolution of a slowly contracting interstellar cloud having some magnetism. The dashed lines represent regions where the field lines are distorted and compressed as the cloud shrinks. As the field lines are compressed, the magnetic field strength increases,



becoming much larger than that normally permeating interstellar space. The primitive solar nebula may have contained a strong magnetic field created in just this manner.

Even small amounts of rotation or magnetism can compete with gravity and greatly alter the evolution of a typical gas cloud, but the interplay among these factors is complex and extremely difficult to study theoretically. In this chapter, we will try to understand the broad outlines of the star-formation process by neglecting these two complicating factors. Bear in mind, however, that both are important in determining the details.


### Gravity and Heat

What determines which interstellar clouds collapse? For that matter, since all clouds exert a gravitational pull, why didn't they all collapse long ago? To answer these questions and understand the processes leading to the stars we see, we must explore in a little more detail the factors that compete with gravity in determining a cloud's fate. By

far the most important of these is the random motion of atoms—or *heat*. *More Precisely 19-1* discusses some other factors that influence—and complicate—the star formation process.

We have already seen numerous instances of the competition between heat and gravity. ∞ (*More Precisely 8-1*) The temperature of a gas is simply a measure of the





If more than a few atoms interact, the group would come together,

start to slide by, but then pause,

and finally contract into a clump.

◀ **FIGURE 19.2 Atomic Motions**

The motions of a few atoms within an interstellar cloud are influenced by gravity so slightly that the atoms' paths are hardly changed (a) before, (b) during, and (c) after an accidental, random encounter.

average speed of the atoms or molecules in it, so the higher the temperature, the greater the average speed of the molecules and hence the higher the pressure of the gas. This is the main reason that the Sun and other stars don't collapse: The outward pressure of their heated gases exactly balances gravity's inward pull. ∞ (Fig. 16.2)

Consider a small portion of a large cloud of interstellar gas. Concentrate first on just a few atoms, as shown in Figure 19.2. Even though the cloud's temperature is very low, each atom still has some random motion because of the cloud's heat. ∞ (More Precisely 3-1) Each atom is also influenced by the gravitational attraction of all its neighbors. The gravitational force is not large, however, because the mass of each atom is so small. When a few atoms accidentally cluster for an instant, as shown in Figure 19.2(b), their combined gravity is insufficient to bind them into a lasting, distinct clump of matter. This accidental cluster will disperse as quickly as it forms. The effect of heat is much stronger than the effect of gravity.

Now consider a larger group of atoms. Imagine, for example, 50, 100, 1000—even a million—atoms, each gravitationally pulling on all the others. With increased mass, the force of gravity is now stronger than before. Will this many atoms exert a combined gravitational attraction strong enough to prevent the clump from dispersing again? The answer—at least under the conditions found in interstellar space—is still no. The gravitational attraction of even this mass of atoms is still far too weak to overcome the effect of heat.

How many atoms must be accumulated in order for their collective pull of gravity to prevent them from dispersing back into interstellar space? The answer, even for a typical cool (100 K) cloud, is a truly huge number. Nearly  $10^{57}$  atoms are required—much more than the  $10^{25}$  grains of sand on all the beaches of the world and even more than the  $10^{51}$  elementary particles that constitute all the atomic nuclei in our entire planet. There is simply nothing on Earth comparable to a star.

## Modeling Star Formation

The next two sections describe the currently accepted theoretical view of star formation, derived in large part from numerical experiments performed on high-speed

computers. The results are mathematical predictions of a multifaceted problem incorporating gravity, heat, nuclear reaction rates, elemental abundances, and other physical processes operating in contracting interstellar clouds (see *More Precisely 19-1*).

Scientific theories always develop in response to experimental or observational data, and theories of star formation are no exception. ∞ (Sec. 1.2) The theory of star formation has evolved to explain innumerable observations of stars and star-forming regions. However, the phenomenology in this case is so complex and diverse that it is helpful to have a theoretical framework to “connect the dots” between phenomena that might otherwise appear unrelated. Accordingly, we present the theory first and then discuss how and where the observational data fit into and support the theoretical picture.

### CONCEPT Check

✓ What basic competitive process controls star formation?

## 19.2 The Formation of Stars Like the Sun

Star formation begins when gravity begins to dominate over heat, causing a cloud to lose its equilibrium and start contracting. Only after the cloud has undergone radical changes in its internal structure is equilibrium finally restored.

In the process of becoming a main-sequence star like the Sun, an interstellar cloud goes through seven basic evolutionary stages, as listed in Table 19.1. The stages are characterized by varying central temperatures, surface temperatures, central densities, and radii of the prestellar object. They trace its progress from a cold, dark interstellar cloud to a hot, bright star. The numbers given in the table and in the following discussion are valid *only* for stars of approximately the same mass as that of the Sun. In the next section, we will relax this restriction and consider the formation of stars with masses different from that of the sun.

**TABLE 19.1** Prestellar Evolution of a Solar-Type Star

Stage	Approximate Time to Next Stage (yr)	Central Temperature (K)	Surface Temperature (K)	Central Density (particles/m <sup>3</sup> )	Diameter* (km)	Object
1	$2 \times 10^6$	10	10	$10^9$	$10^{14}$	Interstellar cloud
2	$3 \times 10^4$	100	10	$10^{12}$	$10^{12}$	Cloud fragment Cloud fragment/protostar
3	$10^5$	10,000	100	$10^{18}$	$10^{10}$	
4	$10^6$	1,000,000	3000	$10^{24}$	$10^8$	Protostar
5	$10^7$	5,000,000	4000	$10^{28}$	$10^7$	Protostar
6	$3 \times 10^7$	10,000,000	4500	$10^{31}$	$2 \times 10^6$	Star
7	$10^{10}$	15,000,000	6000	$10^{32}$	$1.5 \times 10^6$	Main-sequence star

\* Round numbers; for comparison, recall that the diameter of the Sun is  $1.4 \times 10^6$  km, whereas that of the solar system is roughly  $1.5 \times 10^{10}$  km.

Note again the time scales involved in these stages—even the shortest spans a thousand human generations. Astronomers have not gained this insight by watching a single cloud or group of clouds evolve from start to finish. Rather, they combine theory and observation to refine a still evolving mathematical model of how stars form.

## Stage 1: An Interstellar Cloud


The first stage in the star-formation process is a dense interstellar cloud—the core of a dark dust cloud or perhaps a molecular cloud. These clouds are truly vast, sometimes spanning tens of parsecs ( $10^{14}$ – $10^{15}$  km) across. Typical temperatures are about 10 K throughout, with a density of perhaps  $10^9$  particles/m<sup>3</sup>. Stage-1 clouds contain thousands of times the mass of the Sun, mainly in the form of cold atomic and molecular gas. (The dust in a stage-1 cloud both cools the cloud as it contracts and plays a crucial role in planet formation, but it constitutes a negligible fraction of the total mass of the cloud.) [∞ \(Sec. 15.2\)](#)

Despite their low internal temperatures, most observed dark interstellar clouds seem to have enough internal pressure to support themselves against the force of gravity. [∞ \(More Precisely 8-1\)](#) However, if such a cloud is to be the birthplace of stars, it must become unstable, start to collapse under its own gravity, and eventually break up into smaller pieces. Most astronomers think that the process of star formation is triggered when some external event, such as the shock of a nearby stellar explosion or the pressure wave produced

when a nearby O- or B-type star forms and ionizes its surroundings, squeezes a cloud beyond the point where pressure can resist gravity's inward pull. [∞ \(Secs. 16.2, 17.5\)](#) Or perhaps the cloud's supporting magnetic field leaks away as charged particles slowly drift across the confining field lines, leaving the gas unable to support its own weight (*More Precisely 19-1*).

Whatever the cause, theory suggests that once the collapse begins, fragmentation into smaller and smaller clumps of matter naturally follows, as gravitational instabilities continue to operate in the gas. As illustrated in Figure 19.3, a typical cloud can break up into tens, hundreds, or even thousands, of fragments, each imitating the shrinking behavior of the parent cloud and contracting ever faster. The whole process, from a single stable cloud to many collapsing fragments, takes a few million years.

In this way, depending on the precise conditions under which fragmentation takes place, an interstellar cloud can produce either a few dozen stars, each much larger than



**▲ FIGURE 19.3 Cloud Fragmentation** As an interstellar cloud contracts, gravitational instabilities cause it to fragment into smaller pieces. The pieces themselves continue to fall inward and fragment, eventually forming many tens or hundreds of individual stars.

our Sun, or a whole cluster of hundreds of stars, each comparable to or smaller than our Sun. There is little evidence of stars born in isolation, one star from one cloud. Most stars—perhaps even all stars—appear to originate as members of multiple systems or large groups of stars. The Sun, which is now found alone and isolated in space, probably escaped from the larger system in which it formed, perhaps after an encounter with another star or some much larger object (such as a molecular cloud).

## Stage 2: A Collapsing Cloud Fragment

The second stage in our evolutionary scenario represents the physical conditions in just one of the many fragments that develop in a typical interstellar cloud. A fragment destined to form a star like the Sun contains between 1 and 2 solar masses of material at this stage. Estimated to span a few hundredths of a parsec across, this fuzzy, gaseous blob is still about 100 times the size of our solar system. Its central density by this time is roughly  $10^{12}$  particles/m<sup>3</sup>.

Even though the fragment has shrunk substantially, its average temperature is not much different from that of the original cloud. The reason is that the gas constantly radiates large amounts of energy into space. The material of the fragment is so thin that photons produced within it easily escape without being reabsorbed by the cloud, so virtually all the energy released in the collapse is radiated away and does not cause any significant increase in temperature. Only at the center, where the radiation must traverse the greatest amount of material to escape, is there any appreciable temperature rise. The gas there may be as warm as 100 K by this stage. For the most part, however, the fragment stays cold as it shrinks.

The process of continued fragmentation is eventually stopped by the increasing density within the shrinking cloud. As stage-2 fragments continue to contract, they eventually become so dense that radiation cannot get out of the cloud easily. The trapped radiation then causes the temperature to rise, the pressure to increase, and the fragmentation to cease.

## Stage 3: Fragmentation Ceases

By the start of stage 3, several tens of thousands of years after it first began contracting, a typical stage-2 fragment has shrunk to roughly the size of our solar system (still 10,000 times the size of our Sun). The density in the inner regions has just become high enough that the gas is opaque to the radiation it emits, so the core of the fragment begins to heat up considerably, as noted in Table 19.1. The central temperature has reached about 10,000 K—hotter than the hottest steel furnace on Earth. However, the temperature in the fragment's outer parts has not increased much. The gas there is still able to radiate its energy into space and so remains cool. The density increases much faster in the center of the

fragment than near the edge, so the outside is both cooler and thinner than the interior. By this time, the central density is approximately  $10^{18}$  particles/m<sup>3</sup> (still only  $10^{-9}$  kg/m<sup>3</sup> or so).

For the first time, our contracting cloud fragment is beginning to resemble a star. The dense, opaque region at the center is called a **protostar**—an embryonic object at the dawn of star birth. The protostar's mass grows as more and more material rains down on it from the surrounding, still shrinking, fragment. However, the protostar's radius continues to decrease because pressure is still unable to overcome the relentless pull of gravity. After stage 3, we can distinguish a “surface” on the protostar—its *photosphere*. Inside the photosphere, the protostellar material is opaque to the radiation it emits.\* From here on, the surface temperatures listed in Table 19.1 refer to the photosphere of the collapsing fragment and not to its low-density “periphery,” where radiation can easily escape and the temperature remains low.


## Stage 4: A Protostar

As the protostar evolves, it shrinks, its density grows, and its temperature rises, both in the core and at the photosphere. Some 100,000 years after the fragment began to form, it reaches stage 4, where its center seethes at about 1,000,000 K. Electrons and protons ripped from atoms whiz around at hundreds of kilometers per second, yet the temperature is still well short of the  $10^7$  K needed to ignite the proton–proton nuclear reactions that fuse hydrogen into helium. ☞ (Sec. 16.6) Still much larger than the Sun, our gassy heap is now about the size of Mercury's orbit. Heated by the material falling on it from above, it now has a surface temperature of a few thousand kelvins.


Knowing the protostar's radius and surface temperature, we can calculate its luminosity. Surprisingly, it turns out to be several thousand times the luminosity of the Sun. Even though the protostar has a surface temperature only about half that of the Sun, it is hundreds of times larger, making its total luminosity very large indeed—in fact, much greater than the luminosity of most main-sequence stars. Because nuclear reactions have not yet begun, the protostar's luminosity is due entirely to the release of gravitational energy as the protostar continues to shrink and material from the surrounding fragment continues to fall onto its surface.

By the time stage 4 is reached, our protostar's physical properties can be plotted on the Hertzsprung–Russell (H–R) diagram, as shown in Figure 19.4. Recall that an H–R diagram is a plot of two key stellar properties: surface temperature (increasing to the left) and luminosity (increasing

\*Note that this is the same definition of “surface” that we used for the Sun in Chapter 16. ☞ (Sec. 16.1)



▲ **FIGURE 19.4 Protostar on the H–R Diagram** The red arrow indicates the approximate evolutionary track followed by an interstellar cloud fragment before reaching the end of the Kelvin–Helmholtz contraction phase as a stage-4 protostar. The boldface numbers on this and subsequent H–R plots refer to the prestellar evolutionary stages listed in Table 19.1.



▲ **FIGURE 19.5 Interstellar Cloud Evolution** Artist's conception of the changes in an interstellar cloud during the early evolutionary stages outlined in Table 19.1. (Not drawn to scale.) The duration of each stage, in years, is indicated.


upward). ∞ (Sec. 17.5) The luminosity scale in the figure is expressed in terms of the solar luminosity ( $4 \times 10^{26}$  W). Our G2-type Sun is plotted at a temperature of 6000 K and a luminosity of 1 unit. As before, the dashed diagonal lines in the H–R diagram represent an object's radius, allowing us to follow the changes in the protostar's size as it evolves. At each phase of the star's evolution, its surface temperature and luminosity can be represented by a point on the diagram. The motion of that point as the star evolves is known as the star's **evolutionary track**. It is a graphical representation of a star's life.

The red track in Figure 19.4 depicts the approximate path followed by our interstellar cloud fragment since it became a protostar at stage 3 (which itself lies off the right-hand edge of the figure). This early evolutionary track is known as the *Kelvin–Helmholtz contraction phase*, after the two European physicists (Lord Kelvin and Hermann von Helmholtz) who first studied the subject.

Figure 19.5 is an artist's sketch of an interstellar gas cloud proceeding along the evolutionary path outlined so far. As the stage-3 fragment contracts, it spins faster (to conserve angular momentum) and flattens into a rotating **protostellar disk** perhaps 100 AU in diameter, surrounding the central stage-4 protostar. ∞ (More Precisely 6-1) Recall that we first saw this process in Chapter 6, where we referred to the disk as the *solar nebula*. ∞ (Sec. 6.7) If the star is ultimately going to have a planetary system, by stage 4 that process is already well underway. ∞ (Sec. 15.2) However, regardless of whether planets actually form,

astronomers think that protostellar disks are common—the vast majority of protostars (perhaps all) are accompanied by disks at this stage of their evolution.

Our protostar is still not in equilibrium. Even though its temperature is now so high that outward-directed pressure has become a powerful countervailing influence against gravity's continued inward pull, the balance is not yet perfect. The protostar's internal heat gradually diffuses out from the hot center to the cooler surface, where it is radiated away into space. As a result, the overall contraction slows, but it does not stop completely. From our perspective on Earth, this is quite fortunate: If the heated gas were somehow able to counteract gravity completely before the star reached the temperature and density needed to start nuclear burning in its core, the protostar would simply radiate away its heat and never



**Interactive** **FIGURE 19.6** **Newborn Star on the H–R Diagram** The changes in a protostar’s observed properties are shown by the path of decreasing luminosity, from stage 4 to stage 6, often called the Hayashi track. At stage 7, the newborn star has arrived on the main sequence.

become a true star. The night sky would be abundant in faint protostars, but completely lacking in the genuine article. Of course, there would be no Sun either, so it is unlikely that we, or any other intelligent life-form, would exist to appreciate these astronomical subtleties.

After stage 4, the protostar on the H–R diagram moves down (toward lower luminosity) and slightly to the left (toward higher temperature), as shown in Figure 19.6. Its surface temperature remains almost constant, and it becomes less luminous as it shrinks. This portion of our protostar’s evolutionary path, running from point 4 to point 6 in Figure 19.6, is often called the *Hayashi track*, after C. Hayashi, a 20th-century Japanese astrophysicist whose groundbreaking work in the 1960s on the evolution of pre-main-sequence stars still provides the theoretical basis for all studies of star formation.

Protostars on the Hayashi track often exhibit violent surface activity during this phase of their evolution, resulting in extremely strong protostellar winds, much denser than the solar wind that flows from our own Sun. As mentioned previously, this portion of the protostar’s evolution is often called the **T Tauri** phase, after T Tauri, the first “star” (actually protostar) to be observed in that stage of prestellar development. [☞ \(Sec. 15.3\)](#)

## Stage 5: Protostellar Evolution

By stage 5 on the Hayashi track, the protostar approaches the main sequence. It has shrunk to about 10 times the size of the Sun, its surface temperature is about 4000 K, and its luminosity has fallen to about 10 times the solar value. At this point, the central temperature has reached about 5,000,000 K. The gas is completely ionized by now, but the protons still do not have enough thermal energy to overcome their mutual electromagnetic repulsion and enter the realm of the nuclear binding force. [☞ \(Sec. 16.6\)](#) The core is still too cool for nuclear fusion to begin.

Events proceed more slowly as the protostar approaches the main sequence. The initial contraction and fragmentation of the interstellar cloud occurred quite rapidly, but by stage 5, as the protostar nears the status of a full-fledged star, its evolution slows. The cause of this slowdown is heat: Even gravity must struggle to compress a hot object. The contraction is governed largely by the rate at which the protostar’s internal energy can be radiated away into space. The greater this radiation of internal energy—that is, the more rapidly energy moves through the star to escape from its surface—the faster the contraction occurs. As the luminosity decreases, so, too, does the rate of contraction.

## Stage 6: A Newborn Star

Some 10 million years after its first appearance, the protostar finally becomes a true star. By the bottom of the Hayashi track, at stage 6, when our roughly 1-solar-mass object has shrunk to a radius of about 1,000,000 km, the contraction has raised the central temperature to 10,000,000 K, enough to ignite nuclear burning. Protons begin fusing into helium nuclei in the core, and a star is born. As shown in Figure 19.6, the star’s surface temperature at this point is about 4500 K, still a little cooler than the Sun. Even though the newly formed star is slightly larger in radius than our Sun, its lower temperature means that its luminosity is somewhat less than (actually, about two-thirds of) the solar value.

## Stage 7: The Main Sequence at Last

Over the next 30 million years or so, the stage-6 star contracts a little more. In making this slight adjustment, the star’s central density rises to about  $10^{32}$  particles/m<sup>3</sup> (more conveniently expressed as  $10^5$  kg/m<sup>3</sup>), the central temperature increases to 15,000,000 K, and the surface temperature reaches 6000 K. By stage 7, the star finally arrives at the main sequence, just about where our Sun now resides. Pressure and gravity are finally balanced, and the rate at which nuclear energy is generated in the core exactly matches the rate at which energy is radiated from the surface.

The evolutionary events just described occur over the course of some 40 to 50 million years. Although this is a long time by human standards, it is still less than 1 percent of the Sun’s lifetime on the main sequence. Once an object begins fusing hydrogen in its core and establishes a “gravity-in, pressure-out” equilibrium, it is destined to burn steadily for a very long time. The star’s location on the H–R diagram—that is, its surface temperature and luminosity—will remain virtually unchanged for the next 10 billion years.

### CONCEPT Check

✓ What distinguishes a collapsing cloud from a protostar and a protostar from a star?


## 19.3 Stars of Other Masses

The numerical values and the evolutionary track just described are valid only for the case of a 1-solar-mass star. The temperatures, densities, and radii of prestellar objects of other masses exhibit similar trends, but the numbers and the tracks differ, in some cases considerably. Perhaps not surprisingly, the most massive fragments within interstellar clouds tend to produce the most massive protostars and, eventually, the most massive stars. Similarly, low-mass fragments give rise to low-mass stars. Whatever the mass, the end point of the prestellar evolutionary track is the main sequence.

### The Zero-Age Main Sequence

Figure 19.7 compares the theoretical pre-main-sequence track taken by the Sun with the corresponding evolutionary tracks of a 0.3-solar-mass star and a 3-solar-mass star. All three tracks traverse the H–R diagram in the same general manner, but cloud fragments that eventually form stars more massive than the Sun approach the main sequence along a higher track on the diagram, whereas those destined to form less massive stars take a lower track. The *time* required for an interstellar cloud to become a main-sequence star also depends strongly on its mass. The most massive cloud fragments heat up to the required 10 million K and become O-type stars in a mere million years, roughly  $\frac{1}{50}$  the time taken by the Sun. The opposite is the case for prestellar objects having masses less than that of our Sun. A typical M-type star, for example, requires nearly a billion years to form.

A star is considered to have reached the main sequence when hydrogen burning begins in its core and the star’s properties settle down to stable values. The main-sequence line thus predicted by theory is called the **zero-age main sequence** (or ZAMS, for short). The fact that the theoretically derived zero-age main sequence agrees very well with the actual main sequences observed for stars in the vicinity of the Sun and in more distant



▲ **FIGURE 19.7 Prestellar Evolutionary Tracks** Prestellar evolutionary paths for stars more massive and less massive than our Sun.

star clusters (see Section 19.6) provides strong support for the modern theory of star formation and stellar structure. ∞ (Sec. 1.2)

If all gas clouds contained precisely the same elements in exactly the same proportions, mass would be the sole determinant of a newborn star’s location on the H–R diagram and the zero-age main sequence would be a well-defined line rather than a broad band. However, the composition of a star affects its internal structure (mainly by changing the opacity of its outer layers), and this in turn affects both the star’s temperature and its luminosity on the main sequence. Stars with more heavy elements tend to be cooler and slightly less luminous than stars that have the same mass, but contain fewer heavy elements. As a result, differences in composition between stars “blur” the zero-age main sequence into the broad band we observe.

It is important to realize that *the main sequence is itself not an evolutionary track—stars do not evolve along it*. Rather, it is just a “way station” on the H–R diagram where stars stop and spend most of their lives—low-mass stars at the bottom, high-mass stars at the top. Once on the main sequence, a star stays in essentially the same location on the H–R diagram during its whole time as a stage-7 object. (In other words, a star that arrives on the main sequence as, say, a G-type star can never “work its way up” to become

a B- or an O-type main-sequence blue supergiant or move down to become an M-type red dwarf.) As we will see in Chapter 20, the next stage of stellar evolution occurs when a star moves away from the main sequence. A star leaving the main sequence and entering this next stage has pretty much the same surface temperature and luminosity it had when it arrived on the main sequence millions (or billions) of years earlier.

## Failed Stars

Some cloud fragments are too small ever to become stars. Consider, for example, the giant planet Jupiter. It formed in the Sun's protostellar disk (the solar nebula) and contracted under the influence of gravity. The resultant heat is still detectable, but the planet did not have enough mass for gravity to crush its matter to the point of nuclear ignition. ∞ (Sec. 11.3) Instead, Jupiter became stabilized by heat and rotation before the planet's central temperature became hot enough to fuse hydrogen—Jupiter never evolved beyond the protostar stage. If it, or any of the other jovian planets, had continued to accumulate gas from the solar nebula, it might have become a star (almost certainly to the detriment of life on Earth). However, that did not occur—virtually all the matter present during the formative stages of our solar system is now gone, swept away by the solar wind during the Sun's T Tauri phase. ∞ (Sec. 15.2)

Low-mass gas fragments simply lack the mass needed to initiate nuclear burning. Rather than turning into stars, they continue to cool, eventually becoming compact, dark “clinkers”—cold fragments of unburned matter—orbiting a star or moving alone through interstellar space. On the basis of theoretical modeling, astronomers think that the minimum mass of gas needed to generate core temperatures high enough to begin nuclear fusion is about 0.08 solar mass (80 times the mass of Jupiter). Our practical definition of a star requires that it shine via the energy released by nuclear fusion reactions in its core. Thus this mass of 0.08 times the mass of the Sun is a lower limit on the masses of all stars in the universe.

Vast numbers of “substellar” objects may well be scattered throughout the universe—fragments frozen in time somewhere along the Kelvin–Helmholtz contraction phase. Small, faint, and cool (and growing ever colder), they are known collectively as **brown dwarfs**. For reasons discussed in more detail in *Discovery 19-1*, researchers generally reserve the term *brown dwarf* to mean a low-mass prestellar fragment of more than about 12 Jupiter masses (so Jupiter itself is not a brown dwarf, by this definition). Anything smaller is simply called a *planet*.

Observationally, these faint, low-mass objects are difficult to study, be they planets or brown dwarfs associated with stars or interstellar cloud fragments far from any star (see *Discovery 19-1*). Current observations suggest that up

to 100 billion cold, dim substellar objects may lurk in the depths of interstellar space—a number comparable to the total number of “real” stars in our Galaxy.

## CONCEPT Check

✓ Do stars evolve along the main sequence?

## 19.4 Observations of Cloud Fragments and Protostars

How can we verify the theoretical picture just outlined? The age of our entire civilization is much shorter than the time needed for a single interstellar cloud to contract and form a star. We can never observe individual objects proceed through the full panorama of star birth. However, we can do the next best thing: We can observe many different objects—interstellar clouds, protostars, and young stars approaching the main sequence—as they appear today at different stages of their evolutionary paths.

The various evolutionary stages just described draw on evidence from different parts of the electromagnetic spectrum, and each observation is like part of a jigsaw puzzle. ∞ (Sec. 3.3) When properly oriented relative to all the others, the pieces can be used to build up a picture of the full life cycle of a star.

## Evidence of Cloud Contraction

Prestellar objects at stages 1 and 2 are not yet hot enough to emit much infrared radiation, and certainly no optical radiation arises from their dark, cool interiors. The best way to study the early stages of cloud contraction and fragmentation is to observe the radio emission from interstellar molecules within those clouds. Consider again M20, the splendid emission nebula studied in Chapter 18. ∞ (Sec. 18.2) The brilliant region of glowing, ionized gas shown in Figure 18.7 is not our main interest here, however; instead, the youthful O- and B-type stars that energize the nebula alert us to the general environment in which stars are forming. Emission nebulae are indicators of star birth.

The region surrounding M20 contains galactic matter that seems to be contracting. The presence of (optically) invisible gas there was illustrated in Figure 18.20, which showed a contour map of the abundance of the formaldehyde ( $\text{H}_2\text{CO}$ ) molecule. Formaldehyde and many other molecules are widespread in the vicinity of the nebula, especially throughout the dusty regions below and to the right of the emission nebula itself. Further analysis of the observations suggests that this region of greatest molecular abundance is also contracting and fragmenting and is well on its way toward forming a star—or, more likely, a star cluster.

## DISCOVERY 19-1


## Observations of Brown Dwarfs

Cruelly put, brown dwarfs are stellar failures—objects that formed through the contraction and fragmentation of an interstellar cloud, just as stars do, but fell short of the critical mass of about 0.08 solar mass (80 times the mass of Jupiter) needed to start hydrogen fusion in their cores. Interstellar space could contain huge numbers of these dim objects.

Although hundreds of brown dwarfs are now known, detecting them is no easy task, as they are small, cool, and hence very faint. ∞ (Sec. 3.4) We can detect stars by means of telescopes, and we can infer the presence of interstellar atoms and molecules by spectroscopic analysis, but astronomical objects of intermediate size outside our solar system remain hard to see. One place astronomers have looked is in binary-star systems, using many of the same techniques they employ in the search for extrasolar planets. ∞ (Sec. 15.5) The images below show two binary-star systems containing brown dwarf candidates (marked by arrows). Note in each case how much fainter the brown dwarf is than its companion. Very high resolution is usually needed to separate the two.


The first (left) is an image of Gliese 623, which was originally identified as a binary system because of its variations in radial velocity. ∞ (Secs. 15.5, 17.7) From the binary's measured orbital separation and period, the mass of the faint companion appears to be approximately 0.1 solar mass—very close to the limit for a brown dwarf, although astronomers still aren't certain of its exact mass. ∞ (Sec. 2.8) The “rings” in the image are instrumental artifacts. The second image shows the binary-star system Gliese 229. These two objects are 7" apart; the fainter “star” has a luminosity only a few millionths that of the Sun and an estimated mass about 50 times that of Jupiter. (The diagonal streak in the image is caused by an overexposure of the brighter star in the CCD chip used to record it.)

Actually, the dividing line between brown dwarfs and Jupiter-like planets is not completely clear-cut, especially given the varied properties of the many extrasolar planetary systems now known. ∞ (Sec. 15.6) Researchers distinguish between “stellar” objects (stars and brown dwarfs), which form within their own contracting cloud fragment as described in the text, and planets, which form in the nebular disk around a larger parent. For definiteness, many draw the dividing line at about 12 times the mass of Jupiter. Above that mass (but below





80 Jupiter masses), although core temperatures never become high enough for hydrogen fusion to occur, a contracting fragment will experience a brief phase of *deuterium fusion*, as the core becomes hot enough for any deuterium nuclei that are present in the original cloud to combine. The phase ends once the deuterium is consumed, and the fragment's “nuclear” lifetime is over. Below 12 Jupiter masses, no nuclear fusion of any kind is expected. The drawing above compares the sizes of some stars, brown dwarfs, and planets.


Infrared and spectroscopic studies offer other ways of searching for brown dwarfs, especially those that are not in binaries. Infrared observations are particularly effective because brown dwarfs emit most of their radiation in that part of the spectrum, whereas true stars tend to be brightest at the near-infrared and optical ranges. The final image below captures a star cluster just north of the Orion Nebula taken by the *Spitzer Space Telescope*. The bright objects are stars, but many of the faint specks are brown dwarf candidates. Researchers estimate that some 10–15 percent of the “stars” in Orion are actually brown dwarfs.



(NASA)




(NASA)






The interstellar clouds in and around M20 thus provide tentative evidence of three distinct phases of star formation, as shown in Figure 19.8. The huge, dark molecular cloud surrounding the visible nebula is the stage-1 cloud. Both its density and its temperature are low—about  $10^8$  particles/ $m^3$  and 20 K, respectively. Greater densities and temperatures typify smaller regions within this large cloud. The totally



(a)



(b)

(c)

obscured regions labeled A and B, where the molecular emission of radio energy is strongest, are such denser, warmer fragments. Here, the total gas density is observed to be at least  $10^9$  particles/ $m^3$ , and the temperature is about 100 K. The Doppler shifts of the radio lines observed in the vicinity of region B imply that this portion of M20, labeled “contracting fragment” in the figure, is infalling. Recent infrared observations (Figure 19.8c) reveal the candidate protostars themselves, identified by the warmth of their growing embryos tucked inside. Less than a light-year across, the region has a total mass over a thousand times the mass of the Sun—considerably more than the mass of M20 itself. The region lies somewhere between stages 1 and 2 of Table 19.1.


The third star-formation phase shown in Figure 19.8 is M20 itself. The glowing region of ionized gas results directly from a massive O-type star that formed there within the past million years or so. Because the central star is already fully formed, this final phase corresponds to stage 6 or 7 of our evolutionary scenario.

### Evidence of Cloud Fragments

Other parts of our Milky Way Galaxy provide sketchy evidence for prestellar objects in stages 3 through 5. The Orion complex, shown in Figure 19.9, is one such region about 1400 light-years away. Lit from within by several O-type stars, the bright Orion Nebula is partly surrounded by a vast molecular cloud that extends even beyond the roughly  $10 \times 30$ -light-year region bounded by the photograph in Figure 19.9(b).

The Orion molecular cloud harbors several smaller sites of intense radiation emitted by molecules deep within

**◀ FIGURE 19.8 Star Formation Phases** (a) The M20 region shows observational evidence for three broad phases in the birth of a star. The parent cloud is stage 1 of Table 19.1. The region labeled “contracting fragment” likely lies between stages 1 and 2. Finally, the emission nebula (M20 itself) results from the formation of one or more massive stars (stages 6 and 7). (b) A close-up (including *Hubble* inlays) of the area near region B outlines (in drawn ovals) especially dense knots of dusty matter. (c) A *Spitzer Telescope* infrared image of the same scene reveals those cores thought to be stellar embryos (arrows). (*AURA*; *NASA*)




**Interactive FIGURE 19.9 Orion Nebula, Up Close** (a) The constellation Orion, with the region around its famous emission nebula marked by a rectangle. The Orion Nebula is the middle “star” of Orion’s sword (see Figure 1.8). (b) Enlargement of the framed region in part (a), but here shown in the infrared, revealing how the nebula is partly surrounded by a vast molecular cloud. Various parts of this cloud are probably fragmenting and contracting, with even smaller sites forming protostars. The three frames at the right show some of the evidence for those protostars: (c) nearly real-color visible image of embedded nebular “knots” within the Orion Nebula itself, (d) false-color radio image of some intensely emitting molecular sites, and (e) high-resolution image of one of many young protostars surrounded by disks of gas and dust where planets might ultimately form. (*P. Sanz/Alamy; SST; Cfa; NASA*)

the core of the cloud fragment. Their extent, shown in Figures 19.9(d) and (e), measures about  $10^{10}$  km, or  $\frac{1}{1000}$  of a light-year, about the diameter of our solar system. Their density is about  $10^{15}$  particles/ $\text{m}^3$ , much denser than the surrounding cloud. Although the temperature of these smaller regions cannot be estimated reliably, many researchers regard the regions as objects well on their way to stage 3. We cannot determine whether those regions will eventually form stars like the Sun, but it does seem certain that the intensely emitting objects in them are on the threshold of becoming protostars.

## Evidence of Protostars

In the hunt for, and study of, objects at more advanced stages of star formation, radio techniques become less useful, because stages 4, 5, and 6 have increasingly higher temperatures. By Wien’s law, their emission shifts toward shorter wavelengths, so these objects shine most strongly in the infrared.  $\infty$  (Sec. 3.4) One particularly bright infrared emitter, known as the *Becklin–Neugebauer* object, was detected in the core of the Orion molecular cloud in the 1970s. Its luminosity is around a thousand times the




◀ **FIGURE 19.10 Protostars** (a) An edge-on infrared image of a planetary system-sized dusty disk in the Orion region, showing heat and light emerging from its center. On the basis of its temperature and luminosity, this unnamed source appears to be a low-mass protostar on the Hayashi track (around stage 5) in the H–R diagram. (b) An optical, face-on image of a slightly more advanced circumstellar disk surrounding an embedded protostar in Orion. (NASA)

luminosity of the Sun. Most astronomers agree that this warm, dense blob is a high-mass protostar, probably in or around stage 4.

Until the *Infrared Astronomy Satellite (IRAS)* was launched in the early 1980s, astronomers were aware of giant stars forming only in clouds far away. ∞ (Sec. 5.7) But *IRAS* showed that many such stars are forming much closer to home, and some of these protostars have masses comparable to that of our Sun. Figure 19.10 shows two examples of low-mass protostars, both spotted by *HST* in a rich star-forming region in Orion. Their infrared heat signatures are those expected of an object on the Hayashi track, at around stage 5.

The energy sources for some infrared objects seem to be luminous hot stars that are hidden from optical view by surrounding dark clouds. Apparently, these stars are already so hot that they emit large amounts of ultraviolet radiation, which is mostly absorbed by “cocoon” of dust surrounding them. The absorbed energy is then reemitted by the dust as infrared radiation. These bright infrared sources are known as *cocoon nebulae*. Two considerations support the idea that



◀ **FIGURE 19.11 Protostellar Wind** (a) The nebular disk around a protostar can be the site of intense heating and strong outflows, forming a bipolar jet perpendicular to the disk. (b) As the disk is blown away by the wind, the jets fan out, eventually (c) merging into a spherical wind. In contrast to this art, part (d) is an actual infrared image of a hot young star (at right) whose powerful winds are ripping away the disk (at left) surrounding a Sun-like star (at center). This system is located about 750 pc away in the star-forming cloud IC 1396. (SST)

the hot stars heating the dust have only recently ignited: (1) The dust cocoons are predicted to disperse quite rapidly once their central stars form, and (2) they are invariably found in the dense cores of molecular clouds. The central stars probably lie near stage 6.

## Protostellar Winds

Protostars often exhibit strong winds. Radio and infrared observations of hydrogen and carbon monoxide molecules in the Orion molecular cloud have revealed gas expanding outward at velocities approaching 100 km/s. High-resolution interferometric observations have disclosed expanding knots of water emission within the same star-forming region and have linked the strong winds to the protostars themselves. ∞ (Sec. 5.6) These winds may be related to the violent surface activity associated with many protostars.


As mentioned earlier, a young protostar may be embedded in an extensive protostellar disk of nebular material in which planets are forming. ∞ (Sec. 15.2) Strong heating within the turbulent disk and a powerful protostellar wind combine to produce a **bipolar flow**, expelling two “jets” of matter in the directions perpendicular to the disk, as illustrated by the art in Figure 19.11(a)–(c). As the protostellar wind gradually destroys the disk, blowing it away into space, the outflow widens until, with the disk gone, the

wind flows away from the star equally in all directions, as is approximately shown by actual infrared imagery in Figure 19.11(d). Figure 19.12 shows the emission from an especially clear bipolar flow, along with an artist’s conception of the system producing it.

These outflows can be very energetic. Figure 19.13 shows a portion of the Orion molecular cloud, south of the Orion Nebula, where a newborn star is seen still surrounded by a bright nebula, its turbulent wind spreading out into the interstellar medium. Below the star (enlarged in the inset) are twin jets known as HH1 and HH2. (“HH” stands for *Herbig–Haro*, the investigators who first cataloged such objects.) Formed in another (unseen) protostellar disk—the protostar itself is still hidden within the dusty cloud fragment from which it formed—these jets have traveled outward for almost half a light-year before colliding with interstellar matter. More Herbig–Haro objects can be seen in the upper-right portion of the figure. One of them, the oddly shaped “waterfall,” may be due to an earlier outflow from the same protostar responsible for the existence of HH1 and HH2.

### PROCESS OF SCIENCE Check


- ✓ How can a “snapshot” of the universe today test our theories of the evolution of individual objects?



▲ **FIGURE 19.12 Bipolar Jets** (a) This remarkable image shows two jets emanating from the young star system HH30, the result of matter accreting onto (via the disk), and then expelled from (via the jet), an embryonic star near the center. The system is viewed roughly edge-on to the disk. (b) An artist’s conception of a young star system meant to illustrate more clearly what’s happening in (a), showing two jets flowing perpendicular to the disk of gas and dust rotating around the star. (NASA; D. Berry)



◀ **FIGURE 19.13 Protostellar Outflow** This view of the Orion molecular cloud shows the outflow from a newborn star, still surrounded by nebular gas. The inset shows a pair of jets called HH1 and HH2, formed when matter falling onto another protostar (still obscured by the dusty cloud fragment from which it formed) creates a pair of high-speed gas jets perpendicular to the flattened protostellar disk. Several more Herbig–Haro objects can be seen at the top right of the main image—one of them resembling a “waterfall.” (AURA; NASA)




## 19.5 Shock Waves and Star Formation

The subject of star formation is really much more complicated than the preceding discussion suggests. Interstellar space is populated with many kinds of clouds, fragments, protostars, stars, and nebulae, all interacting in a complex fashion and each type of object affecting the behavior of all the other types. For example, the presence of an emission nebula in or near a molecular cloud probably influences the evolution of the entire region. We can easily imagine expanding waves of matter driven outward by the high temperatures and pressures in the nebula. As the waves crash into the surrounding molecular cloud, interstellar gas tends to pile up and become compressed. Such a shell of gas, rushing rapidly through space, known as a **shock wave**, can push ordinarily thin matter into dense sheets, just as a plow pushes snow.

Many astronomers regard the passage of a shock wave through interstellar matter as the triggering mechanism needed to initiate star formation in a galaxy. Calculations show that when a shock wave encounters an interstellar cloud, it races around the thinner exterior of the cloud more rapidly than it can penetrate the cloud’s thicker interior.

Thus, shock waves do not blast a cloud from only one direction, but effectively squeeze it from many directions. Atomic bomb tests have experimentally demonstrated this squeezing: Shock waves created in the blast tend to surround buildings, causing them to be blown together (imploded) rather than apart (exploded). The “contracting fragment” in Figure 19.8 may well have been triggered by the shock wave from the M20 nebula. Note the correspondence between the shock-compressed region at the lower right and the high-density molecular gas revealed by radio studies (Figure 18.19). Once shock waves have begun compressing an interstellar cloud, natural gravitational instabilities take over, dividing the cloud into the fragments that eventually form stars.

Emission nebulae are by no means the only generators of interstellar shock waves. At least four other driving forces are available: the relatively gentle deaths of old stars in the form of planetary nebulae (to be discussed in Chapter 20); the much more violent ends of certain stars in supernova explosions (Chapter 21); the spiral-arm waves that plow through the Milky Way (Chapter 23); and interactions between galaxies (Chapter 24). Supernovae are by far the most energetic, and probably also the most efficient, means of piling up matter into dense clumps. However, they



◀ **FIGURE 19.14 Generations of Star Formation** (a) Star birth and (b) shock waves lead to (c) more star births and more shock waves in a continuous cycle of star formation in many areas of our Galaxy.

are relatively few and far between, so the other mechanisms may be more important overall in triggering star formation. Although the evidence is somewhat circumstantial, the presence of young (and thus fast-forming) O- and B-type stars in the vicinity of supernova remnants does suggest that the birth of stars is often initiated by the violent, explosive deaths of others.

This picture of shock-induced star formation is complicated by the fact that O- and B-type stars form quickly, live briefly, and die explosively. These massive stars, themselves perhaps born of a passing shock wave, may in turn create new shock waves, either through the expanding nebular gas produced by their births or through their explosive deaths. The new shock waves can produce “second-generation” stars, which in turn will explode and give rise to still more shock waves, and so on. As depicted in Figure 19.14, star formation resembles a chain reaction. Other, lighter stars are also formed in the process, of course, but they are largely “along for the ride.” It is the O- and B-type stars that drive the star-formation wave through the cloud.

Observational evidence lends some support to this chain-reaction picture. Groups of stars nearest molecular clouds do indeed appear to be the youngest, whereas those farther away seem to be older. Figure 19.15 shows an *HST* image of a star-forming region in the galaxy NGC 4214, which lies some 13 million light-years from Earth. A series of bright emission nebulae, powered by hot young stars, can be seen, suggesting that a wave of star formation recently swept across the region, triggering the sequence seen here.



▲ **FIGURE 19.15 A Wave of Star Formation?** A group of star-forming regions in the galaxy NGC 4214 possibly displays several generations in a sequential chain of star formation. (NASA)

**CONCEPT Check**

- ✓ Why might we expect multiple episodes of star formation to occur in some locations?

## 19.6 Star Clusters

The end result of the collapse of a cloud is a group of stars, all formed from the same parent cloud and lying in the same region of space. Such a collection of stars is called a **star cluster**. Figure 19.16 shows a spectacular view of a newborn star cluster and (part of) the interstellar cloud from which it came.


Because all the stars formed at the same time out of the same cloud of interstellar gas and under the same environmental conditions, clusters are near-ideal “laboratories” for stellar studies—not in the sense that astronomers can perform experiments on them, but because the properties of the stars are very tightly constrained. The only factor distinguishing one star from another in the same cluster is mass, so theoretical models of star formation and evolution can be compared with reality without the complications introduced by the broad spreads in age, chemical composition, and place of origin found when we consider all stars in our Galactic neighborhood.

## Clusters and Associations

Figure 19.17(a) shows a small star cluster called the Pleiades, or Seven Sisters, a well-known naked-eye object in the constellation Taurus, lying about 120 pc from Earth. This type of loose, irregular cluster, found mainly in the plane of the Milky Way (see Figure 18.4), is called an **open cluster**. Open clusters typically contain from a few hundred to a few tens of thousands of stars and are a few parsecs across.

Figure 19.17(b) shows the H–R diagram of stars in the Pleiades. The cluster contains stars in almost all parts of the main sequence—only the very brightest main-sequence stars are missing. (The brightest six or seven stars in the diagram have just left the main sequence, as will be discussed in Chapter 20.) Thus, even though we have no direct evidence of the cluster’s birth, we can estimate its age as less than about 100 million years, the lifetime of a main-sequence B-type star. (Sec. 17.8) If all the stars in the cluster formed at the same time, then the red stars must be young, too. The wisps of leftover gas evident in the photograph are further evidence of the cluster’s relative youth. In addition, the system is abundant in heavy elements that (as we will see) could have been created only within the cores of many generations of ancient stars long since perished.



Less massive, but more extended, clusters are known as **associations**. These clusters typically contain no more than a few hundred bright stars, but may span many tens of parsecs. Associations tend to be rich in very young stars. Those



▲ **FIGURE 19.16 Newborn Cluster** The star cluster NGC 3603 and part of the larger molecular cloud in which it formed. The cluster contains about 2000 bright stars and lies some 20,000 light-years from Earth. Radiation from its most massive stars has cleared a cavity in the cloud several light-years across. The inset shows the central area more clearly, revealing many small stars less massive than the Sun. (ESO; NASA)





(a)



(b)



(a)



(b)



▲ **FIGURE 19.17 Open Cluster** (a) The Pleiades cluster (also known as the Seven Sisters because only six or seven of its stars can be seen with the naked eye) lies about 400 light-years from the Sun. (b) An H–R diagram for all the stars of this well-known open cluster. (AURA)

▲ **FIGURE 19.18 Globular Cluster** (a) The globular cluster Omega Centauri is approximately 16,000 light-years from Earth and spans some 130 light-years in diameter. (b) A H–R diagram of some of its stars. (P. Seitzer)

containing many pre-main-sequence T Tauri stars are known as *T associations*, whereas those with prominent O- and B-type stars, such as the Trapezium in Orion (see Figure 19.20a on page 491), are called *OB associations*. As a class, associations tend to be very loosely bound—if they are bound at all. Many appear to be expanding freely into space and dissolving following their formation. It is quite likely that the main difference between associations and open clusters is simply the efficiency

(as measured by the fraction of gas that eventually ends up in stars) with which stars formed from the parent cloud.

Figure 19.18(a) shows a very different type of star cluster, called a **globular cluster**. All globular clusters are roughly spherical (which accounts for their name), are generally found away from the Milky Way plane, and contain hundreds of thousands, and sometimes millions, of stars spread out over about 50 pc. Figure 19.18(b) is an H–R diagram of the cluster



shown, which is called Omega Centauri. Notice the many differences between this H–R diagram and that of Figure 19.17(b)—globular clusters present a stellar environment very different from that of open clusters like the Pleiades. The distance to Omega Centauri cluster has been determined by a variation on the method of spectroscopic parallax, applied to the entire cluster rather than to individual stars. ∞ (Sec. 17.6) It lies about 5000 pc from Earth.

The most outstanding spectroscopic feature of globular clusters is their lack of upper-main-sequence stars. Astronomers in the 1920s and 1930s, working with instruments incapable of detecting stars fainter than about 1 solar luminosity at the distances of globular clusters, and having no theory of stellar evolution to guide them, were puzzled by the H–R diagrams they saw when they looked at the globular clusters. Indeed, a comparison of just the top halves of the diagrams (so that the lower main sequences cannot be seen) reveals few similarities between Figures 19.17(b) and 19.18(b).

Most globular clusters contain no main-sequence stars with masses greater than about 0.8 times the mass of the Sun. The more massive O- through F-type stars have long since exhausted their nuclear fuel and disappeared from the main sequence (in fact becoming the red giants and other luminous stars above the main sequence, as we will see in Chapter 20). ∞ (Sec. 17.8) From the theory of stellar evolution (Chapter 20), the A-type stars in Figure 19.18b are now known to be stars at much later stages in their evolution that just happen to be passing through the location of the upper main sequence. On the basis of these and other observations, astronomers estimate that most globular clusters are at least 10 billion years old—they contain the oldest known stars in our Galaxy.

Other observations confirm the great ages of globular clusters. For example, their spectra show few heavy elements, implying that these stars formed in the distant past, when heavy elements were much less abundant than they are today (Chapter 21). Astronomers speculate that the 150 or so globular clusters observed today are just the survivors of a much larger population of clusters that formed long ago.

## Clusters and Nebulae

How many stars form in a cluster, and of what type are they? How much gas is left over? What does the collapsed cloud look like once star formation has run its course? At present, although the main stages in the formation of individual stars (stages 3–7) are becoming clearer, the answers to these more general questions (involving stages 1 and 2) are still sketchy. They await a more thorough understanding of the star-formation process.

In general, the more massive the collapsing region, the more stars are likely to form there. In addition, we know from H–R diagrams of observed stars that low-mass stars are much more common than high-mass ones. ∞ (Sec. 17.8) For every O- or B-type giant, hundreds or even thousands of G-, K-, and M-type dwarfs may form. The precise number of stars of any given mass or spectral type likely depends in a complex (and

poorly understood) way upon conditions within the parent cloud. The same is true of the *efficiency* of star formation—the fraction of the total mass that actually finds its way into stars—which determines the amount of leftover material. However, if, as is usually the case, one or more O- or B-type stars form, their intense radiation and winds will cause the surrounding gas to disperse rapidly, leaving behind a young star cluster.


## The Cluster Environment

In recent years, astronomers have come to realize that physical interactions—close encounters and even collisions—between protostars within a star cluster may be very important in determining the properties of the stars that eventually form. Supercomputer simulations of star-forming clouds suggest that, while the seven stages presented earlier (and listed in Table 19.1) remain a good description of the overall formation process, the sequence of events leading to a main-sequence star can be strongly influenced by events within the cluster itself. Figure 19.19 presents frames from two such simulations, illustrating some of the interactions just described. Note the “clumpy” nature of the star-formation process. Stars tend to form as small groups in dense pockets of gas, and these groups subsequently merge to form the larger cluster.


The simulations reveal that the strong gravitational fields of the most massive protostars give them a competitive advantage over their smaller rivals in attracting gas from the surrounding nebula, causing the giant protostars to grow even faster. However, as the most massive stars grow and heat their surroundings, it becomes more difficult for them to accrete new gas. At the same time, encounters between stars usually disrupt the smaller protostellar disks, terminating the growth of the central protostars and ejecting planets and low-mass brown dwarfs from the disk into intracluster space. In dense clusters these interactions may even lead to mergers and further growth of massive objects. Thus, even before the energetic newborn O and B stars expel the cluster gas, the formation of a few large bodies can significantly inhibit the growth of smaller ones.\*

All these considerations clearly illustrate the important role played by a future star’s environment in the star-formation process and provide important insight into why low-mass stars are so much more common than high-mass stars. The first few massive bodies to form tend to prevent the formation of additional high-mass stars by stealing their “raw material” and ultimately disrupting the environment in which other stars are growing. This tendency also helps explain the existence of brown dwarfs, by providing at least two natural ways (disk destruction and gas dispersion) in which star formation can stop before nuclear fusion begins in a growing stellar core. *Discovery 19-2* describes another

\*Compare this picture of large objects dominating the accretion process at the expense of the smaller bodies around them with the standard view of planet formation presented in Chapter 15. ∞ (Sec. 15.1–3)



(a)




(b)

system in which the gas-dispersal process may be almost complete.

Young star clusters are often shrouded in gas and dust, making them hard to see in visible light. However, infrared observations clearly demonstrate that star clusters really are found within star-forming regions. Figure 19.20 compares optical and infrared views of the central regions of the Orion Nebula. The optical image in Figure 19.20(a) shows the Trapezium, the group of four bright stars responsible for ionizing the nebula; the infrared image in Figure 19.20(b) reveals an extensive cluster of stars within and behind the visible nebula. This remarkable infrared image shows many stages of star formation, with nearly 1000 new stars forming. The speckled green fuzz arises when jets of gas shoot out from those young stars and ram into the surrounding cloud.

**Interactive MA** **FIGURE 19.19 Protostellar Collisions** In the congested environment of a young cluster, star formation is a competitive and violent process. (a) These frames show how stars form in small clumps at different locations in the cloud. The clumps subsequently merge, in this case into a small association a few hundred times the mass of the Sun. (b) Large protostars may grow by “stealing” gas from smaller ones, and the extended disks surrounding most protostars can lead to collisions and mergers. This frame from another simulation shows a small star cluster emerging from an interstellar cloud that originally contained about 50 solar masses of material, distributed over a volume 1 light-year across. (*I. Bonnell and M. Bate*)



**Interactive MA** **FIGURE 19.20 Young Stars in Orion** (a) A short-exposure visible-light image (observed with a filter that is transparent mainly to certain emission lines of oxygen) shows the central regions of the Orion Nebula and four bright O-type stars known as the Trapezium, but few obvious other stars. (b) A *Spitzer Space Telescope* view of the same part of the nebula shows an extensive star cluster containing stars of many masses, possibly including many brown dwarfs (see also Figures 5.29c and d as well as 19.9). (*Lick Observatory; NASA*)

## DISCOVERY 19-2





## Eta Carinae

At the heart of the Carina emission nebula (shown in the main figure below) lies a remarkable object called Eta Carinae (object at bottom right). With an estimated mass of around 100 times the mass of the Sun and a luminosity of 5 million times the solar value, Eta Carinae is one of the most massive stars known. Formed probably only a few hundred thousand years ago, this star has had an explosive, though brief, life. In the mid-19th century, Eta Carinae produced an outburst that made it one of the brightest stars in the southern sky (even though it lies some 2200 pc away from Earth, a very long way compared to most of the bright stars visible in our night sky). During this “Great Eruption,” which peaked in 1843, the star expelled more than 2 solar masses of material in less than a decade and released as much visible energy as a supernova explosion (see Chapter 21), yet it somehow survived the event.


The images at right show the most active part of the nebula: At top is a



(ESO)



(NASA)





*Chandra X-ray Telescope* image that gives a glimpse of the object’s violence, and at bottom a *Hubble Telescope* image that was carefully processed to reveal fine detail, is the highest-resolution view of the explosion obtained to date. Dust lanes, tiny condensations in the outflowing material, and dark radial streaks of unknown origin all appear with exquisite clarity. The star itself is the white dot at the center of the image. The two ends of the “peanut” (at the top right and bottom left) are blobs of material ejected in the 1843 outburst, now racing away from the star at hundreds of kilometers per second—perhaps enough to expel the surrounding nebular gas and convert the Carina Nebula into the Carina Cluster. Perpendicular to the line joining these two blobs is a thin disk of gas, also moving outward at high speed.

The details of the events leading to the Eta Carinae outburst are unclear. Quite possibly, such episodes of violent activity are the norm for supermassive stars. In 2005, astronomers discovered that Eta Carinae has a binary companion, an even hotter, but fainter star orbiting just 11 AU away—not much more than Eta Carinae’s estimated radius of 5 AU.

Many researchers suspect that the interaction between the two stars may have been responsible for the Great Eruption. However, although a few comparable outbursts have been observed in other galaxies, they are so rare that astronomers still do not know what constitutes “typical” behavior for such exotic objects.

## Cluster Lifetimes

Eventually, star clusters dissolve into individual stars. In some cases, the irregular star formation process illustrated in Figure 19.19 simply leaves the newborn cluster gravitationally unbound. In others, the ejection of left-over gas reduces the cluster's mass so much that it becomes unbound and quickly disperses. In clusters that survive the early gas-loss phase, stellar encounters tend to eject the lightest stars from the cluster, just as the gravitational slingshot effect can propel spacecraft around the solar system.  (Discovery 6-1) At the same time, the tidal gravitational field of the Milky Way Galaxy slowly strips outlying stars from the cluster.  (Sec. 7.6) Occasional distant encounters with giant molecular clouds also tend to remove stars from a cluster. Even a near miss may disrupt the cluster entirely.

As a result of all these influences, most open clusters break up in a few hundred million years, although the actual lifetime depends on the cluster's mass and its location in the

Galaxy. Loosely bound associations may survive for only a few tens of millions of years, whereas some very massive open clusters are known from their H-R diagrams to be almost 5 billion years old. In a sense, only when a star's parent cluster has completely dissolved is the star-formation process really complete. The road from a gas cloud to a single, isolated star like the Sun is long and tortuous indeed!

Take another look at the sky one clear, dark evening. Ponder all of the cosmic activity you have learned about as you peer upward at the stars. After studying this chapter, you may find that you have to modify your view of the night sky. Even the seemingly quiet nighttime darkness is dominated by continual change.

### CONCEPT Check

- ✓ If stars in a cluster all form at the same time, how can some influence the formation of others?

**The Big Question** When did the first stars form? We observe stars forming today throughout the Milky Way and myriad other galaxies, and studies of distant stars imply that they formed even more efficiently billions of years ago. Astronomers are pushing back the veil of ancient star formation, trying to understand how and when conditions in the early universe first allowed gas without walls—namely, stars—to ignite as brilliant balls of fire.


# Chapter Review

## SUMMARY

**1** Stars form when an interstellar cloud collapses under its own gravity and breaks up into pieces comparable in mass to our Sun. A cold interstellar cloud may fragment into many smaller clumps of matter, from which stars eventually form. The evolution of the contracting cloud can be represented as an **evolutionary track** (p. 475) on the Hertzsprung–Russell diagram. As a collapsing prestellar fragment heats up and becomes denser, it eventually becomes a **protostar** (p. 474)—a warm, very luminous object that emits mainly infrared radiation. Eventually, a protostar's central temperature becomes high enough for hydrogen fusion to begin, and the protostar becomes a star.




**2** For a star like the Sun, the whole formation process takes about 50 million years. More massive stars pass through similar formation stages, but much more rapidly. Stars less massive than the Sun take much longer to form. The **zero-age main sequence** (p. 477) is the region in the H-R diagram where stars



lie when the formation process is over. Mass is the key property in determining a star's characteristics and life span. The most massive stars have the shortest formation times and main-sequence lifetimes. At the other extreme, some low-mass fragments never reach the point of nuclear ignition and become **brown dwarfs** (p. 478).

**3** Many of the objects predicted by the theory of star formation have been observed in real astronomical objects. The dark interstellar regions near emission nebulae often provide evidence of cloud fragmentation and protostars. Radio telescopes are used in studying the early phases of cloud contraction and fragmentation; infrared observations allow us to see later stages of the process. Many well-known emission nebulae, lit by several O- and B-type stars, are partially engulfed by molecular clouds, portions of which are fragmenting and contracting, with smaller sites forming protostars.



**4** Protostars can produce powerful protostellar winds. These winds encounter less resistance in the directions perpendicular to a star's protostellar disk and often expel two jets of matter in the


directions of the protostar's poles in a **bipolar flow** (p. 483). The protostellar winds gradually destroy the disk, and eventually the wind flows away from the star equally in all directions. **Shock waves** (p. 484) are produced as young hot stars ionize the surrounding gas, forming emission nebulae.



These shock waves can compress other interstellar clouds and trigger more star formation, possibly producing chain reactions of star formation in molecular cloud complexes.

**5** A single collapsing and fragmenting cloud can give rise to hundreds or thousands of stars—a **star cluster** (p. 486). The formation of the most massive stars may play an important role in suppressing the further formation of stars from lower-mass cluster members. **Open clusters** (p. 486), with a few hundred to a

few thousand stars, are found mostly in the plane of the Milky Way. They typically contain many bright blue stars, indicating that they formed relatively recently. **Globular clusters** (p. 487) are found mainly away from the Milky Way plane and may contain millions of stars. They include no main-sequence stars much more massive than the Sun, indicating that they formed long ago. Infrared observations have revealed young star clusters or associations in several emission nebulae. Eventually, clusters break up into individual stars, although the entire process may take hundreds of millions or even billions of years.



**MasteringAstronomy**® For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- LO1** Briefly describe the basic chain of events leading to the formation of a star like the Sun.
- What are the roles of heat, rotation, and magnetism in the process of stellar birth?
- What is an evolutionary track?
- Why does the evolution of a protostar slow down as the star approaches the main sequence?
- LO2** In what ways do the formative stages of high-mass stars differ from those of stars like the Sun?
- What are brown dwarfs?
- What are T Tauri stars?
- LO3 POS** Stars live much longer than we do, so how do astronomers test the accuracy of theories of star formation?
- At what evolutionary stages must astronomers use radio and infrared radiation to study prestellar objects? Why can't they use visible light?
- LO4** What is a shock wave? Of what significance are shock waves in star formation?
- POS** Explain the usefulness of the H-R diagram in studying the evolution of stars. Why can't evolutionary stages 1–3 be plotted on the diagram?
- LO5** What do star clusters and associations have to do with star formation?
- Compare and contrast the observed properties of open star clusters and globular star clusters.
- POS** How can we tell whether a star cluster is young or old?
- In the formation of a star cluster with a wide range of stellar masses, is it possible for some stars to die out before others have finished forming? Do you think this will have any effect on the cluster's formation?

## Conceptual Self-Test: Multiple Choice

- If a newly forming star has an excess of heat, then it will likely have (a) more gravity; (b) less gravity; (c) a slower contraction rate; (d) a rapid contraction rate.
- The gravitational contraction of an interstellar cloud is primarily the result of its (a) mass; (b) composition; (c) diameter; (d) pressure.
- The interstellar cloud from which our Sun formed was (a) slightly larger than the Sun; (b) comparable in size to Saturn's orbit; (c) comparable in mass to the solar system; (d) thousands of times more massive than the Sun.
- A protostar that will eventually turn into a star like the Sun is significantly (a) smaller; (b) more luminous; (c) fainter; (d) less massive than the Sun.
- Prestellar objects in which nuclear fusion never starts are referred to as (a) terrestrial planets; (b) brown dwarfs; (c) protostars; (d) globules.

6. The current theory of star formation is based upon (a) amassing evidence from many different regions of our Galaxy; (b) carefully studying the births of a few stars; (c) systematically measuring the masses and rotation rates of interstellar clouds; (d) observations made primarily at short wavelengths.
7. **VIS** If the initial interstellar cloud in Figure 19.14 (“Generations of Star Formation”) were much more massive, the result would be (a) the formation of more stars; (b) contraction of the cloud due to stronger gravitational attraction; (c) stars forming closer together; (d) stronger shock waves.
8. **VIS** One of the primary differences between the Pleiades cluster, shown in Figure 19.17(a), and Omega Centauri, shown in Figure 19.18(a), is that the Pleiades cluster is much (a) larger; (b) younger; (c) farther away; (d) denser.
9. **VIS** If the H–R diagram shown in Figure 19.18(b) (“Globular Cluster”) were redrawn to illustrate a much younger cluster, the main-sequence turnoff would shift to (a) higher temperature; (b) higher pressure; (c) higher frequency; (d) a spectral classification of K or M.
10. A typical open cluster will dissolve in about the same amount of time as the time since (a) North America was first visited by Europeans; (b) dinosaurs walked on Earth; (c) Earth was formed; (d) the universe formed.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

1. ●● In order for an interstellar gas cloud to contract, the average speed of its constituent particles must be less than half the cloud’s escape speed. ∞ (More Precisely 8-1) Will a (spherical) molecular hydrogen cloud with a mass of 1000 solar masses, a radius of 10 pc, and a temperature of 10 K begin to collapse? Why or why not?
2. ● A protostar on the Hayashi track evolves from a temperature of 3500 K and a luminosity 5000 times that of the Sun to a temperature of 5000 K and a luminosity of 3 solar units. What is the protostar’s radius (a) at the start and (b) at the end of the evolution?
3. ●● Use the radius–luminosity–temperature relation to explain how a protostar’s luminosity changes as it moves from stage 4 (temperature 3000 K, radius  $2 \times 10^8$  km) to stage 6 (temperature 4500 K, radius  $10^6$  km). What is the change in absolute magnitude? ∞ (Sec. 17.2)
4. ● What is the (approximate) absolute magnitude of a stage-5 protostar? (See Figure 19.6.)
5. ● By how many magnitudes does a 3-solar-mass star decrease in brightness as it evolves from stage 4 to stage 6? (See Figure 19.7.)
6. ●● Use the H–R diagrams in this chapter to estimate by what factor a 1000-solar-luminosity, 3000-K protostar is larger than a main-sequence star of the same luminosity.
7. ● What is the luminosity, in solar units, of a brown dwarf whose radius is 0.1 solar radius and whose surface temperature is 600 K (0.1 times that of the Sun)?
8. ●● What is the maximum distance at which the brown dwarf in the previous problem could be observed by a telescope of limiting apparent magnitude (a) 18, (b) 30?

## Activities

### Collaborative

1. Open clusters are generally found in the plane of the Galaxy. If you can see the hazy band of the Milky Way arcing across your night sky—in other words, if you are far from city lights and looking at an appropriate time of night and year—you can simply sweep with your binoculars along the Milky Way. Numerous “clumps” of stars will pop into view. Many will turn out to be open star clusters. For more detail, they are best viewed through a small telescope. The most easily visible clusters are those in the Messier catalog, although there are many others besides. Interesting Messier clusters and associations include M6, M7, M11, M35, M37, M44, M45, M52, M67, and M103. How many of these can you find? How many clusters can you find that are not on Messier’s list?
2. Globular star clusters are harder to find. They are intrinsically larger, but they are also much farther away and therefore appear smaller in the sky. The most famous globular cluster visible from the Northern Hemisphere is M13 in the constellation Hercules, visible on spring and summer evenings. It contains a half million or so of the Galaxy’s most ancient stars. It may

be glimpsed through binoculars as a little ball of light located about one-third of the way from the star Eta to the star Zeta in the Keystone asterism of the constellation Hercules. Telescopes reveal this cluster as a magnificent, symmetrical grouping of stars. Can you find the following well-known globular clusters: M3, M4, M5, M13, M15? Look online or in a star chart for details on how to locate them.

### Individual

1. The constellation Orion the Hunter is prominent in the winter sky. Its most noticeable feature is a short, straight row of three medium-bright stars: the famous belt of Orion. A line of stars extends from the easternmost star of the belt, toward the south. This is Orion’s sword. Toward the bottom of the sword is the sky’s most famous emission nebula, the Orion Nebula (M42). Observe the Orion Nebula with your eye, with binoculars, and with a telescope. What is its color? How can you account for this? With the telescope, try to find the Trapezium, a grouping of four stars in the center of the nebula. These are hot, young stars; their energy causes the Orion Nebula to glow.



# 20

## Stellar Evolution

### THE LIFE AND DEATH OF A STAR

After reaching the main sequence, a newborn star changes little in outward appearance for more than 90 percent of its lifetime. However, at the end of that period, as the star begins to run out of fuel and die, its properties once again change greatly. Aging stars travel along evolutionary tracks that take them far from the main sequence as they end their lives. In this and the next two chapters, we will study the evolution of stars during and after their main-sequence burning stages.

We will find that the ultimate fate of a star depends primarily on its mass—although interactions with other stars can also play a decisive role—and that the final states of stars can be strange, indeed. By continually comparing theoretical calculations with detailed observations of stars and binaries of all types, astronomers have refined the theory of stellar evolution into a precise and powerful tool for understanding the universe.

**The Big Picture** The story of the birth, development, and death of stars is one of the greatest accomplishments of 20th-century science. Yet, ironically, no one has ever seen even a single star pass through all of its many varied changes. Like archaeologists who examine bones and artifacts from long ago to learn more about the evolution of human culture, astronomers observe stars of different ages to construct a consistent model of how stars evolve over billions of years.

**LEFT:** Resembling a cosmic hourglass or celestial butterfly, this striking image captures hot gas released by a dying star about 3800 light-years away. Known as NGC 6302, or informally the Bug Nebula, this complex object is a planetary nebula—an old star shedding its outer layers over light-year dimensions as it ends its life. Its peculiar shape results from a belt of dust (dark lane at center) that obscures the dying star and partially blocks the rolling cauldrons of outwardly expelled gas. (STScI)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Explain why stars evolve off the main sequence.
- 2 Outline the events that occur as a Sun-like star evolves from the main sequence to the giant branch.
- 3 Explain how the Sun will eventually come to fuse helium in its core, and describe what happens when that occurs.
- 4 Summarize the stages in the death of a typical low-mass star, and describe the resulting remnant.
- 5 Contrast the evolutionary histories of high-mass and low-mass stars.
- 6 Present the observations that help verify the theory of stellar evolution.
- 7 Explain how the evolution of stars in binary systems may differ from that of isolated stars.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.



## 20.1 Leaving the Main Sequence

Most stars spend most of their lives on the main sequence. A star like the Sun, for example, after spending a few tens of millions of years in its formative stages (1–6 in Chapter 19), resides on or near the main sequence (stage 7) for 10 billion years before turning into something else. ∞ (Sec. 19.2) That “something else” is the main topic of this chapter.

### Observing Stellar Lifetimes

No one has ever witnessed the complete evolution of any star, from birth to death. Stars take a very long time—millions, billions, even trillions of years—to evolve. ∞ (Secs. 17.8, 19.2) Yet, in less than a century, astronomers devised a comprehensive theory of stellar evolution that is one of the best-tested in all of astronomy. How can we talk so confidently about what took place billions of years in the past, and what will happen billions of years in the future? The answer is that we can observe billions of stars in the universe, enough to see examples of every stage of stellar development, allowing us to test and refine our theoretical ideas. ∞ (Sec. 1.2) Just as we can piece together a picture of the human life cycle by studying a snapshot of all the residents of a large city, so we can construct a picture of stellar evolution by studying the myriad stars we see in the night sky.

Note that astronomers use the term “evolution” here to mean change *during the lifetime of an individual star*. Contrast this with the usage of the term in biology, where it refers to changes in the characteristics of a *population* of plants or animals over many generations. In fact, as we will see in Chapter 21, populations of stars do evolve in the latter “biological” sense, as the overall composition of the interstellar medium (and hence of each new stellar generation) changes slowly over time due to nuclear fusion in stars. However, in astronomical parlance, “stellar evolution” always refers to changes during a single stellar lifetime.

### Structural Change

On the main sequence, a star slowly fuses hydrogen into helium in its core. This process of nuclear fusion is called **core hydrogen burning**. In Chapter 16, we saw how the proton–proton fusion chain powers the Sun. ∞ (Sec. 16.6) Here, by the way, is another instance where astronomers use a fairly familiar term in a quite unfamiliar way: To astronomers “burning” always means nuclear fusion in a star’s core and not the chemical reaction (such as the combustion of wood or gasoline in air) we would normally think of in everyday speech. Chemical burning does not directly affect atomic nuclei.

As discussed in Chapter 16 (see Figure 16.4), a main-sequence star is in a state of *hydrostatic equilibrium*, in which pressure’s outward push exactly counteracts gravity’s inward pull. ∞ (Sec. 16.2) This is a stable balance between

gravity and pressure in which a small change in one always results in a small compensating change in the other. You should keep that figure in mind as you study the various stages of stellar evolution described next. Much of a star’s complex behavior can be understood in these simple terms. Eventually, however, as the hydrogen in the core is consumed, the star’s internal balance starts to shift, and both its internal structure and its outward appearance begin to change more rapidly: The star leaves the main sequence.

Once a star begins to move away from the main sequence, its days are numbered. The post-main-sequence stages of stellar evolution—the end of a star’s life—depend critically on the star’s mass. As a rule of thumb, we can say that low-mass stars die gently, whereas high-mass stars die catastrophically. The dividing line between these two very different outcomes lies around eight times the mass of the Sun, and in this chapter we will refer to stars of more than 8 solar masses as “high-mass” stars. Within both the “high-mass” and the “low-mass” (i.e., less than 8 solar masses) categories, there are substantial variations, some of which we will point out as we proceed.

Rather than dwelling on the many details, we will concentrate on a few representative evolutionary sequences. We begin by considering the evolution of a fairly low-mass star like the Sun. The stages described in the next few sections pertain to the Sun as it nears the end of its fusion cycle 5 billion years from now. The numbers continue the sequence begun in Chapter 19. In fact, most of the qualitative features of the discussion apply to any low-mass star, although the exact numbers vary considerably. Later, we will broaden our discussion to include all stars, large and small.

### PROCESS OF SCIENCE Check

✓ How can astronomers “see” stars evolve in time?

## 20.2 Evolution of a Sun-Like Star

The surface of a main-sequence star like the Sun occasionally erupts in flares and spots, but for the most part the star does not exhibit any sudden, large-scale changes in its properties. Its average surface temperature remains fairly constant, whereas its luminosity increases very slowly with time. The Sun has roughly the same surface temperature as it had when it formed nearly 5 billion years ago, even though it is some 30 percent brighter than it was at that time.

This state of affairs cannot continue indefinitely. Eventually, drastic changes occur in the star’s interior structure. After about 10 billion years of steady core hydrogen burning, a Sun-like star begins to run out of fuel. It’s a little like an automobile cruising effortlessly along a highway at constant speed for many hours, only to have the engine suddenly cough and sputter as the gas gauge reaches empty. Unlike automobiles, though, stars are not easy to refuel.

## Stage 8: The Subgiant Branch

As nuclear fusion proceeds, the composition of the star's interior changes as its hydrogen fuel is depleted. Figure 20.1 illustrates the increase in helium abundance and the corresponding decrease in hydrogen abundance that take place in the stellar core as the star ages. Three cases are shown: (a) the chemical composition of the original core, (b) the composition after 5 billion years, and (c) the composition after 10 billion years. Case (b) represents approximately the present state of our Sun.




The star's helium content increases fastest at the center, where temperatures are highest and the burning is fastest. The helium content also increases near the edge of the core, but more slowly because the burning rate is less rapid there. The inner, helium-rich region becomes larger and more deficient in hydrogen as the star continues to shine. Eventually, about 10 billion years after the star arrived on the main sequence (Figure 20.1c), hydrogen becomes depleted at the center, the nuclear fires there subside, and the location of principal burning moves to higher layers in the core. An inner core of nonburning pure helium starts to grow.

Without nuclear burning to maintain it, the outward-pushing gas pressure weakens in the helium inner core. However, the inward pull of gravity does not. Once the outward push against gravity is relaxed—even a little—structural changes in the star become inevitable. As the hydrogen is consumed, the inner core begins to contract. When all the hydrogen at the center is gone, the process accelerates.

If more heat could be generated, then the core might regain its equilibrium. For example, if helium in the core were to begin fusing into some heavier element, then energy would be created as a by-product of helium burning, and the necessary gas pressure would be reestablished. But the helium at the center cannot burn—not yet, anyway. Despite its high temperature, the core is far too cold to fuse helium into anything heavier.

Recall from Chapter 16 that a minimum temperature of about  $10^7$  K is needed to fuse hydrogen into helium. Only above that temperature do colliding hydrogen nuclei (i.e., protons) have enough speed to overwhelm the repulsive electromagnetic force between them. [∞ \(Sec. 16.6\)](#) Because helium nuclei, with two protons each, carry a greater positive charge, their electromagnetic repulsion is larger, and even higher temperatures are needed to cause them to fuse—at least  $10^8$  K. A core composed of helium at  $10^7$  K thus cannot generate energy through fusion.


The shrinkage of the helium core releases gravitational energy, driving up the central temperature and heating the overlying burning layers. The higher temperatures—now well over  $10^7$  K (but still less than  $10^8$  K)—cause hydrogen nuclei to fuse even more rapidly than before. Figure 20.2 depicts this situation, in which hydrogen is burning at a furious rate in a shell surrounding the nonburning inner core of helium “ash” in the center. This phase is known as the **hydrogen-shell-burning** stage. The hydrogen shell generates



**▲ FIGURE 20.1 Solar Composition Change** Theoretical estimates of the changes in a Sun-like star's composition show how hydrogen (yellow) and helium (blue) abundance vary within the star from birth to death (top to bottom). All these changes speed up as the nuclear burning rate increases with time.

energy faster than did the original main-sequence star's hydrogen-burning core, and the shell's energy production continues to increase as the helium core continues to shrink. Strange as it may seem, the star's response to the disappearance of the fire at its center is to get brighter!

Table 20.1 summarizes the key stages through which a solar-mass star evolves. The table is a continuation of Table 19.1, except that the density units have been changed from particles per cubic meter to the more convenient



**FIGURE 20.2 Hydrogen-Shell Burning** As a star's core converts more and more of its hydrogen into helium, the hydrogen in the shell surrounding the nonburning helium "ash" burns ever more violently. By the time the star has reached the bottom of the giant branch (around stage 8 in Table 20.1), its core has shrunk to a few tens of thousands of kilometers in diameter, and its photosphere is 10 times the star's original size.

kilograms per cubic meter and sizes are expressed as radii rather than diameters. The numbers in the "Stage" column refer to the evolutionary stages noted in the figures and discussed in the text.

After a lengthy stay on the main sequence, the star's temperature and luminosity are once again beginning to change, and we can trace these changes via the star's evolutionary track on the H–R diagram. [∞ \(Sec. 19.2\)](#) Figure 20.3 shows the star's path away from the main sequence, labeled as stage 7. The star first evolves to the right on the diagram, its surface temperature dropping whereas its luminosity increases only slightly. By stage 8, the star's radius has increased to about three times the radius of the Sun. The star at this stage is called a **subgiant**. Its roughly horizontal path from its main-sequence location (stage 7) to stage 8 on the figure is called the **subgiant branch**.

## Stage 9: The Red-Giant Branch


Our aging star is now far from the main sequence and no longer in stable equilibrium. The helium core is unbalanced and shrinking. The rest of the core is also unbalanced, fusing hydrogen into helium at an ever-increasing rate. The gas pressure produced by this enhanced hydrogen burning causes the star's nonburning outer layers to increase in radius. Not even gravity can stop this inexorable change. While the core is shrinking and heating up, the overlying layers are expanding and cooling. The star is on its way to becoming a red giant. The transformation from normal main-sequence star to elderly red giant takes about 100 million years.

By stage 8, the star's surface temperature has fallen to the point at which much of the interior is opaque to the radiation

**TABLE 20.1** Evolution of a Sun-Like Star

Stage	Approximate Time to Next Stage (yr)	Central Temperature ( $10^6$ K)	Surface Temperature (K)	Central Density ( $\text{kg}/\text{m}^3$ )	Radius (km)	Radius (solar radii)	Object
7	$10^{10}$	15	6000	$10^5$	$7 \times 10^5$	1	Main-sequence star
8	$10^8$	50	4000	$10^7$	$2 \times 10^6$	3	Subgiant branch
9	$10^5$	100	4000	$10^8$	$7 \times 10^7$	100	Helium flash
10	$5 \times 10^7$	200	5000	$10^7$	$7 \times 10^6$	10	Horizontal branch
11	$10^4$	250	4000	$10^8$	$4 \times 10^8$	500	Asymptotic-giant branch
12	$10^5$	300	100,000	$10^{10}$	$10^4$	0.01	Carbon core
		—	3000	$10^{-17}$	$7 \times 10^8$	1000	Planetary nebula*
13	—	100	50,000	$10^{10}$	$10^4$	0.01	White dwarf
14	—	Close to 0	Close to 0	$10^{10}$	$10^4$	0.01	Black dwarf

\* Values refer to the envelope.



▲ **FIGURE 20.3 Red Giant on the H–R Diagram** As its helium core shrinks and its outer envelope expands, the star leaves the main sequence (stage 7). At stage 8, the star is well on its way to becoming a red giant. The star continues to brighten and grow as it ascends the red-giant branch to stage 9. As noted in Chapter 17, the dashed diagonal lines are lines of constant radius, allowing us to gauge the changes in the size of the star.

from within. Beyond this point, convection carries the core’s enormous energy output to the surface. One consequence of that convection is that the star’s surface temperature remains nearly constant between stages 8 and 9. The almost vertical path followed by the star between those stages is known as the **red-giant branch** of the H–R diagram. By stage 9, hydrogen shell burning in the still-shrinking core is so ferocious that the giant’s luminosity is many hundreds of times the solar value. Its radius by this time is around 100 solar radii.

The red giant is huge—about the size of Mercury’s orbit. In contrast, its helium core is surprisingly small—only about a thousandth the size of the entire star, making the core just a few times larger than Earth. The central density is enormous: Continued shrinkage of the red giant’s core has compacted its helium gas to approximately  $10^8 \text{ kg/m}^3$ . Contrast this value with the  $10^{-3} \text{ kg/m}^3$  in the giant’s outermost layers, with the  $5000 \text{ kg/m}^3$  average density of Earth, and with the  $150,000 \text{ kg/m}^3$  in the present core of the Sun. About 25 percent of the mass of the entire star is packed into its planet-sized core.

A familiar example of a low-mass star in the red-giant phase is the KIII giant Arcturus (see Figure 17.15), one of the brightest stars in the sky. Its mass is about 1.5 times that of

the Sun. Currently in the hydrogen-shell-burning stage and ascending the red-giant branch, Arcturus has a radius some 21 times that of the Sun and emits about 160 times more energy than the Sun, much of it in the infrared part of the spectrum.


## Stage 10: Helium Fusion

Should the unbalanced state of a red-giant star continue, the core would eventually collapse, and the rest of the star would slowly drift into space. The forces and pressures at work inside a red giant would literally tear it apart. In fact, for stars less than about one-quarter the mass of the Sun, that is precisely what will eventually happen (in a few hundred billion years—see Section 20.3).

However, for a star like the Sun, this simultaneous shrinking and expanding does not continue indefinitely. A few hundred million years after a solar-mass star leaves the main sequence, something else happens: *Helium begins to burn in the core.* By the time the central density has risen to about  $10^8 \text{ kg/m}^3$  (at stage 9), the temperature has reached the  $10^8 \text{ K}$  needed for helium to fuse into carbon, and the central fires reignite.

The reaction that transforms helium into carbon occurs in two steps. First, two helium nuclei come together to form a nucleus of beryllium-8 ( $^8\text{Be}$ ), a highly unstable isotope that would normally break up into two helium nuclei in about  $10^{-12} \text{ s}$ . However, at the high densities found in the core of a red giant, it is possible that the beryllium-8 nucleus will encounter another helium nucleus before breakup occurs, fusing with the helium nucleus to form carbon-12 ( $^{12}\text{C}$ ). This is the second step of the helium-burning reaction. In part, it is because of the electrostatic repulsion between beryllium-8 (containing four protons) and helium-4 (containing two) that the temperature must reach  $10^8 \text{ K}$  before that step can take place.


Symbolically, we can represent this next stage of stellar fusion as follows:



Helium-4 nuclei are traditionally known as *alpha particles*. The term dates from the early days of nuclear physics, when the true nature of these particles, emitted by many radioactive materials, was unknown. Because three alpha particles are required to get from helium-4 to carbon-12, the foregoing reaction is usually called the **triple-alpha process**.

## The Helium Flash

For stars comparable in mass to the Sun, a complication arises when helium fusion begins. At the high densities found in the core, the gas has entered a new state of matter whose properties are governed by the laws of quantum mechanics (the branch of physics describing the behavior of

matter on subatomic scales), rather than by those of classical physics.  (Sec. 4.2)

Up to now, we have been concerned primarily with the nuclei—protons, alpha particles, and so on—that make up virtually all the star’s mass and that participate in the reactions that generate its energy. However, the star contains another important constituent: a vast sea of electrons stripped from their parent nuclei by the ferocious heat in the stellar interior. At this stage in our story, these electrons play an important role in determining the star’s evolution.

Under the conditions found in the stage-9 red-giant core, a rule of quantum mechanics known as the *Pauli exclusion principle* (after Wolfgang Pauli, one of the founding fathers of quantum physics) prohibits the electrons in the core from being squeezed too close together. In effect, the exclusion principle tells us that we can think of the electrons as tiny rigid spheres that can be squeezed relatively easily up to the point of contact, but that become virtually incompressible thereafter. In the language of quantum mechanics, this condition is known as *electron degeneracy*; the pressure associated with the contact of the tiny electron spheres is called **electron degeneracy pressure**.\* It has nothing to do with the *thermal pressure* (due to the star’s heat) that we have been studying up to now. In fact, in our red-giant core, the pressure resisting the force of gravity is supplied almost entirely by degenerate electrons. Hardly any of the core’s support results from “normal” thermal pressure, and this fact has dramatic consequences once the helium begins to burn.

Under normal (“nondegenerate”) circumstances, the core could react to, and accommodate, the onset of helium burning, but in the core’s degenerate state, the burning becomes unstable, with literally explosive consequences. In a star supported by thermal pressure, the increase in temperature produced by the onset of helium fusion would lead to an increase in pressure. The gas would then expand and cool, reducing the burning rate and reestablishing equilibrium, just as discussed earlier.

In the electron-supported core of a solar-mass red giant, however, the pressure is largely *independent* of the temperature. When burning starts and the temperature increases, there is no corresponding rise in pressure, no expansion of the gas, no drop in the temperature, and no stabilization of the core. Instead, the core is unable to respond to the rapidly changing conditions within it. The pressure remains more or less unchanged as the nuclear reaction rates increase, and the temperature rises rapidly in a runaway condition called the **helium flash**.


For a few hours, the helium burns ferociously. Eventually, the flood of energy released by this period of runaway fusion heats the core to the point at which normal thermal pressure once again dominates. Finally able to react to the

\*The term refers to an idealized condition in which all particles (electrons in this case) are in their lowest possible energy states. As a result, the star cannot be compressed into a more compact configuration.


energy dumped into it by helium burning, the core expands, its density drops, and equilibrium is restored as the inward pull of gravity and the outward push of gas pressure come back into balance. The core, now stable, begins to fuse helium into carbon at temperatures well above  $10^8$  K.

The helium flash terminates the giant star’s ascent of the red-giant branch of the H–R diagram. Yet, despite the violent ignition of helium in the core, the flash does *not* increase the star’s luminosity. On the contrary, the energy released in the helium flash expands and cools the core and ultimately results in a *reduction* in the energy output. On the H–R diagram, the star jumps from stage 9 to stage 10, a stable state with steady helium burning in the core. As indicated in Figure 20.4, the surface temperature is now higher than it was on the red-giant branch, but the luminosity is considerably less than at the helium flash. This adjustment in the star’s properties occurs quite quickly—in about 100,000 years.

At stage 10, our star is now stably burning helium in its core and fusing hydrogen in a shell surrounding it. The star resides in a well-defined region of the H–R diagram known as the **horizontal branch**, where core-helium-burning stars remain for a time before resuming their journey around the H–R diagram. The star’s specific position within this region is determined mostly by its mass—not its original mass, but whatever mass remains after its ascent of the red-giant



▲ **FIGURE 20.4 Horizontal Branch** A large increase in luminosity occurs as a star ascends the red-giant branch, ending in the helium flash. The star then settles down into another equilibrium state at stage 10, on the horizontal branch.



◀ **FIGURE 20.5 Helium-Shell Burning** Within a few million years after the onset of helium burning (stage 9), carbon ash accumulates in the star's inner core. Above this core, hydrogen and helium are still burning in concentric shells.

branch. The two masses differ because, during the red-giant stage, strong stellar winds eject large amounts of matter from a star's surface. As much as 20 to 30 percent of the original stellar mass may escape during that period. It so happens that more massive stars have lower surface temperatures at this stage, but all stars have roughly the same luminosity after the helium flash. As a result, stage-10 stars tend to lie along a horizontal line on the H-R diagram, with more massive stars to the right and less massive ones to the left.


### Stage 11: Back to the Giant Branch

The nuclear reactions in our star's helium core burn on, but not for long. Whatever helium exists in the core is rapidly consumed, and the dying star once again ascends the giant branch.

The triple-alpha helium-to-carbon fusion reaction—like the proton-proton and CNO-cycle hydrogen-to-helium reactions before it—proceeds at a rate that increases rapidly with temperature. At the extremely high temperatures found in the horizontal-branch core, the helium fuel doesn't last long—no more than a few tens of millions of years after the initial flash.

As helium fuses to carbon, a new carbon-rich inner core begins to form, and phenomena similar to those that took place during the earlier buildup of helium recur. Helium becomes depleted at the center of the star, and eventually fusion ceases there. The nonburning carbon core shrinks in size—even as its mass increases due to helium fusion—and heats up as gravity pulls it inward, causing the hydrogen- and helium-burning rates in the overlying layer of the core to increase. As illustrated in Figure 20.5, the star now contains a contracting carbon core surrounded by a helium-burning shell, which is in turn surrounded by a hydrogen-burning shell. The outer

envelope of the star—the nonburning layers surrounding the core—expands, much as it did earlier during the first red-giant stage. By the time it reaches stage 11 in Figure 20.6, the star has become a swollen red giant for the second time.



▲ **FIGURE 20.6 Reascending the Giant Branch** A carbon-core star reenters the giant region of the H-R diagram—this time on a track called the *asymptotic-giant branch* (stage 11)—for the same reason it evolved there the first time around: Lack of nuclear fusion at the center causes the core to contract and the overlying layers to expand.

To distinguish the second ascent of the giant branch from the first, the star's track during the second phase is often referred to as the **asymptotic-giant branch**.<sup>\*</sup> The burning rates in the shells around the carbon core are much fiercer this time around, and the star's radius and luminosity increase to values even greater than those reached at the helium flash on the first ascent. The carbon core grows in mass as more and more carbon is produced in the helium-burning shell above it, but continues to shrink in radius, driving the hydrogen-burning and helium-burning shells to higher and higher temperatures and luminosities.

### CONCEPT Check


✓ Why does a star get brighter as it runs out of fuel in its core?

## 20.3 The Death of a Low-Mass Star

Figure 20.7 illustrates the stages through which a G-type star like the Sun will pass over the course of its evolution. As our star moves from stage 10 (the horizontal branch) to stage 11 (the asymptotic-giant branch), its envelope swells, while its inner carbon core, too cool for further nuclear burning, continues to contract. If the central temperature could become high enough for carbon fusion to occur, still heavier products could be synthesized, and the newly generated energy might again support the star, restoring for a time the equilibrium between gravity and heat. However, as we will see in a moment, only high-mass stars reach temperatures high enough for this to occur.

*\*This rather intimidating term is borrowed from mathematics. An asymptote to a curve is a second curve that approaches ever closer to the first as the two are extended to infinity. Theoretically, if the star remained intact, the asymptotic-giant branch would approach the red-giant branch from the left as the luminosity increased and would effectively merge with the red-giant branch near the top of Figure 20.6. However, as we will see in Section 20.3, a Sun-like star will not live long enough for that to occur.*

**Interactive FIGURE 20.7 G-Type Star Evolution** Artist's conception of the relative sizes and changing colors of a normal G-type star (such as our Sun) during its formative stages, on the main sequence, and while passing through the red-giant and white-dwarf stages. At maximum swelling, the red giant is approximately 70 times the size of its main-sequence parent; the core of the giant is about  $\frac{1}{15}$ th the main-sequence size and would be barely discernible if this figure were drawn to scale. The duration of time spent in the various stages—protostar, main-sequence star, red giant, and white dwarf—is roughly proportional to the lengths shown in this imaginary trek through space. The star's brief stay on the horizontal and asymptotic-giant branches are not shown.




For solar-mass stars, the central temperature never reaches the 600 million K needed for a new round of nuclear reactions to occur. The red giant is very close to the end of its nuclear-burning lifetime.

## The Fires Go Out

Before the carbon core can attain the incredibly high temperatures needed for carbon ignition, its density reaches a point beyond which it cannot be compressed further. At about  $10^{10}$  kg/m<sup>3</sup>, the electrons in the core once again become degenerate, the contraction of the core ceases, and the core's temperature stops rising. This stage (stage 12 in Table 20.1) represents the maximum compression that the star can achieve—there is simply not enough matter in the overlying layers to bear down any harder.


The core density at this stage is extraordinarily high. A single cubic centimeter of core matter would weigh 1000 kg on Earth—a ton of matter compressed into a volume about the size of a grape! Yet, despite the extreme compression of the core, the central temperature is “only” about 300 million K. Some oxygen is formed via reactions between carbon and helium at the inner edge of the helium-burning shell—that is,



However, collisions among nuclei are neither frequent nor violent enough to create any heavier elements. For all practical purposes, the central fires go out once carbon has formed.

## Stage 12: A Planetary Nebula

Our aged stage-12 star is now in quite a predicament. Its inner carbon core no longer generates energy. The outer-core shells continue to burn hydrogen and helium, and as more and more of the inner core reaches its final, high-density state, the nuclear burning increases in intensity. Meanwhile, the envelope continues to expand and cool, reaching a



▲ **FIGURE 20.8 Red-Giant Instability** Buffeted by helium-shell flashes from within, and subject to the destabilizing influence of recombination, the outer layers of a red giant become unstable and enter into a series of growing pulsations. Eventually, the envelope is ejected and forms a planetary nebula.

maximum radius of about 300 times that of the Sun—big enough to engulf the planet Mars.

Around this time, the burning becomes quite unstable. The helium-burning shell is subject to a series of explosive *helium-shell flashes*, caused by the enormous pressure in the helium-burning shell and the extreme sensitivity of the triple-alpha burning rate to small changes in temperature. The flashes produce large fluctuations in the intensity of the radiation reaching the star’s outermost layers, causing those layers to pulsate violently as the envelope repeatedly is heated, expands, cools, and then contracts (Figure 20.8). The amplitude of the pulsations grows as the temperature of the core continues to increase and the nuclear burning intensifies in the surrounding shells.

Compounding the star’s problems is the increasing instability of its surface layers. Around the peak of each pulsation, the surface temperature drops below the point at which electrons can recombine with nuclei to form atoms. ∞ (Sec. 4.2) Each recombination produces additional photons, giving the gas a little extra outward “push” and causing some of it to escape. Thus, driven by increasingly intense radiation from within, and accelerated by instabilities in

both the core and the outer layers, virtually all of the star’s envelope is ejected into space in less than a few million years at a speed of a few tens of kilometers per second.


In time, a rather unusual-looking object results. The “star” now consists of two distinct parts, both of which constitute stage 12 of Table 20.1. At the center is a small, well-defined core of mostly carbon ash. Hot, dense, and still very luminous, only the outermost layers of this core still fuse helium into carbon and oxygen. Well beyond the core lies an expanding cloud of dust and cool gas—the ejected envelope of the giant—spread over a volume roughly the size of our solar system.

As the core exhausts its last remaining fuel, it contracts and heats up, moving to the left in the H–R diagram. Eventually, it becomes so hot that its ultraviolet radiation ionizes the inner parts of the surrounding cloud, producing a spectacular display called a **planetary nebula**. Some well-known examples are shown in Figures 20.9 and 20.10. In all, more than 1500 planetary nebulae are known in our Galaxy. The word *planetary* here is misleading, for these objects have no association with planets. The name originated in the 18th century, when, viewed at poor resolution through small telescopes, these shells of glowing gas looked to some astronomers like the circular disks of planets in our solar system.

Note that the mechanism by which planetary nebulae shine is basically the same as that powering the emission nebulae we studied earlier: ionizing radiation from a hot star embedded in a cool gas cloud. ∞ (Sec. 18.2) However, recognize that these two classes of object have very different origins and represent completely separate phases of stellar evolution. The emission nebulae discussed in Chapter 18 are signposts of recent stellar birth. Planetary nebulae, by contrast, indicate impending stellar death.

Astronomers once thought that the escaping giant envelope would be more or less spherical in shape, completely surrounding the core in three dimensions, just as it had while still part of the star. Figure 20.9(a) shows an example where this may well in fact be the case. The “ring” of this planetary nebula is in reality a three-dimensional shell of glowing gas—its halo-shaped appearance is only an illusion. As illustrated in Figure 20.9(b), the nebula looks brighter near the edges simply because there is more emitting gas along the line of sight there, creating the illusion of a bright ring.


However, such cases now seem to be in the minority. There is growing evidence that, for reasons not yet fully understood, the final stages of red-giant mass loss are often decidedly *nonspherical*. For example, the famous Ring Nebula shown in








(a)



(c)



(b)

**◀ FIGURE 20.9 Ejected Envelope** (a) Abell 39, some 6800 light-years away, is a classic planetary nebula shedding a spherical shell of gas. (b) The brightened appearance around the edge of Abell 39 is caused by the thinness of the shell of glowing gas around the central core. Very little gas exists along the line of sight between the observer and the central star (path A), so that part of the shell is invisible. Near the edge of the shell, however, more gas exists along the line of sight (paths B and C), so the observer sees a glowing ring. (c) The Ring Nebula, perhaps the most famous of all planetary nebulae at about 4900 light-years away, is too small and dim to be seen with the naked eye. Astronomers once thought its appearance could be explained in much the same way as that of Abell 39. However, it now seems that the Ring really is ring shaped, but researchers are unsure why. (AURA; NASA)


Figure 20.9(c) may well actually *be* a ring, and not just our view of a glowing spherical shell, and many planetary nebulae are much more complex than that. As illustrated in Figure 20.10, some planetary nebulae exhibit much more complex structures, suggesting that the star's environment—including the existence of a binary companion—can play an important role in determining the nebula's shape and appearance.

The central star fades and eventually cools, and the expanding gas cloud becomes more and more diffuse, eventually dispersing into interstellar space. After just a few tens of thousands of years, the glowing planetary nebula disappears from view. As the cloud rejoins the interstellar medium, it plays a vital role in the evolution of our Galaxy. During the final stages of the red giant's life, nuclear reactions between carbon and unburned helium in the core create oxygen and, in some cases, even heavier elements, such as neon and magnesium. Some of these reactions also release neutrons, which, carrying no electrical charge, have no electrostatic barrier to overcome and hence can interact with


existing nuclei to form still heavier elements (see Chapter 21). All of these elements—helium, carbon, oxygen, and heavier ones—are “dredged up” from the depths of the core into the envelope by convection during the star's final years to enrich the interstellar medium when the giant envelope escapes. The evolution of low-mass stars is the source of virtually all the carbon-rich dust observed throughout the plane of our own and other galaxies. ∞ (Sec. 18.1)

### Stage 13: A White Dwarf

The carbon core—the stellar remnant at the center of the planetary nebula—continues to evolve. Formerly concealed by the atmosphere of the red-giant star, the core becomes visible as the envelope recedes. Several tens of thousands of years are needed for the core to appear from behind the veil of expanding gas. The core is very small. By the time the envelope is ejected as a planetary nebula, the core has shrunk to about the size of Earth. (In some cases, it may be even smaller than our planet.) Its mass is about half the mass of the Sun. Shining only by stored heat, not by nuclear reactions, this small “star” has a white-hot surface when it first becomes visible, although it appears dim because of its small size. The core's temperature



(a) R I V U X G




(b) R I V U X G



(c) R I V U X G

**▲ FIGURE 20.10 Planetary Nebulae** (a) The Eskimo Nebula clearly shows several “bubbles” (or shells) of material being blown into space from this planetary nebula, which resides some 4800 light-years away in the constellation Gemini. (b) The Cat’s Eye Nebula, about 3200 light-years away, is an example of a much more complex planetary nebula, possibly produced by a pair of binary stars (unresolved at center) that have both shed envelopes (red seen in visible light, blue in X-rays). (c) M2-9, some 2000 light-years away, shows surprising twin lobes (or jets) of glowing gas emanating from a central, dying star and racing out at speeds of about 300 km/s. (AURA; NASA)




**▲ Interactive FIGURE 20.11 White Dwarf on the H–R Diagram** A star’s passage from the horizontal branch (stage 10) to the white-dwarf stage (stage 13) by way of the asymptotic-giant branch creates an evolutionary path that cuts across the entire H–R diagram.

and size give rise to its new name: **white dwarf**. This is stage 13 of Table 20.1. The approximate path followed by the star on the H–R diagram as it evolves from stage-11 red giant to stage-13 white dwarf is shown in Figure 20.11.

Not all white-dwarf stars are found as the cores of planetary nebulae: Several hundred have been discovered “naked” in our Galaxy, their envelopes expelled to invisibility (or perhaps stripped away by a binary companion—to be discussed shortly) long ago. Figure 20.12 shows an example of a white dwarf, Sirius B, that happens to lie particularly close to Earth; it is the faint binary companion of the much brighter and better known Sirius A. [∞ \(More Precisely 17-2\)](#) Some properties of Sirius B are listed in Table 20.2. With more than the mass of the Sun

Mass	1.1 solar masses
Radius	0.0073 solar radius (5100 km)
Luminosity (total)	0.0025 solar luminosity ( $10^{24}$ W)
Surface temperature	27,000 K
Average density	$3.9 \times 10^9$ kg/m <sup>3</sup>



▲ **FIGURE 20.12 Sirius Binary System** Sirius B (the speck of light to the right of the much larger and brighter star Sirius A) is a white dwarf star, a companion to Sirius A. The “spikes” on the image of Sirius A are not real; they are caused by the support struts of the telescope. (*Palomar Observatory*)


packed into a volume smaller than Earth, Sirius B has a density about a million times greater than anything familiar to us in the solar system. In fact, Sirius B has an unusually high mass for a white dwarf—it is thought to be the

evolutionary product of a star roughly four times the mass of the Sun. *Discovery 20-1* discusses another possible peculiarity of Sirius B’s evolution.

*Hubble Space Telescope (HST)* observations of nearby globular clusters have revealed the white-dwarf sequences long predicted by theory, but previously too faint to detect at such large distances. Figure 20.13(a) shows a ground-based view of the globular cluster M4, lying 1700 pc from Earth. Part (b) of the figure shows an *HST* closeup of a small portion of the cluster, revealing dozens of white dwarfs among the cluster’s much brighter main sequence, red-giant, and horizontal-branch stars. When plotted on an H–R diagram (see Figure 20.14), the white dwarfs fall nicely along the path indicated in Figure 20.11.

Not all white dwarfs are composed of carbon and oxygen. As mentioned earlier, theory predicts that very low-mass stars (less than about one-quarter the mass of the Sun) will never reach the point of helium fusion. Instead, the core of such a star will become supported by electron degeneracy pressure before its central temperature reaches the 100 million K needed to start the triple-alpha process. The interiors of such stars are completely convective, ensuring that fresh hydrogen continually mixes from the envelope into the core. As a result, unlike the case of the Sun illustrated in Figure 20.2, a nonburning helium inner core never appears, and eventually *all* of the star’s hydrogen is converted to helium, forming a *helium white dwarf*.

The time needed for this kind of transformation to occur is very long—hundreds of billions of years—so no




▲ **FIGURE 20.13 Distant White Dwarfs** (a) The globular cluster M4 (see also Figure 18.13), as seen through a large ground-based telescope at Kitt Peak National Observatory in Arizona, is the closest globular cluster to us, at 5500 light-years away. (b) A peek at M4’s “suburbs” by the *Hubble Space Telescope* shows nearly a hundred white dwarfs within a small 2-square-light-year region. (*AURA; NASA*)



helium white dwarfs have ever actually formed in this way.  $\infty$  (Sec. 17.8) However, if a solar-mass star is a member of a binary system, it is possible for its envelope to be stripped away during the red-giant stage by the gravitational pull of its companion (see Section 20.6), exposing the helium core and terminating the star's evolution before helium fusion can begin. Several such low-mass helium white dwarfs have in fact been detected in binary systems.

Finally, in stars somewhat more massive than the Sun (close to the 8-solar-mass limit on “low-mass” stars at the time the carbon core forms), temperatures in the core may become high enough that an additional reaction,



can occur, ultimately leading to the formation of a rare *neon–oxygen white dwarf*.

## Stage 14: A Black Dwarf

Once an isolated star becomes a white dwarf, its evolution is over. (As we will see in Chapter 21, white dwarfs in binary systems may undergo further activity.) The isolated white dwarf continues to cool and dim with time, following the white–yellow–red track near the bottom of the H–R diagram of Figure 20.11 and eventually becoming a **black dwarf**—a cold, dense, burned-out ember in space. This is stage 14 of Table 20.1, the graveyard of stars.



The cooling dwarf does not shrink much as it fades away. Even though its heat is leaking away into space, gravity does not compress it further. At the enormously high densities in the star (from the white-dwarf stage on), the resistance of electrons to being squeezed together—the same electron degeneracy that prevailed in the red-giant core around the time of the helium flash—supports the star, even as its temperature drops almost to absolute zero. As the dwarf cools, it remains about the size of Earth.

## Comparing Theory with Reality

All the H–R diagrams and evolutionary tracks presented so far are theoretical constructs based largely on computer models of the interior workings of stars. Before continuing our study of stellar evolution, let's take a moment to compare our models with actual observations. Figure 20.14(a) shows the beautiful globular cluster M80, which lies about 8000 pc from Earth. Figure 20.14(b) shows a composite H–R diagram recently constructed by using the stars of a number of other globular clusters of roughly the same age and composition as M80. The diagram spans the entire range of stellar luminosities, from bright red giants to faint red and white dwarfs. Fitting theoretical models of the main-sequence, giant, and horizontal branches (see Section 20.5) implies an age of about 12 billion years, making these clusters among the



(a)




(b)

**▲ FIGURE 20.14 Globular Cluster H–R Diagram** (a) The globular cluster M80, some 26,000 light-years from Earth. (b) Combined H–R diagram, based on ground- and space-based observations, for several globular clusters similar in overall composition to M80. The various evolutionary stages predicted by theory and depicted schematically in Figure 20.11 are clearly visible. Note also the blue stragglers—main-sequence stars that appear to have been “left behind” as other stars evolved into giants. They are probably the result of merging binary systems or actual collisions between stars of lower mass in this remarkably dense stellar system. (See also Figure 20.20.) (NASA; data courtesy W. E. Harris)

## DISCOVERY 20-1

## Learning Astronomy from History

Sirius A, the brighter of the two objects shown in Figure 20.12, appears twice as luminous as any other visible star, excluding the Sun. Its absolute brightness is not very great, but because it is not very far from us (less than 10 light-years away), its apparent brightness is very large.  (Sec. 17.2) Sirius has been prominent in the nighttime sky since the beginning of recorded history. Cuneiform texts of the ancient Babylonians refer to the star as far back as 1000 B.C., and historians know that the star strongly influenced the agriculture and religion of the Egyptians of 3000 B.C.

Even though a star's evolution takes such a long time, we might have a chance to detect a slight change in Sirius because the recorded observations of this star go back several thousand years. The chances for success are improved in this case because Sirius A is so bright that even the naked-eye observations of the ancients should be reasonably accurate. Interestingly, recorded history does suggest that Sirius A has changed in appearance, but the observations are confusing. Every piece of information about Sirius recorded between the years 100 B.C. and A.D. 200 claims that this star was *red*. (No earlier records of its color are known.) In contrast, modern observations now show it to be white or bluish white—definitely *not* red.

If these reports are accurate, then Sirius has apparently changed from red to blue white in the intervening years. According to the theory of stellar evolution, however, no star should be able to change its color in this way in that short a time. Such a color change should take at least several tens of thousands of years and perhaps a lot longer. It should also leave some evidence of its occurrence.

Astronomers have offered several explanations for the rather sudden change in Sirius A, including the suggestions that (1) some ancient observers were wrong and other scribes copied their mistaken writings; (2) a Galactic dust cloud passed between Sirius A and Earth some 2000 years ago, reddening the star much as Earth's dusty atmosphere often reddens our Sun at dusk; and (3) the companion to Sirius A, Sirius B, was a red giant and the dominant star of this double-star system 2000 years ago, but has since expelled its planetary nebular shell to reveal the white-dwarf star that we now observe.

Each of these explanations presents problems. How could the color of the sky's brightest star have been incorrectly recorded for hundreds of years? Where is the intervening galactic cloud now? Where is the shell of the former red giant? We are left with the uneasy feeling that the sky's brightest star doesn't fit particularly well into the currently accepted scenario of stellar evolution.


oldest-known objects in the Milky Way Galaxy and, as such, key indicators of conditions in the early universe.


The great age of this cluster means that stars more massive than about 0.8 solar mass have already evolved beyond the red-giant stage, becoming mainly white dwarfs. The H–R diagram for this cluster can therefore be compared directly with Figure 20.11, as the red-giant-branch, horizontal-branch, and asymptotic-giant-branch stars are all of roughly 1 solar mass. The similarity between theory and observation is striking: Stars in each of the evolutionary stages 7–13 can be seen in numbers consistent with the theoretical models. (See also the acetate inset in this chapter.) Astronomers place great confidence in the theory of stellar evolution precisely because its predictions are so often found to be in excellent agreement with plots of real stars.

Note that the points in Figure 20.14(b) are shifted a little to the left relative to Figure 20.11. This is because of differences in composition between stars like the Sun and stars in globular clusters. For reasons to be discussed more fully in Chapter 21, the old globular cluster stars contain much lower concentrations of “heavy” elements (astronomical jargon for anything more massive than helium). One result of this is that the interiors and atmospheres of those stars tend to be slightly more transparent to radiation

from within, allowing the energy to escape more easily and making the stars slightly smaller and hotter than solar-type stars of the same mass.

The objects labeled as **blue stragglers** in Figure 20.14(b) appear at first sight to contradict the theory just described. They are observed in many star clusters, lying on the main sequence, but in locations suggesting that they should have evolved into white dwarfs long ago, given the parent cluster's age. They are main-sequence stars, but they did not form when the cluster did. Instead, they formed much more recently, through *mergers* of lower mass stars—so recently, in fact, that they have not yet had time to evolve into giants.

In some cases, these mergers are the result of stellar evolution in binary systems, as the component stars evolved, grew, and came into contact (see Section 20.6). In others, the mergers are thought to be the result of actual *collisions* between stars. The core of M80 contains a huge number of stars packed into a relatively small volume. For example, a sphere of radius 2 pc centered on the Sun contains exactly four stars, including the Sun itself.  (Sec. 17.1) At the center of M80, the same 2-pc sphere would contain more than 10 *million* stars—our night sky would be ablaze with thousands of objects brighter than Venus! The dense central cores of globular clusters are among the few places in the entire universe where stellar collisions are likely to occur.



◀ **FIGURE 20.15 Multiple Stellar Generations** The ground-based H–R diagram (b) of the globular cluster NGC 2808 (a) shows an apparently normal main sequence. But more precise observations from *HST* (c) reveal that the main sequence is actually made up of three distinct sequences (as noted), increasing in helium content from right to left. These observations imply multiple generations of star formation shortly after the cluster formed, but no theory can yet explain just how this occurred. (NASA)

## 20.4 Evolution of Stars More Massive than the Sun

High-mass stars evolve much faster than their low-mass counterparts. The more massive a star, the more ravenous is its fuel consumption and the shorter is its main-sequence lifetime. The Sun will spend a total of some 10 billion years on the main sequence, but a 5-solar-mass B-type star will remain there for only a hundred million years. A 10-solar-mass O-type star will depart in just 20 million years or so. This trend toward much faster evolution for more massive stars continues even after the star leaves the main sequence.

All evolutionary changes happen much more rapidly for high-mass stars because their larger mass and stronger gravity generate more heat, speeding up *all* phases of stellar evolution. In fact, helium fusion proceeds so quickly that the high-mass star has a very different evolutionary track. Its envelope swells and cools as the star becomes a *supergiant*.

### Red Supergiants


Stars leave the main sequence for one basic reason: They run out of hydrogen in their cores. As a result, the early stages of stellar evolution beyond the main sequence are qualitatively the same in all cases: Main-sequence hydrogen burning in the core (stage 7) eventually gives way to the formation of a nonburning, collapsing helium core surrounded by a hydrogen-burning shell (stages 8 and 9). A high-mass star leaves the main sequence on its journey toward the red-giant region with an internal structure quite similar to that of its low-mass cousin. Thereafter, however, their evolutionary tracks diverge.

Figure 20.16 compares the post-main-sequence evolution of three stars, respectively, having masses 1, 4, and 10 times the mass of the Sun. Note that, whereas stars like the

High-precision observations from *HST* have revealed a new, and as yet unresolved, mystery about globular clusters that may force astronomers to change significantly their views on how massive star clusters form. Figure 20.15 presents the H–R diagram for the cluster NGC 2808, showing (in part c) what appear to be *three* distinct main sequences undetected in earlier ground-based observations. The stars in the three sequences contain different fractions of helium, carbon, and nitrogen and are thought to be the result of multiple episodes of star formation occurring over the course of about 100 million years. Models suggest that the two more-helium-rich generations formed from gas enriched by stellar evolution in the first-generation stars, but astronomers do not yet know how this could have occurred in the time available. Whatever happened, it seems to have been a common phenomenon, as high-resolution studies of many globular clusters now reveal similar multiple sequences and chemical variations. Indeed, some observers would go so far as to claim that multiple stellar populations like these are the norm in the Galactic globular cluster system.

### CONCEPT Check

✓ Why does fusion cease in the core of a low-mass star?



**▲ FIGURE 20.16 High-Mass Evolutionary Tracks** Evolutionary tracks for stars of 1, 4, and 10 solar masses (shown only up to helium ignition in the low-mass case). Stars with masses comparable to the Sun ascend the giant branch almost vertically, whereas higher-mass stars move roughly horizontally across the H–R diagram from the main sequence into the red-giant region. The most massive stars experience smooth transitions into each new burning stage. No helium flash occurs for stars more massive than about 2.5 solar masses. Some points are labeled with the element that has just started to fuse in the inner core.

Sun ascend the red-giant branch almost vertically, stars of higher mass move nearly horizontally across the H–R diagram after leaving the upper main sequence. Their luminosities stay roughly constant as their radii increase and their surface temperatures drop.

In stars having more than about 2.5 times the mass of the Sun, helium burning begins smoothly and stably, *not* explosively—there is no helium flash. Calculations indicate that the more massive a star, the lower is its core density when the temperature reaches the  $10^8$  K necessary for helium ignition, and the smaller is the contribution to the pressure from degenerate electrons. As a result, above 2.5 solar masses, the unstable core conditions described earlier do not occur. The 4-solar-mass red giant in Figure 20.16 remains a red giant as helium starts to fuse into carbon. There is no sudden jump to the horizontal branch and no subsequent reascend of the giant branch. Instead, the star loops smoothly back and forth near the top of the H–R diagram.

A much more important divergence occurs at approximately 8 solar masses—the dividing line between high

and low mass mentioned in Section 20.1. A low-mass star never achieves the 600 million K needed to fuse carbon nuclei, so it ends its life as a carbon–oxygen (or possibly neon–oxygen) white dwarf. A high-mass star, however, can fuse not only hydrogen and helium, but also carbon, oxygen, and even heavier elements as its inner core continues to contract and its central temperature continues to rise. The rate of burning accelerates as the core evolves.

Evolution proceeds so rapidly in the 10-solar-mass star of Figure 20.16 that the star doesn’t even reach the red-giant region before helium fusion begins. The star achieves a central temperature of  $10^8$  K while it is still quite close to the main sequence. As each element is burned to depletion at the center, the core contracts and heats up, and fusion starts again. A new inner core forms, contracts again, heats again, and so on. The star’s evolutionary track continues smoothly across the supergiant region of the H–R diagram, seemingly unaffected by each new phase of burning. The star’s radius increases as its surface temperature drops, so the star swells to become a red supergiant.  $\infty$  (Sec. 17.4)

With heavier and heavier elements forming at an ever-increasing rate, the high-mass star shown in Figure 20.16 is very close to the end of its life. We will discuss the evolution and ultimate fate of such a star in more detail in the next chapter, but suffice it to say here that it is destined to die in a violent supernova—a catastrophic explosion releasing energy that will most likely literally blow the star to pieces—soon after carbon and oxygen begin to fuse in its core. High-mass stars evolve so rapidly that, for most practical observational purposes, they explode and die shortly after leaving the main sequence.

A good example of a post-main-sequence blue supergiant is the bright star Rigel in the constellation Orion. With a radius some 70 times that of the Sun and a total luminosity of more than 60,000 solar luminosities, Rigel is thought to have had an original mass about 17 times that of the Sun, although a strong stellar wind has probably carried away a significant fraction of its mass since it formed. Although still near the main sequence, Rigel is probably already fusing helium into carbon in its core.


Perhaps the best-known red supergiant is Betelgeuse (shown in Figures 17.8 and 17.11), also in Orion, and Rigel’s rival for the title of brightest star in the constellation. Its luminosity is roughly  $10^4$  times that of the Sun in visible light and perhaps four times that in the infrared. Astronomers think that Betelgeuse is currently fusing helium into carbon and oxygen in its core, but its eventual fate is uncertain. As best we can tell, the star’s mass at formation was between 12 and 17 times the mass of the Sun. However, like Rigel and many other supergiants, Betelgeuse has a strong stellar wind and is known to be surrounded by a huge shell of dust of its own making (see *Discovery 20-2*). It also pulsates, varying in

## DISCOVERY 20-2


## Mass Loss from Giant Stars

Astronomers now know that stars of all spectral types are active and have stellar winds. Consider the highly luminous, hot, blue O- and B-type stars, which have by far the strongest winds. Satellite and rocket observations have shown that their wind speeds may reach 3000 km/s. The result is a yearly mass loss sometimes exceeding  $10^{-6}$  solar mass per year. Over the relatively short span of 1 million years, these stars blow a tenth of their total mass—more than an entire solar mass of material—into space. The powerful stellar winds, driven directly by the pressure of the intense ultraviolet radiation emitted by the stars themselves, hollow out vast cavities in the interstellar gas.

The black-and-white photograph here shows the supergiant star AG Carinae—50 times more massive than the Sun and a million times brighter—shedding its outer atmosphere.



(NASA)



The star is shown puffing out vast clouds of gas and dust. (The star, at the center, is intentionally obscured to show the surrounding faint nebula more clearly; the bright vertical line is also an artifact—an effect of the optical system used to hide the star.)


The four-part accompanying *Hubble* image captures another stellar outburst in the second half of the year 2002, during which a star brightened more than a half-million times our Sun's luminosity. This star, with the tongue-twisting name V838 Monocerotis, is a highly variable (and poorly understood) red supergiant about 20,000 light-years distant. Actually, what we are seeing here is not matter being expelled outward as fast as the images imply; rather, a burst of light—often called a “light echo”—is illuminating shells of gas and dust now surrounding the star, but that had been shed long ago. For scale, the rightmost image is about 7 light-years across.

Observations made with radio, infrared, and optical telescopes have shown that luminous cool stars (e.g., K- and M-type red giants) also lose mass at rates comparable to those at which luminous hot stars lose mass. Red-giant wind velocities, however, are much lower, averaging merely 30 km/s. They carry roughly as much mass into space as do O-type stellar winds, because their densities are generally much greater. Also, because luminous red stars are inherently cool objects (with surface temperatures of only about 3000 K), they emit virtually no ultraviolet radiation, so the mechanism driving the winds must differ from that driving the winds of luminous hot stars. We can only surmise that gas turbulence, magnetic fields, or both in the atmospheres of the red giants are somehow responsible. The surface conditions in red giants are in some ways similar to those in T Tauri protostars, which are also known to exhibit strong winds. Possibly the same basic mechanism—violent surface activity—is responsible for both kinds of winds.

Unlike winds from hot stars, winds from these cool stars are rich in dust particles and molecules. Nearly all stars eventually evolve into red giants, so such winds are a major source of new gas and dust to interstellar space and also provide a vital link between the cycle of star formation and the evolution of the interstellar medium.



(NASA)





**TABLE 20.3** End Points of Evolution for Stars of Different Masses

Initial Mass (Solar Masses)	Final State
Less than 0.08	(Hydrogen) brown dwarf
0.08–0.25	Helium white dwarf
0.25–8	Carbon–oxygen white dwarf
8–12 (approx.)*	Neon–oxygen white dwarf
Greater than 12*	Supernova (Chapter 21)

\* Precise numbers depend on the (poorly known) amount of mass lost while the star is on, and after it leaves, the main sequence.

radius by about 60 percent. The pulsations and strong wind may be related to the huge spots observed on the star’s surface (Figure 17.11). Together, they suggest that Betelgeuse has lost a lot of mass since it formed, but just how much remains uncertain.

## The End of the Road

Protostars and stars evolve because gravity always tends to cause a nonburning stellar core to contract and heat up. The contraction continues until it is halted either by electron degeneracy pressure or by the onset of a new round of nuclear fusion. In the latter case, a new nonburning core builds up, and the process repeats. The more massive the star, the more repetitions occur before the star finally dies. Table 20.3 lists some possible outcomes of stellar evolution for stars of different masses. For completeness, *brown dwarfs*—the end product of low-mass protostars unable even to fuse hydrogen in their cores—are included in the list [∞](#) (Sec. 19.3)

Note that our earlier dividing line of 8 solar masses between “low mass” and “high mass” really refers to the mass at the time the carbon core forms. Since very luminous stars often have strong stellar winds (*Discovery 20-2*), main-sequence stars as massive as 10 to 12 times the mass of the Sun may still manage to avoid going supernova. Unfortunately, we do not know exactly how much mass either Rigel or Betelgeuse has lost, so we cannot yet tell whether they are above or below the threshold for becoming a supernova. Either might explode or instead become a neon–oxygen white dwarf, but for now we can’t say which. We may just have to wait and see—in a million years or so we will know for sure!

### CONCEPT Check

- ✓ **What is the essential evolutionary difference between high-mass and low-mass stars?**

## 20.5 Observing Stellar Evolution in Star Clusters

Star clusters provide excellent test sites for the theory of stellar evolution. Every star in a given cluster formed at almost the same time, from the same interstellar cloud, and with virtually the same composition. Only the mass varies from one star to another, thus allowing us to check the accuracy of our theoretical models in a very straightforward way. Having studied the evolutionary tracks of individual stars in some detail, let’s now consider how their collective appearance changes in time.


In Chapter 19, we saw how astronomers estimate the ages of star clusters by determining which of their stars have already left the main sequence. [∞](#) (Sec. 19.6) In fact, the main-sequence lifetimes that go into those age measurements represent only a tiny fraction of the data obtained from theoretical models of stellar evolution. Starting from the zero-age main sequence, astronomers can predict exactly how a newborn cluster should look at any subsequent time—which stars are on the main sequence, which are becoming giants, and which have already burned themselves out. Although we cannot see into the interiors of stars to test our models, we can compare stars’ outward appearances with theoretical predictions. The agreement—in detail—between theory and observation is remarkably good.

### The Evolving Cluster H–R Diagram

We begin our study shortly after the cluster’s formation, with the upper main sequence already fully formed and burning steadily and with stars of lower mass just beginning to arrive on the main sequence, as shown in Figure 20.17(a). The appearance of the cluster at this early stage is dominated by its most massive stars: the bright blue supergiants. Now let’s follow the cluster forward in time and see how it evolves by using an H–R diagram.

Figure 20.17(b) shows the appearance of our cluster’s H–R diagram after 10 million years. The most massive O-type stars have left the main sequence. Most have already exploded and vanished, as just discussed, but one or two may still be visible as red supergiants. The remaining stars in the cluster are largely unchanged in appearance—their evolution is slow enough that little happens to them in such a relatively short period. The cluster’s H–R diagram shows the main sequence slightly cut off, along with a rather poorly defined red-giant region. Figure 20.18 shows the twin open clusters *h* and  $\chi$  (the Greek letter chi) Persei, along with their combined H–R diagram. Comparing Figure 20.18(b) with such diagrams as those in Figure 20.17, astronomers estimate the age of this pair of clusters to be about 10 million years.

After 100 million years (Figure 20.17c), stars brighter than type B5 or so (about 4–5 solar masses) have left the main sequence, and a few more red supergiants are visible. By this time, most of the cluster’s low-mass stars have finally arrived



◀ **FIGURE 20.17 Cluster Evolution on the H–R Diagram** These are H–R diagrams for a hypothetical, evolving star cluster. (a) Initially, stars on the upper main sequence are already burning steadily while the lower main sequence is still forming. (b) At  $10^7$  years, O-type stars have already left the main sequence (as indicated by the arrows), and a few red giants are visible. (c) By  $10^8$  years, more red giants are visible, and the lower main sequence is almost fully formed. (d) At  $10^9$  years, the subgiant and red-giant branches are just becoming evident, and the formation of the lower main sequence is complete. (e) At  $10^{10}$  years, the cluster’s subgiant, red-giant, horizontal, and asymptotic-giant branches are all discernible, and many white dwarfs have now formed.


on the main sequence, although the dimmest M-type stars may still be in their contraction phase. The appearance of the cluster is now dominated by bright B-type main-sequence stars and brighter red supergiants.

At any time during the cluster’s evolution, the original main sequence is intact up to some well-defined stellar mass, corresponding to the stars that are just leaving the main sequence at that instant. We can imagine the main sequence being “peeled away” from the top down, with fainter and fainter stars turning off and heading for the giant branch as time goes on. Astronomers refer to the high-luminosity end of the observed main sequence as the **main-sequence turnoff**. The mass of a star that is just evolving off the main sequence at any moment is known as the *turnoff mass*.


At 1 billion years, the main-sequence turnoff mass is around 2 solar masses, corresponding roughly to spectral type A2. The subgiant and giant branches associated with the evolution of low-mass stars are just becoming visible, as indicated in Figure 20.17(d). The formation of the lower main sequence is now complete. In addition, the first white dwarfs have just appeared, although they are often too faint to be observed at the distances of most clusters. Figure 20.19 shows the Hyades open cluster and its H–R diagram, which appears to lie between Figures 20.17(c) and 20.17(d), suggesting that the cluster’s age is about 600 million years.

At 10 billion years, the turnoff point has reached solar-mass stars of spectral type G2. The subgiant and giant branches are now clearly discernible (see Figure 20.17e), and the horizontal and asymptotic-giant branches appear as distinct regions in the H–R diagram. Many white dwarfs are also present in the cluster. Although stars in all these evolutionary stages are also present in the 1-billion-year-old cluster shown in Figure 20.17(d), they are few in number then—typically only a few percent of the total number of stars in the cluster. Also, because they evolve so rapidly, these high-mass stars spend very little time in the various regions. Low-mass stars are much more numerous and evolve more slowly, so more of them spend more time in any given region of the H–R diagram, allowing their evolutionary tracks to be more easily discerned.

Figure 20.20 shows the globular cluster 47 Tucanae. By carefully adjusting their theoretical models until the cluster’s main sequence, subgiant, red-giant, and horizontal branches are all well matched, astronomers have determined the age




(a) R I V U X G




(b)

**▲ FIGURE 20.18 Newborn Cluster H–R Diagram** (a) The “double cluster”  $\eta$  and  $\chi$  Persei, two open clusters that apparently formed at the same time, possibly even orbiting one another. (b) The H–R diagram of the pair indicates that the stars are very young—probably only 10–15 million years old. Even so, the most massive stars have already left the main sequence. (NOAO; data from T. Currie)



(a) R I V U X G



(b)


**▲ FIGURE 20.19 Young Cluster H–R Diagram** (a) The Hyades cluster, a relatively young group of stars visible to the naked eye, is found 150 light-years away in the constellation Taurus. (b) The H–R diagram for this cluster is cut off at about spectral type A, implying an age of about 600 million years. A few massive stars have already become white dwarfs. (NOAO; ESA)

of 47 Tucanae to be between 10 and 12 billion years, a little older than our hypothetical cluster in Figure 20.17(e). In fact, globular-cluster ages determined in this way show a remarkably small spread: All the globular clusters in our Galaxy appear to have formed between about 10 and 12 billion years ago.


## The Theory of Stellar Evolution

The modern theory of the lives and deaths of stars is one more excellent example of the scientific method in action.

(Sec. 1.2) Faced with a huge volume of observational data, with little or no theory to organize or explain it, astronomers in the late 19th and early 20th centuries painstakingly classified and categorized the properties of the



**▲ FIGURE 20.20 Old Cluster H–R Diagram** (a) The southern globular cluster 47 Tucanae. (b) Fitting its main-sequence turnoff and its giant and horizontal branches to theoretical models gives 47 Tucanae an age of between 12 and 14 billion years, making it one of the oldest-known objects in the Milky Way Galaxy. The inset is a high-resolution ultraviolet image of 47 Tucanae’s core region, taken with the *Hubble Space Telescope* and showing many blue stragglers—massive stars lying on the main sequence above the turnoff point, resulting perhaps from the merging of binary-star systems. (See also Figure 20.14.) The points representing white dwarfs, some red dwarfs, and blue stragglers have been added to the original dataset, based on *Hubble* observations of this and other clusters. The white-dwarf data are for the cluster M4 (Figure 20.13). Data on the faintest main-sequence stars shown were obtained from ground-based observations. The thickness of the lower main sequence is due almost entirely to observational limitations, which make it difficult to determine accurately the apparent brightnesses and colors of low-luminosity stars. (ESO; NASA)



### PROCESS OF SCIENCE Check

✓ Why are observations of star clusters so important to the theory of stellar evolution?

## 20.6 Stellar Evolution in Binary Systems


stars they observed. ∞ (Sec. 17.5) During the first half of the 20th century, as quantum mechanics began to yield detailed explanations of the behavior of light and matter on subatomic scales, theoretical explanations of many key stellar properties emerged. ∞ (Sec. 4.2) Since the 1950s, a truly comprehensive theory has emerged, tying together the basic disciplines of atomic and nuclear physics, electromagnetism, thermodynamics, and gravitation into a coherent whole. Theory and observation have proceeded hand in hand, each refining and validating the details of the other as astronomers continue to hone their understanding.

Stellar evolution is one of the great success stories of astrophysics. Like all good scientific theories, it makes definite testable predictions about the universe while remaining flexible enough to incorporate new discoveries as they occur. At the start of the 20th century, some scientists despaired of ever knowing even the compositions of the stars, let alone why they shine and how they change. Today, the theory of stellar evolution is a cornerstone of modern astronomy. Its predictions extend our understanding of the cosmos literally to the limits of the observable universe.

We have noted that most stars in our Galaxy are not isolated objects, but are actually members of binary-star systems. However, our discussion of stellar evolution has so far focused exclusively on isolated stars. This narrow focus prompts us to ask how membership in a binary-star system changes the evolutionary tracks we have just described. Indeed, because nuclear burning occurs deep in a star’s core, does the presence of a stellar companion have any significant effect at all? Perhaps not surprisingly, the answer depends on the distance between the two stars in question.

For a binary system whose component stars are very widely separated—that is, the distance between the stars is greater than perhaps a thousand stellar radii—the two stars evolve more or less independently of one another, each following the track appropriate to an isolated star of its particular mass. However, if the two stars are closer, then the gravitational pull of one may strongly influence the envelope of the other. In that case, the physical properties of both may deviate greatly from those calculated for isolated single stars.

As an example, consider the star Algol (Beta Persei, the second-brightest star in the constellation Perseus). By



◀ **FIGURE 20.21 Stellar Roche Lobes** Each star in a binary system can be pictured as being surrounded by a “zone of influence,” or Roche lobe, inside of which matter may be thought of as being “part” of that star. The two teardrop-shaped Roche lobes meet at the Lagrangian point between the two stars. Outside the Roche lobes, matter may flow onto either star with relative ease.


studying its spectrum and the variation in its light intensity, astronomers have determined that Algol is actually a binary (in fact, an eclipsing double-lined spectroscopic binary, as described in Chapter 17), and they have measured its properties very accurately. ∞ (Sec. 17.7) Algol consists of a 3.7-solar-mass main-sequence star of spectral type B8 (a blue giant) with a 0.8-solar-mass red-subgiant companion moving in a nearly circular orbit around it. The stars are 4 million km apart and have an orbital period of about 3 days.

A moment’s thought reveals that there is something odd about these findings. On the basis of our earlier discussion, the more massive main-sequence star should have evolved *faster* than the less massive component. If the two stars formed at the same time (as is assumed to be the case), there should be no way that the 0.8-solar-mass star could be approaching the giant stage first. Either our theory of stellar evolution is seriously in error, or something has modified the evolution of the Algol system. Fortunately for theorists, the latter is the case.

As sketched in Figure 20.21, each star in a binary system is surrounded by its own teardrop-shaped “zone of influence,” inside of which its gravitational pull dominates the effects of both the other star and the overall rotation of the binary.

▶ **FIGURE 20.22 Algol Evolution** (a) Initially, Algol was probably a detached binary made up of two main-sequence stars. (b) As the more massive component (star 1) left the main sequence, it expanded to fill, and eventually overflow, its Roche lobe, transferring large amounts of matter onto its smaller companion (star 2). (c) Today, star 2 is the more massive of the two, but it is on the main sequence. Star 1 is still in the subgiant phase and fills its Roche lobe, causing a steady stream of matter to pour onto its companion.

Any matter within that region “belongs” to the star and cannot easily flow onto the other component or out of the system. Outside the two regions, it is possible for gas to flow toward either star relatively easily. The two teardrop-shaped regions are called **Roche lobes**, after Edouard Roche, the French mathematician who first studied the binary-system problem in the 19th century and whose work we have already encountered in the context of planetary rings. ∞ (Sec. 12.4) The Roche lobes of the two stars meet at a point on the line joining them—the inner Lagrangian point ( $L_1$ ), which we saw in Chapter 14 in discussing asteroid motions in the solar system. ∞ (Sec. 14.1) This Lagrangian point is a place



where the gravitational pulls of the two stars exactly balance the rotation of the binary system. The greater the mass of one component, the larger is its Roche lobe and the farther from its center (and the closer to the other star) is the Lagrangian point.

Astronomers think that Algol started off as a *detached* binary, with both components lying well within their respective Roche lobes. For reference, let us label the component that is now the 0.8-solar-mass subgiant as star 1 and the 3.7-solar-mass main-sequence star as star 2. Initially, star 1 was the more massive of the two, having perhaps three times the mass of the Sun. It thus turned off the main sequence first. Star 2 was originally a less massive star, perhaps comparable in mass to the Sun. As star 1 ascended the giant branch, it overflowed its Roche lobe and gas began to flow onto star 2. This transfer of matter had the effect of reducing the mass of star 1 and increasing that of star 2, which in turn caused the Roche lobe of star 1 to shrink as its gravity decreased. As a result, the rate at which star 1 overflowed its Roche lobe increased, and a period of unstable *rapid mass transfer* ensued, transporting most of star 1's envelope onto star 2. Eventually, the mass of star 1 became less than that of star 2. Detailed calculations show that the rate of mass transfer dropped sharply at that point, and the stars entered the relatively stable state we see today. These changes in Algol's components are illustrated in Figure 20.22.

Being part of a binary system has radically altered the evolution of both stars in the Algol system. The original

high-mass star 1 is now a low-mass red subgiant, whereas the roughly solar mass star 2 is now a massive blue-giant main-sequence star. The removal of mass from the envelope of star 1 may prevent it from ever reaching the helium flash. Instead, its naked core may eventually be left behind as a *helium white dwarf*. In a few tens of millions of years, star 2 will itself begin to ascend the giant branch and fill its own Roche lobe. If star 1 is still a subgiant or a giant at that time, a contact binary system will result. If, instead, star 1 has by then become a white dwarf, a new mass-transfer period—with matter streaming from star 2 back onto star 1—will begin. In that case (as we will see in Chapter 21), Algol may have a very active and violent future in store for it.

Just as molecules exhibit few of the physical or chemical properties of their constituent atoms, binaries can display types of behavior that are quite different from the behavior of either of their component stars. The Algol system is a fairly simple example of binary evolution, yet it gives us an idea of the sorts of complications that can arise when two stars evolve interdependently. We will return to the subject in the next two chapters, when we continue our discussion of stellar evolution and the strange states of matter that may ensue.

### CONCEPT Check

- ✓ Why is it important to understand the evolution of binary stars?


**The Big Question** If stars like the Sun end their lives so quietly and so similarly, why do their scattered remnants look so different on the sky? Planetary nebulae display all sorts of weird shapes and sizes, some with rings and spheres, others with loops and jets. What causes these dissimilar structures? Are they somehow intrinsic to the stars themselves, or are they due to the complex environment through which dying stars expel their contents back into interstellar space?

## Chapter Review

### SUMMARY


**1** Stars spend most of their lives on the main sequence, in the **core-hydrogen-burning** (p. 496) phase of stellar evolution, stably fusing hydrogen into helium at their centers. Stars leave the main sequence when the hydrogen in their cores is exhausted. The Sun is about halfway through its main-sequence lifetime and will reach this stage in about 5 billion years. Low-mass stars evolve much more slowly than the Sun; high-mass stars much faster.

**2** When the central nuclear fires in the interior of a solar-mass star cease, the helium in the star's core is still too cool to fuse into anything heavier. With no internal energy source, the helium



core is unable to support itself against its own gravity and begins to shrink. At this stage, the star is in the **hydrogen-shell-burning** (p. 497) phase, with a nonburning helium core surrounded by a layer of burning hydrogen.



The energy released by the contracting helium core heats the hydrogen-burning shell, greatly increasing the nuclear reaction rates there. The star becomes much brighter, while the envelope expands and cools. A low-mass star like the Sun moves




off the main sequence on the H–R diagram first along the **subgiant branch** (p. 498) and then almost vertically up the **red-giant branch** (p. 499).

3 Eventually, the contracting core of a Sun-like star reaches the point at which helium begins to fuse into carbon, but conditions at the onset of helium burning are such that the electrons in the core can be thought of as tiny, hard spheres that, once brought into contact, present stiff resistance to being compressed any further. This **electron degeneracy pressure** (p. 500) makes the core unable to “react” to the new energy source, and helium burning begins violently in a **helium flash** (p. 500). The flash expands the core and reduces the star’s luminosity, sending the star onto the **horizontal branch** (p. 500) of the H–R diagram. The star now has a core of burning helium surrounded by a shell of burning hydrogen.


4 As helium burns in the core, it forms an inner core of nonburning carbon. The carbon core shrinks and heats the overlying burning layers, and the star once again becomes a red giant, even more luminous than before. It reenters the red-giant region of the H–R diagram along the **asymptotic-giant branch** (p. 502). The core of a low-mass star never becomes hot enough to fuse carbon. Such a star continues to ascend the asymptotic-giant branch until its envelope is ejected into space as a **planetary nebula** (p. 503). At that point, the core becomes visible as a hot, faint, and extremely dense white dwarf, whereas the planetary nebula diffuses into space, carrying helium and some carbon into the interstellar medium. The white dwarf cools and fades, eventually becoming a cold **black dwarf** (p. 507).




5 High-mass stars evolve more rapidly than low-mass stars because larger mass results in higher central temperature. High-mass stars never initiate a helium flash, and they attain central temperatures high enough to fuse carbon. These stars become red supergiants, forming heavier and heavier elements in their cores at an increasingly rapid pace, and eventually die explosively.



6 The theory of stellar evolution can be tested by observing star clusters, all of whose stars formed at the same time. As time goes by, the most massive stars leave the main sequence first, then the intermediate-mass stars, and so on. At any instant, no stars with masses above the cluster’s **main-sequence turnoff** (p. 513) mass remain on the main sequence. Stars below this mass have not yet evolved into giants and so still lie on the main sequence. By comparing the main-sequence turnoff mass with the theoretical predictions, astronomers can measure a cluster’s age.



7 Stars in binary systems can evolve quite differently from isolated stars because of interactions with their companions. Each star is surrounded by a teardrop-shaped **Roche lobe** (p. 516), which defines the region of space within which matter “belongs” to the star. As a binary star evolves into the giant phase, it may overflow its Roche lobe, and gas flows from the giant onto its companion. Stellar evolution in binaries can produce states that are not achievable in single stars. In a sufficiently wide binary, both stars evolve as though they were isolated.



**MasteringAstronomy**® For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information.

**LO** connects to the introduction’s numbered Learning Outcomes.

## Review and Discussion

1. Why don’t stars live forever? Which stars live the longest?
2. **LO1** How long can a star like the Sun keep burning hydrogen in its core? Why is the depletion of hydrogen in the core of a star such an important event?
3. **LO2** Roughly how big (in AU) will the Sun become when it enters the red-giant phase?
4. How long does it take for a star like the Sun to evolve from the main sequence to the top of the red-giant branch?
5. Do all stars eventually fuse helium in their cores?
6. **LO3** What is the helium flash?
7. Describe an important way in which winds from red-giant stars are linked to the interstellar medium.
8. What is the internal structure of a star on the asymptotic-giant branch?
9. **LO4** What is a planetary nebula? Why do many planetary nebulae appear as rings?
10. What are white dwarfs? Why are they hard to observe?
11. **LO5** How do the late evolutionary stages of high-mass stars differ from those of low-mass stars?
12. **LO6 POS** How do astronomers test the theory of stellar evolution?
13. **POS** How can astronomers measure the age of a star cluster?
14. **LO7** What are the Roche lobes of a binary system?
15. Why is it odd that the binary system Algol consists of a low-mass red giant orbiting a high-mass main-sequence star? How did Algol come to be in this configuration?

## Conceptual Self-Test: Multiple Choice

1. A star will evolve off the main sequence when it uses up (a) all of its hydrogen; (b) half of its hydrogen; (c) most of the hydrogen in the core; (d) all of its gas.
2. On the main sequence, massive stars (a) conserve their hydrogen fuel by burning helium; (b) burn their hydrogen fuel more rapidly than the Sun; (c) burn their fuel more slowly than the Sun; (d) evolve into stars like the Sun.
3. Compared to other stars on the H–R diagram, red-giant stars are so named because they are (a) cooler; (b) fainter; (c) denser; (d) younger.
4. When the Sun is on the red-giant branch, it will be found at the (a) upper left; (b) upper right; (c) lower right; (d) lower left of the H–R diagram.
5. After the core of a Sun-like star starts to fuse helium on the horizontal branch, the core becomes (a) hotter; (b) cooler; (c) larger; (d) dimmer with time.
6. **VIS** If the evolutionary track in Overlay 3, showing a Sun-like star, were instead illustrating a significantly more massive star, its starting point (stage 7) would be (a) up and to the right; (b) down and to the left; (c) up and to the left; (d) down and to the right.
7. A white dwarf is supported by the pressure of tightly packed (a) electrons; (b) protons; (c) neutrons; (d) photons.
8. **VIS** When the Sun leaves the main sequence, in Figure 20.3, “Red Giant on the H–R Diagram,” it will become (a) hotter; (b) brighter; (c) more massive; (d) younger.
9. A star like the Sun will end up as a (a) blue giant; (b) white dwarf; (c) binary star; (d) red dwarf.
10. Compared to the Sun, stars plotted near the bottom left of the H–R diagram are much (a) younger; (b) more massive; (c) brighter; (d) denser.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

1. ● The Sun will leave the main sequence when roughly 10 percent of its hydrogen has been fused into helium. Using the data given in Section 16.5 and Table 16.2, calculate the total amount of mass destroyed (i.e., converted into energy) and the total energy released in the process.
2. ● Use the radius–luminosity–temperature relation to calculate the radius of a red supergiant with temperature 3000 K (half the solar value) and total luminosity 10,000 times that of the Sun. ∞ (Sec. 17.4) How many planets of our solar system would this star engulf?
3. ● What would be the luminosity of the Sun if its surface temperature were 3000 K and its radius were (a) 1 AU, (b) 5 AU?
4. ● Use the radius–luminosity–temperature relation to calculate the radius of a 12,000-K (twice the temperature of the Sun) 0.0004-solar-luminosity white dwarf.
5. ●● The Sun will reside on the main sequence for  $10^{10}$  years. If the luminosity of a main-sequence star is proportional to the fourth power of the star’s mass, what is the mass of a star that is just now leaving the main sequence in a cluster that formed (a) 400 million years ago, (b) 2 billion years ago?
6. ●● A main-sequence star at a distance of 20 pc is barely visible through a certain telescope. The star subsequently ascends the giant branch, during which time its temperature drops by a factor of three and its radius increases a hundredfold. What is the new maximum distance at which the star would still be visible in the same telescope?
7. ● Calculate the average density of a red-giant core of 0.25 solar mass and radius 15,000 km. Compare your answer with the average density of the giant’s envelope, if it has a 0.5 solar mass and its radius is 0.5 AU, and with the central density of the Sun. ∞ (Sec. 16.2)
8. ●● The radius of Betelgeuse varies by about 60 percent within a period of 3 years. If the star’s surface temperature remains roughly constant, by how much does its absolute magnitude change during this time?

## Activities

### Collaborative

1. The Ring Nebula (M57) is perhaps the most famous planetary nebula. At magnitude 9, it is faint, but a 6-inch or larger telescope should show its structure. To locate it, find Beta and Gamma Lyrae, the second and third brightest stars in the constellation Lyra. The ring lies between them, about one-third the way from Beta to Gamma. Don’t expect the Ring to look as colorful as the *Hubble* images you may have seen! Can you see any color in it? The Messier catalog contains three other planetary nebulae—M27 (the Dumbbell), M76 (the Little Dumbbell), and M97 (the Owl). Consult an online catalog and see if you can find them. The last two will be the most challenging.

### Individual

1. Can you find the Hyades cluster? It lies about 46 pc away in the constellation Taurus, making up the “face” of the bull. It appears to surround the very bright star Aldebaran, the bull’s eye, which makes it easy to locate in the sky. Aldebaran is a low-mass red giant, about twice the mass of the Sun, probably on the asymptotic-giant branch of its evolution. Despite appearances, it is not part of the Hyades cluster. In fact, it lies only about half as far away—about 20 pc from Earth.





# Stellar Explosions

## NOVAE, SUPERNOVAE, AND THE FORMATION OF THE ELEMENTS

What fate awaits a star when it runs out of fuel? For a low-mass star, the white-dwarf stage is not necessarily the end of the road: The potential exists for further violent activity if a binary companion can provide additional fuel. High-mass stars—whether members of binary stars or not—are also destined to die explosively, releasing vast amounts of energy, creating many heavy elements, and scattering the debris throughout interstellar space.

These cataclysmic explosions may trigger the formation of new stars, continuing the cycle of stellar birth and death. In this chapter, we will study in more detail both the processes responsible for the explosions and the mechanisms that create the elements from which we ourselves are made.

**The Big Picture** There is something philosophically intriguing about the idea that the deaths of some stars cause the birth of others. Build up, break down, change . . . dust to dust is a scientific concept. Many of the elements composing our world and ourselves were created in the violent explosions of long-gone stars. It sounds rather poetic that we are made of mostly stardust, but it happens to be true.

**LEFT:** All elements heavier than iron were created in supernovae—violent stellar explosions that mark the deaths of massive stars. Supernovae have been observed in many locations across the sky, often in galaxies far from our own. This billion-bit mosaic of several images from the *Hubble Space Telescope* shows a much closer example—the Crab Nebula about 6500 light-years away. The debris field seen here is scattered across 5 light-years, the different colors indicating various heavy elements. The explosion was actually seen in the sky about a thousand years ago as a massive star blew itself to smithereens. (*STScI*)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Explain how white dwarfs in binary-star systems can become explosively active.
- 2 Summarize the sequence of events leading to the violent death of a massive star.
- 3 Describe the two types of supernovae, and explain how each is produced.
- 4 Present the observational evidence for the occurrence of supernovae in our Galaxy.
- 5 Explain the origin of elements heavier than helium, and discuss the significance of these elements for the study of stellar evolution.
- 6 Outline how the universe continually recycles matter through stars and the interstellar medium.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.

## 21.1 Life after Death for White Dwarfs

Although most stars shine steadily day after day and year after year, some change dramatically in brightness over very short periods of time. One type of star, called a **nova** (plural: *novae*), may increase enormously in brightness—by a factor of 10,000 or more—in a matter of days and then slowly return to its initial luminosity over a period of weeks or months. The word *nova* means “new” in Latin, and to early observers these stars did indeed seem new because they appeared suddenly in the night sky. Astronomers now recognize that a nova is not a new star at all. It is instead a white dwarf—a normally very faint star—undergoing an explosion on its surface that results in a rapid, temporary increase in the star’s luminosity.


Figures 21.1(a) and (b) illustrate the brightening of a typical nova. Figure 21.1(c) shows a nova light curve, demonstrating how the luminosity rises dramatically in a matter of days and then fades slowly back to normal over the course of several months. On average, two or three novae are observed each year. Astronomers also know of many *recurrent novae*—stars that have been observed to “go nova” several times over the course of a few decades.

What could cause such an explosion on a faint, dead star? The energy involved is far too great to be explained by flares or other surface activity, and as we saw in the previous chapter, there is no nuclear activity in the dwarf’s interior. [∞ \(Sec. 20.3\)](#) To understand what happens, we must consider again the fate of a low-mass star after it enters the white-dwarf phase.


We noted in Chapter 20 that the white-dwarf stage represents the end point of a star’s evolution. Subsequently, the star simply cools, eventually becoming a black dwarf—a burned-out ember in interstellar space. This scenario is quite correct for an *isolated* star, such as our Sun. However, should the star be part of a *binary* system, an important new possibility exists. If the distance between the two stars is small enough, then the dwarf’s tidal gravitational field can pull matter—primarily hydrogen and helium—away from the surface of its main-sequence or giant companion. [∞ \(Sec. 7.6\)](#) The system then becomes a mass-transferring binary, similar to those discussed in Chapter 20. A stream of gas leaves the companion through the inner ( $L_1$ ) Lagrangian point and flows onto the dwarf. [∞ \(Sec. 20.6\)](#)

Because of the binary’s rotation and the white dwarf’s small size, material leaving the companion does not fall


directly onto the dwarf, as indicated in Figure 20.21. Instead, such material “misses” the compact star, loops around behind it, and goes into orbit around it, forming a swirling, flattened disk of matter called an **accretion disk** (shown in Figure 21.2). Due to the effects of viscosity (i.e., friction) within the gas, the orbiting matter in the disk drifts gradually inward, its temperature increasing steadily as it spirals down onto the dwarf’s surface. The inner part of the accretion disk becomes so hot that it radiates strongly in the visible, the ultraviolet, and even the X-ray portions of the electromagnetic spectrum. In many systems, the disk outshines the white dwarf itself and is the main source of the light emitted between nova outbursts. X-rays from the



(a)




(b)



(c)

► **FIGURE 21.1 Nova** Nova Herculis 1934 in (a) March 1935 and (b) May 1935, after brightening by a factor of 60,000. (c) The light curve of a typical nova displays a rapid rise followed by a slow decline in the light received from the star, in good agreement with the explanation of the nova as a nuclear flash on a white-dwarf’s surface. (UC/Lick Observatory)



◀ **FIGURE 21.2 Close Binary System**

If a white dwarf in a semidetached binary system is close enough to its companion (in this case, a main-sequence star), its gravitational field can tear matter from the companion's surface. Compare Figure 20.22, but note that, unlike the scenario shown in that earlier figure, the matter does not fall directly onto the white dwarf's surface. Instead, it forms an accretion disk of gas spiraling down onto the dwarf.

hot disk are routinely observed in many galactic novae. The point at which the infalling stream of matter strikes the accretion disk often forms a turbulent “hot spot,” causing detectable fluctuations in the light emitted by the binary system.

The “stolen” gas becomes hotter and denser as it builds up on the white dwarf's surface. Eventually, its temperature exceeds  $10^7$  K, and the hydrogen ignites, fusing into helium at a furious rate. (Figure 21.3a–d illustrates the sequence of events.) This surface-burning stage is as brief as it is violent: The star suddenly flares up in luminosity and then fades away as some of the fuel is exhausted and the remainder is blown off into space. If the event happens to be visible from Earth, we see a nova. Figure 21.4 shows two novae apparently caught in the act of expelling mass from their surfaces. A nova's decline in brightness results from the expansion


and cooling of the white dwarf's surface layers as they are blown into space. Studies of the details of the brightness curve associated with a nova provide astronomers with a wealth of information about both the dwarf and its binary companion.

A nova represents one way in which a star in a binary system can extend its “active lifetime” well into the white-dwarf stage. Recurrent novae can, in principle, repeat their violent outbursts many dozens, if not hundreds, of times. But even more extreme possibilities exist at the end of stellar evolution. Vastly more energetic events may be in store, given the right circumstances.


### CONCEPT Check

✓ Will the Sun ever become a nova?


*The sequence starts with a small white dwarf at upper left and proceeds to the right while orbiting the big red star, eventually igniting an explosion.*



▲ **FIGURE 21.3 Nova Explosion** In this artist's conception, a white-dwarf star (actually faraway at upper left) orbits a cool red giant (a). As the dwarf swings around in an elliptical orbit, coming closer to the giant, material accretes from the giant to the dwarf and accumulates on the white dwarf's surface (b and c). The dwarf star then ignites in hydrogen fusion as a nova outburst (d). (D. Berry)




(a)



(b)

**▲ FIGURE 21.4 Nova Matter Ejection** (a) The ejection of material from a star's surface can clearly be seen in this image of Nova Persei, taken some 50 years after it suddenly brightened by a factor of 40,000 in 1901. This corresponds approximately to Figure 21.3(d). (b) Nova Cygni, imaged here with a European camera on the *Hubble Space Telescope*, erupted in 1992. At left, more than a year after the blast, a rapidly billowing bubble is seen; at right, 7 months after that, the shell continued to expand and distort. The images are fuzzy because the object is more than 10,000 light-years away. (Palomar Observatory; ESA)



## 21.2 The End of a High-Mass Star


A low-mass star—a star with a mass of less than about 8 solar masses—never becomes hot enough to burn carbon in its core. It ends its life as a carbon–oxygen (or possibly neon–oxygen) white dwarf. [☞ \(Sec. 20.3\)](#) A high-mass star, however, can fuse not just hydrogen and helium, but also carbon, oxygen, and even heavier elements as its inner core continues to contract and its central temperature continues to rise. [☞ \(Sec. 20.4\)](#) The burning rate accelerates as the core evolves. Can anything stop this runaway process? Is there a stable “white-dwarf-like” state at the end of the evolution of a high-mass star? What is the ultimate fate of such a star? To answer these questions, we must look more carefully at fusion in massive stars.

### Fusion of Heavy Elements

Figure 21.5 is a cutaway diagram of the interior of a highly evolved star of large mass. Note the numerous layers in which various nuclei burn. As the temperature increases with depth, the ash of each burning stage becomes the fuel for the next stage. At the relatively cool periphery of the core, hydrogen fuses into helium. In the intermediate layers, shells of helium, carbon, and oxygen burn to form heavier nuclei. Deeper down reside neon, magnesium, silicon, and other heavy nuclei, all produced by nuclear fusion in the layers overlying the core. (Recall that, to astronomers, a “heavy” element is anything more massive than helium.) The core itself is composed of iron. We will study the key reactions in this burning chain in more detail later in the chapter.

As each element is burned to depletion at the center, the core contracts, heats up, and starts to fuse the ash of the previous burning stage. A new inner core forms, contracts again, heats again, and so on. Through each period of stability and instability, the star's central temperature increases, the nuclear reactions speed up,

#### **Interactive FIGURE 21.5 Heavy-Element Fusion**


 Cutaway diagram of the interior of a highly evolved star of mass greater than 8 solar masses. The interior resembles the layers of an onion, with shells of progressively heavier elements burning at smaller and smaller radii and at higher and higher temperatures. The core is actually only a few times larger than Earth, whereas the star is hundreds of times larger than the Sun.

and the newly released energy supports the star for ever-shorter periods of time. For example, in round numbers, a star 20 times more massive than the Sun burns hydrogen for 10 million years, helium for 1 million years, carbon for a thousand years, oxygen for a year, and silicon for a week. Its iron core grows for less than a day.

## Collapse of the Iron Core

Once the inner core begins to change into iron, our high-mass star is in trouble. As illustrated in Figure 21.6, iron is the *most stable* element there is. To understand the figure, imagine fusing four protons to form helium-4. According to the figure, the mass per particle of a helium-4 nucleus is less than the mass of a proton, so mass is lost and (in accordance with the law of conservation of mass and energy) energy is released.  $\infty$  (Sec. 16.6) Similarly, combining three helium-4 nuclei to form carbon results in a net loss of mass, again releasing energy. In other words, the left side of the figure shows how light elements can fuse to release energy. The right side of the figure shows the opposite process, known as **fission**. Here, combining nuclei will increase the total mass per particle and hence absorb energy, so fusion can't occur. However, splitting a heavy nucleus (such as uranium, or plutonium, which lies just off the right edge of the figure) into lighter nuclei does release energy—this is how nuclear reactors and atomic bombs work.

Iron lies at the dividing line between these two types of behavior—at the lowest point of the curve in the figure. Iron nuclei are so compact that energy cannot be extracted either by combining them into heavier elements or by splitting them into lighter ones. In effect, iron plays the role of a fire extinguisher, damping the inferno in the stellar core.




**▲ FIGURE 21.6 Nuclear Masses** This graph shows how the masses (per nuclear particle—proton or neutron) of most known nuclei vary with nuclear mass. When light nuclei fuse (left side of the figure), the mass per particle decreases and energy is released.  $\infty$  (Sec. 16.6) Similarly, when heavy nuclei split apart (right side), the total mass again decreases and energy is again released.

With the appearance of substantial quantities of iron, the central fires cease for the last time, and the star's internal support begins to dwindle. The star's foundation is destroyed, and its equilibrium is gone forever. Even though the temperature in the iron core has reached several billion kelvins by this stage, the enormous inward gravitational pull of matter ensures catastrophe in the very near future. Gravity overwhelms the pressure of the hot gas, and the star implodes, falling in on itself.

The core temperature rises to nearly 10 billion K. According to Wien's law, at that temperature individual photons have tremendously high energies—enough to split iron into lighter nuclei and then to break those lighter nuclei apart until only protons and neutrons remain.  $\infty$  (Sec. 4.2) This process is known as *photo disintegration* of the heavy elements in the core. In less than a second, the collapsing core undoes all the effects of nuclear fusion that occurred during the previous 10 million years! But to split iron and lighter nuclei into smaller pieces requires a lot of energy (Figure 21.6, moving from iron to the left). After all, this splitting is just the opposite of the fusion reactions that generated the star's energy during earlier times. Photodisintegration *absorbs* thermal energy—in other words, it cools the core and thus reduces the pressure there. As nuclei are destroyed, the core of the star becomes even less able to support itself against its own gravity. The collapse accelerates.


Now the core consists entirely of simple elementary particles—electrons, protons, neutrons, and photons—at enormously high densities, and it is still shrinking. As the density of the core continues to rise, the protons and electrons are crushed together, forming neutrons and neutrinos:



This process is sometimes called the *neutronization* of the core. Recall from our discussion in Chapter 16 that the neutrino is an extremely elusive particle that hardly interacts at all with matter.  $\infty$  (Sec. 16.6) Even though the central density by this time may exceed  $10^{12} \text{ kg/m}^3$ , most of the neutrinos produced by neutronization pass through the core as if it weren't there. They escape into space, carrying away energy as they go, further reducing the core's pressure support.


## Supernova Explosion

The disappearance of the electrons and the escape of the neutrinos make matters even worse for the core's stability. There is now nothing to prevent it from collapsing all the way to the point at which the neutrons come into contact with one another, at the incredible density of about  $10^{15} \text{ kg/m}^3$ . At this point, the neutrons in the shrinking core offer rapidly increasing resistance to further compression, producing enormous pressures that finally slow the core's gravitational



◀ **FIGURE 21.7 Supernova 1987A**

A supernova called SN 1987A (arrow) was exploding near this nebula (called 30 Doradus) at the moment the photograph on the right was taken. The photograph on the left is the normal appearance of the star field. (See *Discovery 21-1.*) (AURA)



collapse. By the time the collapse is actually halted, however, the core has overshot its point of equilibrium and may reach densities as high as  $10^{17}$  or  $10^{18}$   $\text{kg}/\text{m}^3$  before turning around and beginning to reexpand. Like a fast-moving ball hitting a brick wall and bouncing back, the core becomes compressed, stops, and then rebounds—with a vengeance!

The events just described do not take long. Only about a second elapses from the start of the collapse to the “bounce” at nuclear densities. At that point, the core rebounds. An enormously energetic shock wave sweeps through the star at high speed, blasting all the overlying layers—including all the heavy elements just formed outside the iron inner core—into space. Although computer models are still somewhat inconclusive, and the details of how the shock reaches the surface and destroys the star remain uncertain, the end result is not: The star explodes, in one of the most energetic events known in the universe (see Figure 21.7). For a period of a few days, the exploding star may rival in brightness the entire galaxy in which it resides. This spectacular death rattle of a high-mass star is known as a **core-collapse supernova**.

### CONCEPT Check

✓ Why does the iron core of a high-mass star collapse?

## 21.3 Supernovae

Let’s compare a supernova with a nova. Like a nova, a **supernova** is a star that suddenly increases dramatically in brightness and then slowly dims again, eventually fading from view. In its unexploded state, a star that will become a supernova is known as the supernova’s *progenitor*. In some cases, supernovae light curves can appear quite similar to those of novae, and a distant supernova can look a lot like a nearby


nova—so much so, in fact, that the difference between the two was not fully appreciated until the 1920s. But novae and supernovae are now known to be quite different phenomena. Supernovae are much more energetic events, driven by very different underlying physical processes.\*

### Novae and Supernovae

Well before they understood the causes of either novae or supernovae, astronomers knew of clear observational differences between them. The most important of these differences is that a supernova is more than a million times brighter than a nova. A supernova produces a burst of light billions of times brighter than the Sun, reaching that level of brightness within just a few hours after the start of the outburst. The total amount of electromagnetic energy radiated by a supernova during the few months it takes to brighten and fade away is roughly  $10^{43}$  J—nearly as much energy as the Sun will radiate during its *entire*  $10^{10}$ -year lifetime! (Enormous as this energy is, however, it pales in comparison with the energy emitted in the form of neutrinos, which may be 100 times greater.)

A second important difference is that the same star may become a nova many times, but a star can become a supernova only once. This fact was unexplained before astronomers knew the precise nature of novae and supernovae, but it is easily understood now that we understand how and why these explosions occur. The nova accretion–explosion cycle described earlier can take place over and over again, but a

\*When discussing novae and supernovae, astronomers tend to blur the distinction between the observed event (the sudden appearance and brightening of an object in the sky), the process responsible for the event (a violent explosion in or on a star), and the object itself (the star itself is called a nova or a supernova, as the case may be). The terms can have any of the three meanings, depending on the context.



**▲ FIGURE 21.8 Supernova Light Curves** The light curves of typical Type I and Type II supernovae both show that their maximum luminosities can sometimes reach that of a billion suns, but there are characteristic differences in the decline of the luminosity after the initial peak. Type I light curves resemble those of novae (see Figure 21.1), but the total energy released is much larger. Type II curves have a characteristic plateau during the declining phase.

supernova destroys the star involved, with no possibility of a repeat performance.

In addition to the distinction between novae and supernovae, there are also important observational differences *among* supernovae. Some supernovae contain very little hydrogen, according to their spectra, whereas others contain a lot. Also, the light curves of the hydrogen-poor supernovae are qualitatively different from those of the hydrogen-rich ones. On the basis of these observations, astronomers divide supernovae into two classes, known simply as Type I and Type II. **Type I supernovae**, the hydrogen-poor kind, have a light curve somewhat similar in shape to that of typical novae; **Type II supernovae**, whose spectra show lots of hydrogen, usually have a characteristic “plateau” in the light curve a few months after the maximum (see Figure 21.8). Observed supernovae are divided roughly equally between these two categories.

## Carbon-Detonation Supernovae

What is responsible for these differences among supernovae? Is there more than one way in which a supernova explosion can occur? The answer is yes. To understand the alternative supernova mechanism, we must return to the processes that cause novae and consider the long-term consequences of their accretion–explosion cycle.

Novae eject matter from a white dwarf’s surface, but they do not necessarily expel or burn all the material that has accumulated since the last outburst. In other words, there is a tendency for the dwarf’s mass to increase slowly with each new nova cycle. As its mass grows and the internal pressure required to support its weight rises, the white

dwarf can enter into a new period of instability—with disastrous consequences.

Recall that a white dwarf is held up not by thermal pressure (heat), but by the degeneracy pressure of electrons that have been squeezed so close together that they have effectively come into contact with one another. [☞ \(Sec. 20.3\)](#) However, there is a limit to the pressure that these electrons can exert. Consequently, there is a limit to the mass of a white dwarf, above which electrons cannot provide the pressure needed to support the star. Detailed calculations show that the maximum mass of a white dwarf is about 1.4 solar masses, a mass often called the *Chandrasekhar mass*, after the Indian-American astronomer Subramanyan Chandrasekhar, whose work in theoretical astrophysics earned him a Nobel Prize in physics in 1983.


If an accreting white dwarf exceeds the Chandrasekhar mass, the pressure of the degenerate electrons in its interior becomes unable to withstand the pull of gravity, and the star immediately starts to collapse. Its internal temperature rapidly rises to the point at which carbon can fuse into heavier elements. Carbon fusion begins everywhere throughout the white dwarf almost simultaneously, and the entire star explodes in another type of supernova—a so-called **carbon-detonation supernova**—comparable in violence to the “implosion” supernova associated with the death of a high-mass star, but born of a very different cause. In an alternative and (many astronomers think) possibly more common scenario, two white dwarfs in a binary system may collide and merge to form a massive, unstable star. The end result is the same: a carbon-detonation supernova.

We can now understand the differences between Type I and Type II supernovae. The explosion resulting from the detonation of a carbon white dwarf, the descendant of a low-mass star, is a supernova of Type I. Because this conflagration stems from a system containing virtually no hydrogen, we can readily see why the spectrum of a Type I supernova shows little evidence of that element. The appearance of the light curve (as we will soon see) results almost entirely from the radioactive decay of unstable heavy elements produced in the explosion itself.


The implosion–explosion of the core of a massive star, described earlier, produces a Type II supernova. Detailed computer models indicate that the characteristic shape of the Type II light curve is just what would be expected from the expansion and cooling of the star’s outer envelope as it is blown into space by the shock wave sweeping up from below. The expanding material consists mainly of unburned gas—hydrogen and helium—so it is not surprising that those elements dominate the supernova’s observed spectrum. (See *Discovery 21-1* for an account of a well-studied Type II supernova that confirmed many basic theoretical predictions, while also forcing astronomers to revise the details of their models.)



(a) Type I Supernova



(b) Type II Supernova



**▲ FIGURE 21.9 Two Types of Supernova** Type I and Type II supernovae have different causes. These sequences depict the evolutionary history of each type. (a) A Type I supernova usually results when a carbon-rich white dwarf pulls matter onto itself from a nearby red-giant or main-sequence companion. (b) A Type II supernova occurs when the core of a high-mass star collapses and then rebounds in a catastrophic explosion.

Figure 21.9 summarizes the processes responsible for the two different types of supernovae. We emphasize that, despite the similarity in the total amounts of energy involved, Type I and Type II supernovae are unrelated to one another. They occur in stars of very different types, under very different circumstances. All high-mass stars become Type II (core-collapse) supernovae, but only a tiny fraction of low-mass stars evolve into white dwarfs that ultimately explode as Type I (carbon-detonation) supernovae. However, there are far more low-mass stars than high-mass stars, so, by a remarkable coincidence, the two types of supernova occur at roughly the same rate.

## Supernova Remnants

We have plenty of evidence that supernovae have occurred in our Galaxy. Occasionally, the explosions themselves are visible from Earth. In many other cases, we can detect their glowing remains, or **supernova remnants**. One of the best-studied supernova remnants is known as the Crab Nebula, shown in Figure 21.10. The Crab has greatly dimmed now, but the original explosion in the year A.D. 1054 was so brilliant that manuscripts of ancient Chinese and Middle Eastern astronomers claim that its brightness greatly exceeded that of Venus and—according to some (possibly exaggerated) accounts—even rivaled

that of the Moon. For nearly a month, this exploded star reportedly could be seen in broad daylight. Native Americans also left engravings of the event in the rocks of what is now the southwestern United States.

The Crab Nebula certainly has the appearance of exploded debris. Even today, the knots and filaments give a strong indication of past violence. In fact, astronomers have proof that this matter was ejected from some central explosion. Doppler-shifted spectral lines indicate that the nebula—the envelope of the high-mass star that exploded to create this Type II supernova—is expanding into space at several thousand kilometers per second. A vivid illustration of the phenomenon is provided by Figure 21.11, which was made by superimposing a positive image of the Crab Nebula taken in 1960 and a negative image taken in 1974. If the gas were not in motion, the positive and negative images would overlap perfectly, but they do not. The gas moved outward in the intervening 14 years. Tracing the motion backward in time, astronomers have found that the explosion must have occurred about nine centuries ago, consistent with the Chinese observations.

The nighttime sky harbors many relics of stars that blew up long ago. Figure 21.12 is another example. It shows the Vela supernova remnant, whose expansion velocities imply that its central star exploded around 9000 B.C. The remnant lies only 1600 light-years from Earth. Given its




**Interactive FIGURE 21.10**  
**Crab Supernova Remnant**

**MA** This remnant of an ancient Type II supernova is called the Crab Nebula (or M1 in the Messier catalog). It resides about 6500 light-years from Earth and has an angular diameter about one-fifth that of the full Moon. The main image was taken with the Very Large Telescope in Chile, the inset by the *Hubble* telescope in orbit. (ESO; NASA)


proximity, the Vela supernova may have been as bright as the Moon for several months. We can only speculate what impact such a bright supernova might have had on the myths, religions, and cultures of Stone Age humans when it first appeared in the sky.

Although hundreds of supernovae have been observed in other galaxies during the 20th century, no astronomer using modern equipment has ever observed a supernova in our own Galaxy. A viewable Milky Way star has not exploded since Galileo first turned his telescope to the



**◀ FIGURE 21.11 The Crab in Motion**

Positive and negative photographs of the Crab Nebula taken 14 years apart do not superimpose exactly, indicating that the gaseous filaments are still moving away from the site of the explosion. The positive image in glowing white was taken first, and then the black (negative) filaments were overlaid later—hence the black (but still glowing) outlying debris is farther from the center of the blast. The scale is roughly the same as in Figure 21.10. (Harvard College Observatory)



**Interactive** **FIGURE 21.12 Vela Supernova Remnant** The glowing gases of the Vela supernova remnant are spread across  $6^\circ$  of the sky. The inset shows more clearly some of the details of the nebula's extensive filamentary structure. (The long diagonal streak was caused by the passage of an Earth-orbiting satellite while the photo was being exposed.) (D. Malin/AAT)

heavens almost four centuries ago. The last supernovae observed in our Galaxy, by Tycho in 1572 and Kepler (and others) in 1604, caused a worldwide sensation in Renaissance times. The sudden appearance and subsequent fading of these very bright objects helped shatter the Aristotelian idea of an unchanging universe.

On the basis of stellar evolutionary theory, astronomers calculate that an observable supernova ought to occur in our Galaxy every 100 years or so. Even at a distance of several kiloparsecs, a supernova would (temporarily) outshine Venus, the brightest planet in our sky, so it seems unlikely that astronomers could have missed any since the last one nearly four centuries ago. Our part of the Milky Way seems long overdue for a supernova. However, a truly nearby supernova—within a few hundred parsecs, say—would be a very rare event, occurring only every 100,000 years or so. Humanity may be destined to see all supernovae from a distance.

### PROCESS OF SCIENCE Check

- ✓ How did astronomers know, even before the mechanisms were understood, that there were at least two distinct physical processes at work in creating supernovae?

## 21.4 Formation of the Elements

Up to now, we have studied nuclear reactions mainly for their role in stellar energy generation. Now let's consider them again, but this time as the processes responsible for creating much of the world in which we live. The evolution of the elements, combining nuclear physics with astronomy, is a complex subject and a very important problem in modern astronomy.

### Types of Matter


We currently know of 115 different elements, ranging from the simplest—hydrogen, containing one proton—to the most complex, first reported in 2004 and known for now as ununpentium, which has 115 protons and 184 neutrons in its nucleus (see Appendix 3, Table 2). In 1999, researchers claimed the discovery of elements 116 and 118, but the experimental findings have not been replicated, and these elements are not “officially” recognized.) All elements exist in different *isotopes*, each having the same number of protons, but a different number of neutrons. We often think of the most common or stable isotope as being the “normal” form of an element. Some elements, and many isotopes, are radioactively unstable, meaning that they eventually decay into other, more stable, nuclei.

The 81 stable elements found on Earth make up the overwhelming bulk of matter in the universe. In addition, 10 radioactive elements—including radon and uranium—also occur naturally on our planet. Even though the half-lives (the time required for half the nuclei to decay into something else) of these elements are very long (typically, millions or even billions of years), their slow, but steady, decay over the 4.5 billion years since the solar system formed means that they are scarce on Earth, in meteorites, and in lunar samples. [\(More Precisely 7-2, Sec. 14.4\)](#) They are not observed in stars—there is just too little of them to produce detectable spectral lines.

Besides these 10 naturally occurring radioactive elements, 19 more radioactive elements have been artificially produced under special conditions in nuclear laboratories on Earth. The debris collected after nuclear weapons tests also contains traces of some of these elements. Unlike the naturally occurring radioactive elements, the artificial ones decay into other elements quite quickly (in much less than a million years). Consequently, they, too, are extremely rare in nature. Two other elements round out our list: Promethium is a stable element that is found on our planet only as a by-product of nuclear laboratory experiments; technetium is an unstable element that is found in stars, but does not exist on Earth—any technetium that existed in our planet at its formation decayed long ago.

## Abundance of Matter

How and where did all these elements form? Were they always present in the universe, or were they created after the universe formed? Since the 1950s, astronomers have come to realize that the hydrogen and most of the helium in the universe are *primordial*—that is, these elements date from the very earliest times (see Chapter 27). All other elements in our universe result from **stellar nucleosynthesis**—that is, they were formed by nuclear fusion in the hearts of stars.



**▲ FIGURE 21.13 Elemental Abundance** This graph summarizes the cosmic abundances of the elements and their isotopes, expressed relative to the abundance of hydrogen. The horizontal axis shows each of the listed elements' atomic number—the number of protons in the nucleus. Notice how many common terrestrial elements are found on “peaks” of the graph, surrounded by elements that are tens or hundreds of times less abundant. Notice also the large peak around the element iron. Reasons for the peaks are discussed in the text.

To test this idea, we must consider not just the different kinds of elements and isotopes, but also their observed abundances, graphed in Figure 21.13. The curve shown is derived largely from spectroscopic studies of stars, including the Sun. The essence of the figure is summarized in Table 21.1, which combines all the known elements into

**TABLE 21.1 Cosmic Abundances of the Elements**

Elemental Group of Particles	Percent Abundance by Number*
Hydrogen (1 nuclear particle)	90
Helium (4 nuclear particles)	9
Lithium group (7–11 nuclear particles)	0.000001
Carbon group (12–20 nuclear particles)	0.2
Silicon group (23–48 nuclear particles)	0.01
Iron group (50–62 nuclear particles)	0.01
Middle-weight group (63–100 nuclear particles)	0.00000001
Heaviest-weight group (over 100 nuclear particles)	0.000000001

\* The total does not equal 100 percent because of uncertainties in the abundance of helium. All isotopes of all elements are included.

## DISCOVERY 21-1

## Supernova 1987A


In 1987, astronomers were treated to a spectacular supernova in the Large Magellanic Cloud (LMC), a small satellite galaxy orbiting our own (see Section 24.2). Observers in Chile first saw the explosion on February 24, and within a few hours, nearly all Southern Hemisphere telescopes and every available orbiting spacecraft were focused on the object. It was officially named SN 1987A. (The SN stands for “supernova,” 1987 gives the year, and A identifies the supernova as the first seen that year.) This was one of the most dramatic changes observed in the universe in nearly 400 years. A 15-solar-mass B-type supergiant star with the catalog name SK-69°202 exploded and outshone all the other stars in the LMC combined for a few weeks, as shown in the “before” and “after” images of Figure 21.7.

Because the LMC is relatively close to Earth and because the explosion was detected so soon after it occurred, SN 1987A has provided astronomers with a wealth of detailed information on supernovae, allowing them to make key comparisons between theoretical models and observational reality. By and large, the theory of stellar evolution described in the text has held up very well. Still, SN 1987A did hold some surprises.

According to its hydrogen-rich spectrum, the supernova was of Type II—the core-collapse type—as expected for a high-mass parent star such as SK-69°202. But according to Figure 20.16 (which was computed for stars in our own Galaxy), the parent star should have been a red supergiant at the time of the explosion—not a blue supergiant, as was actually observed. This unexpected finding caused theorists to scramble in search of an explanation. It now seems that, relative to young stars in the Milky Way, the parent star’s envelope was deficient in heavy elements. This deficiency had little effect on the evolution of the core and on the supernova explosion, but it did change the star’s evolutionary track on the H–R diagram. Unlike a Milky Way star with the same mass, SK-69°202 shrank and looped back toward the main sequence once helium ignited in its core. Following the ignition of carbon, the star, with a surface temperature of around 20,000 K, had just begun to return to the right on the H–R diagram when the rapid chain of events leading to the supernova occurred.

The shape of the light curve of SN 1987A, shown in the first figure, also differed somewhat from the “standard” Type II shape (see Figure 21.8). The peak brightness was less than the expected value. For a few days after its initial detection, the supernova faded as it expanded and cooled rapidly. After about a week, the surface temperature had dropped to about 5000 K, at which point electrons and protons near the expanding

surface recombined into atomic hydrogen, making the surface layers less opaque and allowing more radiation from the interior to leak out. As a result, the supernova brightened rapidly as it grew. The temperature of the expanding layers reached a peak in late May, by which point the radius of the expanding photosphere was about  $2 \times 10^{10}$  km—a little larger than our solar system. Subsequently, the photosphere cooled as it expanded, and the luminosity dropped as the internal supply of heat from the explosion dissipated into space.




Much of the preceding description would apply equally well to a Type II supernova in our own Galaxy. The differences between the SN 1987A light curve shown here and the Type II light curve in Figure 21.8 are mainly the result of the (relatively) small size of SN 1987A’s parent star. The peak luminosity of SN 1987A was less than that of a “normal” Type II supernova because SK-69°202 was small and quite tightly bound by gravity. A lot of the energy emitted in the form of visible radiation (and evident in Figure 21.8) was used up in expanding SN 1987A’s stellar envelope, so far less was left over to be radiated into space. Thus, SN 1987A’s luminosity during the first few months was lower than expected, and the early peak evident in the figure did not occur. The peak in the SN 1987A light curve at about 80 days actually corresponds to the plateau in the Type II light curve in Figure 21.8.

About 20 hours before the supernova was detected optically, a brief (13-second) burst of neutrinos was simultaneously recorded by underground detectors in Japan and the United States. (Sec. 16.7) As discussed in the text, the neutrinos are predicted to arise when electrons and protons in the star’s collapsing core merge to form neutrons. The neutrinos preceded the light because they escaped during the collapse, whereas the first light of the explosion was emitted only after the supernova shock had plowed through the body of the star to the surface. In fact, theoretical models consistent with these observations suggest that vastly more energy was emitted in the form of

eight groups based on the total numbers of nuclear particles (protons and neutrons) that they contain. (All isotopes of all elements are included in both the table and the figure, although only a few elements are marked by dots and labeled in the figure.) Any theory proposed for the creation

of the elements must reproduce these observed abundances. The most obvious feature is that heavy elements are generally much less abundant than lighter elements. However, the many peaks and troughs evident in the figure also represent important constraints.



neutrinos than in any other form. The supernova's neutrino luminosity was many tens of thousands of times greater than its optical energy output.

Despite some unresolved details in SN 1987A's behavior, the detection of the neutrino pulse is considered to be a brilliant confirmation of theory. This singular event—the detection of neutrinos—may well herald a new age of astronomy. For the first time, astronomers have received information from a specific body beyond the solar system by radiation outside the electromagnetic spectrum.

Theory predicts that the expanding remnant of SN 1987A is now on the verge of being resolvable by optical telescopes. The accompanying photographs show the barely resolved remnant (at center) surrounded by a much larger shell of glowing gas (in yellow). Scientists reason that the progenitor of the supernova expelled this shell during its red-giant phase, some 40,000 years before the explosion. The image we see results from the initial flash of ultraviolet light from the supernova hitting the ring and causing

it to glow brightly. As the debris from the explosion itself strikes the ring, it has become a temporary, but intense, source of X-rays. The 2000 *Chandra* X-ray image and diagram at top right show the fastest-moving ejecta impacting the irregular inner edge of the ring, forming the small (1000 AU in diameter) glowing regions on its left side. The six insets to the main image clearly show the ring “lighting up” as the shock wave from the explosion reaches it.

These images also show core debris (purple) moving outward toward the ring. The six insets show material cooling and becoming fainter as it expands at nearly 3000 km/s. The main image also revealed, to everyone's surprise, two additional faint rings that might be caused by radiation sweeping across an hourglass-shaped bubble of gas, itself perhaps the result of a nonspherical “bipolar” stellar wind from the progenitor star before the supernova occurred.


Buoyed by the success of stellar-evolution theory and armed with firm theoretical predictions of what should happen next, astronomers eagerly await future developments in the story of this remarkable object.

## Hydrogen and Helium Burning

Let's begin by reviewing the reactions leading to the production of heavy elements at various stages of stellar evolution. Look again at Figure 21.6 as we discuss the reactions involved. Stellar nucleosynthesis begins with the proton–proton chain

studied in Chapter 16. ∞ (Sec. 16.6) Provided that the temperature is high enough—at least 10 million K—a series of nuclear reactions occurs, ultimately forming a nucleus of ordinary helium ( ${}^4\text{He}$ ) from four protons ( ${}^1\text{H}$ ):






▲ **FIGURE 21.14 Proton Fusion** The basic proton–proton hydrogen-burning reaction combines four protons to form a nucleus of helium-4, releasing energy in the process. (See also Figure 16.27.)

Recall that the positrons immediately interact with nearby free electrons, producing high-energy gamma rays through matter–antimatter annihilation. The neutrinos rapidly escape, carrying energy with them, but playing no direct role in nucleosynthesis. The existence of these reactions has been directly confirmed in nuclear experiments conducted in laboratories around the world during recent decades. In massive stars, an alternate sequence of reactions called the CNO cycle, involving nuclei of carbon, nitrogen, and oxygen, may greatly accelerate the hydrogen-burning process, but the basic four-protons-to-one-helium-nucleus reaction, illustrated in Figure 21.14, is unchanged.


As helium builds up in the core of a star, the burning ceases, and the core contracts and heats up. When the temperature exceeds about 100 million K, helium nuclei can overcome their mutual electrical repulsion, leading to the *triple-alpha reaction*, which we discussed in Chapter 20 (Sec. 20.2):




The net result of this reaction is that three helium-4 nuclei are combined into one carbon-12 nucleus (Figure 21.15), releasing energy in the process.

## Carbon Burning and Helium Capture

At higher and higher temperatures, heavier and heavier nuclei can gain enough energy to overcome the electrical repulsion between them. At about 600 million K (reached only in the cores of stars much more massive than the Sun), carbon nuclei can fuse to form magnesium, as depicted in Figure 21.16(a):




However, because of the rapidly mounting nuclear charges—that is, the increasing number of protons in the nuclei—fusion reactions between any nuclei larger than carbon require such high temperatures that they are actually quite





▲ **FIGURE 21.15 Helium Fusion** The basic triple-alpha helium-burning reaction occurs in post-main-sequence stars, where three helium-4 nuclei combine to form carbon-12.

uncommon in stars. The formation of most heavier elements occurs by way of an easier path. For example, the repulsive force between two carbon nuclei is three times greater than the force between a nucleus of carbon and one of helium. Thus, carbon–helium fusion occurs at a lower temperature than that at which carbon–carbon fusion occurs. As we saw in Section 20.3, at temperatures above 200 million K, a carbon-12 nucleus colliding with a helium-4 nucleus can produce oxygen-16:




If any helium-4 is present, this reaction, shown in Figure 21.16(b), is much more likely to occur than the carbon–carbon reaction.




▲ **FIGURE 21.16 Carbon Fusion** Carbon can form heavier elements (a) by fusion with other carbon nuclei or, more commonly, (b) by fusion with a helium nucleus.

Similarly, the oxygen-16 thus produced may fuse with other oxygen-16 nuclei at a temperature of about 1 billion K to form sulfur-32:



However, it is much more probable that an oxygen-16 nucleus will capture a helium-4 nucleus (if one is available) to form neon-20:



The second reaction is more likely because it occurs at a lower temperature than that necessary for oxygen–oxygen fusion.

Thus, as the star evolves, heavier elements tend to form through **helium capture** rather than by fusion of like nuclei. As a result, elements with nuclear masses of 4 units (i.e., helium itself), 12 units (carbon), 16 units (oxygen), 20 units (neon), 24 units (magnesium), and 28 units (silicon) stand out as prominent peaks in Figure 21.13, our chart of cosmic abundances. Each element is built by combining the preceding element and a helium-4 nucleus as the star evolves.


Helium capture is by no means the only type of nuclear reaction occurring in evolved stars. As nuclei of many different kinds accumulate, a great variety of reactions become possible. In some, protons and neutrons are freed from their parent nuclei and are absorbed by others, forming new nuclei with masses intermediate between those formed by helium capture. Laboratory studies confirm that common nuclei, such as fluorine-19, sodium-23, phosphorus-31, and many

others, are created in this way. However, their abundances are not as great as those produced directly by helium capture, simply because the helium-capture reactions are much more common in stars. For this reason, many of these elements (those with masses not divisible by four, the mass of a helium nucleus) are found in the troughs of Figure 21.13.

## Iron Formation

Around the time silicon-28 appears in the core of a star, a competitive struggle begins between the continued capture of helium to produce even heavier nuclei and the tendency of more complex nuclei to break down into simpler ones. The cause of this breakdown is heat. By now, the star's core temperature has reached the unimaginably large value of 3 billion K, and the gamma rays associated with that temperature have enough energy to break a nucleus apart, as illustrated in Figure 21.17(a). This is the same process of photodisintegration that will ultimately accelerate the star's iron core in its final collapse toward a Type II supernova.


Under the intense heat, some silicon-28 nuclei break apart into seven helium-4 nuclei. Other nearby nuclei that have not yet photodisintegrated may capture some or all of these helium-4 nuclei, leading to the formation of still heavier elements (Figure 21.17b). The process of photodisintegration provides raw material that allows helium capture to proceed to greater masses. Photodisintegration continues, with some heavy nuclei being destroyed and others increasing in mass. In succession, the star forms sulfur-32, argon-36, calcium-40, titanium-44, chromium-48, iron-52, and nickel-56. The chain of reactions building from silicon-28 up to nickel-56 is



This two-step process—photodisintegration followed by the direct capture of some or all of the resulting helium-4 nuclei (or alpha particles)—is often called the *alpha process*.

Nickel-56 is unstable, decaying rapidly first into cobalt-56 and then into a stable iron-56 nucleus. Any unstable nucleus will continue to decay until stability is achieved, and iron-56 is the most stable of all nuclei (Figure 21.6).

◀ **FIGURE 21.17 Alpha Process** (a) At high temperatures, heavy nuclei (such as silicon, shown here) can be broken apart into helium nuclei by high-energy photons. (b) Other nuclei can capture the helium nuclei—or alpha particles—thus produced, forming heavier elements by the so-called alpha process. This process continues all the way to the formation of nickel-56 (in the iron group).





Thus, the alpha process leads inevitably to the buildup of iron in the stellar core.


Another way of describing Figure 21.6 is to say that iron's 26 protons and 30 neutrons are bound together more strongly than the particles in any other nucleus. Iron is said to have the greatest *nuclear binding energy* of any element—more energy per particle is required to break up (unbind) an iron-56 nucleus than the nucleus of any other element. This enhanced stability of iron explains why some of the heavier nuclei in the iron group are more abundant than many lighter nuclei (see Table 21.1 and Figure 21.13): Nuclei tend to “accumulate” near iron as stars evolve.

## Making Elements Beyond Iron


If the alpha process stops at iron, how did heavier elements, such as copper, zinc, and gold, form? To form them, some nuclear process other than helium capture must have been involved. That other process is **neutron capture**: the formation of heavier nuclei by the absorption of neutrons.

Deep in the interiors of highly evolved stars, conditions are ripe for neutron capture to occur. Neutrons are produced as “by-products” of many nuclear reactions, so there are many of them present to interact with iron and other nuclei. Neutrons have no charge, so there is no repulsive barrier for them to overcome in combining with positively charged nuclei. As more and more neutrons join a nucleus, its mass continues to grow.


Adding neutrons to a nucleus—iron, for example—does not change the element. Rather, a more massive isotope of the same element is produced. Eventually, however, so many neutrons have been added to the nucleus that it becomes unstable and then decays radioactively to form a stable nucleus of some other element. The neutron-capture process then continues. For example, an iron-56 nucleus can capture a single neutron to form a relatively stable isotope, iron-57:



This reaction may be followed by another neutron capture:



Thus, another relatively stable isotope, iron-58, is produced, and this isotope can capture yet another neutron to produce an even heavier isotope of iron:



Iron-59 is known from laboratory experiments to be radioactively unstable. It decays in about a month into cobalt-59, which is stable. The neutron-capture process then resumes: Cobalt-59 captures a neutron to form the unstable cobalt-60, which in turn decays to nickel-60, and so on.

Each successive capture of a neutron by a nucleus typically takes about a year, so most unstable nuclei have plenty of time to decay before the next neutron comes along. Researchers usually refer to this “slow” neutron-capture

mechanism as the *s-process*. It is the origin of the copper and silver in the coins in our pockets, the lead in our car batteries, and the gold (and the zirconium) in the rings on our fingers. As mentioned earlier, similar slow neutron-capture processes involving nuclei of lower mass are responsible for many of the elements intermediate between those formed by helium capture. These reactions are thought to be particularly important during the late (asymptotic-giant branch) stages of low-mass stars. [∞ \(Sec. 20.3\)](#)

## Making the Heaviest Elements

The s-process explains the synthesis of stable nuclei up to, and including, bismuth-209, the heaviest-known nonradioactive nucleus, but it cannot account for the heaviest nuclei, such as thorium-232, uranium-238, or plutonium-242. Any attempt to form elements heavier than bismuth-209 by slow neutron capture fails because the new nuclei decay back to bismuth as fast as they form. Accordingly, there must be yet another nuclear mechanism that produces the very heaviest nuclei. This process is called the *r-process* (where r stands for “rapid,” in contrast to the “slow” s-process just described). The r-process operates very quickly, occurring (we think) literally during the supernova explosion that signals the death of a massive star.

During the first 15 minutes of the supernova blast, the number of free neutrons increases dramatically as heavy nuclei are broken apart by the violence of the explosion. Unlike the s-process, which stops when it runs out of stable nuclei, the neutron-capture rate during the supernova is so great that even unstable nuclei can capture many neutrons before they have time to decay. Jamming neutrons into light- and middleweight nuclei, the r-process is responsible for the creation of the heaviest-known elements. The heaviest of the heavy elements, then, are actually born *after* their parent stars have died. However, because the time available for synthesizing these heaviest nuclei is so brief, they never become very abundant. Elements heavier than iron (see Table 21.1) are a billion times less common than hydrogen and helium.

## Observational Evidence for Stellar Nucleosynthesis

The modern picture of the formation of the elements involves many different types of nuclear reactions occurring at many different stages of stellar evolution, from main-sequence stars all the way to supernovae. Elements of the periodic table from hydrogen to iron are built first by fusion and then by alpha capture, with proton and neutron capture filling in the gaps. Elements beyond iron form by neutron capture and radioactive decay. Ultimately, these elements are ejected into interstellar space as the stars in which they form reach the ends of their lives.


Scientific theories must continually be tested and validated by experiment and observation, and the theory of

stellar nucleosynthesis is no exception.  $\infty$  (Sec. 1.2) Yet almost all of the nuclear processes just described take place deep in the hearts of stars, hidden from our view, and the stars responsible for the heavy elements we see today are all long gone. How, then, can we be sure that the sequences of events presented here actually occurred (and are still occurring today)? The answer is that the theory of stellar nucleosynthesis makes many detailed predictions about the numbers and types of elements formed in stars, affording astronomers ample opportunity to observe and test its consequences. We are reassured of the theory's basic soundness by three particularly convincing pieces of evidence.


First, the rates at which various nuclei are captured and the rates at which they decay are known from laboratory experiments. When these rates are incorporated into detailed computer models of the nuclear processes occurring in stars and supernovae, the resulting elemental abundances agree extremely well, point by point, with the observational data presented in Figure 21.13 and Table 21.1. The match is remarkably good for elements up through iron and is still fairly close for heavier nuclei. Although the reasoning is indirect, the agreement between theory and observation is so striking that most astronomers regard it as very strong evidence in support of the entire theory of stellar evolution and nucleosynthesis.

Second, the presence of one particular nucleus—technetium-99—provides direct evidence that heavy elements really do form in the cores of stars. Laboratory measurements show that the technetium nucleus has a radioactive half-life of about 200,000 years, a very short time, astronomically speaking. No one has ever found even traces of naturally occurring technetium on Earth because it all decayed long ago. The observed presence of technetium in the spectra of many red-giant stars implies that it must have been synthesized in their cores through neutron capture—the only known way in which technetium can form—within the past few hundred thousand years and then transported by convection to the surface. Otherwise, we would not observe it. Many astronomers consider the spectroscopic evidence for technetium as proof that the s-process really does operate in evolved stars.


Third, the study of typical light curves from Type I supernovae indicates that radioactive nuclei form as a result of the explosion. Figure 21.18(a) (see also Figure 21.8) displays the dramatic rise in luminosity at the moment of explosion and the characteristic slower decrease in brightness. Depending on the initial mass of the exploded star, the luminosity takes from several months to many years to decrease to its original value, but the *shape* of the decay curve is nearly the same for all exploded stars. These curves have two distinct features: After the initial peak, the luminosity declines rapidly; then



(a)



(b)



(c)


► **FIGURE 21.18 Supernova Energy Emission** (a) The light curve of a Type I supernova, showing not only the dramatic increase and slow decrease in luminosity, but also the characteristic change in the rate of decay about 2 months after the explosion (red arrow). This particular supernova occurred in the faraway galaxy IC 4182 in 1938. (b) Theoretical calculations of the light emitted by the radioactive decay of nickel-56 and cobalt-56 produce a light curve similar to those actually observed in real supernova explosions, lending strong support to the theory of stellar nucleosynthesis. (c) This mostly X-ray image, taken by the orbiting *Chandra* telescope in 2013, shows the aftermath of a titanic stellar explosion known as Kepler's supernova. It was first observed by many people on Earth in 1604 and is named after the famous German astronomer who studied it even before the invention of the telescope. (NASA/CXC)

it decreases at a slower rate. This abrupt change in the rate of luminosity decay invariably occurs about 2 months after the explosion, regardless of the intensity of the outburst.

We can explain the two-stage decline of the luminosity curve in Figure 21.18(a) in terms of the radioactive decay of unstable nuclei, notably nickel-56 and its decay product cobalt-56, produced in abundance during the early moments of the supernova. From theoretical models of the explosion, we can calculate the amounts of these elements expected to form, and we know their half-lives from laboratory experiments. Because each radioactive decay produces a known amount of energy, we can then determine how the light emitted by these unstable elements should vary in time. The result is in very good agreement with the observed light curve in Figure 21.18(b)—the luminosity of a Type I supernova is entirely consistent with the decay of about 0.6 solar mass of nickel-56. More direct evidence for the presence of these unstable nuclei was first obtained in the 1970s, when a gamma-ray spectral feature of decaying cobalt-56 was identified in a supernova observed in a distant galaxy.

### CONCEPT Check

- ✓ Why are the elements carbon, oxygen, neon, and magnesium, whose masses are multiples of four, as well as the element iron, so common on Earth?




## 21.5 The Cycle of Stellar Evolution

The theory of stellar nucleosynthesis can naturally account for the observed differences in the abundances of heavy elements between the old globular-cluster stars and stars now forming in our Galaxy. ∞ (Sec. 20.5) Even though an evolved star continuously creates new heavy elements in its interior, changes in the star's composition are confined largely to the core, and the star's spectrum gives little indication of events within its core. Convection may carry some reaction products (such as the technetium observed in many red giants) from the core into the envelope, but the outer layers largely retain the star's original composition. Only at the end of the star's life are its newly created elements released and scattered into space.

Thus, the spectra of the *youngest* stars show the *most* heavy elements, because each new generation of stars increases the concentration of these elements in the interstellar clouds from which the next generation forms. Accordingly, the photosphere of a recently formed star contains a much greater abundance of heavy elements than that of a star that formed long ago. Knowledge of stellar evolution allows astronomers to estimate the ages of stars from purely spectroscopic studies, even when the stars are isolated and are not members of any cluster. ∞ (Sec. 20.5) In the last three chapters, we have seen all the ingredients that make up the complete cycle of star formation and evolution in our Galaxy. Let's briefly summarize that process, which is illustrated in Figure 21.19:

1. *Stars form* when part of an interstellar cloud is compressed beyond the point at which it can support itself against its own gravity. The cloud collapses and fragments, forming a cluster of stars. The hottest stars heat and ionize the surrounding

**Interactive FIGURE 21.19 Stellar Recycling** The cycle of star formation and evolution continuously replenishes our Galaxy with new heavy elements and provides the driving force for the creation of new generations of stars. Clockwise from the top are an interstellar cloud (Barnard 68), a star-forming region (RCW 38), a massive star ejecting a “bubble” and about to explode (NGC 7635), and a supernova remnant and its heavy-element debris (N49). (ESO; NASA)

- gas, sending shock waves through the surrounding cloud, modifying the formation of lower-mass stars, and possibly triggering new rounds of star formation.  (Sec. 19.6)
2. Within the cluster, *stars evolve*. The most massive stars evolve fastest, creating the heaviest elements in their cores and spewing them forth into the interstellar medium in supernovae. Lower-mass stars take longer to evolve, but they, too, create heavy elements and contribute significantly to the “seeding” of interstellar space when they shed their envelopes as planetary nebulae. Roughly speaking, low-mass stars are responsible for the carbon, nitrogen, and oxygen that make life on Earth possible. High-mass stars produced the iron and silicon that make up Earth itself, as well as the heavier elements on which much of our technology is based.
  3. The creation and *explosive dispersal* of newly formed elements are accompanied by further shock waves, whose passage through the interstellar medium simultaneously enriches the medium and compresses it into further star formation. Each generation of stars increases the concentration of heavy elements in

the interstellar clouds from which the next generation forms. As a result, recently formed stars contain a much greater abundance of heavy elements than do stars that formed long ago.

In this way, although some material is used up in each cycle—turned into energy or locked up in low-mass stars—the galaxy continuously recycles its matter. Each new round of formation creates stars with more heavy elements than the preceding generation had. From the old globular clusters, which are observed to be deficient in heavy elements relative to the Sun, to the young open clusters, containing much larger amounts of these elements, we observe this enrichment process in action. Our Sun is the product of many such cycles. We ourselves are another. Without the elements synthesized in the hearts of stars, neither Earth nor the life it harbors would exist.

### CONCEPT Check


- ✓ Why is stellar evolution important to life on Earth?

**The Big Question** Despite widespread observations of supernovae and their scattered debris, researchers still don't know exactly how these massive stars actually manage to explode. Although many times the mass of the Sun, these stars defy gravity by ripping themselves apart. That they blow their whole being to smithereens is well established, but how do they do it, reversing their catastrophic inward collapse to become outwardly exploding stars?


## Chapter Review

### SUMMARY

**1** A **nova** (p. 522) is a star that suddenly increases greatly in brightness, then slowly fades back to its normal appearance over a period of months. It is the result of a white dwarf in a binary system drawing hydrogen-rich material from its companion. The gas spirals inward in an **accretion disk** (p. 522) and builds up on the white-dwarf's surface, eventually becoming hot and dense enough for the hydrogen to burn explosively, temporarily causing a large increase in the dwarf's luminosity.



**2** Stars more massive than about 8 solar masses form heavier and heavier elements in their cores, at a more and more rapid pace. As they do so, their cores form a layered structure



consisting of burning shells of successively heavier elements. The process stops at iron, whose nuclei can neither be fused together nor split to produce energy. As a star's iron core grows in mass, it eventually becomes unable to support itself against gravity and begins to collapse. At the high temperatures produced during the collapse, iron nuclei are broken down into protons and neutrons. The protons combine with electrons to form more neutrons. Eventually, when the core becomes so dense that the neutrons are effectively brought into physical contact with one another, the collapse stops and the core rebounds, sending a violent shock wave out through the rest of the star. The star explodes in a **core-collapse supernova** (p. 526).


**3** Astronomers classify **supernovae** (p. 526) into two broad categories: Type I and Type II. These classes differ by their light curves and their composition. **Type I supernovae** (p. 527) are hydrogen poor and have a light curve similar in shape to that of a nova. **Type II supernovae** (p. 527) are




hydrogen rich and have a characteristic plateau in the light curve a few months after maximum. A Type II supernova is a core-collapse supernova. A Type I supernova occurs when a carbon–oxygen white dwarf in a binary system gains mass, collapses, and explodes as its carbon ignites. This type of supernova is called a **carbon-detonation supernova** (p. 527).

**4** Theory predicts that a supernova visible from Earth should occur within our Galaxy about once a century, although none has been observed in the last 400 years. We can see evidence of a past supernova in the form of a **supernova remnant** (p. 528)—a shell of exploded debris surrounding the site of the explosion and expanding into space at a speed of thousands of kilometers per second.


**5** All elements heavier than helium are formed by **stellar nucleosynthesis** (p. 531)—the production of new elements by nuclear reactions in the cores of evolved stars. Elements heavier than carbon tend to form by **helium capture** (p. 535),



rather than by the fusion of more massive nuclei. At high enough core temperatures, photodisintegration breaks apart some heavy nuclei, providing helium-4 nuclei for the synthesis of even more massive elements, up to iron. Elements beyond iron form by **neutron capture** (p. 536) in the cores of evolved stars. During a supernova, rapid neutron capture occurs, producing the heaviest nuclei of all. Comparisons between theoretical predictions of element production and observations of element abundances in stars and supernovae provide strong support for the theory of stellar nucleosynthesis.



**6** The processes of star formation, evolution, and explosion form a cycle that constantly enriches the interstellar medium with heavy elements and sows the seeds of new generations of stars. Without the elements produced in supernovae, life on Earth would be impossible.



**MasteringAstronomy**® For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- LO1** Under what circumstances will a binary star produce a nova?
- What is an accretion disk, and how does one form?
- What is a light curve? How can it be used to identify a nova or a supernova?
- LO2** Why does the core of a massive star collapse?
- What are the observational differences between Type I and Type II supernovae?
- LO3** How do the mechanisms responsible for Type I and Type II supernovae explain their observed differences?
- Roughly how often would we expect a supernova to occur in our own Galaxy? How often would we expect to *see* a galactic supernova?
- LO4 POS** What evidence is there that many supernovae have occurred in our Galaxy?
- POS** How can astronomers estimate the age of an isolated star?
- LO5** What proof do astronomers have that heavy elements are formed in stars?
- As a star evolves, why do heavier elements tend to form by helium capture rather than by fusion of like nuclei?
- Why do the cores of massive stars evolve into iron and not heavier elements?
- How and where are nuclei heavier than iron formed?
- POS** Why was supernova 1987A so important? Why are neutrino detectors important to the study of supernovae?
- LO6** How do supernovae help “recycle” galactic matter?

## Conceptual Self-Test: Multiple Choice

- A white dwarf can dramatically increase in brightness only if (a) it has binary companion; (b) fusion restarts in its core; (c) it spins very rapidly; (d) it was the core of a very massive star.
- A nova differs from a supernova in that the nova (a) can occur only once; (b) is much more luminous; (c) involves only high-mass stars; (d) is much less luminous.
- Which of the following stars will become hot enough to form elements heavier than oxygen? (a) A star that is half the mass of the Sun. (b) A star having the same mass as the Sun. (c) A star that is twice as massive as the Sun. (d) A star that is eight times more massive than the Sun.
- A massive star becomes a supernova when it (a) collides with a stellar companion; (b) forms iron in its core; (c) suddenly

increases in surface temperature; **(d)** suddenly increases in mass.

5. **VIS** Figure 21.8 (“Supernova Light Curves”) indicates that a supernova whose luminosity declines steadily in time is most likely associated with a star that is **(a)** without a binary companion; **(b)** more than eight times the mass of the Sun; **(c)** on the main sequence; **(d)** comparable in mass to the Sun.
6. An observable supernova should occur in our Galaxy about once every **(a)** year; **(b)** decade; **(c)** century; **(d)** millennium.
7. Which of the following is *not* evidence for supernovae in our Galaxy? **(a)** The rapid expansion and filamentary structure of the Crab Nebula. **(b)** Historical records from China and Europe. **(c)** The existence of binary stars in our Galaxy. **(d)** The existence of iron on Earth.
8. Nuclear fusion in the Sun will **(a)** never create elements heavier than helium; **(b)** create elements up to and including oxygen; **(c)** create all elements up to and including iron; **(d)** create some elements heavier than iron.
9. Most of the carbon in our bodies originated in **(a)** the Sun; **(b)** the core of a red-giant star; **(c)** a supernova; **(d)** a nearby galaxy.
10. The silver in our jewelry formed in **(a)** the Sun; **(b)** the core of a red-giant star; **(c)** a supernova; **(d)** a nearby galaxy.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

1. • A certain telescope can just detect the Sun at a distance of 10,000 pc. What is the apparent magnitude of the Sun at this distance? (For convenience, take the Sun’s absolute magnitude to be 5.) What is the maximum distance at which the telescope can detect a nova having a peak luminosity of  $10^5$  solar luminosities?
2. • Repeat the previous calculation for a supernova having a peak luminosity  $10^{10}$  times that of the Sun. What would be the apparent magnitude of the explosion if it occurred at a distance of 10,000 Mpc? Would it be detectable by any existing telescope?
3. •• At what distance would a supernova of absolute magnitude  $-20$  look as bright as the Sun? As the Moon? Would you expect a supernova to occur that close to us?
4. • Calculate the orbital speed of matter in an accretion disk just above the surface of a 0.6-solar-mass, 15,000-km-diameter white dwarf.
5. • Using the Sun’s current luminosity, estimate its total energy output, assuming a  $10^{10}$  year main-sequence lifetime. How does this compare with the energy released by a typical supernova?
6. •• The *Hubble Space Telescope* is observing a distant Type I supernova with peak apparent magnitude 24. Using the light curve in Figure 21.8, estimate how long after the peak brightness the supernova will become too faint to be seen.
7. • The Crab Nebula is now about 1 pc in radius. If it exploded in A.D. 1054, roughly how fast is it expanding? (Assume a constant expansion rate. Is that a reasonable assumption?)
8. •• Suppose that stars form in our Galaxy at an average rate of 10 per year. Suppose also that all stars greater than 8 solar masses explode as supernovae. Assuming that 0.36 percent of all stars fall into this category (Figure 17.23), estimate the rate of Type II supernovae in our Galaxy.


## Activities

### Collaborative

1. Look up a table of isotopes in the *Handbook of Chemistry and Physics* (available in the reference section of your library) or its equivalent online. Pick some unstable (radioactive) isotopes mentioned in the text and follow their decay into a final stable isotope. For example, choose nickel-56, iron-59, cobalt-60, and nickel-63, formed by the alpha process and s-process. In each case, note the half-life of all decays, how the isotope and its descendants decay, and what particles and radiation are produced. Repeat this exercise for the fissionable nuclei uranium-235, uranium-238, and plutonium-239.

### Individual

1. In 1758, Charles Messier discovered the sky’s most legendary supernova remnant, now called M1, or the Crab Nebula. It is located northwest of Zeta Tauri, the star that marks the southern tip of the horns of Taurus the Bull. Try to find it. An 8-inch telescope reveals the Crab’s oval shape, but it will appear faint. A 10-inch or larger telescope will show some of its famous filamentary structure.



# Neutron Stars and Black Holes

## STRANGE STATES OF MATTER

Our study of stellar evolution has led us to some very unusual and unexpected objects. Red giants, white dwarfs, and supernovae surely represent extreme states of matter completely unfamiliar to us here on Earth. Yet stellar evolution—and in particular, its end point, the death of a star—can have even more bizarre consequences. The strangest states of all result from the catastrophic implosion–explosion of stars much more massive than our Sun.

Neutron stars and black holes are among the most exotic objects in the universe. They are the end of the road for massive stars, and their bizarre properties boggle the imagination. Yet theory and observation seem to agree that, fantastic or not, they really do exist in space.

**The Big Picture** The almost unimaginable violence of supernova explosions may create objects so extreme in their behavior that they require us to reconsider some of our most cherished laws of physics. They open up a science fiction writer’s dream of fantastic phenomena that border on reality. They may even one day force scientists to construct a whole new theory of the universe.

**LEFT:** This stunning image is actually a composite of three images taken by telescopes in orbit: optical light (in yellow) observed with *Hubble*, X-ray radiation (blue and green) with *Chandra*, and infrared radiation (red) with *Spitzer*. This object is known as Cassiopeia A, the remnant of a supernova whose radiation first reached Earth about 300 years ago. The debris field shown here is about 11,000 light-years away and extends across some 10 light-years. The small turquoise dot at the center may be a neutron star created in the blast, the sole survivor of the explosion. (NASA)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Describe the properties of neutron stars, and explain how these strange objects are formed.
- 2 Explain the nature and origin of pulsars, and account for their characteristic radiation.
- 3 List and explain some of the observable properties of neutron-star binary systems.
- 4 Outline the basic characteristics of gamma-ray bursts and some theoretical attempts to explain them.
- 5 Describe how black holes are formed, and discuss their effects on matter and radiation in their vicinity.
- 6 Present Einstein’s theories of relativity, and discuss how they relate to neutron stars and black holes.
- 7 Relate the phenomena that occur near black holes to the warping of space around them.
- 8 Explain the difficulties in observing black holes, and describe some ways in which a black hole might be detected.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.



## 22.1 Neutron Stars

In Chapter 21 we saw how some stars can explode violently as *supernovae*, scattering debris across large regions of interstellar space. What remains after a supernova? Is the entire progenitor (parent) star blown to bits and dispersed throughout interstellar space, or does some portion of it survive?

### Stellar Remnants

For a Type I (carbon-detonation) supernova, most astronomers regard it as quite unlikely that any central remnant is left after the explosion. The entire star is shattered by the blast. However, for a Type II supernova, involving the implosion and subsequent rebound of a massive star's iron core, theoretical calculations indicate that part of the star may survive. ∞ (Sec. 12.2) The explosion destroys the parent star, but it may leave a tiny ultracompressed **remnant** at its center—all that remains of a star's inner core after stellar evolution has ceased. A white dwarf, the dense end point of the evolution of a low-mass star, is another example of a stellar remnant.\* ∞ (Sec. 20.3) Even by the high-density standards of a white dwarf, though, the matter within this severely compacted core is in a very strange state, unlike anything we are ever likely to find (or create) on Earth.


Recall from Chapter 21 that during the moment of implosion of a massive star—just prior to the supernova itself—the electrons in the core violently smash into the protons there, forming neutrons and neutrinos. ∞ (Sec. 21.2) The neutrinos leave the scene at (or nearly at) the speed of light, accelerating the collapse of the neutron core, which continues to contract until its particles come into contact. At that point, the central portion of the core rebounds, creating a powerful shock wave that races outward through the star, expelling matter violently into space.

The key point here is that the shock wave does not start at the very center of the collapsing core. The innermost part of the core—the region that “bounces”—remains intact as the shock wave it causes destroys the rest of the star. After the violence of the supernova has subsided, this ball of neutrons is all that is left. Researchers colloquially call this core remnant a **neutron star**, although it is not a star in any true sense of the word because all of its nuclear reactions have ceased forever.

### Neutron-Star Properties

Neutron stars are extremely small and very massive. Composed purely of neutrons packed together in a tight ball about 20 km across, a typical neutron star is not much bigger than a small asteroid or a terrestrial city (see Figure 22.1),

\*These remnants are small and compact—no larger than Earth in the case of a white dwarf and far smaller still for a neutron star. They should not be confused with supernova remnants: glowing clouds of debris scattered across many parsecs of interstellar space. ∞ (Sec. 21.3)



▲ **FIGURE 22.1 Neutron Star** Neutron stars are not much larger than many of Earth's major cities. In this fanciful comparison, a typical neutron star sits alongside Manhattan Island. (NASA)

yet its mass is greater than that of the Sun. With so much mass squeezed into such a small volume, neutron stars are incredibly dense. Their average density can reach  $10^{17}$  or even  $10^{18}$  kg/m<sup>3</sup>, nearly a billion times denser than a white dwarf. (For comparison, the density of a normal atomic nucleus is about  $3 \times 10^{17}$  kg/m<sup>3</sup>.) A single thimbleful of neutron-star material would weigh 100 million tons—about as much as a good-sized terrestrial mountain. In a sense, we can think of a neutron star as a single enormous nucleus, with an atomic mass of around  $10^{57}$ ! At these densities, neutrons resist further packing in very much the same way as electrons do (at much lower densities) in a white dwarf—this **neutron degeneracy pressure** supports the neutron star.

Neutron stars are solid objects. Provided that a sufficiently cool one could be found, you might even imagine standing on it. However, doing so would not be easy, as a neutron star's gravity is extremely powerful. A 70-kg (150-pound) human would weigh the Earth equivalent of about 10 trillion kg (10 billion tons). The severe pull of a neutron star's gravity would flatten you much thinner than a piece of paper!

In addition to large mass and small size, newly formed neutron stars have two other very important properties. First, they *rotate* extremely rapidly, with periods measured in fractions of a second. This is a direct result of the law of conservation of angular momentum, which tells us that any rotating body must spin faster as it shrinks. ∞ (More Precisely 6-1) Even if the core of the progenitor star were initially rotating quite slowly (once every couple of weeks, say, as is observed in many upper main-sequence stars), it would be spinning a few times per second by the time it had reached a diameter of 20 km.

Second, newborn neutron stars have very strong *magnetic fields*. The original field of the progenitor star is amplified by the collapse of the core because the contracting material squeezes the magnetic field lines closer together, creating a magnetic field trillions of times stronger than Earth's. [∞ \(More Precisely 19-1\)](#)

In time, theory indicates, our neutron star will spin more and more slowly as it radiates its energy into space, and its magnetic field will diminish. However, for a few million years after its birth, these two properties combine to provide the primary means by which this strange object can be detected and studied.


### CONCEPT Check

✓ Are all supernovae expected to lead to neutron stars?

## 22.2 Pulsars

Can we be sure that objects as strange as neutron stars really exist? The answer is a confident yes. The first observation of a neutron star occurred in 1967, when Jocelyn Bell, a graduate student at Cambridge University, made a surprising discovery. She observed an astronomical object emitting radio radiation in the form of rapid *pulses*. Each pulse consisted of an 0.01-second burst of radiation, after which there was nothing. Then, 1.34 s later, another pulse would arrive. The interval between the pulses was astonishingly uniform—so accurate, in fact, that the repeated emissions could be used as a precise clock. Figure 22.2 is a recording of part of the radio radiation from the pulsating object Bell discovered.

More than 1500 of these pulsating objects are now known in the Milky Way Galaxy. They are called **pulsars**. Each pulsar has its own characteristic pulse period and duration. In some cases, the pulse periods are so stable that they are by far the most accurate natural clocks known in the universe—more accurate even than the best atomic clocks on Earth. In some cases, the period is predicted to change by only a few seconds in a million years. The best current model describes a pulsar as a compact, spinning neutron star that periodically flashes radiation toward Earth.




▲ **FIGURE 22.2 Pulsar Radiation** This recording shows the regular change in the intensity of the radio radiation emitted by the first pulsar discovered, known as CP 1919. Some of the object's pulses are marked by arrows.

## The Lighthouse Model

When Bell made her discovery in 1967, she did not know what she was looking at. Indeed, no one at the time knew what a pulsar was. The explanation of pulsars as spinning neutron stars won Bell's thesis advisor, Antony Hewish, a share of the 1974 Nobel Prize in physics. Hewish reasoned that the only physical mechanism consistent with such precisely timed pulsations is a small rotating source of radiation. Only rotation can cause the high degree of regularity of the observed pulses, and only a small object can account for the sharpness of each pulse. Radiation emitted from different regions of an object larger than a few tens of kilometers across would arrive at Earth at slightly different times, blurring the pulse profile.

Figure 22.3 outlines the important features of this pulsar model. Two “hot spots” on the surface of a neutron star, or in the magnetosphere just above the surface, continuously emit radiation in a narrow “searchlight” pattern. These spots are most likely localized regions near the neutron-star's magnetic poles, where charged particles, accelerated to extremely high energies by the star's rotating magnetic field, emit radiation along the star's magnetic axis. The hot spots radiate more or less steadily, and the resulting beams sweep through space like a revolving lighthouse beacon, as the neutron star rotates. Indeed, this pulsar model is often known as the **lighthouse model**. If the neutron star happens to be oriented such that the beam sweeps across Earth, we see the star as a pulsar. The beams are observed as a series of rapid pulses—each time one of the beams flashes past Earth, a pulse is seen. The period of the pulses is the star's rotation period.

A few pulsars are definitely associated with supernova remnants, although not all such remnants have a detectable pulsar within them. Figure 22.4(c) shows a pair of optical photographs of the Crab pulsar, at the center of the Crab supernova remnant (Figures 22.4a and b). [∞ \(Sec. 21.3\)](#) In the left frame, the pulsar is off; in the right frame, it is on. The rapid variation in the pulsar's light, with a pulse period of about 33 milliseconds, is shown in Figure 22.4(d). The Crab also pulses in the radio and X-ray parts of the spectrum. By observing the speed and direction of the Crab's ejected matter, astronomers have worked backward to pinpoint the location in space at which the explosion must have





**MA** **Narrated FIGURE 22.3 Pulsar Model** The “lighthouse model” of neutron-star emission explains many of the observed properties of pulsars. Charged particles, accelerated by the magnetism of the neutron star, flow along the magnetic field lines, producing radiation that beams outward. At greater distances from the star, the field lines channel these particles into a high-speed outflow in the star’s equatorial plane, forming a pulsar wind. The beam sweeps across the sky as the neutron star rotates. If it happens to intersect Earth, we see a pulsar—much like a lighthouse beacon.

occurred and where the supernova core remnant should be located. ∞ (Sec. 21.3) It corresponds to the location of the pulsar. This is all that remains of the once massive star whose supernova was observed in 1054.

As indicated in Figure 22.3, the neutron star’s strong magnetic field and rapid rotation channel high-energy particles from near the star’s surface into the surrounding nebula (compare the expanding envelope of the 1054 supernova—Figure 22.4a). The result is an energetic *pulsar wind* that flows outward at almost the speed of light, primarily in the star’s equatorial plane. As it slams into the nebula, the wind heats the gas to very high temperatures. Figure 22.4(b) shows this process in action in the Crab—the combined *Hubble/Chandra* image reveals rings of hot X-ray-emitting gas moving rapidly away from the pulsar. Also visible in the image is a jet of hot gas (*not* the beam of radiation from the pulsar) escaping perpendicular to the equatorial plane. Eventually, the energy from the pulsar wind is deposited into the Crab nebula, where


it is radiated into space by the nebular gas, powering the spectacular display we see from Earth. ∞ (Fig. 21.10)

Most pulsars emit pulses in the form of radio radiation, but some (like the Crab) have been observed to pulse in the visible, X-ray, and gamma-ray parts of the spectrum as well. Figure 22.5 shows the Crab and the nearby Geminga pulsar in gamma rays. Geminga is unusual in that, although it pulsates strongly in gamma rays, it is barely detectable in visible light and not at all at radio wavelengths. Whatever types of radiation are produced, these electromagnetic flashes at different frequencies all occur at regular, repeated intervals, as we would expect, since they arise from the same object. However, pulses at different frequencies do not necessarily all occur at the same instant in the pulse cycle. The periods of most pulsars are quite short, ranging from about 0.03 s to 0.3 s (that is, flashing between 3 and 30 times per second). The human eye is insensitive to such rapid flashes, making it impossible to observe the flickering of a pulsar by eye, even




◀ **FIGURE 22.4 Crab Pulsar** In the core of the Crab Nebula (a), the Crab pulsar (c) blinks on and off about 30 times each second. In this pair of optical images, the pulsing can be clearly seen. (b) This more recent *Chandra* X-ray image of the Crab, superimposed on a *Hubble* optical image, shows the central pulsar, as well as rings of hot X-ray-emitting gas in the equatorial plane, driven outward by the pulsar wind. Also visible in the image is a jet of hot gas (*not* the beam of radiation from the pulsar) escaping perpendicular to the equatorial plane. (d) This radio recording shows the main pulse and its precursor, the latter probably related to the beam directed away from Earth. (ESO; NASA; *Lick Observatory*; NRAO)

Here the pulsar is unseen . . .




OFF  
(c)

. . . here it's seen (arrow).



ON




(d)

with a large telescope. Fortunately, instruments can record pulsations of light that the human eye cannot detect.

Most known pulsars are observed (usually by Doppler measurements) to have high speeds—much greater than the typical speeds of stars in our Galaxy. The most likely explanation for these anomalously high speeds is that neutron stars may receive substantial “kicks” due to asymmetries in the supernovae in which they formed. Such asymmetries, which are predicted by theory, are generally not very pronounced, but if the supernova’s enormous energy is channeled even slightly in one direction, the newborn neutron star can recoil in the opposite direction with a speed of many tens or even hundreds of kilometers per second. Thus, observations of pulsar velocities give theorists additional insight into the detailed physics of supernovae.

## Neutron Stars and Pulsars

All pulsars are neutron stars, but not all neutron stars are observed as pulsars, for two reasons. First, the two ingredients that make the neutron star pulse—rapid rotation and a strong magnetic field—both diminish with time, so the pulses gradually weaken and become less frequent. Theory indicates that, within a few tens of millions of years, the beam weakens and the pulses all but stop. Second, even a young, bright neutron star is not necessarily detectable as a pulsar from our vantage point on Earth. The pulsar beam depicted in Figure 22.3 is relatively narrow—perhaps as little as a few degrees across in some cases. Only if the neutron star happens to be oriented in just the right way do we actually see pulses. When we see those pulses from Earth, we call the body a pulsar. Note that



◀ **FIGURE 22.5 Gamma-Ray Pulsars** (a) The Crab and Geminga pulsars lie fairly close to one another in the sky. Unlike the Crab, Geminga is barely visible at optical wavelengths and undetectable in the radio region of the spectrum. (b) Sequence of *Compton Gamma-Ray Observatory* images showing Geminga's 0.24-s pulse period. (NASA)

we are using the term “pulsar” here to mean the pulsing object we observe if the beam crosses Earth. However, many astronomers use the word more generically to mean *any* young neutron star producing beams of radiation as in Figure 22.3. Such an object will be a pulsar as seen from some directions—just not necessarily ours!

Given our current knowledge of star formation, stellar evolution, and neutron stars, our observations of pulsars are consistent with the ideas that (1) *every* high-mass star dies in a supernova explosion, (2) most supernovae leave a neutron star behind (a few result in black holes, as discussed in a moment), and (3) *all* young neutron stars emit beams of radiation, just like the pulsars we actually detect. A few pulsars are definitely associated with supernova remnants, clearly establishing those pulsars’ explosive origin. On the basis of estimates of the rate at which massive stars have formed over the lifetime of the Milky Way, astronomers reason that, for every pulsar we know of, there must be several hundred thousand more neutron stars moving unseen somewhere in our Galaxy. Some formed relatively recently—less than a few million years ago—and simply happen not to be beaming their energy toward Earth. However, the vast majority are old, their youthful pulsar phase long past.

Neutron stars (and black holes too) were predicted by theory long before they were actually observed, although their extreme properties made many scientists doubt that they would ever be found in nature. The fact that we now have strong observational evidence, not just for their existence but also for the vitally important roles they play in many areas of high-energy astrophysics, is yet another testament to the fundamental soundness of the theory of stellar evolution.

### CONCEPT Check

✓ Why don't we see pulsars at the centers of all supernova remnants?


## 22.3 Neutron-Star Binaries

We noted in Chapter 17 that most stars are not single, but instead are members of binary systems. ∞ (Sec. 17.7) Although many pulsars are known to be isolated (i.e., not part of any binary), at least some do have binary companions, and the same is true of neutron stars in general (even the ones not seen as pulsars). One important consequence of this pairing is that the masses of some neutron stars have been determined quite accurately. All the measured masses are fairly close to 1.4 times the mass of the Sun—the Chandrasekhar mass of the stellar core that collapsed to form the neutron-star remnant—although a neutron star with a mass twice that of the Sun has recently been reported.


### X-Ray Sources

The late 1970s saw several important discoveries about neutron stars in binary-star systems. Numerous X-ray sources were found near the central regions of our Galaxy and also near the centers of a few rich star clusters. Some of these sources, known as **X-ray bursters**, emit much of their energy in violent eruptions, each thousands of times more luminous than our Sun, but lasting only a few seconds. A typical burst is shown in Figure 22.6.


This X-ray emission arises on or near neutron stars that are members of binary systems. Matter torn from the



(a)



(b)




◀ **FIGURE 22.6 X-Ray Burster** An X-ray burster produces a sudden, intense flash of X-rays, followed by a period of relative inactivity lasting as long as several hours. Then another burst occurs. (a) An optical photograph of the globular star cluster Terzan 2, showing a 2'' dot (arrow) at the center where the X-ray bursts originate. (b) X-ray images taken before and during the outburst. The most intense X-rays correspond to the position of the black dot shown in (a). (SAO; NASA)

surface of the (main-sequence or giant) companion by the neutron star's strong gravitational pull accumulates on the neutron star's surface. As in the case of white-dwarf accretion (see Chapter 21), the material does not fall directly onto the surface. Instead, as illustrated in Figure 22.7(a), it forms an accretion disk (compare with Figure 21.2, which depicts the white-dwarf equivalent). ∞ (Sec. 21.1) The gas goes into a tight orbit around the neutron star and then spirals slowly inward. The inner portions of the accretion disk become extremely hot, releasing a steady stream of X-rays.

As gas builds up on the neutron star's surface, its temperature rises due to the pressure of overlying material. Soon the temperature becomes hot enough to fuse hydrogen. The result is a sudden period of rapid nuclear burning that releases a huge amount of energy in a brief, but intense, flash of X-rays—an *X-ray burst*. After several hours of renewed accumulation, a fresh layer of matter produces the next burst. Thus, an X-ray burst is much like a nova on a white dwarf, but occurring on a far more violent scale because of the neutron star's much stronger gravity. ∞ (Sec. 21.1)

Not all the infalling gas makes it onto the neutron star's surface, however; in at least one case—an object known as SS 433,\* lying roughly 5000 pc from Earth—we have direct observational evidence that some material is instead shot completely out of the system at enormously high speeds. SS 433





\*The name simply identifies the object as the 433rd entry in a particular catalog of stars with strong optical emission lines.




(a)

Sometimes, jets of hot gas are ejected perpendicular to the disk.

◀ **FIGURE 22.7 X-Ray Emission** (a) Matter flows from a normal star toward a compact neutron-star companion and falls toward the surface in an accretion disk. As the gas spirals inward under the neutron star's intense gravity, it heats up, becoming so hot that it emits X-rays. (b) False-color radiographs of the peculiar object SS 433, made at monthly intervals (left to right), show its jets moving outward and its central source rotating under the gravitational influence of the companion star. (NRAO)



(b)



expels more than one Earth mass of material every year in the form of two oppositely directed narrow jets moving roughly perpendicular to the disk. Observations of the Doppler shifts of optical emission lines produced within the jets themselves imply speeds of almost 80,000 km/s—more than 25 percent of the speed of light! As the jets interact with the interstellar medium, they emit radio radiation, as shown in Figure 22.7(b).

Jets of this sort are apparently quite common in astronomical systems in which an accretion disk surrounds a compact object (such as a neutron star or a black hole). They are thought to be produced by the intense radiation and magnetic fields near the inner edge of the disk, although the details of their formation are still uncertain. Once again, note that these jets are *not* the “lighthouse” beams of radiation from the neutron star itself, shown in Figure 22.3, that can result in a pulsar, nor are they associated with a pulsar wind, as in Figure 22.4(b).

Since the discovery of SS 433, roughly a dozen stellar-mass objects with comparable properties have been discovered in our Galaxy, and we will see examples of similar phenomena on much larger scales in later chapters. Indeed, the current terminology for “stellar-scale” objects like SS 433—**microquasar**—derives from their much more energetic galactic counterparts (called *quasars*; see Sec. 24.4). SS 433 has been particularly important in the study of microquasars because we can actually observe both its disk and its jets, instead of simply having to assume their existence, as we do in more distant cosmic objects.

## Millisecond Pulsars


In the mid-1980s an important new category of pulsars was found: a class of very rapidly rotating objects called **millisecond pulsars**. Some 250 are currently known in the Milky Way Galaxy. These objects spin hundreds of times per second (i.e., their pulse period is a few milliseconds). This speed is about as fast as a typical neutron star can spin without flying apart. In some cases, the star’s equator is moving at more than 20 percent of the speed of light, a speed that suggests a phenomenon bordering on the incredible: a cosmic object of kilometer dimensions, more massive than our Sun, spinning almost at breakup speed and making nearly a thousand complete revolutions *every second!* Yet the observations and their interpretation leave little room for doubt.

The story of these remarkable objects is further complicated because most of them—about two-thirds—are found in globular clusters. This is odd, since globular clusters are known to be very old—10 billion years, at least. ∞ (Sec. 20.5) Yet, Type II supernovae (the kind that create neutron stars) are associated with massive stars that explode within a few tens of *millions* of years after their formation, and no stars have formed in any globular cluster since the cluster itself came into being. Thus, no new neutron star has been produced in a globular cluster in a very long time. But the pulsar produced by a supernova is


expected to slow down in only a few million years, and after 10 billion years its rotation should have all but ceased. Thus, the rapid rotation of the pulsars found in globular clusters cannot be a relic of their birth. Instead, these objects must have been “spun up”—that is, had their rotation rates increased—by some other, much more recent, mechanism.

The most likely explanation for the high rotation rate of pulsars is that the neutron star has been spun up by drawing in matter from a companion star. As matter spirals down onto the star’s surface in an accretion disk, it provides a “push” that makes the neutron star spin faster (see Figure 22.8). Theoretical calculations indicate that this process can spin the star up to breakup speed in about a hundred million years. This general picture is supported by the finding that, of the 150 or so millisecond pulsars seen in globular clusters, roughly half are known to be members of binary systems. The remaining solo millisecond pulsars were probably formed when an encounter with another star ejected the pulsar from the binary or when the pulsar’s own intense radiation destroyed its companion.

Thus, although a pulsar like the Crab is the direct result of a supernova, millisecond pulsars are the product of a two-stage process. First, the neutron star was formed in an ancient supernova, billions of years ago. Second, through a relatively recent interaction with a binary companion, the neutron star then achieved the rapid spin that we observe today. Once again, we see how members of a binary system can evolve in ways quite different from the manner in which single stars evolve. Notice that the scenario of accretion onto a neutron star from a binary companion is the same scenario that we just used to explain the existence of X-ray bursters. In fact, the two phenomena are closely linked. Many X-ray bursters may be on their way to becoming millisecond pulsars, and many millisecond pulsars are X-ray sources, powered by the trickle of material still falling onto them from their binary companions.



▲ **FIGURE 22.8 Millisecond Pulsar** As infalling matter strikes the star, it moves almost parallel to the surface, so it tends to make the star spin faster. Eventually, this process can result in a millisecond pulsar—a neutron star spinning at the incredible rate of hundreds of revolutions per second.




◀ **FIGURE 22.9 Cluster X-Ray Binaries** The dense core of the old globular cluster 47 Tucanae harbors more than 100 separate X-ray sources (shown in the *Chandra* image at bottom right). More than half of these are thought to be binary millisecond pulsars, still accreting small amounts of gas from their companions after an earlier period of mass transfer spun them up to millisecond speeds. (ESO; NASA)

Figure 22.9 shows the globular cluster 47 Tucanae, together with a *Chandra* image of its core showing no fewer than 108 X-ray sources—about 10 times the number that had been known in the cluster prior to *Chandra*'s launch. Roughly half of these sources are millisecond pulsars; the cluster also contains two or three “conventional” neutron-star binaries.


Most of the remaining sources are white-dwarf binaries, similar to those discussed in Chapter 21.

∞ (Sec. 21.1)

The way in which a neutron star can become a member of a binary system is the subject of active research, because the violence of a supernova explosion would be expected to blow the binary apart in many cases. Only if the supernova progenitor lost a lot of mass before the explosion would the binary system be likely to survive. Alternatively, by interacting with an existing binary and displacing one of its components, a neutron star may become part of a binary system *after* it is formed, as depicted in Figure 22.10. Astronomers are eagerly searching the skies for more millisecond pulsars to test their ideas.



## Pulsar Planets



▲ **FIGURE 22.10 Binary Exchange** A neutron star can encounter a binary made up of two low-mass stars, ejecting one of them and taking its place. This mechanism provides a means of forming a binary system with a neutron-star component (which may later evolve into a millisecond pulsar) without having to explain how the binary survived the supernova explosion that formed the neutron star.

Radio astronomers can capitalize on the precision with which pulsar signals repeat themselves to make extremely accurate measurements of pulsar motion. In January 1992, radio astronomers at the Arecibo Observatory found that the pulse period of a recently discovered millisecond pulsar lying some 500 pc from Earth varied in an unexpected, but quite regular, way. Careful analysis of the data has revealed that the period fluctuates on two distinct time scales—one of 67 days, the other of 98 days. The changes in the pulse period are small—less than one part in  $10^7$ —but repeated observations have confirmed their reality.

These fluctuations are caused by the Doppler effect as the pulsar wobbles back and forth in space. ∞ (Sec. 3.5) But what causes the wobble? The Arecibo group thinks that it is the result of the combined gravitational pulls of not one, but *two*, planets, each about three times the mass of Earth! One orbits the pulsar at a distance of 0.4 AU and the other at a distance of 0.5 AU. Their orbital periods are



67 and 98 days, respectively, matching the timing of the fluctuations. In April 1994, the group announced further observations that not only confirmed their earlier findings, but also revealed the presence of a *third* body, with mass comparable to Earth's Moon, orbiting only 0.2 AU from the pulsar.

These remarkable results constituted the first definite evidence of planet-sized bodies outside our solar system. A few other millisecond pulsars have since been found with similar behavior. However, it is unlikely that any of these planets formed in the same way as our own. Any planetary system orbiting the pulsar's progenitor star was almost certainly destroyed in the supernova explosion that created the pulsar. As a result, scientists are still unsure about how these planets came into being. One possibility involves the binary companion that provided the matter necessary to spin the pulsar up to millisecond speeds. Possibly, the pulsar's intense radiation and strong gravity destroyed the companion and then spread its matter out into a disk (a little like the solar nebula) in whose cool outer regions the planets might have condensed.

Astronomers have been searching for decades for planets orbiting main-sequence stars like our Sun, on the assumption that planets are a natural by-product of star formation.  $\infty$  (Sec. 6.7) As we have seen, these searches have now identified many extrasolar planets, although only a few planets comparable in mass to Earth have so far been detected.  $\infty$  (Sec. 15.5) It is ironic that the first Earth-sized planets to be found outside the solar system orbit a dead star and have little or nothing in common with our own world!

### CONCEPT Check

✓ What is the connection between X-ray sources and millisecond pulsars?


## 22.4 Gamma-Ray Bursts

Discovered serendipitously in the late 1960s by military satellites looking for violators of the Nuclear Test Ban Treaty and first made public in the 1970s, **gamma-ray bursts** consist of bright, irregular flashes of gamma rays typically lasting only a few seconds (Figure 22.11a). Until the 1990s, it was thought that gamma-ray bursts were basically “scaled-up” versions of X-ray bursters in which even more violent nuclear burning resulted in the release of the more energetic gamma rays. However, this is not the case.


### Distances and Luminosities

Figure 22.11(b) shows an all-sky plot of the positions of 2704 bursts detected by the *Compton Gamma-Ray Observatory* (CGRO) during its nine-year operational lifetime.  $\infty$  (Sec. 5.7) On average, CGRO detected gamma-ray bursts at the rate of about one a day. Note that the bursts are distributed uniformly across the sky (their distribution is said to be *isotropic*), rather than being confined to the relatively narrow band of the Milky Way (compare Figure 5.37). The bursts seemingly never repeat at the same location, show no obvious clustering, and appear unaligned with any known large-scale structure, near or far. Although CGRO was unable to measure distances to any of the bursts it observed, the isotropy of the data convinced most astronomers that the bursts originate far outside our own Galaxy—at so-called *cosmological distances*, comparable to the scale of the universe itself.

In fact, measuring the distance to a gamma-ray burst is no easy task. The gamma-ray observations do not provide enough information to tell us how far away the burst is, so astronomers must instead associate the burst with some other object in the sky—called the burst *counterpart*—whose distance can be measured by other means. The techniques for studying counterparts generally involve observations in the optical or



▲ **FIGURE 22.11 Gamma-Ray Bursts** (a) Plots of intensity versus time (in seconds) for three gamma-ray bursts. Note the substantial differences between them. Some bursts are irregular and spiky, whereas others are much more smoothly varying. (b) Positions on the sky of all the gamma-ray bursts detected by *Compton Observatory* during its nearly 9-year operating lifetime. The bursts appear to be distributed uniformly across the entire sky, which is mapped here with the plane of the Milky Way running horizontally across the center of the plot. (NASA)



▲ **FIGURE 22.12 Gamma-Ray Burst Counterparts** The long-duration gamma-ray burst, GRB 080319B, was one of the brightest yet observed. Its light that reached Earth on March 19, 2008, was emitted 7.5 billion years ago, yet for a few seconds the flash would have been visible to the unaided eye—if anyone had been looking in just the right spot! Only moments after the detonation, it was observed in X-rays (a) and in visible light (b). (NASA; ESO)

X-ray parts of the electromagnetic spectrum. The problem is that the resolution of a gamma-ray telescope is quite poor, so the burst positions may be uncertain by up to a degree, and a relatively large region of the sky must be scanned in search of a counterpart. ∞ (Sec. 5.7) In addition, the “afterglow” of a burst at X-ray or optical wavelengths fades rapidly, severely limiting the time available to complete the search.

The most successful searches for burst counterparts have been carried out by satellites combining gamma-ray detectors with X-ray and/or optical telescopes. NASA’s *Swift* mission, launched in 2004 and still operational, combines a wide-angle gamma-ray detector (to monitor as much of the sky as possible) with two telescopes: one X-ray and one optical/ultraviolet instrument. The gamma-ray detector system pinpoints the burst to an accuracy of about 4 arc minutes, and within seconds the onboard computer automatically repositions the satellite to point the X-ray and optical telescopes in that direction. At the same time, the craft relays the burst position to other instruments in space and on the ground. *Swift* detects burst counterparts at the rate of about one per week and has played a pivotal role in advancing our understanding of these violent phenomena. Figures 22.12(a) and (b) show *Swift* X-ray and optical images of GRB 080319B, one of the brightest bursts to date. Automated observations at many wavelengths began within seconds of *Swift*’s detection, making this burst one of the most intensively studied on record.

The first direct measurement of the distance to a gamma-ray burst was made in 1997, when astronomers succeeded in obtaining a spectrum of the visible afterglow of a particularly energetic burst. The spectrum contained absorption lines of iron and magnesium, but they were redshifted by almost a factor of two in wavelength. Such redshifts, as we will see in

Chapter 24, are the result of the expansion of the universe, and they are clear proof—a “smoking gun,” if you will—that this particular gamma-ray burst, and presumably all others, really does lie at cosmological distances. (This one occurred more than 2 billion parsecs from Earth.) To date, the distances to hundreds of gamma-ray bursts have been measured from their afterglows. All are very large, implying that the bursts must be extremely energetic, otherwise they wouldn’t be detectable by our equipment.

If we assume that the gamma rays are emitted equally in all directions, we can easily calculate the total energy of a burst using the inverse-square law. ∞ (Sec. 17.2) Doing so, we would find that each burst apparently generates more energy—and in some cases hundreds of times more energy—than a

typical supernova explosion, all in a matter of seconds! Such enormous energies would defy theoretical explanation. Fortunately for theorists, the above simple estimate greatly *overestimates* the actual amount of energy produced. Very probably, the radiation is emitted in the form of a rather narrow *jet*, so the energy we see is representative of only a small fraction of the sky.

As an analogy, consider a handheld laser pointer of the sort commonly used in talks and lectures. It radiates only a few milliwatts of power—far less than a household lightbulb—but it appears enormously bright if you happen to look directly into the beam. (*Don’t* do this, by the way!) Like the laser beam, a gamma-ray burst appears to be so bright because all of its energy is concentrated in almost a single direction instead of being radiated in all directions into space. Taking this into account reduces the total emission to much more understandable—but still “supernova-scale”—levels.

## What Causes the Bursts?

Not only are gamma-ray burst sources extremely energetic, they are also very *small*. The millisecond flickering in the bursts implies that whatever their origin, all of their energy must come from a volume no larger than a *few hundred kilometers* across. The reasoning is as follows: If the emitting region were, say, 300,000 km—1 light-second—across, even an instantaneous change in intensity at the source would be smeared out over a time interval of 1 s as seen from Earth, because light from the far side of the object would take 1 s longer to reach us than light from the near side. For the gamma-ray variation not to be blurred by the light travel time, the source cannot be more than 1 light-millisecond, or just 300 km, in diameter.

Theoretical models of gamma-ray bursts describe the burst as a *relativistic fireball*—an expanding region, probably a jet, of superhot gas radiating furiously in the gamma-ray part of the spectrum. (The term “relativistic” here means that particles are moving at nearly the speed of light and that Einstein’s theory of relativity is needed to describe them—see Section 22.6.) The complex burst structure and afterglows are produced as the fireball expands, cools, and interacts with its surroundings.


Two leading models for the energy source have emerged, as sketched in Figure 22.13. The first (Figure 22.13a) is the “true” end point of a binary-star system—the *merger* of the component stars. Suppose that both members of the binary evolve to become neutron stars. As the system continues to evolve, gravitational radiation (see *Discovery 22-2*) is released, and the two ultradense stars spiral in toward each other. Once they are within a few kilometers of one another, coalescence is inevitable. Such a merger will likely produce a violent explosion comparable in energy to that generated by a supernova and energetic enough to explain the flashes of gamma rays we observe. The overall rotation of the binary system may channel the energy into a high-speed, high-temperature jet.

The second model (Figure 22.13b), sometimes called a *hypernova*, is a “failed” supernova—but what a failure! In this picture, a very massive star undergoes core collapse much as described earlier for a Type II supernova, but instead

of forming a neutron star, the core collapses to a black hole (see Section 22.5). [∞ \(Sec. 21.2\)](#) At the same time, the blast wave racing outward through the star stalls. Instead of being blown to pieces, the inner part of the star begins to implode, forming an accretion disk around the black hole and creating a relativistic jet. The jet punches its way out of the star, producing a gamma-ray burst as it slams into the surrounding shells of gas expelled from the star during the final stages of its nuclear-burning lifetime. [∞ \(Discovery 20-2\)](#) At the same time, intense radiation from the accretion disk may restart the stalled supernova, blasting what remains of the star into space.

The idea of a relativistic fireball has become widely accepted among workers in this branch of astrophysics, and thanks to the “rapid response network” provided by *Swift* and other instruments, astronomers have detailed observations of many afterglows from both long and short bursts. Can we tell which, if either, of the two models just described is correct? In fact, experts in the field would say that the answer is probably *both*.

The neutron-star merger model naturally accounts for the short gamma-ray bursts, and the rapidly fading X-ray afterglows from the short bursts are consistent with the detailed predictions of the merger scenario. Recent observations also reveal that a few of these bursts may involve the theoretically predicted, but rare, merger between a neutron star and a black hole, which should have its own characteristic light signature.




**▲ FIGURE 22.13 Gamma-Ray Burst Models** Two models have been proposed to explain gamma-ray bursts. Part (a) depicts the merger of two neutron stars; part (b) shows the collapse of a single massive star. Both models predict a relativistic fireball, perhaps releasing energy in the form of jets, as illustrated at right.

The hypernova model predicts bursts of relatively long duration and is the leading explanation of the long bursts. Figure 22.14(a) and (b) show the afterglow of the long-burst GRB 030329, as seen by the 8.2-m VLT in Chile. [∞ \(Sec. 5.2\)](#) Both the spectrum and the light curve were consistent with what astronomers expected from the supernova of a very massive (roughly 25-solar-mass) star. [∞ \(Sec. 21.3\)](#) Figure 22.14(c) shows a simplified light curve of another long burst, illustrating how the prompt “burst” and later “hypernova” components can be distinguished.

### CONCEPT Check

✓ **What are gamma-ray bursts, and why have they posed such a challenge to current theory?**



▲ **FIGURE 22.14 Hypernova?** The gamma-ray burst GRB 030329 may prove crucial to theorists’ understanding of the physical processes underlying these violent phenomena. First detected by the *High Energy Transient Explorer 2* satellite and subsequently observed at radio, optical, and X-ray wavelengths, the burst counterpart has all the hallmarks of a high-mass supernova, lending strong support to the hypernova model. Here, the counterpart is shown, (a) near the moment of the burst and (b) fading a month later. (c) This simplified schematic shows the emitted radiation from another, similar, gamma-ray burst. (ESO)

## 22.5 Black Holes

Table 22.1 lists the properties of some of the dense stellar remnants we have encountered in this text. Brown, white, and black dwarfs are held up by electron degeneracy—the resistance of tightly packed electrons to further compression. [∞ \(Secs. 20.2, 20.3\)](#) The much denser neutron stars, as we have just seen, are supported by a similar mechanism involving neutrons instead. Squeezed together, the neutrons in a neutron star form a hard ball of matter that not even gravity can compress further. Or can it? Is it possible that, given enough matter packed into a small enough volume, the collective pull of gravity can eventually crush any opposing pressure? Can gravity continue to compress a massive star into an object the size of a planet, a city, a pinhead—even smaller? The answer, apparently, is yes.

### The Final Stage of Stellar Evolution

Although the precise figure is uncertain, mainly because the behavior of matter at very high densities is not well enough understood, most researchers concur that the mass of a neutron star cannot exceed about 3 solar masses. That is the neutron-star equivalent of the white dwarf’s Chandrasekhar mass limit discussed in the previous chapter. [∞ \(Sec. 21.3\)](#) Above this limit, not even tightly packed neutrons can withstand the star’s gravitational pull. In fact, we know of *no* force that can counteract gravity once neutron degeneracy pressure is overwhelmed. If enough material is left behind after a supernova such that the central core exceeds the 3-solar-mass limit, gravity wins the battle with pressure once and for all, and the star’s central core collapses forever. Stellar evolution theory indicates that this is the fate of any star whose main-sequence mass exceeds about 25 times the mass of the Sun.

The limit of 3 solar masses is uncertain, in part because it ignores the effects of magnetism and rotation, both of which are surely present in the cores of evolved stars. Because these effects can compete with gravity, they influence stellar evolution. [∞ \(Sec. 19.1\)](#) In addition, we do not know precisely how the basic laws of physics might change in regions of very dense matter that is both rapidly spinning and strongly magnetized. Generally speaking, theorists expect that the neutron-star mass limit increases when magnetism and rotation are included because even larger amounts of mass will then be needed for gravity to

TABLE 22.1 Properties of Stellar Remnants

Remnant	Typical Mass (solar masses)	Typical Radius (km)	Typical Density (kg/m <sup>3</sup> )	Support	Context (Section)
brown dwarf	less than 0.08	70,000	10 <sup>5</sup>	electron degeneracy	H fusion never started (19.3)
white dwarf	less than 1.4	10,000	10 <sup>9</sup>	electron degeneracy	stellar core after fusion stops at C/O (20.3)
black dwarf	less than 1.4	10,000	10 <sup>9</sup>	electron degeneracy	“cold” white dwarf (20.3)
neutron star	1.4–3 (approx.)	10	10 <sup>18</sup>	neutron degeneracy	remnant of a core collapse supernova (22.1)
black hole	more than 3	10	infinite at the center	none	remnant of a core collapse supernova with massive progenitor (22.5)

compress stellar cores into neutron stars or black holes, but the amount of the increase is not currently known.

As the stellar core shrinks, the gravitational pull in its vicinity eventually becomes so great that even light itself is unable to escape. The resultant object therefore emits no light, no radiation, and no information whatsoever. Astronomers call this bizarre end point of stellar evolution, in which a massive core remnant collapses in on itself and vanishes forever, a **black hole**.

## Escape Speed

Newtonian mechanics—up to now our reliable and indispensable tool in understanding the universe—cannot adequately describe conditions in or near black holes. ∞ (Sec. 2.8) To comprehend these collapsed objects, we must turn instead to the modern theory of gravity: Einstein’s *general theory of relativity*, discussed in Section 22.6. Still, we can usefully discuss some aspects of these strange bodies in more or less Newtonian terms. Let’s consider again the familiar Newtonian concept of escape speed—the speed needed for one object to escape from the gravitational pull of another—supplemented by two key facts from relativity: (1) Nothing can travel faster than the speed of light, and (2) all things, *including light*, are attracted by gravity.

A body’s escape speed is proportional to the square root of the body’s mass divided by the square root of its radius. ∞ (Sec. 2.8) Earth’s radius is 6400 km, and the escape speed from Earth’s surface is just over 11 km/s. Now consider a hypothetical experiment in which Earth is squeezed on all sides by a gigantic vise. As our planet shrinks under the pressure, its mass remains the same, but its escape speed increases because the planet’s radius is decreasing. For example, suppose Earth were compressed to one-fourth its present size. Then the proportionality mentioned in the first sentence of this paragraph predicts that our planet’s escape speed would double. To escape from this compressed Earth, an object would need a speed of at least 22 km/s.

Imagine compressing Earth some more. Squeeze it by an additional factor of, say, a thousand, making its radius hardly more than a kilometer. Now a speed of about

630 km/s would be needed to escape from the planet’s gravitational pull. Compress Earth still further, and the escape speed continues to rise. If our hypothetical vise were to squeeze Earth hard enough to crush its radius to about a centimeter, then the speed needed to escape the planet’s surface would reach 300,000 km/s. But this is no ordinary speed—it is the speed of light, the fastest speed allowed by the laws of physics as we currently know them.

Thus, if, by some fantastic means, the entire planet Earth could be compressed to less than the size of a grape, the escape speed would exceed the speed of light. However, because nothing can in fact exceed that speed, the compelling conclusion is that *nothing—absolutely nothing—could escape from the surface of such a compressed body*.

## Black Hole Properties

The origin of the term *black hole* now becomes clear: No form of radiation—radio waves, visible light, X-rays, indeed, photons of any wavelength—would be able to escape the intense gravity of our grape-sized Earth. With no photons leaving, our planet would be invisible and uncommunicative. No signal of any sort could be sent to the universe beyond. For all practical purposes, such a supercompact Earth could be said to have disappeared from the universe! Only its gravitational field would remain behind, betraying the presence of its mass, now shrunk to a point.

The “one-way” nature of the flow of energy and matter into a black hole means that almost all information about the matter falling into it—gas, stars, spaceships, or people—is lost. Only a few scraps survive. In fact, we now know that, regardless of the composition, structure, or history of the objects that formed the hole, only three physical properties can be measured from the outside: the hole’s *mass*, *charge*, and *angular momentum*. All other information is lost once matter enters the hole. Thus, *just three numbers* are required to completely describe a black hole’s outward appearance and interaction with the rest of the universe.

In this chapter, we will concentrate on black holes that formed from nonrotating, electrically neutral matter. Such objects are *completely specified* once their masses are known.

## The Event Horizon


Astronomers have a special name for the critical radius at which the escape speed from an object would equal the speed of light and within which the object could no longer be seen. It is the **Schwarzschild radius**, after Karl Schwarzschild, the German scientist who first studied its properties. The Schwarzschild radius of any object is simply proportional to the object's mass. For Earth, the Schwarzschild radius is 1 cm; for Jupiter, which is about 300 Earth masses, it is approximately 3 m; for the Sun, with a mass of 300,000 Earth masses, it is 3 km. For a 3-solar-mass stellar core remnant, the Schwarzschild radius is about 9 km. As a convenient rule of thumb, the Schwarzschild radius of an object is simply 3 km, multiplied by the object's mass, measured in solar masses. Every object has a Schwarzschild radius; it is the radius to which the object would have to be compressed for it to become a black hole. Put another way, a black hole is an object that happens to lie within its own Schwarzschild radius.

The surface of an imaginary sphere with radius equal to the Schwarzschild radius and centered on a collapsing star is called the **event horizon**. It defines the region within which no event can ever be seen, heard, or known by anyone outside. Even though there is no matter of any sort associated with it, we can think of the event horizon as the “surface” of a black hole.

A 1.4-solar-mass neutron star has a radius of about 10 km and a Schwarzschild radius of 4.2 km. If we were to keep increasing the star's mass, the star's Schwarzschild radius would grow, although its actual physical radius would not. In fact, the radius of a neutron star *decreases* slightly with increasing mass. By the time the neutron star's mass exceeded about 3 solar masses, it would lie just within its own event horizon, and it would collapse of its own accord. It would not stop shrinking at the Schwarzschild radius: The event horizon is not a physical boundary of any kind—just a communications barrier. The remnant would shrink right past the Schwarzschild radius to ever-diminishing size on its way to being crushed to a point.

Thus, provided that at least 3 solar masses of material remain behind after a supernova explosion, the remnant core will collapse catastrophically, diving below the event horizon in less than a second. The core simply “winks out,” disappearing and becoming a small dark region from which nothing can escape—a literal black hole in space. Theory indicates that this is the likely fate of stars having more than about 20 to 25 times the mass of the Sun.

## 22.6 Einstein's Theories of Relativity

The objects we have been studying in this and the last few chapters have taken us far beyond the scope of Newtonian mechanics and gravitation discussed in Chapter 2.  (Sec. 2.8) Now, in the face of extreme states of matter, speeds comparable to that of light, and gravitational fields so intense that

not even light can escape, these “workhorse” theories must give way to more refined tools. These tools are the theories of *special* and *general relativity*.

### Special Relativity


By the latter part of the 19th century, physicists were well aware of the special status of the speed of light,  $c$ . It was, they knew, the speed at which all electromagnetic waves traveled, and, as best they could tell, it represented an upper limit on the speeds of *all* known particles. Scientists struggled without success to construct a theory of mechanics and radiation in which  $c$  was a natural speed limit.

In 1887, a fundamental experiment carried out by the American physicists A. A. Michelson and E. W. Morley compounded theorists' problems further by demonstrating another important and unique aspect of light: The measured speed of a beam of light is *independent* of the motion of either the observer or the source (see *Discovery 22-1*). No matter what our motion may be relative to the source of the radiation, we always measure precisely the same value for  $c$ : 299,792.458 km/s.


A moment's thought tells us that this is a decidedly non-intuitive statement. For example, if we were traveling in a car moving at 100 km/h and we fired a bullet forward with a speed of 1000 km/h relative to the car, an observer standing at the side of the road would see the bullet pass by at  $100 + 1000 = 1100$  km/h, as illustrated in Figure 22.15(a). However, the Michelson–Morley experiment tells us that if we were traveling in a rocket ship at one-tenth the speed of light,  $0.1c$ , and we shone a searchlight beam ahead of us (Figure 22.15b), an outside observer would measure the speed of the beam not as  $1.1c$ , as the example of the bullet would suggest, but as  $c$ . The rules that apply to particles moving at or near the speed of light are different from those we are used to in everyday life.

The **special theory of relativity** (or just *special relativity*) was proposed by Einstein in 1905 to deal with the preferred status of the speed of light. The theory is the mathematical framework that allows us to extend the familiar laws of physics from low speeds (i.e., speeds much less than  $c$ , which are often referred to as *nonrelativistic*) to very high (or *relativistic*) speeds, comparable to  $c$ . The essential features of the theory are as follows:

1. The speed of light,  $c$ , is the maximum possible speed in the universe, and all observers measure the same value for  $c$ , regardless of their motion. Einstein broadened this statement into the *principle of relativity*: The basic laws of physics are *the same* to all unaccelerated observers.
2. There is no absolute frame of reference in the universe; that is, there is no “preferred” observer relative to whom all other velocities can be measured. Put another way, there is no way to tell who is moving and who is not. Instead, only relative velocities between observers matter (hence the term “relativity”).



◀ **FIGURE 22.15 Speed of Light**  
 (a) A bullet fired from a speeding car is measured by an outside observer to have a speed equal to the sum of the speeds of the car and of the bullet. (b) A beam of light shining forward from a high-speed spacecraft is still observed to have speed  $c$ , regardless of the speed of the spacecraft. The speed of light is thus independent of the speed of the source or of the observer.



gravity into special relativity took Einstein another decade. The result once again overturned scientists' conception of the universe.

In 1915, Einstein illustrated the connection between special relativity and gravity with the following famous "thought experiment." Imagine that you are enclosed in an elevator with no windows, so that you cannot directly observe the outside world, and the elevator is floating in space.

- Neither space nor time can be considered independently of one another. Rather, they are each components of a single entity: **spacetime**. There is no absolute, universal time—observers' clocks tick at different rates, depending on the observers' motions relative to one another.


Special relativity is equivalent to Newtonian mechanics in describing objects that move much more slowly than the speed of light, but it differs greatly in its predictions at relativistic velocities. (See *Discovery 22-1*) Yet, despite their often nonintuitive nature, all of the theory's predictions have been repeatedly verified to a high degree of accuracy. Today, special relativity lies at the heart of modern science. No scientist seriously doubts its validity.

You are weightless. Now suppose that you begin to feel the floor press up against your feet. Weight has apparently returned. There are two possible explanations for this, as shown in Figure 22.16. A large mass could have come nearby, and you are feeling its downward gravitational attraction (Figure 22.16a), or the elevator has begun to accelerate upward, and the

*A person inside a windowless elevator could not distinguish between these two cases.*

## General Relativity

Einstein's special theory of relativity is cast in terms of frames of reference ("observers") moving at *constant* speeds with respect to one another. In constructing his theory, Einstein rewrote the laws of motion expounded by Newton more than two centuries previously. ∞ (Sec. 2.7) But Newton's other great legacy—the theory of gravitation—does not deal with observers moving at constant relative velocities. Rather, gravity causes observers to *accelerate* relative to one another, making for a much more complex mathematical problem. Fitting



▶ **FIGURE 22.16 Einstein's Elevator** Einstein reasoned that no experiment conducted entirely within an elevator can tell the passenger whether the force he feels is (a) due to the gravity of a nearby massive object or (b) caused by the acceleration of the elevator itself.

## DISCOVERY 22-1

## Special Relativity

In 1887 the Michelson–Morley experiment attempted to determine Earth's motion relative to the “absolute” space through which light supposedly moved. As illustrated in the first figure, Michelson and Morley expected the measured speed of a beam of light to change as their equipment moved due to Earth's rotation and orbit around the Sun—faster when the beam was moving opposite to Earth's motion (to the left in the figure) and slower when Earth was “catching up” on it (to the right). In fact, they measured precisely the *same* speed of light for any orientation of their apparatus. This meant either that Earth was not moving through space—which conflicts with the fact that we see stellar parallax—or that Newtonian thinking and human intuition somehow go awry when light is involved. Far from measuring the properties of absolute space, the Michelson–Morley experiment ultimately demolished the entire concept—and with it, the 19th-century view of the universe.

With the special theory of relativity, Einstein explained the Michelson–Morley experiment and elevated the speed of light to the status of a constant of nature. He rewrote the laws of mechanics to reflect that new fact and thereby opened the door to a flood of new physics and a much deeper understanding of the universe. But many commonsense ideas had to be abandoned in the process and replaced with some decidedly less intuitive concepts.

Imagine that you are an observer watching a rocket ship fly past at relative velocity  $v$  and that the craft is close enough for you to make detailed observations inside its cabin. If  $v$  is much less than the speed of light,  $c$ , you would see nothing out of the ordinary—special relativity is consistent with familiar Newtonian mechanics at low velocities. As the ship's velocity increases, however, you begin to notice that it appears to *contract* in the direction in which it is moving. A meterstick on board, identical at launch to the one in your laboratory, is now shorter than its twin. This is called **Lorentz contraction** (or Lorentz–Fitzgerald contraction). The graph shows the stick's measured length aboard the moving ship: At low speeds (bottom) the meterstick measures 1 meter, but at high speeds (top) the stick is shortened considerably. A meterstick moving at 90 percent of the speed of light would shrink to a little less than half a meter.

At the same time, the ship's clock, synchronized prior to launch with your own, now ticks *more slowly*. This phenomenon, known as *time dilation*, has been observed many times in laboratory experiments in which fast-moving radioactive particles are observed to decay more slowly than if they were at rest in the lab. Their internal clocks—their half-lives—are slowed by their rapid motion. [∞ \(More Precisely 7-2\)](#) Although no material particle can actually reach the speed of light, Einstein's theory implies



that, as approaches  $c$ , the measured length of the meterstick will shrink to nearly zero and the clock will slow to a virtual stop.

Of course, from the point of view of an astronaut on board the spaceship, *you* are the one moving rapidly. Seen from the ship, *you* appear to be compressed in the direction of motion, and *your* clock runs slowly! How can this be? The answer is that, in relativity, the familiar concept of *simultaneity*—the idea that two events happen “at the same time”—is no longer well defined, but depends on the observer.

When measuring the length of the moving meterstick, you note the positions of the two ends *at the same time, according to your clock*. But those two events—the two measurements—do *not* occur at the same time as seen by the astronaut on the spaceship. From her viewpoint, your measurement of the leading end of the meterstick occurs *before* your measurement of the trailing end, resulting in the Lorentz contraction you observe. A similar argument applies to measurements of time, such as the period between two clock ticks. Time dilation occurs because the measurements occur at the same location and different times in one frame, but *at different places and times* in the other.

Further experimentation would show that the *mass* of the rocket ship also rises as the ship accelerates, becoming nearly infinite as the ship's speed approaches that of light. Finally, perhaps the best-known prediction of special relativity is that the rocket ship's *energy* and mass are proportional to one another, connected by the famous equation  $E = mc^2$ .

Einstein's revolutionary ideas required physicists to abandon some long-held, cherished, and “obvious” facts about the universe. Perhaps not surprisingly, they encountered initial opposition from many of Einstein's colleagues, but the gain in scientific understanding soon overcame the price in unfamiliarity. Within just a few years, special relativity had become almost universally accepted, and Einstein was on his way to becoming the best-known scientist on the planet.





force you feel is that exerted by the elevator as it accelerates you at the same rate (Figure 22.16b). The crux of Einstein’s argument is this: There is *no* experiment that you can perform within the elevator (without looking outside) that will let you distinguish between these two possibilities.

Thus, Einstein reasoned, there is no way to tell the difference between a gravitational field and an accelerated frame of reference (such as the rising elevator in the thought experiment). This statement is known more formally as the **equivalence principle**. Using it, Einstein set about incorporating gravity into special relativity as a general acceleration of all particles. However, he found that another major modification to the theory of special relativity had to be made. As we have just seen, a central concept in relativity is the notion that space and time are not separate quantities, but instead must be treated as a single entity, spacetime. To incorporate the effects of gravity, the mathematics forced Einstein to the unavoidable conclusion that spacetime had to be *curved*. The resulting theory, the result of including gravity within the framework of special relativity, is called **general relativity**.

The central concept of general relativity is this: Matter—all matter—tends to “warp” or curve space in its vicinity. Objects such as planets and stars react to this warping by changing their paths. In the Newtonian view of gravity, particles move on curved trajectories because they are acted upon by a gravitational force.  $\infty$  (Sec. 2.7) In Einsteinian relativity, those same particles move on curved trajectories because they are falling freely through space, following the curvature of spacetime produced by some nearby massive object. The more the mass, the greater is the warping. Thus, in general relativity, there is no such thing as a “gravitational force” in the Newtonian sense. Objects move as they do because they follow the curvature of spacetime, which is determined by the amount of matter present. Stated more loosely, as summed up by the renowned physicist John Archibald Wheeler, “Spacetime tells matter how to move, and matter tells spacetime how to curve.”

Some props may help you visualize these ideas. Bear in mind, however, that these props are not real, but only tools to help you grasp some exceedingly strange concepts. Imagine a pool table with the tabletop made of a thin rubber sheet rather than the usual hard felt. As Figure 22.17 suggests, such a rubber sheet becomes distorted when a heavy weight (e.g., a rock) is placed on it. The heavier the rock (Figure 22.17a), the larger is the distortion.

Trying to play pool on this table, you would quickly find that balls passing near the rock were deflected by the curvature of the tabletop (Figure 22.17b). The pool balls are not attracted to the rock in any way; rather, they respond to the curvature of the sheet produced by the rock’s presence. In much the same way, anything that moves through space—matter *or* radiation—is deflected by the curvature of spacetime near a star. For example, Earth’s orbital path is the trajectory that results as our planet falls freely in the relatively gentle curvature of space created by our Sun. When the curvature is small


(i.e., gravity is weak), both Einstein and Newton predict the same orbit—the one we observe. However, as the gravitating mass increases, the two theories begin to diverge.

## Curved Space and Black Holes


Modern notions about black holes rest squarely on the general theory of relativity. Although white dwarfs and (to a lesser extent) neutron stars can be adequately described by the classical Newtonian theory of gravity, only the modern Einsteinian theory of relativity can properly account for the bizarre physical properties of black holes.

In Figure 22.17, we saw how the distortion of space (the rubber sheet in our analogy) increases as the mass of the object causing the distortion increases. In these terms, a *black hole*

The amount of mass determines the amount of curvature and hence the amount of deflection.




(a)



(b)

**Interactive FIGURE 22.17 Curved Space** (a) A pool table made of a thin rubber sheet sags when a weight is placed on it. Likewise, space is bent, or warped, in the vicinity of any massive object. (b) A ball rolling across the table is deflected by the curvature of the surface, in much the same way that a planet’s curved orbit is determined by the curvature of spacetime produced by the Sun.



**▲ FIGURE 22.18 Space Warping** Mass causes a rubber sheet (or space) to be curved. As people assemble at a fixed spot on the sheet (marked by an X), the curvature grows larger, as shown in frames (a), (b), and (c). The blue arrows represent some directions in which information can be sent from place to place. (d) The people are eventually sealed inside the bubble, forever trapped and cut off from the outside world.

is a region of space where the gravitational field becomes overwhelming and the curvature of space extreme. At the event horizon itself, the curvature is so great that space “folds over” on itself, causing objects within to become trapped and disappear.

Let’s consider another analogy. Imagine a large extended family of people living on a huge rubber sheet—a sort of gigantic trampoline. Deciding to hold a reunion, they converge on a given place at a given time. As shown in Figure 22.18, one person remains behind, not wishing to attend. She keeps in touch with her relatives by means of “message balls” rolled out to her (and back from her) along the surface of the sheet. These message balls are the analog of radiation carrying information through space.

As the people converge, the rubber sheet sags more and more. Their accumulating mass creates an increasing amount of space curvature. The message balls can still reach the lone person far away in nearly flat space, but they arrive less frequently as the sheet becomes more and more warped and stretched—as shown in Figures 22.18(b) and (c)—and the balls have to climb out of a deeper and deeper well. Finally, when enough people have arrived at the appointed spot, the mass becomes too great for the rubber to support them. As illustrated in Figure 22.18(d), the sheet pinches off into a “bubble,” compressing the people into oblivion and severing their communications with the lone survivor outside. This final stage represents the formation of an event horizon around the reunion party.

Right up to the end—the pinching off of the bubble—two-way communication is possible. Message balls can reach the outside from within (but at a slower and slower rate as the rubber stretches), and messages from outside can get in without difficulty. However, once the event horizon (the bubble) forms, balls from the outside can still fall in, but they can no longer be sent back out to the person left behind, no matter how fast they are rolled. They cannot make it past the “lip” of the bubble in Figure 22.18(d). This analogy (very) roughly depicts how a black hole warps space completely around on itself, isolating its interior from the rest of the universe. The essential ideas—the slowing down and eventual cessation of outward-going signals and the one-way nature of the event horizon once it forms—all have parallels in the case of stellar black holes.

### PROCESS OF SCIENCE Check

- ✓ How do Newton’s and Einstein’s theories differ in their descriptions of gravity?

## 22.7 Space Travel Near Black Holes


Black holes are *not* cosmic vacuum cleaners. They don’t cruise around interstellar space, sucking up everything in sight. The orbit of an object near a black hole is basically the same as its orbit near a star of the same mass. Only if the object happens to pass within a few Schwarzschild radii (perhaps 50 or 100 km for a typical 5- to 10-solar-mass black hole formed in a supernova) of the event horizon is there any significant difference between its actual orbit and the one predicted by Newtonian gravity and described by Kepler’s laws. Of course, if some matter does happen to fall into a black hole—if the object’s orbit happens to take it too close to the event horizon—it will be unable to get out. Black holes are like turnstiles, permitting matter to flow in only one direction: inward.

Because a black hole will accrete at least a little material from its surroundings, its mass, and hence also the radius of its event horizon, tends to increase slowly over time.

### Tidal Forces

Matter flowing into a black hole is subject to great tidal stress. An unfortunate person falling feet first into a solar-mass black hole would find himself stretched enormously in height and squeezed unmercifully laterally. He would be torn apart even before he reached the event horizon, for the pull of gravity would be much stronger at his feet (which are closer to the hole) than at his head. The tidal forces at work in and near a black hole are the same basic phenomenon that is responsible for ocean tides on Earth and the spectacular volcanoes on Io. The only difference is that the tidal forces near a black hole are far stronger than any other force we know in the solar system.

As illustrated (with some artistic license) in Figure 22.19, a similar fate awaits any kind of matter falling into a black hole. Whatever falls in—gas, people, space probes—is vertically stretched and horizontally squeezed and accelerated to high speeds in the process. The net result of all this stretching



▲ **FIGURE 22.19 Black-Hole Heating** Any matter falling into the clutches of a black hole will become severely distorted and heated. This sketch shows an imaginary planet being pulled apart by a black hole's gravitational tides.

and squeezing is numerous, and violent collisions among the torn-up debris cause a great deal of frictional heating of the infalling matter. Material is simultaneously torn apart and heated to high temperatures as it plunges into the hole.

So efficient is the heating that, before reaching the hole's event horizon, matter falling into the hole emits radiation of its own accord. For a black hole of mass comparable to the Sun, the energy is expected to be emitted in the form of X-rays. In effect, the gravitational energy of matter outside the black hole is converted into heat as that matter falls toward the hole. Thus, contrary to what we might expect from an object whose defining property is that nothing can escape from it, the region surrounding a black hole is expected to be a *source* of energy. Of course, once the hot matter falls below the event horizon, its radiation is no longer detectable—it never leaves the hole.

## Approaching the Event Horizon

One safe way to study a black hole would be to go into orbit around it well beyond the disruptive influence of the hole's strong tidal forces. After all, Earth and the other planets of our solar system orbit the Sun without falling into it and without being torn apart. The gravity field around a black hole is basically no different; however, even from a stable circular orbit, a close investigation of the hole would be unsafe for humans. Endurance tests conducted on astronauts of the United States and the former Soviet Union indicate that the human body


cannot withstand stress greater than about 10–20 times the pull of gravity on Earth's surface. This breaking point would occur about 3000 km from a 10-solar-mass black hole (which, recall, would have a 30-km event horizon). Closer than that, the tidal effect of the hole would tear a human body apart.

Let's instead send an imaginary indestructible astronaut—a mechanical robot, say—in a probe toward the center of the hole. Watching from a safe distance in our orbiting spacecraft, we can then examine the nature of space and time near the hole. Our robot will be a useful explorer of theoretical ideas, at least down to the event horizon. After that boundary is crossed, there is no way for the robot to return any information about its findings.

Suppose, for example, our robot has an accurate clock and a light source of known frequency mounted on it. From our safe vantage point far outside the event horizon, we could use telescopes to read the clock and measure the frequency of the light we receive. What might we discover? We would find that the light from the robot would become more and more redshifted as the robot neared the event horizon. Even if the robot used rocket engines to remain motionless, a redshift would still be detected. The redshift is not caused by motion of the light source, nor is it the result of the Doppler effect arising as the robot falls into the hole. Rather, it is a redshift induced by the black hole's gravitational field, predicted by Einstein's general theory of relativity and known as **gravitational redshift**.

We can explain gravitational redshift as follows: According to general relativity, photons are attracted by gravity. As a result, in order to escape from a source of gravity, photons must expend some energy. They have to do work to get out of the gravitational field. They don't slow down at all—photons always move at the speed of light—they just lose energy. Because a photon's energy is proportional to the frequency of its radiation, light that loses energy must have its frequency reduced (or, equivalently, its wavelength lengthened). In other words, as illustrated in Figure 22.20, radiation coming from the vicinity of a massive object will be redshifted to a degree depending on the strength of the object's gravitational field.

As photons traveled from the robot's light source to the orbiting spacecraft, they would become gravitationally redshifted. From our standpoint on the orbiting spacecraft, a green light, say, would become yellow and then red as the robot astronaut neared the black hole. From the robot's perspective, the light would remain green. As the robot got closer to the event horizon, the radiation from its light source would become undetectable with optical telescopes. The radiation reaching us in the orbiting spacecraft would by then be lengthened so much that infrared and then radio telescopes would be needed to detect it. When the robot probe got closer still to the event horizon, the radiation it emitted as visible light would be shifted to wavelengths even longer than conventional radio waves by the time it reached us.



**Interactive** **FIGURE 22.20 Gravitational Redshift** Photons escaping from the strong gravitational field close to a black hole must expend energy to overcome the hole's gravity. As a result, the photons change wavelength; their color changes, and their frequency lessens. This figure shows the effect on two beams of radiation, one of visible light and one of X-rays, emitted from a space probe as it nears the event horizon of a 1-solar-mass black hole.

Light emitted from the event horizon itself would be gravitationally redshifted to infinitely long wavelengths. In other words, each photon would use all its energy trying to escape from the edge of the hole. What was once light (on the robot) would have no energy left upon its arrival at the safely orbiting spacecraft. Theoretically, this radiation would reach us—still moving at the speed of light—but with zero energy. Thus, the light radiation originally emitted would be redshifted beyond our perception.

Now, what about the robot's clock? Assuming that we could read it, what time would it tell? Would there be any observable change in the rate at which the clock ticked as it moved deeper into the hole's gravitational field? From the safely orbiting spacecraft, we would find that any clock close to the hole would appear to tick more *slowly* than an equivalent clock on board the spacecraft. The closer the clock came to the hole, the slower it would appear to run. On reaching the event horizon, the clock would seem to stop altogether. It would be as if the robot astronaut had found immortality! All action would become virtually frozen in time. Consequently, an external observer would never actually witness an infalling astronaut sink below the event horizon. Such a process would appear to take forever.

This apparent slowing down of the robot's clock is known as **time dilation**. It is another clear prediction of

general relativity and in fact is closely related to the gravitational redshift. To see this connection, imagine that we use our light source as a clock, with the passage of (say) a wave crest constituting a "tick." The clock thus ticks at the frequency of the radiation. As the wave is redshifted, the frequency drops, and fewer wave crests pass the distant observer each second—the clock appears to slow down. This thought experiment demonstrates that the redshift of the radiation and the slowing of the clock are one and the same.

From the point of view of the indestructible robot, however, relativity theory predicts no strange effects at all. To the infalling robot, the light source hasn't reddened, and the clock keeps perfect time. In the robot's frame of reference, everything is normal. Nothing prohibits it from coming within the Schwarzschild radius of the hole. No law of physics constrains an object from passing through an event horizon. There is no barrier at the event horizon and no sudden lurch as it is crossed; it is only an imaginary boundary in space. Travelers passing through the event horizon of a sufficiently massive hole (such as might lurk in the heart of our own Galaxy, as we will see) might not even know it—at least until they tried to get out!

The gravitational fields of most astronomical objects are far too weak to produce any significant gravitational redshift, although in many cases the effect can still be measured. Delicate laboratory experiments on Earth and on satellites in near-Earth orbits have succeeded in detecting the tiny gravitational redshift produced by even our own planet's weak gravity. Sunlight is redshifted by only about a thousandth of a nanometer. A few white-dwarf stars do show some significant gravitational reddening of their emitted light, however. Their radii are much smaller than that of our Sun, so their surface gravity is very much stronger than the Sun's. Neutron stars should show a substantial shift in their radiation, but it is difficult to disentangle the effects of gravity, magnetism, and environment on the signals we observe.

## Deep Down Inside

No doubt you are wondering what lies within the event horizon of a black hole. The answer is simple: No one really knows. However, the question is of great interest to theorists, as it raises some fundamental issues that lie at the forefront of modern physics.

Can an entire star simply shrink to a point and vanish? General relativity predicts that, without some agent to compete with gravity, the core remnant of a high-mass star will collapse all the way to a point at which both its density and its gravitational field become infinite. Such a point is called a **singularity**. We should not take this prediction of infinite density too literally, however. Singularities are not physical—rather, they always signal the breakdown of the theory producing them. In other words, the present laws of physics

are simply inadequate to describe the final moments of a star's collapse.

As it stands today, the theory of gravity is incomplete because it does not incorporate a proper (i.e., a quantum mechanical) description of matter on very small scales. As our collapsing stellar core shrinks to smaller and smaller radii, we eventually lose our ability even to describe, let alone predict, its behavior. Perhaps matter trapped in a black hole never actually reaches a singularity. Perhaps it just approaches this bizarre state in a manner that we will someday understand as the subject of *quantum gravity*—the merger of general relativity with quantum mechanics—develops.

Having said that, we can at least estimate how small the core can get before current theory fails. It turns out that by the time that stage is reached, the core is already much smaller than any elementary particle. Thus, although a complete description of the end point of stellar collapse may well require a major overhaul of the laws of physics, for all practical purposes the prediction of collapse to a point is valid. Even if a new theory somehow succeeds in doing away with the central singularity, it is unlikely that the external appearance of the hole or the existence of its event horizon will change. Any modifications to general relativity are expected to occur only on submicroscopic scales, not on the macroscopic (kilometer-sized) scale of the Schwarzschild radius.

Singularities are places where the rules break down, and some very strange things may occur near them. Many possibilities have been envisaged—gateways into other universes, time travel, the creation of new states of matter—but none of them has been proved, and certainly none of them has ever been observed. Because these regions are places where science fails, their presence causes serious problems for many of our cherished laws of physics, from causality (the idea that cause should precede effect, which runs into immediate and severe problems if time travel is possible) to energy conservation (which is violated if material can hop from one universe to another through a black hole). It is currently unclear whether the removal of the central singularity by some future all-encompassing theory would necessarily also eliminate all of these problematic side effects.

Disturbed by the possibility of such chaos in science, some researchers have even proposed a “principle of cosmic censorship”: Nature always hides *any* singularity, such as that found at the center of a black hole, inside an event horizon. In that case, even though physics fails, its breakdown cannot affect us outside, so we are safely insulated from any effects the singularity may have. What would happen if we one day found a so-called *naked singularity* somewhere—a singularity uncloaked by an event horizon? Would relativity theory still hold there? For now, we just don't know.

What sense are we to make of black holes? Do black holes and all the strange phenomena that occur in and around them really exist? The basis for understanding these weird objects is the relativistic concept that mass warps spacetime—which

has already been found to be a good representation of reality, at least for the weak gravitational fields produced by stars and planets (see *More Precisely 22-1* and *Discovery 22-2*). The larger the concentration of mass, the greater is the spacetime warping and, apparently, the stranger are the observational consequences. These consequences are part and parcel of general relativity, and black holes are one of its most striking predictions. As long as general relativity stands as the correct theory of gravity in the universe, black holes are real.

### CONCEPT Check

- ✓ Why would you never actually witness an infalling object crossing the event horizon of a black hole?

## 22.8 Observational Evidence for Black Holes

Theoretical ideas aside, is there any observational evidence for black holes? Can we prove that these strange invisible objects really do exist?


### Stellar Transits?

One way in which we might think we would detect a black hole is if we observed it transiting (passing in front of) a star. Unfortunately, such an event would be extremely hard to see. The approximately 12,000-km-diameter planet Venus is barely noticeable when it transits the Sun, so a 10-km-wide object moving across the image of a faraway star would be completely invisible with either current equipment or any equipment available in the foreseeable future.

Actually, we are even worse off than the previous paragraph suggests. Suppose we were close enough to the star to resolve the disk of the transiting black hole. Then the observable effect would not be a black dot superimposed on a bright background; instead, the background starlight would be deflected as it passed the black hole on its way to Earth, as indicated in Figure 22.21. The effect is the same as the bending of distant starlight around the edge of the Sun, a phenomenon that has been repeatedly measured during solar eclipses throughout the last several decades (see *More Precisely 22-1*). With a black hole, much larger deflections would occur. As a result, the image of a black hole in front of a bright companion star would show not a neat, well-defined black dot, but rather a fuzzy image virtually impossible to resolve, even from nearby.

### Black Holes in Binary Systems

A much better way to find black holes is to look for their effects on other objects. Our Galaxy harbors many binary-star systems in which only one object can be seen. Recall



**▲ FIGURE 22.21 Gravitational Light Deflection** The gravitational bending of light around the edges of a small, massive black hole makes it impossible to observe the hole as a black dot superimposed against the bright background of its stellar companion.

from our study of binary-star systems in Section 17.7 that we need to observe the motion of only one star to infer the existence of an unseen companion and measure some of its properties. In the majority of cases, the invisible companion is simply small and dim, nothing more than an M-type star hidden in the glare of an O- or B-type partner or perhaps shrouded by dust or other debris, making it invisible to even the best available equipment. In either case, the invisible object is *not* a black hole.


A few close binary systems, however, have peculiarities suggesting that one of their members may be a black hole. Some of the most interesting observations, made during the 1970s and 1980s by Earth-orbiting satellites, revealed binary systems in which the invisible member emits large amounts of X-rays. The mass of the emitting object is measured as several solar masses, so we know that it is not simply a small, dim star. Nor is it likely that visible radiation from the X-ray source is obscured by dusty circumstellar debris—in the cases of interest, intense radiation from the

**► FIGURE 22.22 Cygnus X-1** (a) The brightest star in this photograph (marked with its catalog number) is a member of a binary system whose unseen companion, called Cygnus X-1, is a leading black hole candidate. (b) An X-ray image of the field of view outlined by the rectangle in part (a). Since X-rays cannot be seen directly, those emitted by Cygnus X-1 were captured in space by a detector aboard a satellite, changed into radio signals for transmission to the ground, and changed again into electronic signals that were then viewed on a video screen, from which this picture was taken. (*Harvard-Smithsonian Center for Astrophysics*)

binary components would have dispersed the debris into interstellar space long ago.

One particular binary system drawing much attention lies in the constellation Cygnus. Figure 22.22(a) shows the area of the sky in Cygnus where astronomers have reasonably good evidence for a black hole. The rectangle outlines the celestial system of interest, some 6200 light-years from Earth. The black-hole candidate is an X-ray source called Cygnus X-1, studied in detail by the *Uhuru* satellite in the early 1970s. The main observational features of this binary system are as follows:

1. The visible companion of the X-ray source—a blue B-type supergiant with the catalog name HDE 226868—was identified a few years after Cygnus X-1 was discovered. Assuming that the companion lies on the main sequence, we know that its mass must be around 25 times the mass of the Sun.
2. Spectroscopic observations indicate that the binary system has an orbital period of 5.6 days. Combining this information with further spectroscopic measurements of the visible component's orbital speed, astronomers estimate the total mass of the system to be around



## MORE PRECISELY 22-1

### Tests of General Relativity

Special relativity is the most thoroughly tested and most accurately verified theory in the history of science. General relativity, however, is on somewhat less-firm experimental ground.


The problem with verifying general relativity is that its effects on Earth and in the solar system—the places where we can most easily perform tests—are very small. Just as special relativity produces major departures from Newtonian mechanics only when velocities approach the speed of light, general relativity predicts large departures from Newtonian gravity only when extremely *strong* gravitational fields are involved—when orbital speeds and escape velocities become relativistic.

We will encounter other experimental and observational tests of general relativity elsewhere in this chapter. (See *Discovery 22-2*.) Here, we consider just two “classical” tests of the theory—the deflection of light by the Sun and the effect of relativity on the orbit of Mercury. These tests are solar system observations that helped ensure the acceptance of Einstein’s theory. Later, more accurate measurements confirmed and strengthened the test results. Bear in mind, however, that there are currently no tests of general relativity in the “strong-field” regime—that part of the theory that predicts black holes, for example—so the full theory has never been experimentally tested. Scientists hope that the experiments described in *Discovery 22-2* will be able to test that part of the theory.

At the heart of general relativity is the premise that

everything, including light, is affected by gravity because of the curvature of spacetime. Shortly after he published his theory in 1915, Einstein noted that light from a star should be deflected by a measurable amount as it passes the Sun. According to the theory, the closer to the Sun the light comes, the more it is deflected. Thus, the maximum deflection should occur for a ray that just grazes the solar surface. Einstein calculated that the deflection angle should be  $1.75''$ —a small, but detectable, amount. Of course, it is normally impossible to see stars close to the Sun. However, during a solar eclipse, when the Moon blocks the Sun’s light, the observation becomes possible, as illustrated in the first (highly exaggerated) figure.

In 1919, a team of observers led by the British astronomer Sir Arthur Eddington succeeded in measuring the deflection of starlight during an eclipse. The results were in excellent agreement with the prediction of general relativity. Virtually overnight Einstein became world famous. His previous major accomplishments notwithstanding, this single prediction assured him a permanent position as the best-known scientist



35 solar masses, implying that Cygnus X-1 has a mass about 10 times the mass of the Sun. ∞ (Sec. 17.7)

3. Other detailed studies of Doppler-shifted spectral lines suggest that hot gas is flowing from the bright star toward an unseen companion. ∞ (Sec. 4.5)
4. X-ray radiation emitted from the immediate neighborhood of Cygnus X-1 implies the presence of very high temperature gas, perhaps as hot as several million kelvins (see Figure 22.22b).
5. Rapid time variations of this X-ray radiation imply that the size of the X-ray-emitting region of Cygnus X-1 must be extremely small—in fact, less than a few hundred kilometers across. The reasoning is basically the same as in the discussion of gamma-ray bursts in Section 22.4: X-rays from Cygnus X-1 have been observed to vary in intensity on time scales as short as a millisecond. For

this variation not to be blurred by the travel time of light across the source, Cygnus X-1 cannot be more than 1 light-millisecond, or 300 km, in diameter.

These properties suggest that the invisible X-ray-emitting companion could be a black hole. The X-ray-emitting region is likely an accretion disk formed as matter drawn from the visible star spirals down onto the unseen component. The rapid variability of the X-ray emission indicates that the unseen component must be compact—a neutron star or a black hole. The mass limit of the dark component argues for the latter, for a neutron star’s mass cannot exceed about 3 solar masses. Figure 22.23 is an artist’s conception of this intriguing object. Note that most of the gas drawn from the visible star ends up in a doughnut-shaped accretion disk of matter. As the gas flows toward the black hole, it becomes superheated and emits

on Earth! The high-precision *Hipparcos* satellite observed shifts in the apparent positions of many stars, even those whose line of sight is far from the Sun. ∞ (Sec. 17.1) The shifts are exactly as predicted by Einstein's theory.

**EXAMPLE** According to general relativity, a beam of light passing an object of mass  $M$  at distance  $R$  is deflected through an angle (in radians) of  $4GM/Rc^2$ , where  $G = 6.67 \times 10^{-11} \text{ N m}^2/\text{kg}^2$  is the gravitational constant and  $c = 3.00 \times 10^8 \text{ m/s}$  is the speed of light. Putting in the numbers for the Sun, we obtain  $M = 1.99 \times 10^{30} \text{ kg}$  and  $R = 696,000 \text{ km}$ , and remembering that 1 radian =  $57.3^\circ$ ; we can calculate the deflection to be  $(4 \times 6.67 \times 10^{-11} \times 1.99 \times 10^{30}) / (6.96 \times 10^8 \times [3.00 \times 10^8]^2) \times 57.3$  (degrees per radian)  $\times 3600$  (arc seconds per degree) =  $1.75''$ , as previously stated. ∞ (More Precisely 1-2) In more convenient units, we can write



$$\text{deflection (arc seconds)} = 1.75 \frac{M \text{ (solar masses)}}{R \text{ (solar radii)}}.$$

Note that the deflection is proportional to the mass  $M$  and inversely proportional to the distance  $R$ . Thus, Earth, with mass  $M = 3.0 \times 10^{-6}$  and radius  $R = 9.2 \times 10^{-3}$  solar units, would produce a deflection of just 0.57 milli-arc second (thousandths of an arc second), whereas a white dwarf such as Sirius B, with  $M = 1.1$  and  $R = 0.0073$  in the same units would deflect the beam by 4.4 arc minutes. ∞ (Sec. 20.3) (Neutron stars and black holes produce even greater effects, but the preceding simple formula is valid only when the deflection is small—less than a few degrees.)

The second prediction of general relativity testable within the solar system is that planetary orbits deviate slightly from the perfect ellipses of Kepler's laws. Again, the effect is greatest where

gravity is strongest—that is, closest to the Sun. Thus, the largest relativistic effects are found in the orbit of Mercury. Relativity predicts that Mercury's orbit is not a closed ellipse. Instead, its orbit should rotate slowly, as shown in the second (again exaggerated) diagram. The amount of rotation is very small—only  $43''$  per century—but Mercury's orbit is so well charted that even this tiny effect is measurable.

In fact, the observed rotation rate is  $540''$  per century, much greater than that predicted by relativity. However, when other (nonrelativistic) gravitational influences—primarily the perturbations due to the other planets—are taken into account, the rotation is in complete agreement with the foregoing prediction.



### Interactive FIGURE 22.23 Stellar Black Hole

MA Artist's conception of a binary system containing a large, bright, visible star and an invisible, X-ray-emitting black hole. (Compare with Figure 21.2.) This painting is based on data obtained from detailed observations of Cygnus X-1. (L. Chaisson)

the X-rays we observe just before they are trapped forever below the event horizon.

A few other black hole candidates are known. For example, the third X-ray source discovered in the Large Magellanic Cloud—LMC X-3—is an invisible object that, like Cygnus X-1, orbits a bright companion star. Reasoning similar to that applied to Cygnus X-1 suggests that LMC X-3 has 10 times more mass than the Sun—too massive to be anything but a black hole. Similarly, the X-ray binary



## DISCOVERY 22-2

## Gravity Waves: A New Window on the Universe


Electromagnetic waves are common everyday phenomena. They involve periodic changes in the strengths of electric and magnetic fields. [∞ \(Sec. 3.2\)](#) Electromagnetic waves move through space and transport energy. Any accelerating charged particle, such as an electron in a broadcasting antenna or on the surface of a star, generates electromagnetic waves.

The modern theory of gravity—Einstein’s theory of relativity—also predicts waves that move through space. A *gravity wave* is the gravitational counterpart of an electromagnetic wave. *Gravitational radiation* results from changes in the strength of a gravitational field. Any time an object of any mass accelerates, a gravity wave should be emitted at the speed of light. The wave should produce small distortions in the space through which it passes. Gravity is an exceedingly weak force compared with electromagnetism, so these distortions are expected to be very small—in fact, much smaller than the diameter of an atomic nucleus. [∞ \(More Precisely 16-1\)](#) Yet many researchers think that these tiny distortions are measurable.

The objects most likely to produce gravity waves detectable on Earth are close binary systems containing black holes, neutron stars, or white dwarfs. As these massive components orbit one another, their acceleration results in rapidly changing gravitational fields and the emission of gravitational radiation. As energy escapes in the form of gravity waves, the two objects spiral toward one another, orbiting more rapidly and emitting even more gravitational radiation. As we saw in Section 22.4, neutron-star mergers may well also be the origin of some gamma-ray bursts, so gravitational radiation might provide an alternative means of studying these violent and mysterious phenomena.

Such a slow but steady decay in the orbit of a binary system has in fact been detected. In 1974, radio astronomer Joseph Taylor and his student Russell Hulse at the University of Massachusetts discovered an unusual binary system. Both components are neutron stars, and one is observable from Earth as a pulsar. This system has become known as the *binary pulsar*. Measurements of the periodic Doppler shift of the pulsar’s radiation prove that its orbit is shrinking at exactly the rate predicted by relativity theory if the energy were being carried off by gravity waves. The two neutron stars should merge in an energetic burst of gravitational radiation and gamma rays in less than 300 million years (although most of the radiation will be emitted during the last few seconds). Even though the waves themselves have not been detected, the binary pulsar is regarded by most astronomers as very strong evidence in favor of general relativity. Taylor and Hulse received the 1993 Nobel Prize in physics for their discovery. The first accompanying figure illustrates the scale of, and predicted orbital changes in, the binary pulsar’s orbit.


In 2004, radio astronomers announced the discovery of a *double-pulsar* binary system with an even shorter period than the binary pulsar, implying stronger relativistic effects and a shorter merger time—about 85 million years. Because both components are pulsars, and since the system, by pure luck, also happens to be seen almost exactly edge on by observers on Earth, leading to



eclipses, this system has provided a wealth of detailed information on both neutron stars and gravitational physics.

The second figure shows part of an ambitious gravity-wave observatory called *LIGO*—short for *Laser Interferometric Gravity-Wave Observatory*—which became operational in 2003. Twin detectors, one (shown here) in Hanford, Washington, the other in Livingston, Louisiana, use interference between two laser beams to measure the tiny distortions of space produced by gravitational radiation in the lengths of the 4-km-long arms should a gravity wave pass by. [∞ \(Discovery 3-1\)](#) The instrument is in theory capable of detecting gravity waves from many galactic and extragalactic sources, although, so far, no gravity waves have actually been detected, despite a 2007 upgrade that greatly increased the system’s sensitivity. Additional upgrades boosting the detector’s sensitivity by a further factor of 10 are planned to come on line around 2014.

If these experiments are successful, the discovery of gravity waves could herald a new age in astronomy, in much the same way that invisible electromagnetic waves, virtually unexplored a century ago, revolutionized classical astronomy and led to the field of modern astrophysics.




(NSF)

system A0620–00 contains an invisible compact object of mass 3.8 times that of the Sun. In total, perhaps two dozen known objects in or near our Galaxy may be black holes. Cygnus X-1, LMC X-3, and A0620–00 have the strongest claims.

## Black Holes in Galaxies

Perhaps the strongest evidence for black holes comes not from binary systems in our own Galaxy, but from observations of the *centers* of many galaxies, including our own. Using high-resolution observations at wavelengths ranging from radio to ultraviolet, astronomers have found that stars and gas near the centers of many galaxies are moving extremely rapidly, orbiting some very massive, unseen object. Masses inferred from Newton’s laws range from millions to billions of times the mass of the Sun. [∞ \(More Precisely 2-2\)](#)

The intense energy emission from the centers of these galaxies and the short-timescale fluctuations in that emission suggest the presence of massive, compact objects. In addition, as in the *radio galaxy* shown in Figure 22.24, these objects are also observed to have extended *jets* reminiscent of—but vastly larger than—those associated with neutron stars and black holes. The leading (and at present, the only) explanation is that these energetic objects are powered by huge central **supermassive black holes** accreting stars and gas from their



**▲ FIGURE 22.24 Active Galaxy** Many galaxies are thought to harbor massive black holes at their centers. Indeed, the best evidence for black holes anywhere in the universe is found in galactic nuclei. Shown here in false color is the galaxy 3C296. Blue color shows the distribution of stars in the central elliptical galaxy; red shows huge jets of radio emission extending 500,000 light-years across. (See also the opening image of Chapter 24.) (NRAO)


surroundings. The radio-emitting jets are rooted in the parsec-sized accretion disks surrounding the black holes themselves. We will return to the observations, and the question of how these black holes might have formed, in Chapters 23–24.

Thus, astronomers know of “stellar-mass” black holes, comparable in mass to the Sun, and supermassive black holes of millions or billions of solar masses. The former are the result of stellar evolution, as discussed in the last few chapters; the latter have grown in the centers of galaxies, as we will see in Part 4 of the text. Is there anything in between? In 2000, X-ray astronomers reported the first evidence for the long-sought, but elusive, missing link between these two classes of black holes. Figure 22.25 shows an unusual-looking galaxy called M82, currently the site of an intense and widespread burst of star formation (see Chapter 24); the red plumes are winds of hot gas escaping from numerous star-forming sites in the otherwise quiescent part of the galaxy (shown in blue). The inset shows a *Chandra* image of the innermost few thousand parsecs of M82, revealing a number of bright X-ray sources close to—but *not* at—the center of the galaxy. Their spectra and X-ray luminosities suggest that some may be accreting compact objects with masses ranging from 100 to almost 1000 times the mass of the Sun. If confirmed, they will be the first *intermediate-mass black holes* ever observed.

Too large to be remnants of normal stars and too small to warrant the “supermassive” label, these objects present a puzzle to astronomers. Where did they come from? One possible origin is suggested by follow-up infrared observations from the Subaru and Keck telescopes on Mauna Kea, indicating that some of the X-ray sources are apparently associated with dense, young star clusters. [∞ \(Secs. 5.2, 19.6\)](#) Theorists speculate that collisions between high-mass main-sequence stars in the congested cores of such clusters could lead to the runaway growth of extremely massive and highly unstable stars, which could then collapse to form intermediate-mass black holes. Figure 22.26 shows the current best candidate for a “nearby” star cluster harboring an intermediate-mass black hole—the globular cluster G1 in the Andromeda Galaxy. [∞ \(Sec. 3.1\)](#) The peculiar orbits of stars near the center of this cluster suggest a black hole of mass 20,000 times that of the Sun, and observations of the cluster in both radio and X-rays are consistent with theoretical expectations of the emission from such a massive object in the cluster’s core. However, both the theory and this observation remain controversial.

## Do Black Holes Exist?

You may have noticed that the identification of an object as a black hole really proceeds by elimination. Loosely stated, the argument goes as follows: “Object X is compact and very massive. We don’t know of anything else that can be that small and that massive. Therefore, object X is a black hole.” For the very massive compact objects observed (or inferred to be) in the centers of galaxies, the absence of viable alternatives



◀ **FIGURE 22.25 Intermediary-Mass Black Holes?** X-ray observations (inset below) of the center of the starburst galaxy M82 (at top, about 100,000 light-years across and 12 million light-years away) reveal a collection of bright sources thought to be the result of matter accreting onto intermediary-mass black holes. The black holes are probably young, have masses between 100 and 1000 times the mass of the Sun, and lie relatively far (about 600 light-years) from the center of M82. The brightest (and possibly most massive) black hole candidate is marked by an arrow. (*Subaru; NASA*)

means that the black-hole hypothesis has become widely accepted among astronomers.

However, Cygnus X-1 and the other suspected stellar-mass black holes in binary systems all have masses relatively close to the dividing line separating neutron stars from black holes. Given the present uncertainties in both observation and theory, might they conceivably be merely dim, dense neutron stars and not black holes at all?


Most astronomers do not think this likely but it highlights a problem. It is difficult to unambiguously distinguish a 10-solar-mass black hole from, say, a 10-solar-mass neutron star (if one could somehow exist). Both objects would affect a companion star's orbit in the same way; both would tear mass from its surface, and both would form an accretion disk around themselves that would emit intense X-rays (although many researchers think that the accretion disks may differ sufficiently in detail that the nature of the central object might be identifiable from observations).

We have stressed throughout this text that scientific theories unsupported by observational or experimental evidence are destined not to survive. ∞ (Sec. 1.2) Black holes are a clear prediction of Einstein's general theory of relativity, which is widely regarded as the correct description of gravity in the presence of strong fields and orbital speeds comparable to the speed of light. But we have also seen that general relativity has been tested most thoroughly in situations where gravity is weak and velocities are relatively low, and not at all under the extreme conditions expected near a black hole. So we can legitimately ask, "Do we have any unambiguous evidence that the massive, compact objects just described really are black holes?"

The short answer—at least, if measurements of mass and size alone are insufficient to convince you of a black hole's reality—is no. Detailed measurements of black-hole properties are hard to make and even harder to interpret. Black holes tend to live in very messy astrophysical environments. Astronomers

have uncovered tantalizing hints of black-hole event horizons (rather than hard neutron star surfaces) in several systems, but none have as yet proved conclusive. However, as technology continues to improve, we can expect many more such observations, with increasing precision, as astronomers test a key prediction of Einstein's theory.

So have black holes really been discovered? Despite the uncertainties, the answer is probably yes. Skepticism is healthy in science, but only the most stubborn astronomers (and some do exist!) would take serious issue with the many lines of theoretical reasoning that support the case for black holes. The crucial role played by black holes in the theories of stellar evolution, gamma-ray bursts, and (as we will see in Chapters 24 and 25) the structure and evolution of galaxies is a clear indication of their widespread acceptance in astronomy.



▲ **FIGURE 22.26 Black-Hole Host?** Astronomers have found that stars near the center of the massive globular cluster G1 do not move as expected if the cluster's mass is as smoothly distributed as its light. Instead, the observations suggest that an intermediary-mass black hole resides at the cluster's center. (*NASA*)

Can we guarantee that future modifications to the theory of compact objects will not invalidate some or all of our arguments? No, but similar statements could be made in many areas of astronomy—indeed, about any theory in any area of science. We conclude that, strange as they are, black holes have been detected, both in our Galaxy and beyond. Perhaps someday future generations of space travelers will visit Cygnus X-1 or the center of our Galaxy and (carefully!) test these conclusions firsthand.

Until then, we will have to continue to rely on improving theoretical models and observational techniques to guide our discussions of the mysterious objects known as black holes.

### PROCESS OF SCIENCE Check


✓ How do astronomers “see” black holes?

**The Big Question** Are black holes for real? Decades ago, many astronomers regarded black holes as a kind of cosmic cop-out—the last resort of researchers who couldn’t decipher the bizarre phenomena they observed on the sky. But now, improved observations really do point toward stellar remnants that are indeed denser, brighter, and much more puzzling than the already peculiar neutron stars. Still, might there be other weird collapsed remnants—quark stars, perhaps—that stop short of being genuine black holes?


# Chapter Review

## SUMMARY


1 A core-collapse supernova may leave behind a **remnant** (p. 544)—an ultracompressed ball of material called a **neutron star** (p. 544). Neutron stars are extremely dense and, at formation, are predicted to be very hot, strongly magnetized, and rapidly rotating. They cool down, lose much of their magnetism, and slow down as they age.



2 According to the **lighthouse model** (p. 545), neutron stars, because they are magnetized and rotating, send regular bursts of electromagnetic energy into space. The beams are produced by charged particles confined by the strong magnetic fields. When we can see the beams from Earth, we call the source a **pulsar** (p. 545). The pulse period is the rotation period of the neutron star. Because the pulse energy is beamed into space and because neutron stars slow down as they radiate energy into space, not all neutron stars are seen as pulsars.




3 A neutron star in a close binary system can draw matter from its companion, forming an accretion disk. The material in the disk heats up before it reaches the




neutron star, making the disk a strong source of X-rays. As gas builds up on the star’s surface, the star eventually becomes hot enough to fuse hydrogen. When hydrogen burning starts, it does so explosively, and an **X-ray burster** (p. 548) results. The rapid rotation of the inner part of the accretion disk causes the neutron

star to spin faster as new gas arrives on its surface. The eventual result is a very rapidly rotating neutron star—a **millisecond pulsar** (p. 550). Many millisecond pulsars are found in the hearts of old globular clusters. They cannot have formed recently, so they must have been spun up by interactions with other stars. Analysis of the radiation received shows that some pulsars are orbited by planet-sized objects.


4 **Gamma-ray bursts** (p. 552) are very energetic flashes of gamma rays observed about once per day, distributed uniformly over the entire sky. In some cases, their distances have been measured, placing them far away from us and implying that they are extremely luminous. The leading theoretical models for these explosions postulate the violent merger of neutron stars in a distant binary system or the recollapse and subsequent violent explosion following a “failed” supernova in a very massive star.



5 Einstein’s special theory of relativity deals with the behavior of particles moving at speeds comparable to the speed of light. It agrees with Newton’s theory at low velocities, but makes many very different predictions for high-speed motion. All of its predictions have been repeatedly verified by experiment. The modern replacement for Newtonian gravity is Einstein’s **general theory of relativity** (p. 558), which describes gravity in terms of the warping, or bending, of **spacetime** (p. 558) by the presence of mass. The more mass, the greater the warping. All particles—including photons—respond to that warping by moving along curved paths.




**6** The upper limit on the mass of a neutron star is about 3 solar masses. Beyond that mass, the star can no longer support itself against its own gravity, and it collapses to form a **black hole** (p. 556), a region of space from which nothing can escape.




Very massive stars, after exploding as supernovae, form black holes rather than neutron stars. Conditions in and near black holes can only be described by general relativity. The radius at which the escape speed from a collapsing star equals the speed of light is called the **Schwarzschild radius** (p. 557). The surface of an imaginary sphere, of radius equal to the Schwarzschild radius, surrounding a black hole is called the **event horizon** (p. 557).

**7** To a distant observer, light leaving a spaceship that is falling into a black hole would be subject to **gravitational redshift** (p. 562) as the light climbed out of the hole's intense gravitational field. At the same time, a clock on the spaceship would show **time dilation** (p. 563)—the



clock would appear to slow down as the ship approached the event horizon. The observer would never see the ship reach the surface of the hole. Once within the event horizon, no known force can prevent a collapsing star from contracting all the way to a point-like **singularity** (p. 563), at which point both the density and the gravitational field of the star become infinite. This prediction of relativity theory has yet to be proved. Singularities are places where the known laws of physics break down.

**8** Once matter falls into a black hole, it can no longer communicate with the outside. However, on its way in, it can form an accretion disk and emit X-rays, just as in the case of a neutron star. The best candidates for black holes are binary systems in which one component is a compact X-ray source. Cygnus X-1, a well-studied X-ray source in the constellation Cygnus, is a long-standing black-hole candidate. Studies of orbital motions imply that some binaries contain compact objects too massive to be neutron stars, leaving black holes as the only alternative. There is also strong evidence for **supermassive black holes** (p. 569) in the centers of many galaxies, including our own.



**MasteringAstronomy**® For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- LO1** How does the way in which a neutron star forms determine some of its most basic properties?
- What would happen to a person standing on the surface of a neutron star?
- LO2** What are pulsars, and how are they related to neutron stars? Why aren't all neutron stars seen as pulsars?
- What are X-ray bursters?
- LO3** What is the favored explanation for the rapid spin rates of millisecond pulsars?
- LO4** Why do astronomers think that gamma-ray bursts are very distant and very energetic?
- Describe two leading models for gamma-ray bursts.
- LO5** Use your knowledge of escape speed to explain why black holes are said to be "black."
- LO6** According to special relativity, what is special about the speed of light?
- POS** Why is it so difficult to test the predictions of general relativity? Describe two tests of the theory.
- LO7** What is an event horizon? What would happen to someone falling into a black hole as they approach the event horizon?
- POS** What is the principle of cosmic censorship? Do you think it is a sound scientific principle?
- What makes Cygnus X-1 a good black-hole candidate?
- LO8 POS** What evidence is there for black holes much more massive than the Sun?
- POS** Do you think that planet-size objects orbiting a pulsar should be called planets? Why or why not?

## Conceptual Self-Test: Multiple Choice

- A neutron star is about the same size as (a) a school bus; (b) a U.S. city; (c) the Moon; (d) Earth.
- A neutron star's immense gravitational attraction is due primarily to its small radius and (a) rapid rotation rate; (b) strong magnetic field; (c) large mass; (d) high temperature.
- The most rapidly "blinking" pulsars are those that (a) spin fastest; (b) are oldest; (c) are most massive; (d) are hottest.
- The X-ray emission from a neutron star in a binary system comes mainly from (a) the hot surface of the neutron star

itself; (b) heated material in an accretion disk around the neutron star; (c) the neutron star's magnetic field; (d) the surface of the companion star.

5. **VIS** According to Figure 22.11, gamma-ray bursts are observed to occur (a) mainly near the Sun; (b) throughout the Milky Way Galaxy; (c) approximately uniformly over the entire sky; (d) near pulsars.
6. Black holes result from stars having initial masses (a) less than the mass of the Sun; (b) between 1 and 2 times the mass of the Sun; (c) up to 8 times the mass of the Sun; (d) more than 25 times the mass of the Sun.
7. If the Sun were magically to turn into a black hole of the same mass, (a) Earth would start to spiral inward; (b) Earth's orbit would remain unchanged; (c) Earth would fly off into space; (d) Earth would be torn apart.
8. **VIS** According to the second figure in *Discovery 22-1*, a meterstick in a spaceship traveling at half the speed of light would appear to have a length of (a) 1 meter; (b) 0.87 meter; (c) 0.50 meter; (d) 0.15 meter.
9. The best place to search for black holes is in a region of space that (a) is dark and empty; (b) has recently lost some stars; (c) has strong X-ray emission; (d) is cooler than its surroundings.
10. The best evidence for supermassive black holes in the centers of galaxies is (a) the absence of stars there; (b) rapid gas motion and intense energy emission; (c) gravitational redshift of radiation emitted from near the center; (d) unknown visible and X-ray spectral lines.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

1. ● The angular momentum of a solid body is proportional to the angular velocity of the body times the square of its radius. ∞ (More Precisely 6-1) Using the law of conservation of angular momentum, estimate how fast a collapsed stellar core would spin if its initial spin rate was 1 revolution per day and its radius decreased from 10,000 km to 10 km.
2. ● What would your mass be if you were composed entirely of neutron-star material of density  $3 \times 10^{17} \text{ kg/m}^3$ ? (Assume that your average density is  $1000 \text{ kg/m}^3$ .) Compare your answer with the mass of a typical 10-km-diameter rocky asteroid.
3. ● Calculate the surface gravitational acceleration and escape speed of a 1.4-solar-mass neutron star with a radius of 10 km. What would be the escape speed from a neutron star of the same mass and radius 4 km?
4. ●● Use the radius–luminosity–temperature relation to calculate the luminosity of a 10-km-radius neutron star for temperatures of  $10^5 \text{ K}$ ,  $10^7 \text{ K}$ , and  $10^9 \text{ K}$ . At what wavelengths does the star radiate most strongly in each case? Could the brightest of them be plotted on an H–R diagram?
5. ●● A gamma-ray burst 5000 Mpc away releases  $10^{45}$  joules of energy isotropically in the form of gamma rays, each of energy 250 keV. ∞ (More Precisely 4-1) Some of the rays are detected by an instrument in Earth orbit with an effective collecting area of  $0.75 \text{ m}^2$ . How many gamma-ray photons strike the detector?
6. ● Supermassive black holes are thought to exist in the centers of many galaxies. What would be the Schwarzschild radii of black holes of 1 million and 1 billion solar masses, respectively? How does the 1-million-solar-mass black hole compare in size with the Sun? How does the 1-billion-solar-mass black hole compare in size with the solar system?
7. ●● Use the information presented in *More Precisely 22-1* to estimate the deflection of a beam of light that just grazes the surface of (a) the Moon, (b) Jupiter, and (c) Sirius B. (d) A future generation of space astrometry missions may be able to measure angles as small as  $10^{-6}$  arcsec accurately. At what distance from the Sun would this deflection occur?
8. ●● Calculate the tidal acceleration on a 2-m-tall human falling feet-first into a 1-solar-mass black hole; that is, compute the difference in the accelerations (forces per unit mass) on their head and their feet just as the feet cross the event horizon. ∞ (Sec. 7.6) Repeat the calculation for a 1-million-solar-mass black hole and for a 1-billion-solar-mass black hole (see question 6). Compare these accelerations with the acceleration due to gravity on Earth ( $g = 9.8 \text{ m/s}^2$ ).

## Activities


### Collaborative

1. The text focuses on the simplest kind of black hole—the uncharged, nonrotating *Schwarzschild* black hole—but rotating *Kerr* black holes are extremely important in astronomy. Divide your group in two and research online the properties of Schwarzschild and Kerr black holes. Combine your research to make a joint presentation on the similarities and differences between the two. Focus on properties like the event horizon, the singularity, and the orbits of light and matter near the hole. How fast can a black hole rotate? Which kind of black hole is thought to be most common in nature?

### Individual

1. Find the ninth magnitude companion to Cygnus X-1, the sky's most famous black hole candidate. Even without a telescope, it's easy to locate the region of the heavens where Cygnus X-1 resides. The constellation Cygnus contains a recognizable star pattern, or asterism, in the shape of a large cross. This asterism is called the Northern Cross. The star in the center of the crossbar is called Sadr. The star at the bottom of the cross is called Albireo. Approximately midway between Sadr and Albireo lies the star Eta Cygni. Cygnus X-1 is located slightly less than  $0.5^\circ$  from this star. With or without a telescope, sketch what you see.

Andromeda Galaxy as observed in 1890 (*J. Roberts*)



## PART FOUR

# Galaxies and Cosmology



Harlow Shapley working at his rotating octagonal desk (*Harvard*)


**It is hard to imagine**, but less than 100 years ago the Sun was considered the center of the universe. Earlier studies by Copernicus, Kepler, Galileo, and others had dethroned Earth from a central position, but the Sun itself, at the center of our solar system, was still assumed also to be the center of the Milky Way—which a century ago was identical to the universe. Our true place in the cosmos, and even the existence of countless other galaxies comparable to our own, were completely unknown.

Enter the American astronomer Harlow Shapley (1885–1972), who, by studying variable stars in globular clusters, was able to deduce the size and scale of the Milky Way Galaxy, as well as our position in it. His results, announced in 1918, showed not only that our extended home in space was immensely larger than had previously been realized—about 100,000 light-years across—but also that Earth resided in what he called the "galactic suburbs," now known to be about 25,000 light-years from the center of


the Galaxy. Shapley demonstrated that our Sun is not central, unique, or special in any way. His work was a milestone in our understanding of our place in the universe, certainly one of the most important astronomical discoveries of the 20th century.

Ironically, Shapley's dramatic discovery of the increased size and scale of the Milky Way led him astray regarding another, even more profound, advance in our knowledge at that time: the realization that our Galaxy is only one of many galaxies in the universe. The sheer size of the Milky Way implied by his observations caused him to oppose the idea of a vastly larger universe—he found it hard to believe that there could be other, distant galaxies as huge as our own. Even among eminent scientists, personal biases can sometimes cloud scientific judgment.

The stage was set for a "Great Debate" that occurred at the National Academy of Sciences in Washington in 1920. At issue were the fuzzy "spiral nebulae" (which today we call galaxies): Were they close enough to be part of our own Milky Way, or were they sufficiently distant to be whole galaxies unto themselves? Shapley held that, given that his research had revised upward the size of the Milky Way, the spiral nebulae must be part of our own Galaxy. His opponent, Heber Curtis of California's Lick Observatory, while incorrectly



Edwin Hubble's discovery of variable stars in Andromeda (*Carnegie Institute*)




Andromeda Galaxy as observed today (*R. Gendler*)

rejecting the great size of our Galaxy, correctly argued that the spirals were remote aggregates of stars similar to the Milky Way. Both men presented other scientific arguments supporting their views (see Section 23.2), but both also let personal feelings affect their comprehension of our home Galaxy. With no objective measurements of the true distances to the nebulae, the debate ended in a draw.

Shapley's rival, the Caltech astronomer Edwin Hubble (1884–1953), broke the stalemate just a few years later by using the premier optical telescope of the day, the 2.5-m (100-inch) reflector atop Mt. Wilson. He first resolved the Andromeda Nebula into individual stars and then carefully measured its variable stars, thereby proving that Andromeda was a genuine galaxy millions of light-years distant, well outside our Milky Way. Ironically, Hubble used the same basic technique that Shapley and his Harvard colleagues had pioneered. It was yet another milestone along the road extending the Copernican principle: Neither Earth nor the Sun is special in any way, and even the Galaxy in which we live is just one of myriad galaxies in a much, much larger cosmos.

**Today**, astronomers have extensively mapped the distribution of variable stars in Andromeda. Curiously, we are still struggling to determine that galaxy's distance to better than 10 percent accuracy. Even in the decades-long lifetime of this textbook, Andromeda's quoted distance has fluctuated from about 2.2 to as much as 2.9 million light-years; in this edition, we have averaged the most recent measurements to arrive at a value of 2.5 million light-years. The correct value is important, for upon it rests a key rung in the so-called distance ladder. This cosmic yardstick is used to measure the ranges to billions of other, more distant galaxies and hence to gauge the vastly larger realm of the universe itself.

The Shapley-Curtis debate of yesteryear, together with our current struggles to pin down accurate distances to the truly faraway galaxies, constitute good case studies of how the scientific method actually works. Science is practiced by human beings, and scientists are no different from others who have strong emotions and personal values. Yet, over the course of time, and through much criticism and debate, scientific issues eventually gain a measure of objectivity. By demanding tests and proven facts, the scientific community gradually damps the subjectivity of individuals and arrives at a more objective view among a community of critical thinkers. Reasoned skepticism and repeated testing are hallmarks of the modern scientific method.



Hubble Ultra Deep Field (*STScI*)





# The Milky Way Galaxy

## A SPIRAL IN SPACE

Looking up on a dark, clear night, we are struck by two aspects of the night sky. The first is that the individual stars we see are roughly uniformly distributed in all directions. They all lie relatively close to us, mapping out the local Galactic neighborhood within a few hundred parsecs of the Sun. But this is only a local impression. Ours is a rather provincial view. Beyond those nearby stars, the second thing we notice is a fuzzy band of light—the Milky Way—stretching across the heavens.

The perspective we have is that of an insider's view of the galaxy in which we live—the blended light of countless distant stars. As we consider much larger volumes of space, on scales far, far greater than the distances between neighboring stars, a new level of organization becomes apparent as the large-scale structure of the Milky Way Galaxy is revealed.

**The Big Picture** Our Milky Way Galaxy is just one among nearly a hundred billion other galaxies in the observable universe—a hundred billion galaxies! For astronomers, the Milky Way plays much the same role for galaxies as the Sun does for stars. Our understanding of galaxies throughout the cosmos rests squarely on our knowledge of the size, scale, structure, and dynamics of our own grand system of many varied stars.

**LEFT:** Galaxies like this one, known as NGC 1232 and shown in true color, contain roughly a hundred billion stars bound together by gravity. As we now enter the realm of big dimensions, the graceful winding arms of this majestic spiral galaxy sweep across some 100,000 light-years of space; the whole object is about 60 million light-years distant. Its size, shape, and mass approximate those of our own Galaxy, which has never been photographed in its full grandeur because we live inside it. If this were our Galaxy, the Sun would reside in one of its spiral arms, two-thirds of the way out from the center. (ESO)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Describe the overall structure of the Milky Way Galaxy, and say how the various regions differ from one another.
- 2 Explain the importance of variable stars for measuring the size and shape of our Galaxy.
- 3 Compare and contrast the orbital motions of stars in different regions of the Galaxy.
- 4 Interpret the differences between disk and halo stars in terms of our current understanding of how our Galaxy formed.
- 5 Present some possible explanations for the spiral arms observed in our own and many other galaxies.
- 6 Explain what studies of Galactic rotation reveal about the size and mass of our Galaxy, and discuss the possible nature of dark matter.
- 7 Describe the evidence for a supermassive black hole and some of the other phenomena observed at the center of our Galaxy.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.

## 23.1 Our Parent Galaxy

A **galaxy** is a gargantuan collection of stellar and interstellar matter—stars, gas, dust, neutron stars, black holes—isolated in space and held together by its own gravity. Astronomers are aware of literally billions of galaxies beyond our own. The particular galaxy we happen to inhabit is known as the **Milky Way Galaxy**, or just *the Galaxy*, with a capital *G*.

Our Sun lies in a part of the Galaxy known as the **Galactic disk**—an immense, circular, flattened region containing most of our Galaxy’s luminous stars and interstellar matter (and virtually everything we have studied so far in this book). Figure 23.1 illustrates how, viewed from within, the Galactic disk appears as a band of light stretching across our night sky, a band known as the *Milky Way*. As indicated in the figure, if we look in a direction away from the Galactic disk (red arrows), we see relatively few stars in our field of view. However, if our line of sight happens to lie within the disk (white and blue arrows), we see so many stars that their light merges into a continuous blur.


Paradoxically, although we can study individual stars and interstellar clouds that lie near the Sun in great detail, our location within the Galactic disk makes deciphering our Galaxy’s large-scale structure from Earth a very difficult task—a little like trying to unravel the layout of paths, bushes, and trees in a city park without being able to leave one particular park bench. In some directions, the interpretation

of what we see is ambiguous and inconclusive. In others, foreground objects completely obscure our view of what lies beyond, but we cannot move around them to get a better look. As a result, astronomers who study the Milky Way Galaxy are often guided in their efforts by comparisons with more distant, but much more easily observable, systems.


Figures 23.2 and 23.3 show three galaxies thought to resemble our own in overall structure. Figure 23.2 is the Andromeda Galaxy, the nearest major galaxy to the Milky Way Galaxy, lying nearly 800 kpc (roughly 2.5 million light-years) away. Andromeda’s apparent elongated shape is a consequence of the angle at which we happen to view it. In fact, our Galaxy, like this one, consists of a circular galactic disk of matter that fattens to a **Galactic bulge** at the center. The disk and bulge are embedded in a roughly spherical ball of faint old stars known as the **Galactic halo**. These three basic galactic regions are indicated on the figure. (The halo stars are so faint that they cannot be discerned here.) Figures 23.3(a) and (b) show views of two other galaxies—one seen face-on, the other edge-on—that illustrate these points more clearly.

### CONCEPT Check

- ✓ Why do we see the Milky Way as a band of light across the night sky?




(a)



(b)

### ◀ FIGURE 23.1 Galactic

**Plane** (a) Gazing from Earth toward the Galactic center (white arrow) in this artist’s conception, we see myriad stars stacked up within the thin band of light known as the Milky Way. In the opposite direction (blue arrow), we see little of our Galaxy, much as when looking perpendicular to the disk (red arrows), where far fewer stars exist. (b) This real optical view (from a very dark place on Earth) of the sky in the direction of the white arrow shows the fuzzy (mostly white and “milky”) band or disk of our Milky Way Galaxy (see also Figure 18.1). (A. Mellinger)




▲ **FIGURE 23.2 Andromeda Structure** (a) The Andromeda Galaxy closely resembles the overall layout of our own Milky Way. Its disk and bulge are clearly visible in this image, but the faint halo stars, completely surrounding the disk and bulge, cannot easily be seen here. The white stars sprinkled all across this image are foreground stars in our own Galaxy, lying in the same region of the sky as Andromeda, but about a thousand times closer. (b) More detail within the inner parts of this galaxy, including (c) its peculiar—and still unexplained—double core. (R. Gendler; Palomar/Caltech; NASA)



(a)




(b)



## 23.2 Measuring the Milky Way

Before the 20th century, astronomers' conception of the cosmos differed markedly from the modern view. The fact that we live in just one of many enormous "islands" of matter separated by even larger tracts of apparently empty space was completely unknown, and the clear distinction between "our Galaxy" and "the universe" did not exist. The twin ideas that (1) *the Sun is not at the center of the Galaxy* and (2) *the Galaxy is not at the center of the universe* required both time and hard observational evidence before they gained widespread acceptance. The growth in our knowledge of our Galaxy, as well as the realization that there are many other distant galaxies similar to our own, has gone hand in hand with the development of the cosmic distance scale.

◀ **FIGURE 23.3 Disk Galaxies** (a) This galaxy, called M101 and seen nearly face-on, is somewhat similar in its overall structure to our own Milky Way and Andromeda. (b) The galaxy NGC 4565 is oriented edge-on, allowing us to see clearly its disk and central bulge. (NASA)



◀ **FIGURE 23.4 Herschel's Galaxy Model**

Eighteenth-century astronomer William Herschel constructed this “map” of the Galaxy by counting the numbers of stars he saw in different directions of the sky. Our Sun (marked by the yellow dot) appears to lie near the center of the distribution. The long axis of the diagram roughly parallels the plane of the Galactic disk. (The scale is not Herschel's, but rather is a modern estimate of what he had in mind.)

## Star Counts

In the late 18th century, long before the distances to any stars were known, the English astronomer William Herschel tried to estimate the shape of our Galaxy simply by counting how many stars he could see in different directions in the sky. Assuming that all stars were of about equal brightness, he concluded that the Galaxy was a somewhat flattened, roughly disk-shaped collection of stars lying in the plane of the Milky Way, with the Sun at or near its *center* (Figure 23.4). Subsequent refinements to this approach led to much the same picture. Herschel was unable to estimate the *size* of the Galaxy by this method, but early in the 20th century, with improved knowledge of the properties of stars, some astronomers went so far as to estimate the dimensions of this “Galaxy” as about 10 kpc in diameter by 2 kpc thick.

Today we know that the Milky Way Galaxy is several tens of kiloparsecs across, and the Sun lies far from the center. How could the older picture have been so flawed? The answer is that the earlier observations were made at visible wavelengths, and astronomers failed to take into account the (then-unknown) absorption of visible light by interstellar gas and dust. ∞ (Sec. 18.1) Only in the 1930s did astronomers begin to realize the true extent and importance of the interstellar medium.

Any objects in the Galactic disk more than a few kiloparsecs away from us are hidden from our view (in visible light) by the effects of interstellar dust. The apparent falloff in the density of stars with distance in the plane of the Milky Way is thus not a real thinning of their numbers in space, but simply a consequence of the murky environment in the Galactic disk. The long “fingers” in Herschel's map are directions in which the obscuration happens to be a little less severe than in others. However, because some obscuration occurs in all directions in the disk, the falloff is roughly similar no matter which way we look, so the Sun appears to be more or less at the center. The horizontal extent of Figure 23.4 corresponds approximately to the span of the blue and white arrows in Figure 23.1.

Radiation coming to us from above or below the plane of the Galaxy, where there is less gas and dust along the line of sight, arrives on Earth relatively unscathed. There is still some patchy obscuration, but the Sun happens to be located where the view out of the disk is largely unimpeded by nearby interstellar clouds.

## Spiral Nebulae and Globular Clusters

We have just seen how astronomers' attempts to probe the Galactic disk by optical means are frustrated by the effects of the interstellar medium. However, looking in other directions, out of the Milky Way plane, we can see to much greater distances. During the first quarter of the 20th century, studies of the large-scale structure of our Galaxy focused on two particularly important classes of objects, both found mainly away from the Milky Way. The first is *globular clusters*, those tightly bound swarms of old, reddish stars we met in Chapter 19. ∞ (Sec. 19.6) About 150 are now known in our own Galaxy. The second class consisted of objects that were known at the time as *spiral nebulae*. Examples are shown in Figures 23.2(a) and 23.3(a). We know them today as **spiral galaxies**, comparable in size to our own.

Early 20th-century astronomers had no means of determining the distances to any of these objects. They are too far away to have any observable parallax, and with the technology of the day, main-sequence stars (after the discovery of the main sequence in 1911) could not be clearly identified and measured. For these reasons, neither of the techniques discussed in Chapter 17—trigonometric and spectroscopic parallax—was applicable. ∞ (Secs. 17.1, 17.6) As a result, even the most basic properties—size, mass, and stellar and interstellar content—of globular clusters and spiral nebulae were unknown. It was assumed that the globular clusters lay within our own Galaxy, which was thought at the time to be relatively small (using the size estimates just mentioned). The locations of the spiral nebulae were much less clear.

Knowing the distance to an object is vitally important to understanding its true nature. As an example, consider again the Andromeda “nebula” (Figure 23.2). In the late 19th century, when improved telescopes and photographic techniques allowed astronomers to obtain images showing detail comparable to that in Figure 23.2(a), the newly released photographs caused great excitement among astronomers, who thought they were seeing the formation of a star from a swirling gaseous disk! Comparing Figure 23.2(a) with the figures in Chapter 15 (especially Figure 15.2b), we can perhaps understand how such a mistake could be made—if we

thought that we were looking at a relatively close, star-sized object. Far from demonstrating that Andromeda was distant and large, the new observations seemed to confirm that it was just a small part of our own Galaxy.

Further observations soon made it clear that Andromeda was not a star-forming region. Andromeda's parallax is too small to measure, indicating that it must be at least several hundred parsecs from Earth, and, even at 100 pc—which we now know is vastly less than Andromeda's true distance—an object the size of the solar nebula would be impossible to resolve and simply would not look like Figure 23.2(a). (See Section 22.4 for another, more recent example of how distance measurements directly affect our theoretical understanding of observational data.)

During the first quarter of the 20th century, both the size of our Galaxy and the distances to the spiral nebulae were hotly debated in astronomical circles (see below, and also the discussion in the Part 4 Opener on p. 574). One school of thought maintained that the spiral nebulae were relatively small systems contained within our Galaxy. Other astronomers held that the spirals were much larger objects, lying far outside the Milky Way Galaxy and comparable to it in size. However, with no firm distance information, both arguments were inconclusive. Only with the discovery of a new distance-measurement technique—which we discuss next—was the issue finally settled in favor of the latter view. However, in the process, astronomers' conception of our own Galaxy changed radically and forever.

## A New Yardstick

An important by-product of the laborious effort to catalog stars around the turn of the 20th century was the systematic study of **variable stars**—stars whose luminosity changes with time, some quite erratically, others more regularly. Only a small fraction of stars fall into this category, but those that do are of great astronomical significance.

We have encountered examples of variable stars in earlier chapters. In an eclipsing binary, for example, the total brightness varies because one star in a binary system periodically blocks the light of the other. ∞ (Sec. 17.7) Binary membership has much more violent consequences in novae, also called *cataclysmic variables* because of their sudden, large changes in brightness. ∞ (Sec. 21.3)



In other instances, however, the variability is a basic trait of a star and is not dependent on its being a part of a binary system. We call such a star an *intrinsic variable*. A particularly important class of intrinsic variables is the **pulsating variable stars**, which vary cyclically in luminosity in very characteristic ways (Figure 23.5). Two types of pulsating **variable stars** that have played central roles in revealing both the true extent of our Galaxy and the distances to our galactic neighbors are the **RR Lyrae** and

**Cepheid variables**. Following long-standing astronomical practice, the names come from the first star of each class to be discovered—in this case, the variable star labeled RR in the constellation Lyra and the variable star Delta Cephei, the fourth brightest star in the constellation Cepheus.


RR Lyrae and Cepheid variable stars are recognizable by the characteristic shapes of their light curves. RR Lyrae stars all pulsate similarly (Figure 23.5b), with only small



(a)



▲ **FIGURE 23.5 Variable Stars** (a) The Cepheid variable star WW Cygni is shown here (boxed) on successive nights, near its maximum and minimum brightness; two photos, one from each night, were superimposed and then slightly displaced. (b) Light curve of the pulsating variable star RR Lyrae. All RR Lyrae-type variables have essentially similar light curves, with periods of less than a day. (c) The light curve of WW Cygni, with a period of about 3 days. (Harvard College Observatory)



▲ **FIGURE 23.6 Variable Stars on the H–R Diagram** Pulsating variable stars are found in the instability strip of the H–R diagram. As a high-mass star evolves through the strip, it becomes a Cepheid variable. Low-mass horizontal-branch stars in the instability strip are RR Lyrae variables.

differences in period between them. Observed periods range from about 0.5 to 1 day. Cepheid variables also pulsate in distinctive ways (the regular “sawtooth” pattern in Figure 23.5c), but different Cepheids can have quite different pulsation periods, ranging from about 1 to 100 days. The period of any given RR Lyrae or Cepheid variable is, to a high degree of accuracy, the same from one cycle to the next. The key point is that pulsating variable stars can be recognized and identified *just by observing the variations in the light they emit*.

Note, by the way, that pulsating variable stars have *nothing* whatsoever to do with the pulsars discussed in the previous chapter! Pulsars are rapidly rotating neutron stars beaming energy into space as they spin; as we will see in a moment, pulsating variable stars are “normal” stars undergoing a temporary period of instability as they evolve. ☞ (Sec. 22.2)

Why do Cepheids and RR Lyrae variables pulsate? The basic mechanism was first suggested by the British astrophysicist Sir Arthur Eddington in 1941. The structure of any star is determined in large part by how easily radiation can travel from the core to the photosphere—that is, by the *opacity* of the interior, the degree to which the gas hinders the passage of light through it. If the opacity

rises, the radiation becomes trapped, the internal pressure increases, and the star “puffs up.” If the opacity falls, radiation can escape more easily, and the star shrinks. According to theory, under certain circumstances a star can become unbalanced and enter a state in which the flow of radiation causes the opacity first to rise—making the star expand, cool, and diminish in luminosity—and then to fall, leading to the pulsations we observe.

The conditions necessary to cause pulsations are not found in main-sequence stars. Rather, they occur in evolved post-main-sequence stars as they pass through a region of the H–R diagram known as the *instability strip* (Figure 23.6). When a star’s temperature and luminosity place it in this strip, the star becomes internally unstable. Both its temperature and its radius vary in a regular way, causing the pulsations we observe: For the reasons just described, as the star brightens, its radius shrinks and its surface becomes hotter; as its luminosity decreases, the star expands and cools. As we learned in Chapter 20, high-mass stars evolve across the upper part of the H–R diagram. When their evolutionary tracks take them into the instability strip, they become Cepheid variables. ☞ (Sec. 20.4) RR Lyrae variables are low-mass horizontal-branch stars that lie within the lower portion of the instability strip. ☞ (Sec. 20.2) Thus, pulsating variables are normal stars passing through a brief—roughly million-year—phase of instability as a natural part of stellar evolution.


## The Cosmic Distance Scale

The importance of these stars to Galactic astronomy lies in the fact that once we recognize a star as being of the RR Lyrae or Cepheid type, we can infer its luminosity, and that in turn allows us to measure its distance. The distance calculation is precisely the same as that presented in Chapter 17 during our discussion of spectroscopic parallax. ☞ (Sec. 17.6) Comparing the star’s (known) luminosity with its (observed) apparent brightness yields an estimate of its distance by the inverse-square law: ☞ (Sec. 17.2)

$$\text{apparent brightness} \propto \frac{\text{luminosity}}{\text{distance}^2}.$$

In this way, astronomers can use pulsating variables as a means of determining distances, both within our own Galaxy and far beyond.

How do we infer a variable star’s luminosity? For RR Lyrae stars, doing so is simple. As we saw in Chapter 20, all horizontal branch stars have basically the same luminosity (averaged over a complete pulsation cycle)—about 100 times that of the Sun. ☞ (Sec. 20.2) Thus, once a variable star is recognized as being of the RR Lyrae type, its luminosity is immediately known. For Cepheids, we make use of a close




▲ **FIGURE 23.7 Period–Luminosity Plot** A plot of pulsation period versus average absolute brightness (that is, luminosity) for a group of Cepheid variable stars. The two properties are quite tightly correlated. The pulsation periods of some RR Lyrae variables are also shown.

correlation between average luminosity and pulsation period, discovered in 1908 by Henrietta Leavitt of Harvard University (see *Discovery 23-1*) and known simply as the **period–luminosity relationship**. Cepheids that vary slowly—that is, that have long periods—have large luminosities; conversely, short-period Cepheids have low luminosities.

Figure 23.7 illustrates the period–luminosity relationship for Cepheids found within a thousand parsecs or so of Earth. Astronomers can plot such a diagram for relatively nearby stars because they can measure their distances by using stellar or spectroscopic parallax. Once the distances are known, the luminosities of those stars can be calculated. We know of no exceptions to the period–luminosity relationship, and it is consistent with theoretical calculations of pulsations in evolved stars. Consequently, we assume that it holds for all Cepheids, near and far. Thus, a simple measurement of a Cepheid variable’s pulsation period immediately tells us its luminosity—we just read it off the plot in Figure 23.7. (The roughly constant luminosities of the RR Lyrae variables are also indicated in the figure.)

This distance-measurement technique works well, provided that the variable star can be clearly identified and its pulsation period measured. With Cepheids, the method allows astronomers to estimate distances out to about 25 million parsecs, enough to take us all the way to the nearest galaxies. The less-luminous RR Lyrae stars are not so easily seen as Cepheids, so their useful range is not as great. However, they are much more common, so, within their limited range, they are actually more useful than Cepheids.

Figure 23.8 extends our cosmic distance ladder, begun in Chapter 2 with radar ranging in the solar system and expanded in Chapter 17 to include stellar and spectroscopic parallax, by adding variable stars as a fourth method of



▲ **FIGURE 23.8 Variable Stars on Distance Ladder** Application of the period–luminosity relationship for Cepheid variable stars allows estimates of distances out to about 25 Mpc with reasonable accuracy.

determining distance. Note that because the period–luminosity relationship is calibrated by using nearby stars, this latest rung inherits any and all uncertainties and errors present in the lower levels. Uncertainties also arise from the “scatter” of data points shown in Figure 23.7. Although the overall connection between period and luminosity is unmistakable, the individual data points do not quite lie on a straight line; instead, a range of possible luminosities corresponds to any measured period.

## The Size and Shape of Our Galaxy

Many RR Lyrae variables are found in globular clusters. Early in the 20th century, the American astronomer Harlow Shapley used observations of variable stars to make two very important discoveries about the Galactic globular cluster system. First, he showed that most globular clusters reside at great distances—many thousands of parsecs—from the Sun. Second, by measuring the direction and distance of each cluster, he was able to determine the three-dimensional distribution of the clusters in space (Figure 23.9). In this way, Shapley demonstrated that the globular clusters map out a truly gigantic, and roughly *spherical*, volume of space, about 30 kpc across.\* However, the center of the distribution lies nowhere near our Sun; rather, it is located nearly 8 kpc away from us, in the direction of the constellation Sagittarius.

\*The Galactic globular cluster system and the Galactic halo, of which it is a part, are somewhat flattened in the direction perpendicular to the disk, but the degree of flattening is quite uncertain. The halo is certainly much less flattened than the disk, however.



## DISCOVERY 23-1

## Early “Computers”

A large portion of the early research in observational astronomy focused on monitoring stellar luminosities and analyzing stellar spectra. Much of this pioneering work was done using photographic methods. What is not so well known is that most of the labor was accomplished by women. Around the turn of the 20th century, a few dozen dedicated women—assistants at the Harvard College Observatory—created an enormous database by observing, sorting, measuring, and cataloging photographic information that helped form the foundation of modern astronomy. Some of them went far beyond their duties in the lab to make several of the basic astronomical discoveries often taken for granted today.

The first photograph below taken in 1910, shows several of those women carefully examining star images and measuring variations in luminosity or wavelengths of spectral lines. In the cramped quarters of the Harvard Observatory, they inspected image after image to collect a vast body of data on millions of measurements of hundreds of thousands of stars. Note the plot of stellar luminosity changes pasted on the wall at the left. The pattern is so regular that it likely belongs to a Cepheid variable. Known as “computers” (there were no electronic devices then), these women were paid 25 cents an hour.

The second photograph, taken in 1913, shows a more formal portrait of another group of staff members, along with their director, E.C. Pickering. Though looking rather stern here, Pickering was often described as a true Victorian gentleman who championed a policy, unique at the time, of admitting women to the staff. Also prominent here (and symmetrically positioned to Pickering’s left) is Annie Cannon, perhaps the most accomplished of the early group of women who, beginning in 1880, undertook a survey of the skies that lasted for more than half a century—work that netted Cannon the first Oxford honorary degree awarded to a woman.

The first major result of this work was a record of the brightnesses and spectra of tens of thousands of stars, published in 1890 under the direction of Williamina Fleming (seen standing in the photograph below). On the basis of this compilation, several of these women made fundamental contributions to




astronomy. In 1897 Antonia Maury (who is also pictured in the first photo at left rear) undertook the most detailed study of stellar spectra to that time, enabling Hertzsprung and Russell independently to develop what is now called the H–R diagram. In 1898 Annie Cannon proposed the spectral classification system (described in Chapter 17) that is now the international standard for categorizing stars. [∞ \(Sec. 17.5\)](#) And in 1908 Henrietta Leavitt discovered the period–luminosity relationship for Cepheid variable stars, which later allowed Pickering’s successor as director, Harlow Shapley (see the introductory essay for Part 4), to recognize our Sun’s true position in the universe.

All was not work, however, and socializing was common among this generation of astronomers. The third photograph (below) shows a 1920s scene from a humorous play portraying life at the Observatory, starring (at center) the then youngest of the “lady computers,” Cecilia Payne, who would go on to become one of the foremost astronomers of the 20th century (see the introductory essay for Part 3).



(Harvard College Observatory)



### Narrated **FIGURE 23.9** Globular Cluster Distribution

**MA** Our Sun does not coincide with the center of the very large collection of globular clusters (indicated by the pink dots). Instead, more globular clusters are found in one direction than in any other. The Sun resides closer to the edge of the collection, which measures roughly 30 kpc across. The globular clusters outline the true distribution of stars in the Galactic halo.


In a brilliant intellectual leap, Shapley realized that the distribution of globular clusters maps out the true extent of stars in the Milky Way Galaxy—the region that we now call the Galactic halo. The hub of this vast collection of matter, 8 kpc from the Sun, is the **Galactic center**. Figure 23.9 shows

the distribution, based on modern data, of the 138 globular clusters lying within 20 kpc of the center. As illustrated in Figure 23.10, we live in the “suburbs” of this huge ensemble—in the Galactic disk, the thin sheet of young stars, gas, and dust that cuts through the center of the halo. Since Shapley’s time, astronomers have identified many individual stars—that is, stars not belonging to any globular cluster—within the Galactic halo.

Shapley’s bold interpretation of the globular clusters as defining the overall distribution of stars in our Galaxy was an enormous step forward in human understanding of our place in the universe. Five hundred years ago, Earth was considered the center of all things. Copernicus argued otherwise, demoting our planet to an undistinguished location removed from the center of the solar system. In Shapley’s time, as we have just seen, the prevailing view was that our Sun was the center not only of the Galaxy, but also of the universe. Shapley showed otherwise. With his observations of globular clusters, he simultaneously increased the size of our Galaxy by almost a factor of 10 over earlier estimates and banished our parent Sun to its periphery, virtually overnight!

## The Shapley–Curtis Debate

Curiously, Shapley’s dramatic revision of the size of the Milky Way Galaxy and our place in it only strengthened his erroneous opinion that the spiral nebulae were part of our Galaxy and that our Galaxy was essentially the entire universe. He regarded as beyond belief the idea that there could be other structures as large as our Galaxy. The scientific issues involved in understanding the nature of the spiral nebulae were clearly drawn in a famous 1920 debate



### Interactive **FIGURE 23.10** Stellar Populations in Our

**MA** **Galaxy** Based on observations of young stars and gas in the disk, and the old stars and globular clusters in the halo, astronomers have constructed a comprehensive picture of the structure of our Galaxy. This artist’s conception of a (nearly) edge-on view of the Milky Way Galaxy shows schematically the distributions of young blue stars, open clusters, old red stars, and globular clusters. (The brightness and size of our Sun are greatly exaggerated for clarity.)

between Shapley and Lick Observatory astronomer Heber Curtis. (See also p. 574.) We list here some key elements of the debate, illustrating the sometimes erratic process by which scientific knowledge grows: ∞ (Sec. 1.2)

1. *Size of the Milky Way.* Shapley correctly asserted that the diameter of the Milky Way Galaxy was much larger than the “conventional” scale based on star counts, but then incorrectly concluded that similar-sized galaxies beyond our own could not exist. Curtis incorrectly accepted the smaller size for our Galaxy, but correctly argued that similar galaxies might exist beyond our own.
2. *Distribution of the nebulae on the sky.* Curtis noted that the observed spiral nebulae were found away from the plane of the Galaxy, and he suggested that our Galaxy had a “ring” of occulting material in its plane, like those observed in many edge-on spirals, preventing us from seeing the nebulae in the plane. Shapley simply had to assume that spiral nebulae were, for some unknown reason, not found in the Galactic plane. Curtis was almost correct in this point. Note, however, that the effects of absorption by interstellar dust were unknown at the time. ∞ (Sec. 18.1)
3. *Observations of novae.* Shapley argued (correctly) that the observed apparent brightnesses of some “novae” seen in spiral nebulae implied enormous luminosities if the nebulae lay at large distances. ∞ (Secs. 17.2, 21.1) Curtis suggested (also correctly) that these anomalous events might be members of a second, much brighter, class of nova—today we call them *supernovae*. ∞ (Sec. 21.3)
4. *Brightness and spectra of the nebulae.* Shapley pointed out that the measured brightnesses and colors of spiral nebulae differed from what he would have expected to see if our Galaxy were observed from afar, suggesting that the nebulae were somehow fundamentally different from the Milky Way. Curtis had no answer. We know today that these differences exist because of interstellar absorption and reddening, which prevent astronomers from getting a comparable view of our own Galaxy. ∞ (Sec. 18.1) Curtis did correctly note that spectral lines seen in spiral nebulae were generally consistent with the nebulae’s being assemblages of large numbers of stars, supporting his argument that they were stellar systems comparable to our own Galaxy. ∞ (Sec. 4.2)
5. *Rotation of the nebulae.* Shapley cited published measurements of the angular rotation speeds of some spiral nebulae, which implied that the nebulae would have to be spinning faster than the speed of light if they were very distant and hence very large. ∞ (More Precisely 1-2) Curtis simply responded that the observations were in error. Curtis was right, but he couldn’t prove it at the time.

We see that both men made some correct and some incorrect statements (or conclusions) about the problem. However, with the observations of the day, their disagreements could

not be resolved, and the debate was inconclusive. But technology marches on, and just a few years later, in 1925, American astronomer Edwin Hubble reported that he had observed Cepheids in the Andromeda Galaxy and finally succeeded in measuring its distance. His work firmly established Andromeda as a separate galaxy lying far beyond our own, finally extending the Copernican principle to the Galaxy itself.

### CONCEPT Check

- ✓ Can variable stars be used to map out the structure of the Galactic disk?

## 23.3 Galactic Structure

Based on optical, infrared, and radio observations of stars, gas, and dust, Figure 23.10 illustrates the different spatial distributions of the disk, bulge, and halo of the Milky Way Galaxy. The extent of the halo is based largely on optical observations of globular clusters and other halo stars. However, as we have seen, optical techniques can cover only a small portion of the dusty Galactic disk. Much of our knowledge of the structure of the disk on larger scales is based on radio observations, particularly of the 21-cm radio emission line produced by atomic hydrogen. ∞ (Sec. 18.4)

The center of the gas distribution coincides roughly with the center of the globular cluster system, lying about 8 kpc from the Sun. In fact, the location of the Galactic center is determined most accurately from radio observations of Galactic gas. The densities of both stars and gas in the disk decline quite rapidly beyond about 15 kpc from the Galactic center (although some radio-emitting gas has been observed out to at least 50 kpc).

### The Spatial Distribution of Stars

Perpendicular to the Galactic plane, the disk in the vicinity of the Sun is relatively thin—“only” 300 pc thick, or about one hundredth of the 30-kpc Galactic diameter. Don’t be fooled, though: Even if you could travel at the speed of light, it would take you a thousand years to traverse the thickness of the Galactic disk. The disk may be thin compared with the Galactic diameter, but it is huge by human standards.

Actually, the thickness of the Galactic disk depends on the kinds of objects measured. Young stars and interstellar gas are more tightly confined to the plane than are stars like the Sun, and solar-type stars in turn are more tightly confined than are older K- and M-type dwarfs. The reason for these differences is that stars form in interstellar clouds close to the plane of the disk, but then tend to drift out of the disk over time, mainly due to their interactions with other stars and molecular clouds. Thus, *as stars age, their abundance above and below the plane of the disk slowly increases*. Note that these considerations do not apply to the Galactic halo,

whose ancient stars and globular clusters extend far above and below the Galactic plane. As we will see in a moment, the halo is a remnant of an early stage of our Galaxy’s evolution and predates the formation of the disk.

Recently, improved observational techniques have revealed an intermediate category of Galactic stars, midway between the old halo and the younger disk, both in age and in spatial distribution. Consisting of stars with estimated ages in the range of 7–10 billion years, this *thick-disk* component of the Milky Way Galaxy measures some 2–3 kpc from top to bottom. Its thickness is too great to be explained by the slow drift just described. Like the halo, it appears to be a vestige of our Galaxy’s distant past.

Also shown in Figure 23.10 is our Galaxy’s central bulge, measuring roughly 6 kpc across in the plane of the Galactic disk by 4 kpc perpendicular to the plane. Obscuration by interstellar dust makes it difficult to study the detailed structure of the Galactic bulge in optical images of the Milky Way. (See, for example, Figure 18.4, which would clearly show a large portion of the bulge were it not for interstellar absorption.) However, at longer wavelengths, which are less affected by interstellar matter, a much clearer picture emerges (Figure 23.11; see also Figure 23.3b). Detailed measurements of the motion of gas and stars in and near the bulge imply that it is actually football shaped, about half as wide as it is long, with the long axis of the “football” lying in the Galactic plane. On the basis of these observations, astronomers speculate that the central part of our Galaxy may have a distinctly elongated, or barlike, appearance and that we may live in a galaxy of the “barred-spiral” type, as discussed further in Chapter 24.

## Stellar Populations

Aside from their distributions in space, the three components of the Galaxy—disk, bulge, and halo—have several other properties that distinguish them from one another. First, the halo contains almost *no* gas or dust—just the opposite of the disk and bulge, in which interstellar matter is common. Second, there are clear differences in both *appearance* and *composition* among disk, bulge, and halo stars: Stars in the Galactic bulge and halo are found to be distinctly *redder* than stars found in the disk. Observations of other spiral galaxies also show this trend—the blue-white tint of the disk and the yellowish coloration of the bulge are evident in Figures 23.2(a) and 23.3(a).



All the bright, blue stars visible in our sky are part of the Galactic disk, as are the young, open star clusters and star-forming regions. In contrast, the cooler, redder stars—including those found in the old globular clusters—are more uniformly distributed throughout the disk, bulge, and halo. Galactic disks appear bluish because main-sequence O- and B-type blue supergiants are very much brighter

than G-, K-, and M-type dwarfs, even though the dwarfs are present in far greater numbers.

The explanation for the marked difference in stellar content between disk and halo is this: Whereas the gas-rich Galactic disk is the site of ongoing star formation and so contains stars of all ages, all the stars in the Galactic halo are *old*. The absence of dust and gas in the halo means that no new stars are forming there, and star formation apparently ceased long ago—at least 10 billion years in the past, judging from the types of halo stars we now observe. (Recall from Chapter 20 that most globular clusters are thought to be between 10 and 12 billion years old.) ∞ (Sec. 20.5) The gas density is very high in the inner part of the Galactic bulge, making that region the site of vigorous ongoing star formation, and both very old and very young stars mingle there. The bulge’s gas-poor outer regions have properties more similar to those of the halo.

Support for this picture comes from studies of the spectra of halo stars, which indicate that these stars are far less abundant in heavy elements (i.e., elements heavier than helium) than are nearby stars in the disk. In Chapter 21, we saw how each successive cycle of star formation and evolution enriches the interstellar medium with the products of stellar nucleosynthesis, leading to a steady increase in heavy elements with time. ∞ (Sec. 21.5) Thus, the scarcity of these elements in halo stars is consistent with the view that the halo formed long ago.

Astronomers often refer to young disk stars as *Population I* stars and to old halo stars as *Population II* stars. The idea of two stellar “populations” dates from the 1940s, when the differences between disk and halo stars first became clear. The names are something of an oversimplification, as there is actually a continuous variation in stellar ages throughout the Milky Way Galaxy, not a simple division of stars into two distinct “young” and “old” categories. Nevertheless, the terminology is widely used.



**Interactive FIGURE 23.11 Infrared View of the Milky Way** A wide-angle infrared image of the disk and bulge of the Milky Way Galaxy, as observed by the Two Micron All Sky Survey. Compare Figure 23.3(b). (UMass/Caltech)


## Orbital Motion

Now let's turn our attention to the *dynamics* of the Milky Way Galaxy—that is, to the motion of the stars, dust, and gas it contains. Are the internal motions of our Galaxy's members disordered and random, or are they part of some gigantic “traffic pattern”? The answer depends on our perspective. The motion of stars and clouds we see on small scales (within a few tens of parsecs from the Sun) seems random, but on larger scales (hundreds or thousands of parsecs) the motion is much more orderly.

As we look around the Galactic disk in different directions, a clear pattern of motion emerges (Figure 23.12). Radiation received from stars and interstellar gas clouds in the upper-right and lower-left quadrants of the figure is generally *blueshifted*. At the same time, radiation from stars and gas sampled in the upper-left and lower-right quadrants tends to be *redshifted*. In other words, some regions of the Galaxy (those in the blueshifted directions) are approaching the Sun, whereas others (the redshifted ones) are receding from us. ∞ (Sec. 3.5)

Careful study of the positions and velocities of stars and gas clouds near the Sun leads us to two important

The curved arrows denote the speed of the disk material, which is greater closer to the center.



▲ **FIGURE 23.12 Orbital Motion in the Galactic Disk** Stars and interstellar clouds in the neighborhood of the Sun show systematic Doppler motions, implying that the disk of the Galaxy spins in a well-ordered way. These four Galactic quadrants are drawn to intersect not at the Galactic center, but at the Sun, the location from which observations are made. Because the Sun orbits faster than stars and gas at larger radii, it moves away from material at top left and gains on that at top right, resulting in the Doppler shifts indicated. Likewise, stars and gas in the bottom left quadrant are gaining on us, while material at bottom right is pulling away.

conclusions about the motion of the Galactic disk. First, the entire disk is *rotating*—stars, gas, and dust all move in roughly circular paths around the Galactic center, their orbits governed by the Galaxy's gravitational pull. The orbital speed in the vicinity of the Sun is about 220 km/s. Thus, at the Sun's distance of 8 kpc from the Galactic center, material takes about 225 million years (an interval of time sometimes called 1 *Galactic year*) to complete one circuit.

Second, the Galactic rotation period depends on distance from the Galactic center, being shorter closer to the center and longer at greater distances. In other words, the Galactic disk rotates not as a solid object, but *differentially*. ∞ (Secs. 11.1, 16.1) Accurate measurements, made by the *Hipparcos* satellite, of stars within a few hundred parsecs of the Sun have proved particularly valuable in measuring these important Galactic properties. ∞ (Sec. 17.1) Similar *differential rotation* is observed in Andromeda and many other spiral galaxies.

This picture of orderly circular motion about the Galactic center applies only to the disk: Stars in the Galactic halo and bulge are not so well behaved. The old globular clusters in the halo and the faint, reddish individual stars in both the halo and the bulge do *not* share the disk's well-defined rotation. Instead, their orbital orientations are largely random.\* Although these objects do orbit the Galactic center, they move in all directions, their paths filling an entire three-dimensional volume rather than a nearly two-dimensional disk.

Figure 23.13 contrasts the motion of bulge and halo stars with the much more regular orbits of stars in the Galactic disk. At any given distance from the Galactic center, bulge, or halo, stars move at speeds comparable to the disk's rotation speed at that radius, but in *all* directions, not just one. Their orbits carry these stars repeatedly through the plane of the disk and out the other side. (They don't collide with stars in the disk because interstellar distances are huge compared with the diameters of individual stars—a star or even an entire star cluster passes through the disk almost as though it weren't there—see Section 25.2.) Some well-known stars in the vicinity of the Sun—the bright giant Arcturus, for example—are actually halo stars that are “just passing through” the disk on orbits that take them far above and below the Galactic plane.

Recently, astronomers have detected numerous *tidal streams* in the Galactic halo—groups of stars thought to be the remnants of globular clusters and even small satellite galaxies (see Sec. 24.1) torn apart by our Galaxy's tidal field. Just as micrometeoroid swarms in our solar system follow


\* Halo stars do, in fact, have some net rotation about the Galactic center, but the rotational component of their motion is overwhelmed by the larger random component. The motion of bulge stars also has a rotational component, larger than that of the halo, but still smaller than the random component of stellar motion in the bulge.

**TABLE 23.1** Overall Properties of the Galactic Disk, Halo, and Bulge

Galactic Disk	Galactic Halo	Galactic Bulge
highly flattened	roughly spherical—mildly flattened	somewhat flattened and elongated in the plane of the disk (“football shaped”)
contains both young and old stars	contains old stars only	contains young and old stars; more old stars at greater distances from the center
contains gas and dust	contains no gas and dust	contains gas and dust, especially in the inner regions
site of ongoing star formation	no star formation during the last 10 billion years	ongoing star formation in the inner regions
gas and stars move in circular orbits in the Galactic plane	stars have random orbits in three dimensions	stars have random orbits but some net rotation about the Galactic center
spiral arms	no obvious substructure	central regions probably elongated into a bar; ring of gas and dust near center
overall white coloration, with blue spiral arms	reddish in color	yellow-white

the orbit of their disrupted parent comet long after the comet itself is gone, stars in a tidal stream are now spread out around the entire original orbit of their parent cluster or galaxy. [∞ \(Sec. 14.4\)](#) We will discuss the processes responsible for them in more detail in Chapter 25.

Table 23.1 compares some key properties of the three basic components of the Galaxy.



**▲ FIGURE 23.13 Stellar Orbits in Our Galaxy** Stars in the Galactic disk (blue curves) move in orderly, circular orbits about the Galactic center. In contrast, halo stars (orange curves) move randomly around the center. The orbit of a typical halo star takes it high above the Galactic disk, then down through the disk plane, and then out the other side and far below the disk. The orbital properties of bulge stars are intermediate between those of disk and halo stars.

### CONCEPT Check


✓ Why do astronomers regard the disk and the halo as different components of our Galaxy?

## 23.4 Formation of the Milky Way

Is there some evolutionary scenario that can naturally account for the Galactic structure we see today? The answer is that there is, and it takes us all the way back to the birth of our Galaxy, more than 10 billion years ago. [∞ \(Sec. 20.5\)](#) Not all the details are agreed upon by all astronomers, but the overall picture is now fairly widely accepted. For simplicity, we confine our discussion here to the Galactic disk and halo; in many ways, the bulge is intermediate in its properties between these two extremes. Figure 23.14 illustrates the current view of our Galaxy’s evolution, starting (not unlike the star-formation scenario outlined in Chapter 19) from a contracting cloud of pregalactic gas. [∞ \(Sec. 19.1\)](#)

When the first Galactic stars and globular clusters formed, the gas in our Galaxy had not yet accumulated into a disk. Instead, it was spread out over an irregular and quite extended region of space, spanning many tens of kiloparsecs in all directions (Figure 23.14b). When the first stars formed, they were distributed throughout that volume. Their distribution today (the Galactic halo) reflects that fact—it is an imprint of their birth. Many astronomers think that the very first stars formed even earlier, in smaller systems that later merged to create our Galaxy (Figure 23.14a). Probably, many more stars were born during the mergers themselves, as interstellar gas clouds collided and began to collapse. [∞ \(Sec. 19.5\)](#) Whatever the details, the present-day halo would look much the same in either case.

Since those early times, rotation has flattened the gas in our Galaxy into a relatively thin disk (Figure 23.14c).



◀ **FIGURE 23.14 Milky Way**

**Formation** (a) The Milky Way Galaxy likely formed by a merger of several smaller systems. (b) Early on, our Galaxy was irregularly shaped, with gas distributed throughout its volume. When stars formed during this stage, their orbits carried them throughout an extended three-dimensional volume surrounding the newborn Galaxy. (c) In time, the gas and dust fell to the Galactic plane and formed a spinning disk. The stars that had already formed were left behind in the halo. (d) New stars forming in the disk inherit its overall rotation and so orbit the Galactic center on ordered, circular orbits.

Physically, the process is similar to the flattening of the solar nebula during the formation of the solar system, as described in Chapters 6 and 15, except on a vastly larger scale. ∞ (Secs. 6.7, 15.2) Star formation in the halo ceased billions of years ago when the raw materials—the gas and dust—cooled and fell toward the Galactic plane. Ongoing star formation in the disk gives it its bluish tint, but the halo’s short-lived bright blue stars have long since burned out, leaving only the long-lived red stars that give the halo its characteristic pinkish glow. The Galactic halo is ancient, whereas the disk is full of youthful activity. The thick disk, with its intermediate-age stars, may represent an intermediate stage of star formation that occurred while the gas was still flattening into the plane.

Recent studies of the composition of stars in the Galactic disk suggest that the infall of halo gas is still going on today. The best available models of star formation and stellar nucleosynthesis predict that the fraction of heavy elements in disk stars should be significantly *greater* than is actually observed, unless the gas in the disk is steadily being “diluted” by relatively unevolved gas arriving from the halo at a rate of perhaps 5–10 solar masses per year. ∞ (Sec. 21.5) This may not sound like much mass, but accumulated over billions of years it actually amounts to a significant fraction of the total mass of the disk (see Section 23.6).

This theory also explains the randomly oriented orbits of the halo stars and the more ordered motion of the disk (Figure 23.14d). When the halo developed, the irregularly shaped Galaxy was rotating only very slowly, so there was no strongly preferred direction in which matter tended to move. As a result, halo stars were free to travel along nearly any path once they formed (or when their parent systems merged), leading to the random halo orbits we observe today. After the Galactic disk formed, however, stars that formed from its gas and dust inherited its rotational motion and so move on well-defined, circular orbits. Again, the thick disk’s orbital properties are consistent with the idea that it formed while gas was still sinking to the Galaxy’s midplane.

In principle, the structure of our Galaxy bears witness to the conditions that created it. In practice, however, the interpretation of the observations is made difficult by the sheer complexity of the system we inhabit and by the many competing physical processes that have modified its appearance since it formed. As a result, the early stages of the Milky Way are still quite poorly understood. We will return to the subject of galaxy formation in Chapters 24 and 25.

### CONCEPT Check

✓ Why are there no young halo stars?

## 23.5 Galactic Spiral Arms

If we want to look beyond our immediate neighborhood and study the full extent of the Galactic disk, we cannot rely on optical observations, as interstellar absorption severely limits our vision. In the 1950s, astronomers developed a very important tool for exploring the distribution of gas in our Galaxy: spectroscopic radio astronomy.

### Radio Maps of the Milky Way


The keys to observing Galactic interstellar gas are the 21-cm radio emission line produced by atomic hydrogen and the many radio molecular lines formed in molecular cloud complexes. [∞ \(Sec. 18.4\)](#) Long-wavelength radio waves are largely unaffected by interstellar dust, so they travel more or less unimpeded through the Galactic disk, allowing us to “see” to great distances. Because hydrogen is by far the most abundant element in interstellar space, the 21-cm signals are strong enough that a large portion of the disk can be observed in this way. As noted in Chapter 18, observations of spectral lines from “tracer” molecules, such as carbon monoxide, allow us to study the distribution of the densest interstellar clouds. [∞ \(Sec. 18.5\)](#)

Earlier, we noted that observations of stars within several hundred parsecs of the Sun have allowed astronomers to measure the rotation rate of the Galaxy in the solar neighborhood. As indicated in Figure 23.15, in order to probe to greater distances, astronomers often turn to radio

observations (illustrated here with 21-cm radiation), because long-wavelength radio waves are largely unaffected by interstellar dust, allowing astronomers to study virtually the entire Galactic disk. [∞ \(Sec. 18.4\)](#)

However, the distances to the clouds emitting the radio radiation are often poorly known. To determine a cloud’s location in the disk, astronomers use all available data, coupled with knowledge of Newtonian mechanics, to construct a *mathematical model* of the rotation of stars and gas throughout the Galactic disk. [∞ \(Sec. 2.8\)](#) Assuming circular orbits, the model allows us to turn a measured radial velocity into a distance along the line of sight. As in so many areas of astronomy, theory and observations complement one another: The data refine the theoretical model, and the model in turn provides the framework needed to interpret further observations. [∞ \(Sec. 1.2\)](#)

Radio astronomers couple their observations with this Galactic model to turn their measurements into detailed information about the distribution of gas along the line of sight. Because of the differential rotation described in Section 23.3, the measured velocity of a cloud depends on its *distance* from the Sun (Figure 23.15), and the Galactic model provides the connection between the two. Furthermore, the strength of the signal is a measure of the *density* of gas in the cloud—denser clouds contain more gas and emit more radiation. Thus, knowing direction, distance, and density, astronomers can use observations along different lines of sight to map out the radio-emitting gas in our Galaxy.




### Spiral Structure

Interstellar gas in the Galactic disk exhibits an organized pattern on a grand scale. Near the center, the gas in the disk fattens markedly in the Galactic bulge. Radio-emitting gas

**▲ FIGURE 23.15 Gas in the Galactic Disk** Because the disk of our Galaxy is rotating differentially (inside faster than the outside), 21-cm radio signals from different clumps of hydrogen matter along any line of sight are Doppler shifted by different amounts. Repeated observations in many different directions allow astronomers to map out the distribution of gas in our Galaxy.

*Typical radio spectrum showing how cloud 2 is blueshifted more than cloud 1.*





**Interactive** **FIGURE 23.16 Milky Way Spiral Structure** This face-on artist's conception of our Milky Way Galaxy illustrates the spiral structure of the Galactic disk. It is based on data collected by many teams of astronomers during the past few decades, including radio and infrared maps of stars, gas, and dust. Painted from the perspective of an observer 100 kpc above the Galactic plane, the spiral arms are at their best-determined positions, apparently emanating from a bar whose length is twice its width. Everything is drawn to scale (except for the oversized yellow dot near the top, which represents our Sun). The two small blotches to the left are dwarf galaxies, called the Magellanic Clouds, which are studied in Chapter 24. (Adapted from JPL)

interstellar clouds. The obvious conclusion is that the spiral arms are the part of the Galactic disk where star formation takes place. The brightness of the young stellar objects just listed is the main reason that the spiral arms of other galaxies are easily seen from afar (e.g., Figure 23.3a).

A central problem in understanding spiral structure is explaining how that structure persists over long periods of time. The basic issue is simple: Differential rotation makes it impossible for any large-scale structure “tied” to the disk material to survive. Figure 23.17 shows how a spiral pattern consisting always of the same group of stars and gas clouds would necessarily disappear within a few hundred million years. How, then, do the Galaxy's spiral arms retain their structure over long periods in spite of differential rotation?

has been observed out to at least 50 kpc from the Galactic center. Over much of the inner 20 kpc or so of the disk, the gas is confined within about 100 pc of the Galactic plane. Beyond that distance, the gas distribution spreads out somewhat, to a thickness of several kiloparsecs, and shows definite signs of being “warped,” possibly because of the gravitational influence of a pair of nearby galaxies (to be discussed in Chapter 24; see also Figure 23.16).

Radio studies provide perhaps the best direct evidence that we live in a spiral galaxy. Figure 23.16 is an artist's conception (based on observational data) of the appearance of our Galaxy as seen from far above the disk. The figure clearly shows our Galaxy's **spiral arms**, pinwheel-like structures originating close to the Galactic bulge and extending outward throughout much of the Galactic disk. Our Sun lies near the edge of one of these arms, which wraps around a large part of the disk. Notice, incidentally, the scale markers on Figures 23.9, 23.10, and 23.16: The Galactic globular-cluster distribution (Figure 23.9), the luminous stellar component of the disk (Figure 23.10), and the known spiral structure (Figure 23.16) all have roughly the *same* diameter—about 30 kpc. This scale is fairly typical of spiral galaxies observed elsewhere in the universe.


## Survival of the Spiral Arms

The spiral arms in our Galaxy are made up of much more than just interstellar gas and dust. Studies of the Galactic disk within a kiloparsec or so of the Sun indicate that young stellar and prestellar objects—emission nebulae, O- and B-type stars, and recently formed open clusters—are also distributed in a spiral pattern that closely follows the distribution of


A leading explanation for the existence of spiral arms is that they are **spiral density waves**—coiled waves of gas compression that move through the Galactic disk, squeezing clouds of interstellar gas and triggering the process of star formation as they go. [☞ \(Sec. 19.5\)](#) The spiral arms we observe are defined by the denser-than-normal clouds of gas the density waves create and by the new stars formed as a result of the spiral waves' passage.

This explanation of spiral structure avoids the problem of differential rotation, because the wave pattern is not tied to any particular piece of the Galactic disk. The spirals we see are merely patterns moving through the disk, not great masses of matter being transported from place to place. The density wave moves through the collection of stars and gas making up the disk just as a sound wave moves through air or an ocean wave passes through water, compressing different parts of the disk at different times. Even though the rotation rate of the disk material varies with distance from the Galactic center, the wave itself remains intact, defining the Galaxy's spiral arms.

In fact, over much of the visible portion of the Galactic disk (within about 15 kpc of the center), the spiral wave pattern is predicted to rotate *more slowly* than the stars and gas. Thus,




**Interactive** **FIGURE 23.17 Differential Galactic Rotation** The disk of our Galaxy rotates differentially, as depicted by the small white arrows that represent the angular speed of the disk. If spiral arms were somehow tied to the material of the Galactic disk, such an uneven rotation would cause the spiral pattern to wind up and disappear in a few hundred million years. Spiral arms would be too short-lived to be consistent with the numbers of spiral galaxies observed today.



as shown in Figure 23.18, Galactic material catches up with the wave, is temporarily slowed down and compressed as it passes through, and then continues on its way. (For a more down-to-earth example of an analogous process, see *Discovery 23-2*.)

*This idea of spiral arm construction is probably correct . . . but no one knows for sure.*



As material enters the density wave from behind, the gas is compressed and forms stars. Dust lanes mark the regions of highest-density gas. The most prominent stars—the bright O- and B-type blue giants—live for only a short time, so young stellar associations, emission nebulae, and open clusters with long main sequences are found only within the arms, near their birth sites, just ahead of the dust lanes. The brightness of these young systems emphasizes the spiral structure. Further downstream, ahead of the spiral arms, we see mostly older stars and star clusters. These objects have had enough time since their formation to outdistance the wave and pull away from it. Over millions of years, their random individual motions, superimposed on their overall rotation around the Galactic center, distort and eventually destroy their original spiral configuration, and they become part of the general disk population.

Note, incidentally, that although the spirals shown in Figure 23.18 have two arms each, astronomers are not certain how many arms make up the spiral structure in our

**Interactive** **FIGURE 23.18 Spiral Density Waves** Density-wave theory holds that the spiral arms seen in our own and many other galaxies are waves of gas compression and star formation moving through the material of the galactic disk. Gas enters an arm from behind, is compressed, and forms stars. The spiral pattern is delineated by dust lanes, regions of high gas density, and newly formed O- and B-type stars. The inset at right shows the spiral galaxy NGC 1566, which displays many of the features just described. (*AURA*)


## DISCOVERY 23-2

### Density Waves

In the late 1960s, American astrophysicists C. C. Lin and Frank Shu proposed a way in which spiral arms in the Galaxy could persist for many Galactic rotations. They argued that the arms themselves contain no “permanent” matter. They should thus not be viewed as assemblages of stars, gas, and dust moving intact through the disk—those would quickly be destroyed by differential rotation. Instead, a spiral arm should be envisaged as a density wave—a wave of compression and expansion sweeping through the Galaxy.

A wave in water builds up material temporarily in some places (crests) and lets it down in others (troughs). The wave *pattern* moves across the water, even though the water comprising the peaks and troughs does not. ∞ (Sec. 3.1) Similarly, as the spiral density wave encounters galactic matter, the gas is compressed to form a region of slightly higher than normal density. Galactic material enters the wave, is temporarily slowed down and compressed as it passes through, and then continues on its way. The compression triggers the formation of new stars and nebulae. In this way, the spiral arms are formed and re-formed repeatedly, without disappearing completely. Lin and Shu showed that the process can in fact maintain a spiral pattern for very long periods.

The accompanying figure illustrates the formation of a density wave in a much more familiar context: a traffic jam on a highway, triggered by the presence of a repair crew moving slowly down the road. As cars approach the crew, they slow down temporarily. Then they speed up again as they pass the work site and continue on their way. The result, as might be reported by a high-flying traffic helicopter, is a region of high traffic density, concentrated around the location of the work crew and moving with it. An



own Galaxy (see Figure 23.16). The theory makes no strong predictions on this point.

An alternative possibility is that the formation of stars drives the waves, instead of the other way around. Imagine a row of newly formed massive stars somewhere in the disk. The emission nebula created when these stars form, and the supernovae when they die, send shock waves through the surrounding gas, triggering new star formation. ∞ (Secs. 19.5, 21.5) Thus, as illustrated in Figure 23.19(a), the formation of one group of stars provides the mechanism for


observer on the side of the road, however, sees that the jam never contains the same cars for very long. Cars constantly catch up to the bottleneck, move slowly through it, and then speed up again, only to be replaced by more cars arriving from behind.

The traffic jam is analogous to the region of high stellar density in a Galactic spiral arm. Just as the traffic density wave is not tied to any particular group of cars, the spiral arms are not attached to any particular piece of disk material. Stars and gas enter a spiral arm, slow down for a while, then continue on their orbits around the Galactic center. The result is a moving region of high stellar and gas density, involving different parts of the disk at different times. Notice also that, just as in our Galaxy, the wave moves more slowly than, and independently of, the overall traffic flow.

We can extend our traffic analogy a little further. Most drivers are well aware that the effects of such a tie-up can persist long after the road crew responsible for it has stopped work and gone home for the night. Similarly, spiral density waves can continue to move through the disk even after the disturbance that originally produced them has long since subsided. According to spiral density wave theory, that is precisely what has happened in the Milky Way. Some disturbance in the past produced the wave, which has been moving through the Galactic disk ever since.

the creation of others. Computer simulations suggest that it is possible for the “wave” of star formation created in this manner to take on the form of a partial spiral and for the pattern to persist for some time. However, the process, sometimes known as **self-propagating star formation**, can produce only pieces of spirals, as are seen in some galaxies (Figure 23.19b). It apparently cannot produce the galaxy-wide spiral arms seen in other galaxies and present in our own. It may well be that there is more than one process at work in the spectacular spirals we see.

Again, time unfolds left to right, creating new stars over and over . . .



**▲ FIGURE 23.19 Self-Propagating Star Formation** (a) In this theory of the formation of spiral arms, the shock waves produced by the formation and later evolution of a group of stars provide the trigger for new rounds of star formation. Supernova explosions are used to illustrate the point here, but the formation of emission nebulae and planetary nebulae is also important. (b) This process may well be responsible for the partial spiral arms seen in some galaxies, such as NGC 4314, shown here in true color. (R. Gendler)

## Origin of Spiral Structure

An important question (but one that unfortunately is not answered by either of the two theories just described) is Where do these spirals come from? What was responsible for generating the density wave in the first place or for creating the line of newborn stars whose evolution drives the advancing spiral arm? Scientists speculate that (1) the gravitational effects of our satellite galaxies (the Magellanic Clouds, to be discussed in Chapter 24), (2) instabilities in the gas near the Galactic bulge, or (3) the possible barlike asymmetry within the bulge itself may have had a big enough influence on the disk to get the process going.

The first possibility is supported by growing evidence that many other spiral galaxies seem to have been affected by gravitational interactions with neighboring systems in the relatively recent past (see Chapter 24). However, many astronomers still regard the other two possibilities as equally likely. For example, they point to *isolated* spirals, whose structure clearly cannot be the result of an external interaction. The fact is that we still don't know for sure how galaxies—including our own—acquire such beautiful spiral arms.

### CONCEPT Check

- ✓ Why can't spiral arms simply be clouds of gas and young stars orbiting the Galactic center?


## 23.6 The Mass of the Milky Way Galaxy

We can measure our Galaxy's mass by studying the motions of gas clouds and stars in the Galactic disk. Recall from Chapter 2 that Newton's law of gravity (in the form of the modified version of Kepler's third law) connects the period, orbital size, and masses of any two objects in orbit around each other: ∞ (Sec. 2.8)

$$\text{total mass (solar masses)} = \frac{\text{orbital size (AU)}^3}{\text{orbital period (years)}^2}.$$

As we saw earlier, the distance from the Sun to the Galactic center is about 8 kpc, and the Sun's orbital period is 225 million years. Substituting these numbers into the preceding equation, we obtain a mass of  $(8000 \times 206,000)^3 / (225,000,000)^2$ , or almost  $9 \times 10^{10}$  solar masses—90 billion times the mass of our Sun!

But what mass have we just measured? When we performed the analogous calculation in the case of a planet orbiting the Sun, there was no ambiguity: The result of our calculation was the mass of the Sun. ∞ (More Precisely 2-2) However, the Galaxy's matter is not concentrated at the Galactic center (as the Sun's mass is concentrated at the center of the solar system); instead, Galactic matter is distributed over a large volume of space. Some of it lies inside the Sun's orbit (i.e., within 8 kpc of the Galactic center), and some lies outside, at



▲ **FIGURE 23.20 Weighing the Galaxy** The orbital speed of a star or gas cloud moving around the Galactic center is determined only by the mass of the Galaxy lying inside the orbit (within the gray-shaded sphere). Thus, to measure the Galaxy's total mass, we must observe objects orbiting at large distances from the center.

large distances from both the Sun and the center of the Galaxy. What portion of the Galaxy's mass controls the Sun's orbit? Isaac Newton answered this question three centuries ago: The Sun's orbital period is determined by the portion of the Galaxy that lies *within the orbit of the Sun* (Figure 23.20). This is the mass computed in the foregoing equation.

## Galactic Rotation

The Sun's motion around the Galactic center tells us that the total Galactic mass within the Sun's orbit is about 90 billion solar masses, but it says nothing about the mass lying outside that orbit—that is, more than 8 kpc from the center. To determine the mass of the Galaxy on larger scales, we must measure the orbital motion of stars and gas at greater distances from the Galactic center. Astronomers have found that the most effective way to do this is to make radio observations of gas in the Galactic disk, because radio waves are relatively unaffected by interstellar absorption and allow us to probe to great distances, far beyond the Sun's orbit. On the basis of these studies, radio astronomers have determined our Galaxy's rotation rate at various distances from the Galactic center. The resultant plot of rotation speed versus distance from the center (Figure 23.21) is called the Galactic **rotation curve**.


Knowing the Galactic rotation curve, we can now repeat our earlier calculation to compute the total mass that lies within any given distance from the Galactic center. We find, for example, that the mass within about 15 kpc from the center—the volume defined by the globular clusters and the known spiral structure—is roughly  $2 \times 10^{11}$  solar masses, about twice the mass contained within the Sun's orbit. Does the distribution of matter in the Galaxy “cut off” beyond 15 kpc, where the luminosity drops off sharply? Surprisingly, it does not.

Newton's laws of motion predict that if all of the mass of the Galaxy were contained within the edge of the visible structure, then the orbital speed of stars and gas beyond 15 kpc would decrease with increasing distance from the Galactic center, just as the orbital speeds of the planets diminish as we move outward from the Sun. The dashed line in Figure 23.21 indicates what the rotation curve would look like in that case. However, the true rotation curve is quite different: Far from falling off at larger distances, it *rises* slightly, out to the limits of our measurement capabilities. This slight rise implies that the amount of mass contained within successively larger radii continues to grow beyond the orbit of the Sun, apparently out to a distance of at least 40 or 50 kpc.

According to the equation presented at the beginning of this section, the amount of mass within 40 kpc is approximately  $6 \times 10^{11}$  solar masses. Since  $2 \times 10^{11}$  solar masses lie within 15 kpc of the Galactic center, we have to conclude that at least twice as much mass lies *outside* the luminous part of our Galaxy—the part made up of stars, star clusters, and spiral arms—as lies inside!

## Dark Matter

On the basis of these observations of the Galactic rotation curve, astronomers now regard the luminous portion of the Milky Way Galaxy—the region outlined by the globular clusters and by the spiral arms—as merely the “tip of the Galactic iceberg.” Our Galaxy is in reality very much larger. The luminous region is surrounded by an extensive, invisible **dark halo**, which dwarfs the inner halo of stars and globular clusters and extends well beyond the 15-kpc radius once thought to represent the limit of our Galaxy. But what is the composition of



▲ **FIGURE 23.21 Galaxy Rotation Curve** The rotation curve for the Milky Way Galaxy plots rotation speed against distance from the Galactic center. The dashed curve is the rotation curve expected if the Galaxy “ended” abruptly at a radius of 15 kpc, the limit of most of the known spiral structure. The fact that the red curve does not follow this dashed line, but instead stays well above it, indicates that additional unseen matter must be beyond that radius.

this dark halo? We do not detect enough stars or interstellar matter to account for the mass that our computations tell us must be there. We are inescapably drawn to the conclusion that most of the mass in our Galaxy exists in the form of invisible **dark matter**, which we presently do not understand.

Note, incidentally, that even the “visible” portion of our Galaxy contains substantial amounts of dark matter. The total mass of stars and gas within 15 kpc of the center is estimated, from direct measurements of the luminosities of stars and radio emission from the interstellar medium, to be around  $6 \times 10^{10}$  solar masses. Most of this mass lies in the disk. Comparing this mass with that inferred from the Galactic rotation curve, we see that, even inside this luminous region, dark matter accounts for roughly two-thirds of the Galaxy’s total mass.

The term *dark* here does not refer just to matter that is undetectable in visible light: The material has (so far) escaped detection at *all* wavelengths, from radio to gamma rays. Only by its gravitational pull do we know of its existence. Dark matter is not hydrogen gas (atomic or molecular), nor is it made up of ordinary stars. Given the amount of matter that must be accounted for, we would have been able to detect it with present-day instruments if it were in either of those forms. Its nature and its consequences for the evolution of galaxies and the universe are among the most important questions in astronomy today.

Many candidates have been suggested for this dark matter, although none is proven. Stellar-mass black holes may supply some of the unseen mass, but given that they are the evolutionary products of (relatively rare) massive stars, it is unlikely that there could be enough of them to hide large amounts of Galactic matter. ∞ (Sec. 22.8)


Currently among the strongest “stellar” contenders are brown dwarfs—low-mass prestellar objects that never reached the point of core nuclear burning—white dwarfs, and faint, low-mass red dwarfs. ∞ (Secs. 19.3, 20.3) In the jargon of the field, these objects are collectively known as *M*Assive Compact Halo Objects, or MACHOs for short. In principle, they could exist in great numbers throughout the Galaxy, yet would be exceedingly hard to see because they are so faint.

*Hubble Space Telescope* observations of globular clusters seem to argue against at least the last of the three possibilities listed for MACHOs. Figure 23.22 shows a *Hubble* image of a relatively nearby globular cluster—one close enough that very faint red dwarfs could have been detected if any existed. The *Hubble* data suggest that there is a cutoff at about 0.2 solar

mass, below which stars form much less frequently than had previously been supposed. As a result, stars with very low mass may be unexpectedly rare, at least in the Galactic halo.

A radically different alternative is that the dark matter is made up of exotic *subatomic particles* that pervade the entire universe. To account for the properties of dark matter, these particles must have mass (to produce the observed gravitational effects), but also must interact hardly at all with “normal” matter (because otherwise we would be able to see them). One class of candidate particles satisfying these requirements has been dubbed *Weakly Interacting Massive Particles*, or WIMPs. Many astrophysicists think that such “dark-matter particles” could have been produced in abundance during the very earliest moments of our universe. If they survived to the present day, there might be enough of them to account for all the dark matter apparently out there. We will discuss this possibility and its far-reaching implications in more detail in Chapter 27. These ideas are hard to test, however, because these particles would necessarily be very difficult to detect. Several detection experiments on Earth have been attempted, so far without success.

A few astronomers have proposed a very different explanation for the “dark matter problem,” suggesting that its resolution may lie not in the nature of dark matter, but rather in a modification to Newton’s law of gravity that increases the gravitational force on very large (galactic) scales, doing away with the need for dark matter in the first place. We emphasize that the vast majority of scientists do *not* accept



▲ **FIGURE 23.22 Missing Red Dwarfs** Sensitive visible-light observations with the *Hubble Space Telescope* have apparently ruled out faint red-dwarf stars as candidates for dark matter. The object shown here, the globular cluster 47 Tucanae, is one of many regions searched in the Milky Way. The inset is a high-resolution *Hubble* image of part of the cluster. The red dwarfs that would be expected if they existed in sufficient numbers to account for the dark matter in the Galaxy are not found. (The red stars that are seen are giants.) (AAT; NASA)

this view in any way. However, the very fact that such an unorthodox solution has even been proposed is a testament to our current level of uncertainty. Dark matter is one of the great unsolved mysteries in astronomy today.


## The Search for Stellar Dark Matter

Recently, researchers have obtained insight into the distribution of stellar dark matter by using a key element of Albert Einstein's theory of general relativity: the prediction that a beam of light can be deflected by a gravitational field, which has already been verified in the case of starlight that passes close to the Sun.  $\infty$  (Sec. 22.6, *More Precisely 22-1*) The effect is small in the case of light grazing the Sun, but it has the potential for making distant and otherwise invisible stellar objects observable from Earth. Here's how.

Imagine looking at a distant star as a faint foreground object (a MACHO, such as a brown or white dwarf) happens to cross your line of sight. As illustrated in Figure 23.23, the

intervening object deflects a little more starlight than usual toward you, resulting in a temporary, but quite substantial, *brightening* of the distant star. In some ways, the effect is like the focusing of light by a lens, so the process is known as **gravitational lensing**. The foreground object is referred to as a *gravitational lens*. The amount of brightening and the duration of the effect depend on the mass, distance, and speed of the lensing object. Typically, the apparent brightness of the background star increases by a factor of two to five for a period of several weeks. Thus, even though the foreground object cannot be seen directly, its effect on the light of the background star makes it detectable. (In Chapter 25, we will encounter other instances of gravitational lensing in the universe, but on very much larger scales.)

Of course, stars are very small compared with the distance scale of the Galaxy, and the probability that one star will pass almost directly in front of another, as seen from Earth, is extremely low. But by observing millions of stars every few days over a period of years (using automated



**▲ FIGURE 23.23 Gravitational Lensing** (a) Gravitational lensing by a faint foreground object (such as a brown dwarf) can temporarily cause a background star to brighten significantly, providing a means of detecting otherwise invisible stellar dark matter. (b) These two images show the brightening of a star during a lensing event, this one implying that a massive, but unseen, object passed in front of the unnamed star at the center of the two boxes imaged 6 months apart. (AURA)

telescopes and high-speed computers to reduce the burden of coping with so much data), astronomers have been able to see enough of these events to estimate the amount of stellar dark matter in the Galactic halo. The technique represents an exciting new means of probing the structure of our Galaxy. The observations are consistent with lensing by low-mass white dwarfs and suggest that such stars could account for a significant fraction—perhaps as much as 20 percent—but apparently *not* all, of the dark matter inferred from dynamical studies.

Bear in mind, though, that the identity of the dark matter is not necessarily an all-or-nothing proposition. It is perfectly conceivable—and, in fact, most astronomers think it likely—that more than one type of dark matter exists. For example, it is quite possible that most of the dark matter in the inner (visible) parts of galaxies is in the form of brown dwarfs and very low mass stars, whereas the dark matter farther out may be primarily in the form of exotic particles. We will return to this perplexing problem in later chapters, when we discuss some theories of how galaxies form and evolve, and how matter in the universe may have come into being.

### PROCESS OF SCIENCE Check

✓ The nature of dark matter particles is unknown, yet most scientists regard these particles as the best solution to the dark matter problem. How do you think these statements square with the experimental scientific method presented in Section 1.2?

## 23.7 The Galactic Center


Theory predicts that the Galactic bulge should be densely populated with billions of stars, with the highest densities found closest to the Galactic center. However, we are unable to see this central region of our Galaxy—the interstellar medium in the Galactic disk shrouds what otherwise would be a stunning view. Figure 23.24 shows the optical view we do have of the part of the Milky Way toward the Galactic center, in the general direction of the constellation Sagittarius. Here, the Galactic plane is nearly vertical.

Observations at other wavelengths allow us to peer more deeply into the congested central regions of our Galaxy. The inset to Figure 23.24 is an adaptive-optics infrared image of the innermost parsec. ∞ (Sec. 5.4) It shows a dense central cluster containing roughly 1 million stars. That's a stellar density some 10 *million* times greater than in our solar neighborhood, high enough that stars must experience frequent close encounters and even collisions with one another.

Over the past two decades, combined radio, infrared, and X-ray observations have allowed astronomers to paint a detailed—and intriguing—picture of the Galactic center. They reveal complex structure on many scales, and violent activity in our Galaxy's core.


### Galactic Activity

Figure 23.25(a) is an infrared view of part of Figure 23.24, with the Galactic plane now horizontal. On this scale, infrared radiation has been detected from what appear to be huge clouds



**Interactive** **FIGURE 23.24 Galactic Center** Photograph of stellar and interstellar matter in the direction of the Galactic center. Because of heavy obscuration, even the largest optical telescopes can see no farther than one-tenth the distance to the center. To connect with previous figures, the M8 nebula can be seen at extreme top center. ∞ (Sec. 18.2) The field is roughly 10° vertically across and is a continuation of the bottom part of Figure 18.5. The overlaid box outlines the location of the center of our Galaxy. The inset at right shows an adaptive-optics infrared view of the dense stellar cluster surrounding the Galactic center, whose very core is indicated by the twin arrows. (AURA; ESO)





**▲ FIGURE 23.25 Galactic Center Close-Up** (a) An infrared image around the center of our Galaxy (within the white box) shows many bright stars packed into a relatively small volume. The average density of matter in this boxed region is estimated to be about a million times that in the solar neighborhood. (b) The central portion of our Galaxy, as observed in the radio part of the spectrum, shows a region about 100 pc across surrounding the Galactic center (which lies within the orange-yellow blob at bottom right). The long-wavelength radio emission cuts through the Galaxy's dust, providing a view of matter in the immediate vicinity of the Galaxy's center. (c) This *Chandra* image shows the relation of a hot supernova remnant (red) and Sgr A\*, the suspected black hole at the very center of our Galaxy. (d) The spiral pattern of radio emission arising from Sagittarius A itself suggests a rotating ring of matter only a few parsecs across. All images are false-color, since they lie outside the visible spectrum. (SST; NRAO; NASA)

rich in dust. Radio observations indicate a ring of molecular gas nearly 400 pc across, containing hundreds of thousands of solar masses of material and rotating around the Galactic center at about 100 km/s. The origin of this ring is unclear, although researchers think that the gravity of the Galaxy's central rotating bar may deflect gas from farther out into the dense central regions.

Higher-resolution radio observations reveal further structure on smaller scales. Figure 23.25(b) shows a region called Sagittarius A. (The name simply means that it is the

brightest radio source in the constellation Sagittarius.) It lies at the center of the boxed region in Figure 23.24 and Figure 23.25(a)—and, we think, at the center of our Galaxy. On a scale of about 25 pc, extended filaments can be seen. Their presence suggests to many astronomers that strong *magnetic fields* operate in the vicinity of the center, creating structures similar in appearance to (but much larger than) those observed on the active Sun. ∞ (Sec. 16.5)

On even smaller scales (Figure 23.25c), *Chandra* observations indicate an extended region of hot X-ray-emitting gas,

apparently associated with a supernova remnant, in addition to many other individual bright X-ray sources. And within that lies a rotating ring or disk of molecular gas only a few parsecs across, with streams of matter spiraling inward toward the center (shown, again in the radio, in Figure 23.25d). Note that the scale of this disk is comparable to that of the dense central cluster shown in the inset to Figure 23.24.

What could cause all this activity? An important clue comes from the Doppler broadening of infrared spectral lines emitted from the central swirling whirlpool of gas. [∞ \(Sec. 4.5\)](#) The extent of the broadening indicates that the gas is moving very rapidly. In order to keep this gas in orbit, whatever is at the center must be extremely massive—more than a million solar masses. Given the twin requirements of large mass and small size, a leading contender is a *supermassive black hole*. [∞ \(Sec. 22.8\)](#)

The hole itself is not the source of the energy, of course. Instead, the vast accretion disk of matter drawn toward the hole by the enormous gravity emits the energy as it falls in, just as we saw (on a much smaller scale) in Chapter 22 when we discussed X-ray emission from neutron stars and stellar-mass black holes. [∞ \(Secs. 22.3, 22.8\)](#) The strong magnetic fields, thought to be generated within the accretion disk as matter spirals inward, may act as “particle accelerators,” creating extremely high-energy particles detected on Earth as *cosmic rays*. In the late 1990s, the *Compton Gamma Ray Observatory* found indirect evidence for a fountain of high-energy particles, possibly produced by violent processes close to the event horizon, gushing outward from the hole into the halo more than a thousand parsecs beyond the Galactic center. [∞ \(Sec. 5.7\)](#) Astronomers have reason to suspect that similar events are occurring at the centers of many other galaxies.

## The Central Black Hole


Astronomers have identified a candidate for the supermassive black hole at the Galactic center. At the very heart of Sagittarius A, is a remarkable object with the odd-sounding name **Sgr A\*** (pronounced “saj ay star”). By the standards of the active galaxies to be studied in Chapter 24, this compact **Galactic nucleus** is not particularly energetic. Still, radio observations made during the past two decades, along with more recent X- and gamma-ray observations, suggest that it is nevertheless a pretty violent place. Its total energy output (at all wavelengths) is estimated to be  $10^{33}$  W, more than a million times that of the Sun.

VLBI observations using radio telescopes arrayed from Hawaii to Massachusetts imply that Sgr A\* cannot be much larger than 10 AU, and it is probably a good deal smaller than that. [∞ \(Sec. 5.6\)](#) This size is consistent with the view that the energy source is a massive black hole. Figure 23.26 is perhaps the strongest evidence to date supporting the black-hole picture. It shows a high-resolution infrared image of the innermost 0.04 pc (or 8000 AU across) near the Galactic center, centered on Sgr A\*. Using advanced adaptive-optics techniques on the Keck telescopes and the VLT, U.S. and European researchers have created the first-ever diffraction-limited (0.05” resolution) images of the region. [∞ \(Sec. 5.4\)](#)


Remarkably, the image quality is good enough that the *proper motions* of several of the stars—their orbits around the Galactic center—can clearly be seen. The inset shows a series of observations of one of the brightest stars—called S2—over a 10-year period. The motion is consistent with an orbit around a massive object at the location of Sgr A\*, in accordance with Newton’s laws of motion. [∞ \(Sec. 2.7\)](#) The solid curve on the figure shows the elliptical orbit that best fits the observations: a 15-year orbit with a semimajor axis of 950 AU, corresponding

(from Kepler’s third law, as modified by Newton) to a central mass of approximately 4 million solar masses. The small size of the central object is very clearly demonstrated by the motion of another star in the group (S16), whose extremely eccentric orbit brings it within just 45 AU of the center.

Other observations, using adaptive-optics infrared-imaging techniques, have revealed a bright source very close to Sgr A\* that seems to



◀ **FIGURE 23.26 Orbits Near the Galactic Center** This extremely close-up map of the Galactic center (left) was obtained by infrared adaptive optics, resulting in an ultra-high-resolution image of the innermost 0.1 pc of the Milky Way. The inset shows the orbit of the innermost star in the frame, labeled S2, between 1992 and 2003. The solid line shows the best-fitting orbit for S2 around a black hole of 4 million solar masses, located at Sgr A\* (marked with a cross). (ESO)



**▲ FIGURE 23.27 Galactic Center Zoom** This series of artist’s conceptions of the Galactic center depicts each frame increasing in resolution by a factor of 10. Frame (a) is the same scene as Figure 23.16. Frame (f) is a rendition of a vast whirlpool within the innermost 0.5 parsec of our Galaxy. The data imaged in Figure 23.25 do not closely match these artistic renderings because the Figure 23.25 view is parallel to the Galactic disk—along the line of sight from the Sun to the Galactic center—whereas these six paintings portray a simplified view perpendicular to the disk, while progressively zooming down onto that disk. (*Adapted from L. Chaisson*)

vary with a 10-minute period.  $\infty$  (Sec. 5.4) The source could be a hot spot on the accretion disk that circles the purported hole. Note that, even with the large mass just mentioned, if Sgr A\* is a genuine black hole, the size of its event horizon is still only 0.02 AU.  $\infty$  (Sec. 22.5) Such a small region, 8 kpc away, is currently unresolvable, although radio astronomers are hopeful that improving VLBI techniques will allow them to “see” the event horizon and study the surrounding accretion disk within the next decade.

Figure 23.27 places these findings into a simplified perspective. Each frame is centered on the Galaxy’s core, increasing in resolution by a factor of 10 from one to the next. Frame (a) renders the Galaxy’s overall shape, as painted in Figure 23.16. This frame measures about 50 kpc across. Frame (b) spans a distance of 5 kpc from side to side and is nearly filled by the Galactic bar and the great sweep of the innermost spiral arm. Moving in to a 500-pc span, frame (c) depicts part of the 400-pc ring of gas mentioned earlier. The dark blobs represent giant molecular clouds and the pink patches emission nebulae associated with star formation within those clouds. In parts (b) and (c), the artist has peeled away the bright bulge, enabling us to “see” better into the central regions.

In frame (d), at 50 pc, a pinkish (thin, warm) region of ionized gas surrounds the reddish (denser, warmer) heart of the Galaxy, corresponding approximately to the images shown in Figure 23.25(b) and (c). The energy responsible for this vast ionized region comes from frequent supernovae and other violent phenomena in the Galactic center. Recent multi-wavelength observations reveal that this intense activity has


blasted huge (10 kpc long) magnetized jets of high-energy particles out of the Galactic center, roughly perpendicular to the disk. The total energy in the jets exceeds that of a typical supernova by about a factor of a million. Also shown in frame (d) are numerous young dense star clusters, further evidence of recent bursts of star formation near the Galactic center.

Frame (e), spanning 5 pc, depicts the central star cluster (diluted in the painting for clarity) and the surrounding star-forming ring, along with the tilted, spinning whirlpool of hot ( $10^4$  K) gas surrounding the center of our Galaxy. The innermost part of this gigantic whirlpool is shown in frame (f), in which a swiftly spinning, white-hot disk of gas with temperatures in the millions of kelvins nearly engulfs the central black hole (marked by as the black dot). Two rings of stars, possibly the remains of disrupted star clusters, can also be seen. The black hole itself, and the stellar orbits shown in Figure 23.26, are far too small to be pictured on this scale.

The last decade has seen an explosion in our knowledge of the innermost few parsecs of our Galaxy, and astronomers are working hard to decipher the clues hidden within its invisible radiation. Still, we are only now beginning to appreciate the full complexity of this strange new realm deep in the heart of the Milky Way.

### CONCEPT Check

- ✓ What is the most likely explanation for the energetic events observed at the Galactic center?



**The Big Question** How big is the Milky Way? How well do we know the size, shape, and mass of our colossal home in the cosmos? In recent years, astronomers have upsized our system of stars, gas, and dark matter. Its total mass has been upgraded by nearly a factor of ten, its extended halo might reach halfway to the nearest galaxy, and its dark matter outweighs its normal matter by at least a factor of five. Even so, we may yet have seriously underestimated the scale of this grand system.

## Chapter Review

### SUMMARY

**1** A **galaxy** (p. 578) is a huge collection of stellar and interstellar matter isolated in space and bound together by its own gravity. Because we live within it, the **Galactic disk** (p. 578) of our own Milky Way Galaxy appears as a broad band of light across the sky, a band called the Milky Way. Near the center, the disk thickens into the **Galactic bulge** (p. 578). The disk is surrounded by a roughly spherical **Galactic halo** (p. 578) of old stars and star clusters. Like many others visible in the sky, our Galaxy is a **spiral galaxy** (p. 580). Disk and halo stars differ in their spatial distributions, ages, colors, and orbital motion. The luminous portion of our Galaxy has a diameter of about 30 kpc. In the vicinity of the Sun, the Galactic disk is about 300 pc thick.




**2** The halo can be studied using **variable stars** (p. 581), whose luminosity changes with time. **Pulsating variable stars** (p. 581) vary in brightness in a repetitive and predictable way. Of particular importance to astronomers are **RR Lyrae variables** (p. 581) and **Cepheid variables** (p. 581). All RR Lyrae stars

have roughly the same luminosity. For Cepheids, the luminosity can be determined using the **period–luminosity relationship** (p. 583). Knowing the luminosity, astronomers can apply the inverse-square law to determine the distance. The brightest Cepheids can be seen at distances of millions of parsecs, extending the cosmic distance ladder well beyond our own Galaxy. In the early 20th century, Harlow Shapley used RR Lyrae stars to determine the distances to many of the Galaxy's globular clusters and found that they have a roughly spherical distribution in space, but the center of the sphere lies far from the Sun. The center of their distribution is close to the **Galactic center** (p. 585), about 8 kpc away.




**3** Stars and gas within the Galactic disk move on roughly circular orbits around the Galactic center. Stars in the halo and bulge move on largely random




three-dimensional orbits that pass repeatedly through the disk plane but have no preferred orientation.

**4** The Galactic halo lacks gas and dust, so no new stars are forming there. All halo stars are old. The gas-rich disk is the site of current star formation and contains many young stars. Halo stars appeared early on, before the Galactic disk took shape, when there was no preferred orientation for their orbits. After the disk formed, stars born there inherited its overall spin and so move on circular orbits in the Galactic plane.




**5** Radio observations clearly reveal the extent of our Galaxy's **spiral arms** (p. 592), regions of the densest interstellar gas where star formation is taking place. The spirals cannot be “tied” to the disk material, as the disk's differential rotation would have wound them up long ago. Instead, they may be **spiral density waves** (p. 592) that move through the disk, triggering star formation as they pass by. Alternatively, the spirals may arise from **self-propagating star formation** (p. 594), when shock waves produced by the formation and evolution of one generation of stars trigger the formation of the next.




**6** The Galactic **rotation curve** (p. 596) plots the orbital speed of matter in the disk against distance from the Galactic center. By applying Newton's laws of motion, astronomers can determine

the mass of the Galaxy. They find that the Galactic mass continues to increase beyond the radius defined by the globular clusters and the spiral structure we observe. Our Galaxy, like many others, has an invisible **dark halo** (p. 596) containing far more mass than can be accounted for in the form of luminous matter. The **dark matter** (p. 597) making up these dark halos is of unknown composition. Leading candidates include low-mass stars and exotic subatomic particles. Recent attempts to detect stellar dark matter have used the fact that a faint foreground object can occasionally pass in front of a more distant star, deflecting the star's light and causing its apparent brightness to increase temporarily. This deflection is called **gravitational lensing** (p. 598).



**7** Astronomers working at infrared and radio wavelengths have uncovered evidence for energetic activity within a few parsecs of the Galactic center. The leading explanation is that a black hole roughly 4 million times more massive than the Sun resides there. The hole lies at the center of a dense star cluster containing millions of stars, which is in turn surrounded by a star-forming disk of molecular gas. The observed activity is thought to be powered by accretion onto the black hole, as well as by supernova explosions in the cluster surrounding it.



**MasteringAstronomy**® For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- POS** What evidence do we have that we live in a disk galaxy?
- Why is it difficult to map out our Galaxy from our vantage point on Earth?
- LO1 POS** What do globular clusters tell us about our Galaxy and our place within it?
- How are Cepheid variables used in determining distances? How far away can they be used?
- LO2** What important discoveries were made early in the 20th century by using RR Lyrae variables?
- Of what use is radio astronomy in the study of Galactic structure?
- LO3** Contrast the motions of disk and halo stars.
- LO4** What do the red stars in the Galactic halo tell us about the history of the Milky Way?
- LOS POS** Explain why galactic spiral arms are thought to be regions of recent and ongoing star formation.
- What is self-propagating star formation?
- LO6** What does the rotation curve of our Galaxy tell us about the Galaxy's total mass?
- POS** What is the evidence for dark matter in the Galaxy? Describe some candidates for Galactic dark matter.
- What is gravitational lensing, and can astronomers use it to search for dark matter?
- Why can't optical astronomers easily study the center of our Galaxy?
- LO7 POS** Why do astronomers think that a supermassive black hole lies at the center of the Milky Way Galaxy?

## Conceptual Self-Test: Multiple Choice

- Most of the bright stars in our Galaxy are located in the Galactic (a) center; (b) bulge; (c) halo; (d) disk.
- VIS** According to Figure 23.7 (“Period–Luminosity Plot”), a Cepheid variable star with luminosity 1000 times that of

the Sun has a pulsation period of roughly (a) 1 day; (b) 3 days; (c) 10 days; (d) 50 days.

3. Globular clusters are found mainly in (a) the Galactic center; (b) the Galactic disk; (c) spiral arms; (d) the Galactic halo.
4. Shapley measured the distances to globular clusters by using (a) trigonometric parallax; (b) a comparison of the absolute and apparent magnitudes of variable stars; (c) spectroscopic parallax; (d) radar ranging.
5. In the Milky Way Galaxy, our Sun is located (a) near the Galactic center; (b) about halfway out from the center; (c) at the outer edge; (d) in the halo.
6. A telescope searching for newly formed stars would make the most discoveries if it were pointed (a) directly away from the Galactic center; (b) perpendicular to the Galactic disk; (c) within a spiral arm; (d) between spiral arms.
7. The first stars that formed in the Milky Way now (a) have random orbits in the halo; (b) orbit in the Galactic plane; (c) orbit closest to the Galactic center; (d) orbit in the same direction as the Milky Way spins.
8. **VIS** Figure 23.21 (“Galaxy Rotation Curve”) says (a) the Galaxy rotates like a solid body; (b) far from the center, the Galaxy rotates more slowly than expected based on the light we see; (c) far from the center, the Galaxy rotates more rapidly than expected based on the light we see; (d) there is no matter beyond 15 kpc from the Galactic center.
9. Most of the mass of the Milky Way exists in the form of (a) stars; (b) gas; (c) dust; (d) dark matter.
10. The main evidence for a black hole at the Galactic center is that (a) stars near the center are disappearing; (b) no stars can be seen in the vicinity of the center; (c) stars near the center are orbiting some unseen object; (d) the Galaxy rotates faster than astronomers expect.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

1. ● Calculate the angular diameter of a prestellar nebula of radius 100 AU lying 100 pc from Earth. Compare this with the roughly  $6^\circ$  diameter of the Andromeda Galaxy (Figure 23.2a).
2. ●● How close would the nebula in the previous question have to be in order to have the same angular diameter as Andromeda? Calculate the apparent magnitude of the central star if it had a luminosity 10 times that of the Sun.
3. ● What is the greatest distance at which an RR Lyrae star of absolute magnitude 0 could be seen by a telescope capable of detecting objects as faint as 20th magnitude?
4. ● A typical Cepheid variable is 100 times brighter than a typical RR Lyrae star. How much farther away than RR Lyrae stars can Cepheids be used as distance-measuring tools?
5. ●● Calculate the proper motion (in arc seconds per year) of a globular cluster with a transverse velocity (relative to the Sun) of 200 km/s and a distance of 3 kpc. Do you think that this motion is measurable?
6. ●● Calculate the total mass of the Galaxy lying within 20 kpc of the Galactic center if the rotation speed at that radius is 240 km/s.
7. ●● Using the data presented in Figure 23.21, estimate the distance from the Galactic center at which matter takes (a) 100 million years and (b) 500 million years to complete one orbit.
8. ●● Material at an angular distance of  $0.2''$  from the Galactic center is observed to have an orbital speed of 1200 km/s. If the Sun’s distance to the Galactic center is 8 kpc, and the material’s orbit is circular and is seen edge-on, calculate the radius of the orbit and the mass of the object around which the material is orbiting.

## Activities

### Collaborative

1. Construct your own version of the Messier Catalog, listing the names, types, and coordinates of each of the 110 Messier objects. Plot the celestial coordinates—right ascension and declination, akin to latitude and longitude on Earth—of all objects. Color-code them to distinguish among emission nebulae, open star clusters, globular clusters, and galaxies. What do you notice about the distributions of these objects on the sky? It may help to sketch the location of the Milky Way (more research!) on your plot. Why do you think the galaxies seem to avoid the Galactic plane?

### Individual

1. Observe the Andromeda Galaxy, M31. It’s the most distant object visible to the naked eye, but don’t expect to see anything like Figure 23.2(a)! To locate M31, find Polaris, the pole star, and the constellations Cassiopeia and Andromeda. Follow a line from Polaris through the second “V” in the “W” of Cassiopeia and continue south. That line will pass through M31 before you reach the northern arc of stars in Andromeda. To the unaided eye, from all but the darkest sites, only its nucleus will be visible, looking like a slightly fuzzy star. Use binoculars or a wide-angle eyepiece to view the galaxy and its disk. Switch to higher magnification to view the nucleus and the small satellite galaxies M32, just to its south, and M110, to the northwest.



# Galaxies

## BUILDING BLOCKS OF THE UNIVERSE

As our field of view expands to truly cosmic scales, the focus of our studies shifts dramatically. Planets become inconsequential, stars themselves mere points of hydrogen consumption. Now entire galaxies become the “atoms” from which the universe is built—distant realms completely unknown to scientists just a century ago.

We know of literally millions of galaxies beyond our own. All are vast, gravitationally bound assemblages of stars, gas, dust, dark matter, and radiation separated from us by almost incomprehensibly large distances. Most galaxies are smaller than the Milky Way, some comparable in size, a few much larger. Many seem “normal,” like our own Galaxy—the collective output of billions of stars. But some are sites of explosive events far more energetic than anything ever witnessed in our Galaxy. Such “active” galaxies are probably powered by supermassive black holes.

**The Big Picture** Light collected tonight from the most distant galaxies was emitted by those objects long before Earth even formed. Racing for billions of years across the darkened realms of the cosmos, a minute fraction of their radiation is now intercepted by our telescopes and spacecraft. Captured in the many images of this book, that radiation tells us about not only the properties of faraway galaxies but also a few things about the history of our Galaxy and the universe in which we live.

**LEFT:** Active galaxies, such as this one cataloged as NGC 1316, are much more energetic than normal galaxies like our Milky Way. This is a double image, mixing optical light (acquired by the *Hubble Space Telescope* in Earth orbit) with radio emission (captured by the Very Large Array in New Mexico). At center (in white) is a giant, visible elliptical galaxy that extends about 100,000 light-years across and is probably devouring its small northern neighbor. The result is the complex radio emission (in orange), called Fornax A, spanning more than a million light-years end to end. (*INRAO/STScI*)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 List the basic properties of normal galaxies.
- 2 Outline the distance-measurement techniques that enable astronomers to map the universe beyond the Milky Way.
- 3 Describe how galaxies clump into groups and clusters.
- 4 State Hubble’s law and explain how it is used to derive distances to the most remote objects in the observable universe.
- 5 Specify the basic differences between active and normal galaxies.
- 6 Describe some important features of active galaxies.
- 7 Explain what drives the central engine thought to power all active galaxies.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.



## 24.1 Hubble's Galaxy Classification

Figure 24.1 shows a vast expanse of space lying about 100 million pc from Earth. Almost every patch or point of light in this figure is a separate galaxy—hundreds can be seen in just this one photograph. Over the years, astronomers have accumulated similar images of many millions of galaxies. We begin our study of these enormous accumulations of matter simply by considering their appearance on the sky.

Seen through even a small telescope, images of galaxies look distinctly nonstellar. They have fuzzy edges, and many are quite elongated—not at all like the sharp, point-like images normally associated with stars. Although it is difficult to tell from the photograph, some of the blobs of light in Figure 24.1 are spiral galaxies like the Milky Way Galaxy and Andromeda. Others, however, are definitely not spirals—no disks or spiral arms can be seen. Even when we take into account their different orientations in space, galaxies do *not* all look the same.


The American astronomer Edwin Hubble was the first to categorize galaxies in a comprehensive way. Working with the then recently completed 2.5-m optical telescope on Mount Wilson in California in 1924, he classified the galaxies he saw into four basic types—*spirals*, *barred spirals*,

*ellipticals*, and *irregulars*—solely on the basis of their visual appearance. Many modifications and refinements have been incorporated over the years, but the basic **Hubble classification scheme** is still widely used today.


### Spirals

We saw several examples of **spiral galaxies** in Chapter 23—for example, our own Milky Way Galaxy and our neighbor Andromeda. ∞ (Sec. 23.1) All galaxies of this type contain a flattened galactic disk in which spiral arms are found, a central galactic bulge with a dense nucleus, and an extended halo of faint, old stars. ∞ (Sec. 23.3) The stellar density (i.e., the number of stars per unit volume) is greatest in the *galactic nucleus*, at the center of the bulge. However, within this general description, spiral galaxies exhibit a wide variety of shapes, as illustrated in Figure 24.2.

In Hubble's scheme, a spiral galaxy is denoted by the letter S and classified as type a, b, or c according to the size of its central bulge. Type Sa galaxies have the largest bulges, Type Sc the smallest. The tightness of the spiral pattern is quite well correlated with the size of the bulge (although the correspondence is not perfect). Type Sa spiral galaxies tend to have tightly wrapped, almost circular,



**▲ FIGURE 24.1 Coma Cluster** (a) A collection of many galaxies, each consisting of hundreds of billions of stars. Called the Coma Cluster, this group of galaxies lies about 100 million pc from Earth. (The blue spiked object at top right is a nearby star; virtually every other object in this image is a galaxy.) (b) A recent *Hubble Space Telescope* image of part of the cluster. (AURA; NASA)



▲ **FIGURE 24.2 Spiral Galaxy Shapes** Variation in shape among spiral galaxies. As we progress from type Sa to Sb to Sc, the bulges become smaller and the spiral arms tend to become less tightly wound. (NASA; D. Malin/AAT)

spiral arms, Type Sa galaxies typically have more open spiral arms, and Type Sc spirals often have a loose, poorly defined spiral structure. The arms also tend to become more “knotty,” or clumped, in appearance as the spiral pattern becomes more open.

The bulges and halos of spiral galaxies contain large numbers of reddish old stars and globular clusters, similar to those observed in our own Galaxy and in Andromeda. Most of the light from spirals, however, comes from A-through G-type stars in the galactic disk, giving these galaxies an overall whitish glow. We assume that thick disks exist, too, but their faintness makes this assumption hard to confirm—the thick disk in the Milky Way contributes only a percent or so of our Galaxy’s total light. ∞ (Sec. 23.3)



Like the disk of the Milky Way, the flat disks of typical spiral galaxies are rich in gas and dust. Type Sc galaxies contain the most interstellar matter, Sa galaxies the least. The 21-cm radio radiation emitted by spirals betrays the presence of the gas, and obscuring dust lanes are clearly visible in many systems (see Figures 24.2b and c). ∞ (Sec. 18.4) Stars are forming within the spiral arms, which contain numerous emission nebulae and newly formed O- and B-type stars. ∞ (Secs. 18.2, 23.5) The arms appear bluish because of the presence of bright blue O- and B-type stars there. The photo of the Sc galaxy NGC 2997 shown in Figure 24.2(c) reveals the preponderance of interstellar

gas, dust, and young blue stars tracing the spiral pattern particularly clearly. Spirals are not necessarily young galaxies, however: Like our own Galaxy, they are simply rich enough in interstellar gas to provide for continued stellar birth.

Most spirals are not seen face-on, as they are shown in Figure 24.2. Many are tilted with respect to our line of sight, making their spiral structure hard to discern. However, we do not need to see spiral arms to classify a galaxy as a spiral. The presence of the disk, with its gas, dust, and newborn stars, is sufficient. For example, the galaxy shown in Figure 24.3 is classified as a spiral because of the clear line of obscuring dust seen along its midplane. (Incidentally, this relatively nearby galaxy was another of the “nebulae” figuring in the Shapley–Curtis debate discussed in Chapter 23. ∞ (Sec. 23.2) The visible dust lane was interpreted by Curtis as an obscuring “ring” of material, leading him to suggest that our Galactic plane might contain a similar feature.)

## Barred Spirals

A variation of the spiral category in Hubble’s classification scheme is the **barred-spiral galaxy**. Barred spirals differ from ordinary spirals mainly by the presence of an elongated “bar” of stellar and interstellar matter passing through the




◀ **FIGURE 24.3 Sombrero Galaxy** The Sombrero Galaxy (M104), a spiral system seen edge-on, has a dark band composed of interstellar gas and dust. The large size of this galaxy’s central bulge marks it as type Sa, even though its spiral arms cannot be seen from our perspective. The inset shows this galaxy in the infrared part of the spectrum, highlighting its dust content in false-colored pink. (NASA)

center and extending beyond the bulge, into the disk. The spiral arms project from near the ends of the bar rather than from the bulge (their origin in normal spirals). Barred spirals are designated by the letters SB and are subdivided, like the ordinary spirals, into categories SBa, SBb, and SBc, depending on the size of the bulge. Again, like ordinary spirals, the tightness of the spiral pattern is correlated with the size of the

bulge. Figure 24.4 shows the variation among barred-spiral galaxies. In the case of the SBc category, it is often hard to tell where the bar ends and the spiral arms begin.


Frequently, astronomers cannot distinguish between spirals and barred spirals, especially when a galaxy happens to be oriented with its galactic plane nearly edge-on toward Earth, as in Figure 24.3. Because of the physical



(a) NGC 1300 Type SBa

(b) NGC 1365 Type SBb

(c) NGC 6872 Type SBc



▲ **FIGURE 24.4 Barred-Spiral Galaxy Shapes** Variation in shape among barred-spiral galaxies from SBa to SBc is similar to that for the spirals in Figure 24.2, except that here the spiral arms begin at either end of a bar through the galactic center. In frame (c), the bright star is a foreground object in our own Galaxy; the object at top center is another galaxy that is probably interacting with NGC 6872. (NASA; D. Malin/AAT; ESO)

and chemical similarities of spiral and barred-spiral galaxies, some researchers do not even bother to distinguish between them. Others, however, regard the differences in their structures as very important, arguing that these differences suggest basic dissimilarities in the ways the two types of galaxies formed and evolved.

Based on all available evidence, the Milky Way seems to be a barred spiral galaxy, most likely of type SBb. [∞ \(Sec. 23.3\)](#)

## Ellipticals

Unlike the spirals, **elliptical galaxies** have no spiral arms and, in most cases, no obvious galactic disk—in fact, other than possessing a dense central nucleus, they often exhibit little internal structure of any kind. As with spirals, the stellar density increases sharply in the nucleus. Denoted by the letter E, these systems are subdivided according to how elliptical they appear on the sky. The most circular are designated E0, slightly flattened systems are labeled E1, and so on, all the way to the most elongated ellipticals, of type E7 (Figure 24.5).

Note that an elliptical galaxy's Hubble type depends both on its intrinsic three-dimensional shape and on its orientation relative to the line of sight. A spherical galaxy, a cigar-shaped galaxy seen end-on, and a disk-shaped galaxy seen face-on, would all appear to be circular on the sky and be classified as E0. It can be difficult to decipher a galaxy's true shape solely from its visual appearance.


There is a large range in both the size and the number of stars contained in elliptical galaxies. The largest elliptical galaxies are much larger than our own Milky Way Galaxy. These *giant ellipticals* can range up to hundreds of kiloparsecs across and contain trillions of stars. At the other extreme, *dwarf*

*ellipticals* may be as small as 1 kpc in diameter and contain fewer than a million stars. Their many differences suggest to astronomers that giant and dwarf ellipticals represent distinct galaxy classes, with quite dissimilar formation histories and stellar content. The dwarfs are by far the most common type of ellipticals, outnumbering their brighter counterparts by about 10 to 1. However, most of the *mass* that exists in the form of elliptical galaxies is contained in the larger systems.


The absence of spiral arms is not the only difference between spirals and ellipticals: Most ellipticals also contain little or no cool gas and dust. The 21-cm radio emission from neutral hydrogen gas is, with few exceptions, completely absent, and no obscuring dust lanes are seen. In most cases, there is no evidence of young stars or ongoing star formation. Like the halo of our own Galaxy, ellipticals are made up mostly of old, reddish, low-mass stars. Also, as in the halo of our Galaxy, the orbits of stars in ellipticals are disordered, exhibiting little or no overall rotation; objects move in all directions, not in regular, circular paths as in our Galaxy's disk. Ellipticals differ from our Galaxy's halo in at least one important respect, however: X-ray observations reveal large amounts of very *hot* (several million kelvins) interstellar gas distributed throughout their interiors, often extending well beyond the visible portions of the galaxies (Figure 24.5a,b).

Some giant ellipticals are exceptions to many of these general statements, as they have been found to contain disks of gas and dust in which stars are forming. Astronomers think that these systems may be the results of collisions among gas-rich galaxies (see Section 25.2). Indeed, galactic collisions may have played an important role in determining the appearance of many of the systems we observe today.

Intermediate between the E7 ellipticals and the Sa spirals in the Hubble classification is a class of galaxies




**▲ FIGURE 24.5 Elliptical Galaxy Shapes** (a) The E2 elliptical galaxy M49 is nearly circular in appearance. (b) M84 is slightly more elongated and classified as E3. Both galaxies lack spiral structure, and neither shows evidence of cool interstellar dust or gas, although each has an extensive X-ray halo of hot gas that extends far beyond the visible portion of the galaxy. (c) M110 is a dwarf elliptical companion to the much larger Andromeda Galaxy. (AURA; SAO; R. Gendler)



◀ **FIGURE 24.6 S0 Galaxies** (a) S0 (or lenticular) galaxies contain a disk and a bulge, but no interstellar gas and no spiral arms. Their properties are intermediate between E7 ellipticals and Sa spirals. (b) SB0 galaxies are similar to S0 galaxies, except for a bar of stellar material extending beyond the central bulge. (*Palomar/Caltech*)

that show evidence of a thin disk and a flattened bulge, but that contain no gas and no spiral arms. Two such objects are shown in Figure 24.6. These galaxies are known as **S0 galaxies** if no bar is evident and **SB0 galaxies** if a bar is present. They are also known as *lenticular* galaxies, because of their lens-shaped appearance. They look a little like spirals whose dust and gas have been stripped away, leaving behind just a stellar disk. Observations in recent years have shown that many normal elliptical galaxies have faint disks within them, like the S0 galaxies. As with the S0s, the origin of these disks is uncertain, but some researchers suspect that S0s and ellipticals may be closely related.



## Irregulars

The final class of galaxies identified by Hubble is a catch-all category—**irregular galaxies**—so named because their visual appearance excludes them from the other categories just discussed. Irregulars tend to be rich in interstellar matter and young, blue stars, but they lack any regular structure, such as well-defined spiral arms or central bulges. They are divided into two subclasses: Irr I galaxies and Irr II galaxies. The Irr I galaxies often look like misshapen spirals.

Irregular galaxies tend to be smaller than spirals, but somewhat larger than dwarf ellipticals. They typically contain between  $10^8$  and  $10^{10}$  stars. The smallest such galaxies are called *dwarf irregulars*. As with elliptical galaxies, the dwarf type is the most common. Dwarf ellipticals and dwarf irregulars occur in approximately equal numbers and together make up the vast majority of galaxies in the universe. They are often found close to a larger “parent” galaxy.

Figure 24.7 shows the **Magellanic Clouds**, a famous pair of Irr I galaxies that orbit the Milky Way Galaxy. They

◀ **FIGURE 24.7 Magellanic Clouds** The Magellanic Clouds are prominent features of the night sky in the Southern Hemisphere. Named for the 16th-century Portuguese explorer Ferdinand Magellan, whose around-the-world expedition first brought word of these fuzzy patches of light to Europe, these dwarf irregular galaxies orbit our Galaxy and accompany it on its trek through the cosmos. (a) The Clouds’ relationship to one another in the southern sky reveals both the Small (b) and the Large (c) Magellanic Cloud to have distorted, irregular shapes. (*Mount Stromlo & Siding Spring Observatory; Harvard College Observatory; Royal Observatory, Edinburgh*)

are shown to proper scale in Figure 23.16. Studies of Cepheid variables within the Clouds show them to be approximately 50 kpc from the center of our Galaxy.  $\infty$  (Sec. 23.2) The Large Cloud contains about 6 billion solar masses of material and is a few kiloparsecs across. Both Clouds contain lots of gas, dust, and blue stars (and the recent, well-documented supernova discussed in *Discovery 21-1*), indicating ongoing star formation. Both also contain many old stars and several old globular clusters, so we know that star formation has been going on in them for a very long time.


Radio studies hint at a possible bridge of hydrogen gas connecting the Milky Way to the Magellanic Clouds, although more observational data are still needed to establish this link beyond doubt. It is possible that the tidal force of the Milky Way tore a stream of gas from the Clouds the last time their orbits brought them close to our Galaxy. Of course, gravity works both ways, and many researchers reason that the forces exerted by the Clouds may in turn be responsible for distorting our Galaxy, warping and thickening the outer parts of the Galactic disk.  $\infty$  (Sec. 23.5)

The much rarer Irr II galaxies (Figure 24.8), in addition to their irregular shape, have other peculiarities, often exhibiting a distinctly explosive or filamentary appearance. Their appearance once led astronomers to suspect that violent events had occurred within them. However, it now seems more likely that, in some (but probably not all) cases, we are seeing the result of a close encounter or collision between two previously “normal” systems.

## The Hubble Sequence

Table 24.1 summarizes the basic characteristics of the various types of galaxies. When he first developed his classification scheme, Hubble arranged the galaxies into the “tuning fork” diagram shown in Figure 24.9. The variation in types across the diagram, from ellipticals to spirals to irregulars, is often referred to as the *Hubble sequence*.


Hubble's primary aim in creating this diagram was to indicate similarities in appearance among galaxies. However, he also regarded the tuning fork as an evolutionary sequence from left to right, with E0 ellipticals evolving into flatter ellipticals and S0 systems and ultimately forming disks and spiral arms. Indeed, Hubble's terminology referring to ellipticals as “early-type” and spirals as “late-type” galaxies is still widely used today. However, as far as modern astronomers can tell, there is no direct evolutionary connection of this sort along the Hubble sequence. Isolated normal galaxies do *not* evolve from one type to another. Spirals are not ellipticals that have grown arms, nor are ellipticals spirals that have somehow expelled their star-forming disks. Some astronomers do suspect that bars may be transient features and that barred-spiral galaxies may therefore evolve into ordinary spirals, but, in general, astronomers know of no simple parent-child relationship among Hubble types.



(a) NGC 4449



(b) NGC 1569



**▲ FIGURE 24.8 Irregular Galaxy Shapes** (a) The strangely shaped galaxy NGC 4449 resides within a group of galaxies nearly 4 million parsecs away; it's likely that its peculiar shape results from interactions with its companions that have caused huge rearrangements of its stars, gas, and dust. (b) The galaxy NGC 1569 seems to show an explosive appearance, probably the result of a recent galaxywide burst of star formation. (NASA)

**TABLE 24.1** Galaxy Properties by Type


	<b>Spiral/Barred Spiral (S/SB)</b>	<b>Elliptical* (E)</b>	<b>Irregular (Irr)</b>
Shape and structural properties	Highly flattened disk of stars and gas, containing spiral arms and thickening central bulge. Sa and SBa galaxies have the largest bulges, the least obvious spiral structure, and roughly spherical stellar halos. SB galaxies have an elongated central “bar” of stars and gas.	No disk. Stars smoothly distributed through an ellipsoidal volume ranging from nearly spherical (E0) to very flattened (E7) in shape. No obvious substructure other than a dense central nucleus.	No obvious structure. Irr II galaxies often have “explosive” appearances.
Stellar content	Disks contain both young and old stars; halos consist of old stars only.	Contain old stars only.	Contain both young and old stars.
Gas and dust	Disks contain substantial amounts of gas and dust; halos contain little of either.	Contain hot X-ray–emitting gas, little or no cool gas and dust.	Very abundant in gas and dust.
Star formation	Ongoing star formation in spiral arms.	No significant star formation during the last 10 billion years.	Vigorous ongoing star formation.
Stellar motion	Gas and stars in disk move in circular orbits around the galactic center; halo stars have random orbits in three dimensions.	Stars have random orbits in three dimensions	Stars and gas have highly irregular orbits.

\*As noted in the text, some giant ellipticals appear to be the result of collisions between gas-rich galaxies and are exceptions to many of the statements listed here.

However, the key word in the previous paragraph is *isolated*. As described in Section 25.2, there is now strong observational evidence that collisions and tidal interactions *between* galaxies are commonplace and that these encounters are the main physical processes driving the evolution of galaxies. We will return to this important subject in Chapter 25.

### CONCEPT Check

✓ In what ways are large spirals like the Milky Way and Andromeda not representative of galaxies as a whole?



▲ **FIGURE 24.9 Galactic “Tuning Fork”** The placement of the four basic types of galaxies—ellipticals, spirals, barred spirals, and irregulars—in Hubble’s “tuning-fork” diagram is suggestive of evolution, but this galaxy classification scheme has no known physical meaning. As we will see in Chapter 25, galaxies do evolve, but not (in either direction) along the “Hubble sequence” defined by this figure.

## 24.2 The Distribution of Galaxies in Space

Now that we have seen some of their basic properties, let us ask how galaxies are spread through the expanse of the universe beyond the Milky Way. Galaxies are not distributed uniformly in space. Rather, they tend to clump into still larger agglomerations of matter. As we will see, this uneven distribution is crucial in determining both their appearance and their evolution. As always in astronomy, our understanding hinges on our ability to tell how far away an object lies. We therefore begin by looking more closely at the means used by astronomers to measure distances to galaxies.

### Extending the Distance Scale


Astronomers estimate that some 40 billion galaxies as bright as (or brighter than) our own exist in the observable universe. Some reside close enough for the Cepheid variable technique to work—astronomers have detected and measured the periods of Cepheids in galaxies as far away as 25 Mpc (see Figure 24.10).  $\infty$  (Sec. 23.2) However, some galaxies contain no Cepheid stars (can you think of some reasons that this might be?), and, in any case, most known galaxies lie much farther away than 25 Mpc. Cepheid variables in very distant galaxies simply cannot be observed well enough, even through the world's most sensitive telescopes,

to allow us to measure their apparent brightnesses and periods. To extend our distance-measurement ladder, we must find some new class of object to study.

One way in which researchers have tackled this problem is through observations of **standard candles**—intrinsically bright, easily recognizable astronomical objects whose luminosities are confidently known. The basic idea is very simple. Once an object is identified as a standard candle—by its appearance or by the shape of its light curve, say—its luminosity can be estimated. Comparison of the luminosity with the apparent brightness then gives the object's distance and, hence, the distance to the galaxy in which it resides.  $\infty$  (Sec. 17.2) Note that, apart from the way in which the luminosity is determined, the Cepheid variable technique relies on identical reasoning.


To be most useful, a standard candle must (1) have a *well-defined luminosity*, so that the uncertainty in estimating its brightness is small, and (2) be *bright enough* to be seen at large distances. Over the years, astronomers have explored the use of many types of objects as standard candles—novae, emission nebulae, planetary nebulae, globular clusters, Type I (carbon-detonation) supernovae, and even entire galaxies have been employed. Not all have been equally useful, however: Some have larger intrinsic spreads in their luminosities than others, making them less reliable for measuring distances.

In recent years, planetary nebulae and Type I supernovae have proved particularly reliable as standard candles.  $\infty$  (Secs. 20.3, 21.3) The latter have remarkably consistent peak luminosities and are very bright, allowing them to be identified and measured out to distances of many hundreds of megaparsecs. The small luminosity spread of Type I supernovae is a direct consequence of the circumstances in which these violent events occur. As discussed in Chapter 21, an accreting white dwarf explodes when it reaches the well-defined critical mass at which carbon fusion begins.  $\infty$  (Sec. 21.3) The magnitude of the explosion is relatively insensitive to the details of how the white dwarf formed or how it subsequently reached critical mass, with the result that all such supernovae have quite similar



**▲ FIGURE 24.10 Cepheid in Virgo** This sequence of six snapshots chronicles the periodic changes in a Cepheid variable star in the spiral galaxy M100, a member of the Virgo Cluster of galaxies. The Cepheid appears at the center of each inset, taken at the different times indicated during 1994. The star looks like a square because of the high magnification of the digital CCD camera—we are seeing individual pixels of the image. The 24th-magnitude star varies by about a factor of two in brightness every 7 weeks. (NASA)





**Narrated FIGURE 24.11 Galaxy Rotation** A galaxy's rotation causes some of the radiation it emits to be blueshifted and some to be redshifted. From a distance, when all the radiation from the galaxy is combined into a single beam and analyzed spectroscopically, the redshifted and blueshifted components produce a broadening of the galaxy's spectral lines. The amount of broadening is a direct measure of the rotation speed of the galaxy, such as NGC 4603 shown here. (NASA)

properties.\* Thus, when a Type I supernova is observed in a distant galaxy (we assume that it occurs *in the galaxy*, not in the foreground), astronomers can quickly obtain an accurate estimate of the galaxy's distance.

An important alternative to standard candles was discovered in the 1970s, when astronomers found a close correlation between the rotational speeds and the luminosities of spiral galaxies within a few tens of megaparsecs of the Milky Way Galaxy. Rotation speed is a measure of a spiral galaxy's total mass, so it is perhaps not surprising that this property should be related to luminosity. [∞ \(Sec. 23.5\)](#) What is surprising, though, is how tight the correlation is. The **Tully-Fisher relation**, as it is now known (after its discoverers), allows us to obtain a remarkably accurate estimate of a spiral galaxy's luminosity simply by observing how fast the galaxy rotates. As usual, comparing the galaxy's (true) luminosity with its (observed) apparent brightness yields its distance.

To see how the method is used, imagine that we are looking edge-on at a distant spiral galaxy and observing one particular emission line, as illustrated in Figure 24.11. Radiation from the side of the galaxy where matter is generally approaching us is blueshifted by the Doppler effect. Radiation from the other side, which is receding from us, is redshifted by a similar amount. The overall effect is that line radiation from the galaxy is “smeared out,” or broadened, by the galaxy's rotation. The faster the rotation, the greater the amount of broadening (see Figure 4.18 for the stellar equivalent). By

\*Recall from Chapter 21 that a Type II supernova also occurs when a growing stellar core—this time at the center of a massive star—reaches a critical mass. [∞ \(Sec. 21.2\)](#) However, the outward appearance of the explosion can be significantly modified by the amount of stellar material through which the blast wave must travel before it reaches the star's surface, resulting in a greater spread in observed luminosities. [∞ \(Discovery 21-1\)](#)

measuring the amount of broadening, we can therefore determine the galaxy's rotation speed. Once we know that, the Tully-Fisher relation tells us the galaxy's luminosity.

The particular line normally used in these studies actually lies in the radio part of the spectrum. It is the 21-cm line of cold, neutral hydrogen in the galactic disk. [∞ \(Sec. 18.4\)](#) This line is used in preference to optical lines because (1) optical radiation is strongly absorbed by dust in the disk under study and (2) the 21-cm line is normally very narrow, making the broadening easier to observe. In addition, astronomers often use *infrared*, rather than optical, luminosities, to avoid absorption problems caused by dust, both in our own Galaxy and in others.

The Tully-Fisher relation can be used to measure distances to spiral galaxies out to about 200 Mpc, beyond which the line broadening becomes increasingly difficult to measure accurately. A somewhat similar connection, relating line broadening to a galaxy's *diameter*, exists for elliptical galaxies. Once the galaxy's diameter and angular size are known, its distance can be computed from elementary geometry. [∞ \(More Precisely 1-2\)](#) These methods bypass many of the standard candles often used by astronomers and so provide independent means of determining distances to faraway objects.



As indicated in Figure 24.12, standard candles and the Tully-Fisher relation form the fifth and sixth rungs of our cosmic distance ladder, introduced in Chapter 1 and expanded in Chapters 17 and 23. [∞ \(Secs. 1.6, 17.1, 17.6, 23.2\)](#) In fact, they stand for perhaps a dozen or so related, but separate, techniques that astronomers have employed in their quest to map out the universe on large scales. Just as with the lower rungs, we calibrate the properties of these new techniques by using distances measured by more local means. In this way, the distance-measurement process “bootstraps” itself to greater and greater distances. However, at the same time, the errors and

► **FIGURE 24.12 Extragalactic Distance Ladder** An inverted pyramid summarizes the distance techniques used to study different realms of the universe. The techniques shown in the bottom four rungs of the ladder—radar ranging, stellar parallax, spectroscopic parallax, and variable stars—take us as far as the nearest galaxies. To go farther, we must use other techniques—for example, the Tully-Fisher relation and the use of standard candles—based on distances determined by the four lowest techniques.


uncertainties in each step accumulate, so the distances to the farthest objects are the least well known.

### Clusters of Galaxies

Figure 24.13 sketches the locations of all the known major astronomical objects within about 1 Mpc of the Milky Way. Our Galaxy appears with its dozen or so satellite galaxies—including the two Magellanic Clouds discussed earlier and a small companion (labeled “Sagittarius dwarf” in the figure) lying almost within our own Galactic plane. The Andromeda Galaxy, lying 800 kpc



▲ **FIGURE 24.13 Local Group** The Local Group is made up of more than 50 galaxies within approximately 1 Mpc of our Milky Way Galaxy. Only a few are spirals; most are dwarf elliptical or irregular galaxies, only some of which are shown here. Spirals are colored blue, ellipticals pink, and irregulars white—all of them depicted roughly to scale. The inset map (top right) shows the Milky Way in relation to some of its satellite galaxies. The photographic insets (top left) show two well-known neighbors of the Andromeda Galaxy (M31): the spiral galaxy M33 and the dwarf elliptical galaxy M32 (also visible in Figure 23.2a, a larger-scale view of the Andromeda system). (M. BenDaniel; NASA)



◀ **FIGURE 24.14 Virgo Cluster**

In the central region of the Virgo Cluster of galaxies, about 17 Mpc from Earth, many large spiral and elliptical galaxies can be seen. The inset shows several galaxies surrounding the giant elliptical M86. An even bigger elliptical galaxy, M87, noted at the bottom, will be discussed later in the chapter. (M. BenDaniel; AURA)


from us, is also shown, surrounded by satellites of its own. Two of Andromeda's galactic neighbors are shown in insets. M33 is a spiral, and M32 is a dwarf elliptical, easily seen in Figure 23.2(a) below and to the right of Andromeda's central bulge.

All told, nearly 55 galaxies are known to populate our Galaxy's neighborhood. Three of them (the Milky Way, Andromeda, and M33) are spirals; the remainder are dwarf irregulars and dwarf ellipticals. Together, these galaxies form the **Local Group**—a new level of structure in the universe above the scale of our Galaxy. As indicated in Figure 24.13, the Local Group's diameter is a little over 1 Mpc. The Milky Way Galaxy and Andromeda are by far its largest members, and most of the smaller galaxies are gravitationally bound to one or the other of them. The combined gravity of the galaxies in the Local Group binds them together, like stars in a star cluster, but on a millionfold larger scale. More generally, a collection of galaxies held together by their mutual gravitational attraction is called a **galaxy cluster**.

▶ **FIGURE 24.15 Distant Galaxy Cluster** The galaxy cluster Abell 1689 contains huge numbers of galaxies and resides nearly a billion parsecs from Earth. Virtually every patch of light in this photograph is a separate galaxy. With the most powerful telescopes, astronomers can now discern, even at this great distance, spiral structure in some of the galaxies. We also see many galaxies colliding—some tearing matter from one another, others merging into single systems. (NASA)

Moving beyond the Local Group, the next large concentration of galaxies we come to is the *Virgo Cluster* (Figure 24.14), named after the constellation in which it is found. Lying some 17 Mpc from the Milky Way, the Virgo Cluster does not contain a mere 50 galaxies, however. Rather, it houses more than 2500 galaxies, bound by gravity into a tightly knit group about 3 Mpc across.

Wherever we look in the universe we find galaxies, and most galaxies are members of groups or clusters of galaxies. In practice, the distinction between a “group” and “cluster” is mainly a matter of convention. Groups generally contain only a few bright galaxies (such as the Milky Way and Andromeda) and are quite irregular in shape, whereas large, “rich” clusters like Virgo may contain thousands of individual galaxies distributed fairly smoothly in space. The Coma cluster, shown in Figure 24.1 and lying approximately 100 Mpc away, is another example of a rich cluster. Figure 24.15




is a long-exposure photograph of a much more distant rich cluster, lying about 700 Mpc from Earth. A sizeable minority of galaxies (possibly as many as 40 percent) are not members of any group or cluster, but are apparently isolated systems, moving along through intercluster space. (For simplicity, we will use the term “cluster” below to refer to any gravitationally bound collection of galaxies, large or small.)

We will return to the large-scale distribution of matter in the universe in Chapters 25 and 26.

### PROCESS OF SCIENCE Check

- ✓ What are some of the problems astronomers encounter in measuring the distances to faraway galaxies?



◀ **FIGURE 24.16 Galaxy Spectra** Optical spectra, tallied at left, of several galaxies shown at right. Both the extent of the redshift (denoted by the horizontal red arrows) and the distance from the Milky Way Galaxy to each galaxy (numbers in center column) increase from top to bottom. The vertical yellow arrows denote a pair of dark absorption lines in the observed spectra. The many vertical white lines at the top and bottom of each spectrum are laboratory references. (Adapted from Palomar/Caltech)


## 24.3 Hubble's Law

Now that we have seen some basic properties of galaxies throughout the universe, let's turn our attention to the large-scale *motions* of galaxies and galaxy clusters. Within a galaxy cluster, individual galaxies move more or less randomly. You might expect that, on even larger scales, the clusters themselves would also have random, disordered motion—some clusters moving this way, some that. In fact, that is not the case: On the largest scales, galaxies and galaxy clusters alike move in a very *ordered* way.

### Universal Recession

In 1917, the American astronomer Vesto M. Slipher, working under the direction of Percival Lowell, reported that virtually every spiral galaxy he observed had a redshifted spectrum—it was *receding* from our Galaxy. ∞ (Sec. 3.5) It is now known that, except for a few nearby systems, *every* galaxy takes part in a general motion away from us in all directions. Individual galaxies that are not part of galaxy clusters are steadily receding. Galaxy clusters, too, have an overall recessional motion, although their individual member galaxies move randomly with respect to one another. (Consider a jar full of fireflies that has been thrown into the air. The fireflies within the jar, like the galaxies within the cluster, have random motions due to their individual whims, but the jar as a whole, like the galaxy cluster, has some directed motion as well.)

Figure 24.16 shows the optical spectra of several galaxies, arranged in order of increasing distance from the Milky Way Galaxy. The spectra are redshifted, indicating that the associated galaxies are receding, and the extent of the redshift increases from top to bottom in the figure. There is a connection between Doppler shift and distance:



◀ **FIGURE 24.17 Hubble's Law**  
Plots of recessional velocity against distance (a) for the galaxies shown in Figure 24.16 and (b) for numerous other galaxies within about 1 billion pc of Earth.

The greater the distance, the greater the redshift. This trend holds for nearly all galaxies in the universe. (Two galaxies within our Local Group, including Andromeda, and a few galaxies in the Virgo Cluster display blueshifts and so are moving toward us, but this results from their local motions within their parent clusters—recall the fireflies in the jar.)

Figure 24.17(a) shows recessional velocity plotted against distance for the galaxies of Figure 24.16. Figure 24.17(b) is a similar plot for some more galaxies within about 1 billion parsecs of Earth. Plots like these were first made by Edwin Hubble in the 1920s and now bear his name: *Hubble diagrams*. The data points generally fall close to a straight line, indicating that the rate at which a galaxy recedes is *directly proportional* to its distance from us. This rule is called **Hubble's law**. We can construct such a diagram for any collection of galaxies, provided that we can determine their distances and velocities. The universal recession described by the Hubble diagram is sometimes called the *Hubble flow*.

The recessional motions of the galaxies prove that the cosmos is neither steady nor unchanging on the largest scales. The universe (actually, *space itself*—see Section 26.2) is expanding! But let's be clear on just *what* is expanding and what is not. Hubble's law does not mean that humans, Earth, the solar system, or even individual galaxies and galaxy clusters are physically increasing in size. These groups of atoms, rocks, planets, stars, and galaxies are held together by their own internal forces and are not themselves getting bigger. Only the largest framework of the universe—the vast distances separating the galaxy clusters—is expanding.

To distinguish recessional redshift from redshifts caused by motion *within* an object—for example, galactic orbits within a cluster or explosive events in a galactic nucleus—the redshift resulting from the Hubble flow is called the **cosmological redshift**. Objects that lie so far

away that they exhibit a large cosmological redshift are said to be at *cosmological distances*—distances comparable to the scale of the universe itself.

Hubble's law has some dramatic implications. If nearly all galaxies show recessional velocity according to Hubble's law, then doesn't that mean that they all started their journey from a single point? If we could run time backward, wouldn't all the galaxies fly back to this one point, perhaps the site of some violent event in the remote past? The answer is yes—but not in the way you might expect! In Chapters 26 and 27, we will explore the ramifications of the Hubble flow for the past and future evolution of our universe. For now, however, we set aside its cosmic implications and use Hubble's law simply as a convenient distance-measuring tool.

## Hubble's Constant

The constant of proportionality between recessional velocity and distance in Hubble's law is known as **Hubble's constant**, denoted by the symbol  $H_0$ . The data shown in Figure 24.17 then obey the equation

$$\text{recessional velocity} = H_0 \times \text{distance}.$$

The value of Hubble's constant is the slope of the straight line—recessional velocity divided by distance—in Figure 24.17(b). Reading the numbers off the graph, we get roughly 70,000 km/s divided by 1000 Mpc, or 70 km/s/Mpc (kilometers per second per megaparsec, the most commonly used unit for  $H_0$ ). Astronomers continually strive to refine the accuracy of the Hubble diagram and the resulting estimate of  $H_0$  because Hubble's constant is one of the most fundamental quantities of nature; it specifies the rate of expansion of the entire cosmos.

Hubble's original value for  $H_0$  was about 500 km/s/Mpc, far higher than the currently accepted value. This overestimate was due almost entirely to errors in the cosmic distance scale at the time, particularly the calibrations of Cepheid variables and standard candles. The measured value dropped rapidly as various observational errors were recognized and resolved and distance measurements became more reliable. Published estimates of  $H_0$  entered the "modern" range (within, say, 20 percent of the current value) in roughly the mid-1960s.

As measurement techniques have continued to improve, the uncertainty in the Hubble constant has steadily decreased to the point that now, early in the 21st century, all leading measurements of  $H_0$ , by a variety of different techniques—Tully-Fisher measurements, studies of Cepheid variables in the Virgo Cluster, and observations of standard candles, such as Type I supernovae—are remarkably consistent with one another. We will adopt a rounded-off value of  $H_0 = 70$  km/s/Mpc (a choice roughly in the middle of all recent results, and also in line with some precise cosmological measurements to be discussed in more detail in Chapter 27) as the best current estimate of Hubble's constant for the remainder of the text.

the infrared part of the spectrum! *More Precisely 24-1* discusses in more detail the meaning and interpretation of such large redshifts, apparently implying recessional velocities comparable to the speed of light. According to Hubble's law, the objects that exhibit these redshifts lie more than 9000 Mpc away from us, as close to the limits of the observable universe as astronomers have yet been able to probe.

The speed of light is finite. It takes time for light—or, for that matter, any kind of radiation—to travel from one point in space to another. The radiation that we now see from these most distant objects originated long ago. Incredibly, that radiation was emitted roughly 13 billion years ago (see Table 24.2), well before our planet, our Sun, and perhaps even our Galaxy came into being!


### CONCEPT Check

- ✓ How does the use of Hubble's law differ from the other extragalactic distance-measurement techniques we have seen in this text?

## Top of the Distance Ladder

Using Hubble's law, we can derive the distance to a remote object simply by measuring the object's recessional velocity and dividing by Hubble's constant. Hubble's law thus tops our inverted pyramid of distance-measurement techniques (Figure 24.18). This seventh method simply assumes that Hubble's law holds. If that assumption is correct, then Hubble's law enables us to measure great distances in the universe—so long as we can obtain an object's spectrum, we can determine how far away it is.

Many redshifted objects have recessional motions that are a substantial fraction of the speed of light. The most distant objects thus far observed in the universe—some young galaxies and quasars (Section 24.4)—have redshifts (fractional increases in wavelength) of around 8, meaning that their radiation has been stretched in wavelength not by just a few percent, as with most of the objects we have discussed, but *ninefold*. Their ultraviolet spectral lines are shifted all the way into



▲ **FIGURE 24.18 Cosmic Distance Ladder** Hubble's law tops the hierarchy of distance-measurement techniques. It is used to find the distances of astronomical objects all the way out to the limits of the observable universe.

## MORE PRECISELY 24-1

### Relativistic Redshifts and Look-Back Time

In discussing very distant objects, astronomers usually talk about their *redshifts* rather than their distances. Indeed, it is common for researchers to speak of an event occurring “at” a certain redshift—meaning that the light received today from that event is redshifted by the specified amount. Of course, because of Hubble’s law, redshift and distance are equivalent to one another. However, redshift is the preferred quantity because it is a directly observable property of an object, whereas distance is derived from redshift with the use of Hubble’s constant, whose value is not accurately known. (In Chapter 26 we will see another, much more fundamental, reason why astronomers favor the use of redshift in studies of the cosmos.)

The redshift of a beam of light is, by definition, the *fractional* increase in the wavelength of the light resulting from the recessional motion of the source.  $\infty$  (Sec. 3.5) Thus, a redshift of 1 corresponds to a *doubling* of the wavelength. From the formula for the Doppler shift given previously, the redshift of radiation received from a source moving away from us with speed  $v$  is given by

$$\begin{aligned} \text{redshift} &= \frac{\text{observed wavelength} - \text{true wavelength}}{\text{true wavelength}} \\ &= \frac{\text{recessional velocity, } v}{\text{speed of light, } c}. \end{aligned}$$

**EXAMPLE** Let’s illustrate this relationship with two examples, rounding the speed of light,  $c$ , to 300,000 km/s. A galaxy at a distance of 100 Mpc has a recessional speed (by Hubble’s law) of  $70 \text{ km/s/Mpc} \times 100 \text{ Mpc} = 7000 \text{ km/s}$ . Its redshift is therefore  $7000 \text{ km/s} \div 300,000 \text{ km/s} = 0.023$ . Conversely, an object that has a redshift of 0.05 has a recessional velocity of  $0.05 \times 300,000 \text{ km/s} = 15,000 \text{ km/s}$  and hence a distance of  $15,000 \text{ km/s} \div 70 \text{ km/s/Mpc} = 210 \text{ Mpc}$ .

Unfortunately, although the foregoing equation is correct for low speeds, it does not take into account the effects of relativity. As we saw in Chapter 22, the rules of everyday physics have to be modified when speeds begin to approach the speed of light.  $\infty$  (Discovery 22-1) The formula for the Doppler shift is no exception. In particular, although the formula is valid for speeds much less than the speed of light, when  $v = c$  the redshift is not unity, as the equation suggests, but is in fact *infinite*. That is, radiation received from an object moving away from us at nearly the speed of light is redshifted to almost infinite wavelength.

Thus, do not be alarmed to find that many galaxies and quasars have redshifts greater than unity. This does not mean that they are receding faster than light! It simply means that the preceding simple formula is not applicable. In fact, the real connection between redshift and distance is quite complex, requiring us to make key assumptions about the past history of the universe (see Chapter 26). In place of a formula, we can use Table 24.2, which presents a conversion chart relating redshift and distance. All of the values shown are based on reasonable assumptions and are usable even for large redshifts. We take Hubble’s constant to be 70 km/s/Mpc and assume a flat universe in which matter (mostly dark) contributes roughly 30 percent of the total density (see Section 26.6). The conversions in the table are used consistently throughout this text. The column headed “ $v/c$ ” gives equivalent recessional velocities based on the Doppler effect, taking relativity properly into account. Even though this is *not* the correct interpretation of the redshift (see Section 26.2), we include it here for comparison simply because it is so often quoted in the popular media.

Because the universe is expanding, the “distance” to a galaxy is not very well defined. Do we mean the distance to the galaxy when it emitted the light we see today, the present distance to the galaxy (as presented in the table, even though we do not see the galaxy as it is today), or some other, more appropriate measure? Largely because of this ambiguity, astronomers prefer to work in terms of a quantity known as the *look-back time* (shown

## 24.4 Active Galactic Nuclei

The galaxies described in Section 24.1—those falling into the various Hubble classes—are generally referred to as **normal galaxies**. As we have seen, their luminosities range from a million or so times that of the Sun for dwarf ellipticals and irregulars to more than a trillion solar luminosities for the largest giant ellipticals. For comparison, in round numbers, the luminosity of the Milky Way Galaxy is  $2 \times 10^{10}$  solar luminosities, or roughly  $10^{37}$  W.

In these last two sections we focus our attention on “bright” galaxies, conventionally taken to mean galaxies

with luminosities more than about  $10^{10}$  times the solar value. In these terms, our Galaxy is bright, but not abnormally so.

### Galactic Radiation

A substantial fraction of bright galaxies—perhaps as many as 40 percent—don’t fit well into the “normal” category. Their *spectra* differ significantly from those of their normal cousins, and their *luminosities* can be extremely large. Known collectively as **active galaxies**, they are of great interest to astronomers. The brightest among them are the most energetic objects known in the universe, and all may

in the last column of Table 24.2), which is simply how long ago an object emitted the radiation we see today. Astronomers talk frequently about redshifts and sometimes about look-back times, but they hardly ever talk of distances to high-redshift objects (and *never* about recession velocities, despite what you hear on the news!). Bear in mind, however, that redshift is the only unambiguously measured quantity in this discussion. Statements about “derived” quantities, such as distances and look-back times, all require that we make specific assumptions about how the universe has evolved with time.

For nearby sources, the look-back time is numerically equal to the distance in light-years: The light we receive tonight from a galaxy at a distance of 100 million light-years was emitted 100 million years ago. However, for more distant objects, the look-back time and the present distance in light-years differ because of the expansion of the universe, and the divergence increases dramatically with increasing redshift.

As a simple analogy, imagine an ant crawling across the surface of an expanding balloon at a constant speed of 1 cm/s relative to the balloon’s surface. After 10 seconds, the ant may think it has traveled a distance of 10 cm, but an outside observer with a tape measure will find that it is actually more than 10 cm from its starting point (measured along the surface of the balloon) because of the balloon’s expansion. In exactly the same way, the present distance to a galaxy with a given redshift depends on how the universe expanded in the past. For example, a galaxy now located 15 billion light-years from Earth was much closer to us when it emitted the light we now see. Consequently, its light has taken considerably less than 15 billion years—in fact, about 10 billion years—to reach us.

**TABLE 24.2** Redshift, Distance, and Look-Back Time

Redshift	v/c	Present Distance		Look-Back Time (millions of years)
		(Mpc)	( $10^6$ light-years)	
0.000	0.000	0	0	0
0.010	0.010	43	139	139
0.025	0.025	107	347	343
0.050	0.049	212	691	674
0.100	0.095	419	1370	1300
0.200	0.180	820	2670	2440
0.250	0.220	1010	3300	2950
0.500	0.385	1910	6210	5080
0.750	0.508	2680	8750	6650
1.000	0.600	3350	10,900	7820
1.500	0.724	4450	14,500	9420
2.000	0.800	5300	17,300	10,400
3.000	0.882	6520	21,300	11,600
4.000	0.923	7370	24,000	12,200
5.000	0.946	8000	26,100	12,600
6.000	0.960	8490	27,700	12,800
7.000	0.969	8890	29,000	13,000
8.000	0.976	9220	30,100	13,100
9.000	0.980	9500	31,000	13,200
10.000	0.984	9740	31,800	13,300
50.000	0.999	12,400	40,400	13,700
100.000	1.000	13,000	42,500	13,800
$\infty$	1.000	14,700	47,800	13,800


represent an important, if intermittent, phase of galactic evolution (see Section 25.4). At optical wavelengths, active galaxies often *look* like normal galaxies—familiar components such as disks, bulges, stars, and dark dust lanes can be identified. At other wavelengths, however, their unusual properties are much more apparent.

Most of a normal galaxy’s energy is emitted in or near the visible portion of the electromagnetic spectrum, much like the radiation from stars. Indeed, to a large extent, the light we see from a normal galaxy is just the accumulated light of its many component stars (once the effects of interstellar dust are taken into account), each described

approximately by a blackbody curve.  $\infty$  (Sec. 3.4) By contrast, as illustrated schematically in Figure 24.19, the radiation from active galaxies does *not* peak in the visible. Most active galaxies do emit substantial amounts of visible radiation, but far more of their energy is emitted at invisible wavelengths, both longer and shorter than those in the visible range. Put another way, the radiation from active galaxies is *inconsistent* with what we would expect if it were the combined radiation of myriad stars. Their radiation is said to be *nonstellar*.

Many luminous galaxies with nonstellar emission are known to be *starburst galaxies*—previously normal systems






▲ **FIGURE 24.19 Galaxy Energy Spectra** The energy emitted by a normal galaxy differs significantly from that emitted by an active galaxy. This plot illustrates the general spread of intensity for all galaxies of a particular type and does not represent any one individual galaxy.

currently characterized by widespread episodes of star formation, most likely as a result of interactions with a neighbor. The irregular galaxy NGC 1569 shown in Figure 24.8 is a prime example. We will study these important systems and their role in galaxy evolution in Chapter 25. For purposes of this text, however, we will use the term “active galaxy” to mean a system whose abnormal activity is related to violent events occurring in or near the galactic nucleus. The nuclei of such systems are called **active galactic nuclei**.

Even with this restriction, there is still considerable variation in the properties of galaxies, and astronomers have identified and cataloged a bewildering array of systems falling into the “active” category. For example, Figure 24.20 shows an active galaxy exhibiting both nuclear activity and widespread star formation, with a blue-tinted ring of newborn stars surrounding an extended 1-kpc-wide core of intense emission. Rather than attempting to describe the entire “zoo” of active galaxies, we will instead discuss three basic species: the energetic *Seyfert galaxies* and *radio galaxies* and the even more luminous *quasars*. Although these objects all lie toward the “high-luminosity” end of the active range and represent perhaps only a few percent of the total number of active galaxies, their properties will allow us to identify and discuss features common to active galaxies in general.

The association of galactic activity with the central nucleus is reminiscent of the discussion in Chapter 23 of the center of the Milky Way. ☞ (Sec. 23.7) In our own Galaxy, it seems clear that the activity in the nucleus is associated with the central *supermassive black hole*, whose presence is inferred from observations of stellar orbits in the innermost fraction of a parsec. As we will see, most astronomers think that basically the same thing is going on in the nuclei



▲ **FIGURE 24.20 Active Galaxy** This image of the galaxy NGC 7742 resembles a fried egg, with a ring of blue star-forming regions surrounding a very bright yellow core that spans about 1 kpc. This active galaxy combines star formation with intense emission from its central nucleus and lies roughly 24 Mpc away. (NASA)

of active galaxies and that “normal” and “active” galaxies may differ principally in the degree to which the nonstellar nuclear component of the radiation outshines the light from the rest of the galaxy. This is a powerful unifying theme for understanding the evolution of galaxies, and we will return to it in Chapter 25. For the remainder of this chapter, we concentrate on describing the properties of active galaxies and the black holes that power them.

## Seyfert Galaxies

In 1943, Carl Seyfert, an American optical astronomer studying spiral galaxies from Mount Wilson Observatory, discovered the type of active galaxy that now bears his name. **Seyfert galaxies** are a class of astronomical objects whose properties lie between normal galaxies and the most energetic active galaxies known.

Superficially, Seyferts resemble normal spiral galaxies (Figure 24.21a). Indeed, the stars in a Seyfert’s galactic disk and spiral arms produce about the same amount of visible radiation as do the stars in a normal spiral galaxy. However, most of a Seyfert’s energy is emitted from the galactic nucleus—the center of the overexposed white patch in the figure. The nucleus of a Seyfert galaxy is some 10,000 times brighter than the center of our own Galaxy. In fact, the brightest Seyfert nuclei are 10 times more energetic than the entire Milky Way.

Some Seyferts produce radiation spanning a broad range in wavelengths, from the infrared all the way through ultraviolet and even X-rays. However, the majority (about




▲ **FIGURE 24.21 Seyfert Galaxy** (a) The Circinus galaxy, a Seyfert with a bright compact core, lies some 4 Mpc away. It is one of the closest active galaxies. (b) This graph illustrates the irregular variations in luminosity of the Seyfert galaxy 3C 84 over three decades. These observations were made in the radio part of the electromagnetic spectrum; the optical and X-ray luminosities vary as well. (NASA; NRAO)

75 percent) emit most of their energy in the infrared. Scientists think that much of the high-energy radiation in these Seyferts is absorbed by dust in or near the nucleus and then reemitted as infrared radiation.

Seyfert spectral lines have many similarities to those observed toward the center of our own Galaxy. ∞ (Sec. 23.7) Some of the lines are very broad, most likely indicating rapid (5000 km/s or more) internal motion within the nuclei. ∞ (Sec. 4.5) However, not all of the lines are broad, and some Seyferts show no broad lines at all. In addition, their energy emission often varies in time (Figure 24.21b). A Seyfert's luminosity can double or halve within a fraction of a year. These rapid fluctuations in luminosity lead us to conclude that the source of energy emissions in Seyfert galaxies must be quite compact—simply put, as we saw in Chapter 22, an object cannot “flicker” in less time than radiation takes to cross it. ∞ (Sec. 22.4) The emitting region must therefore be less than 1 light-year across—an extraordinarily small region, considering the amount of energy emanating from it.

Together, the rapid time variability and large radio and infrared luminosities observed in Seyferts imply violent nonstellar activity in their nuclei. As just mentioned, this activity is most likely similar in *nature* to processes occurring at the center of our own Galaxy, but its *magnitude* is thousands of times greater than the comparatively mild events within our own Galaxy's heart. ∞ (Sec. 23.7)



(b)

## Radio Galaxies

As the name suggests, **radio galaxies** are active galaxies that emit large amounts of energy in the radio portion of the electromagnetic spectrum. They differ from Seyferts not only in the wavelengths at which they radiate, but also in both the appearance and the extent of their emitting regions.


Figure 24.22(a) shows the radio galaxy Centaurus A, which lies about 4 Mpc from Earth. Almost none of this galaxy's radio emission comes from a compact nucleus. Instead, the energy is released from two huge extended regions called **radio lobes**—roundish clouds of gas spanning about half a megaparsec and lying well beyond the visible galaxy.\* Undetectable in visible light, the radio lobes of radio galaxies are truly enormous. From end to end, they typically span more than 10 times the size of the Milky Way Galaxy, comparable in scale to the entire Local Group.

Figure 24.22(b) shows the relationship between the galaxy's visible, radio, and X-ray emissions. In visible light, Centaurus A is apparently a large E2 galaxy some 500 kpc in diameter, bisected by an irregular band of dust. Centaurus A is a member of a small cluster of galaxies, and numerical simulations suggest that this peculiar galaxy is probably the result of a collision between an elliptical galaxy and a smaller

\*The term “visible galaxy” is commonly used to refer to those components of an active galaxy that emit visible “stellar” radiation, as opposed to the nonstellar and invisible “active” component of the galaxy's emission.



(a)




spiral galaxy about 500 million years ago. In the crowded confines of a cluster, such collisions may be commonplace (see Section 25.2). The radio lobes are roughly symmetrically placed, jutting out from the center of the visible galaxy and roughly perpendicular to the dust lane, suggesting that they consist of material ejected in opposite directions from the galactic nucleus. This conclusion is strengthened by the presence of a pair of smaller secondary lobes closer to the visible galaxy and by the presence of a roughly 1-kpc-long jet of matter in the galactic center, all aligned with the main lobes (and marked in the figure).


If the material was ejected from the nucleus at close to the speed of light and has subsequently slowed, then Centaurus A's outer lobes were created a few hundred million years ago, quite possibly around the time of the collision thought to be responsible for the galaxy's odd optical appearance. The secondary lobes were expelled more recently. Apparently, some violent process at the center of Centaurus A—most probably triggered by the collision—started up around that time and has been intermittently firing jets of matter out into intergalactic space ever since.

Centaurus A is a relatively low-luminosity source that happens to lie very close to us, astronomically speaking, making it particularly easy to study. Figure 24.23 shows a much more powerful emitter, called Cygnus A, lying roughly 250 Mpc from Earth. The high-resolution radio map in Figure 24.23(b) clearly shows two narrow, high-speed jets joining the radio lobes to the center of the visible galaxy (the dot at the center of the radio image). Notice that, as with Centaurus A, Cygnus A is a member of a small group of galaxies, and the optical image (Figure 24.23a) appears to show two galaxies colliding.

◀ **FIGURE 24.22 Centaurus A Radio Lobes** Radio galaxies, such as Centaurus A (a), often have giant radio-emitting lobes (b) extending a million light-years or more beyond the central galaxy. This entire object might be the result of a collision between two galaxies that took place about 500 million years ago. The lobes cannot be imaged in visible light and must be observed with radio telescopes; they are shown here in false colors, with decreasing intensity from red to yellow to green to blue. The inset (at right) shows a *Chandra* X-ray image of one of the lobes, showing that the jets in the inner parts of the lobes do emit higher-energy radiation. (ESO; NRAO; SAO)




(b)



The radio lobes of the brightest radio galaxies (such as Cygnus A) emit roughly 10 times more energy than the Milky Way Galaxy does at all wavelengths, coincidentally about the same amount of energy emitted by the most luminous Seyfert nuclei. However, despite their names, radio galaxies actually radiate far more energy at shorter wavelengths. Their total energy output can be a hundred (or more) times greater than their radio emission. Most of this energy comes from the nucleus of the visible galaxy. With total luminosities up to a thousand times that of the Milky Way, bright radio galaxies are among the most energetic objects known in the universe. Their radio emission lets us study in detail the connection between the small-scale nucleus and the large-scale radio lobes.

Not all radio galaxies have obvious radio lobes. Figure 24.24 shows a *core-dominated* radio galaxy, most of whose energy is emitted from a small central nucleus (which radio astronomers refer to as the *core*) less than 1 pc across. Weaker radio emission comes from an extended region surrounding the nucleus. It is likely that all radio galaxies have jets and lobes, but what we observe depends on our perspective. As illustrated in Figure 24.25, when a radio galaxy is viewed from the side, we see the jets and lobes. However, if we view the jet almost head-on—in other words, looking *through* the lobe—we see a core-dominated system.




▲ **FIGURE 24.23 Cygnus A** (a) A visible-light image of Cygnus A appears to show two galaxies in collision. (b) On much larger scales, it displays radio-emitting lobes (mapped in blue) on both sides of the visible image. The galaxy in (a) is about the size of the small dot at the center of (b). (NOAO; NRAO)

Our precise location with respect to the jet can also radically affect the *type* of radiation we see. The theory of relativity tells us that radiation emitted by particles moving close to the speed of light is strongly concentrated, or beamed, in the direction of motion. ∞ (Discovery 22-1) As a result, if the observer in Figure 24.25 happens to be directly in line with the beam, the radiation she receives is both very intense and Doppler shifted toward short wavelengths. ∞ (Sec. 3.5) The resulting object is called a **blazar** (Figure 5.35b). Much of the luminosity of the several hundred known blazars is received in the form of X-rays or gamma rays.


Jets are a fairly common feature of active galaxies of all types. Figure 24.26 presents several images of


► **FIGURE 24.25 Radio Galaxy** A central energy source produces high-speed jets of matter that interact with intergalactic gas to form radio lobes. The system may appear to us as either radio lobes or a core-dominated radio galaxy, depending on our location with respect to the jets and lobes.



▲ **FIGURE 24.24 Core-Dominated Radio Galaxy** As shown by this radio contour map of the galaxy M86, the radio emission comes from a bright central nucleus, which is surrounded by an extended region of less-intense emission. The radio map is superimposed on an optical image of the galaxy and some of its neighbors, a wider-field version of which was shown previously in Figure 24.14. (Harvard-Smithsonian CfA)

the giant elliptical galaxy M87, a prominent member of the Virgo Cluster (Figure 24.14). A long-time exposure (Figure 24.26a) shows a large, fuzzy ball of light—a fairly normal-looking E1 galaxy about 100 kpc across. A shorter exposure of M87 (Figure 24.26b), capturing only the galaxy's bright inner regions, reveals a long (2 kpc) thin jet of





**Interactive FIGURE 24.26 M87 Jet** The giant elliptical galaxy M87 (also called Virgo A) is displayed here in a series of zooms. (a) A long optical exposure of the galaxy's halo and embedded central region. (b) A short optical exposure of its core and an intriguing jet of matter, on a smaller scale. (c) An infrared image of M87's jet, examined more closely than in (b). The bright point at left in (c) marks the bright nucleus of the galaxy; the bright blob near the center of the image corresponds to the bright "knot" visible in the jet in (b). (NOAO; NASA)

matter ejected from the galactic center at nearly the speed of light. Computer enhancement shows that the jet is made up of a series of distinct "blobs" more or less evenly spaced along its length, suggesting that the material was ejected during bursts of activity. The jet has also been imaged in the radio, infrared (Figure 24.26c), and X-ray regions of the spectrum.

### CONCEPT Check

- ✓ The energy emission from an active galactic nucleus does not resemble a blackbody curve. Why is this important?


## Quasars

In the early days of radio astronomy, many radio sources were detected for which no corresponding visible object was known. By 1960, several hundred such sources were listed in the *Third Cambridge Catalog*, and astronomers were scanning the skies in search of visible counterparts to these radio sources. Their job was made difficult both by the low resolution of the radio observations (which meant that the observers did not know exactly where to look) and by the faintness of the objects at visible wavelengths.

In 1960, astronomers detected what appeared to be a faint blue star at the location of the radio source 3C 48 (the 48th object on the third *Cambridge* list) and obtained its spectrum. Containing many unknown and unusually broad emission lines, the object's peculiar spectrum defied interpretation. 3C 48 remained a unique curiosity until 1962, when another similar-looking—and similarly mysterious—faint blue object with "odd" spectral lines was discovered and identified with the radio source 3C 273 (Figure 24.27).

The following year saw a breakthrough when astronomers realized that the strongest unknown lines in 3C 273's spectrum were simply familiar spectral lines of hydrogen redshifted by a very unfamiliar amount—about 16 percent, corresponding to a recession velocity of 48,000 km/s! Figure 24.28 shows the spectrum of 3C 273. Some prominent emission lines and the extent of their redshift are marked on the diagram. Once the nature of the strange spectral lines was known, astronomers quickly found a similar explanation for the spectrum of 3C 48, whose 37 percent redshift implied that it was receding from Earth at the astonishing rate of almost one-third the speed of light!

Their huge speeds mean that neither of these two objects can be members of our Galaxy. In fact, their large redshifts indicate that they are very far away indeed. Applying Hubble's law (with our adopted value of the Hubble constant,  $H_0 = 70$  km/s/Mpc), we obtain distances of 650 Mpc for 3C 273 and 1450 Mpc for 3C 48. (See again *More Precisely 24-1* for more information of how these distances are determined and what the large redshifts really mean.)



▲ **FIGURE 24.27 Quasar 3C 273** (a) The bright quasar 3C 273 displays a luminous jet of matter, but the main body of the quasar is starlike in appearance. (b) The jet extends for about 30 kpc and can be seen better in this high-resolution image. (AURA)

However, this explanation of the unusual spectra created an even deeper mystery. A simple calculation using the inverse-square law reveals that, despite their unimpressive optical appearance (see Figure 24.29), these faint “stars” are in fact among the brightest-known objects in the universe! Object 3C 273, for example, has a luminosity of about  $10^{40}$  W, comparable to 20 trillion Suns or a thousand Milky Way Galaxies. More generally, quasars range in luminosity from around  $10^{38}$  W—about the same as the brightest Seyferts—up to nearly  $10^{42}$  W. A value of  $10^{40}$  W (comparable to the luminosity of a bright radio galaxy) is fairly typical.

Clearly not stars (because of their enormous luminosities), these objects became known as *quasi-stellar radio sources* (“quasi-stellar” means “starlike”), or **quasars**. (The name persists even though we now know that not all such highly redshifted, starlike objects are strong radio sources.) More than 200,000 quasars are now known, and the numbers are increasing rapidly as large-scale surveys probe deeper and deeper into space (see *Discovery 25-1*). The distance to the *closest* quasar is 250 Mpc; the farthest lies more than 9000 Mpc away. Most quasars lie well over 1000 Mpc from Earth.


Since light travels at a finite speed, these faraway objects represent the universe as it was in the distant past. The implication is that most quasars date back to much earlier periods of galaxy formation and evolution, rather than more recent times. The prevalence of these energetic objects at great distances tells us that the universe was once a much more violent place than it is today.

Quasars share many properties with Seyferts and radio galaxies. Their radiation is nonstellar and may vary irregularly in brightness over periods of months, weeks, days, or (in some cases) even hours, and some quasars show evidence of jets and extended emission features. Note the jet of luminous matter in 3C 273 (Figure 24.27), reminiscent of the jet in M87 and extending nearly 30 kpc from the center of the quasar. Figure 24.30 shows a quasar with radio lobes similar to those seen in Cygnus A (Figure 24.23b). Quasars have been observed in all parts of the electromagnetic spectrum, although many emit most of their energy in the optical and infrared. About 10–15 percent of quasars (called “radio-loud” quasars) also emit substantial amounts of energy at radio wavelengths, presumably as a result of unresolved jets.

Astronomers once distinguished between active galaxies and quasars on the basis of their appearance, spectra, and distance from us, but today most astronomers think that quasars *are* in fact just the intensely bright nuclei of distant active galaxies lying too far away for the galaxies themselves to be seen. (Figure 25.19 presents *Hubble Space Telescope* observations of several relatively nearby quasars in which the surrounding galaxies are clearly visible.)

### PROCESS OF SCIENCE Check


- ✓ How did the determination of quasar distances change astronomers’ understanding of these objects?



▲ **FIGURE 24.28 Quasar Spectrum** Optical spectrum of the distant quasar 3C 273. (This is a negative, as the lines are actually in emission.) Notice both the redshift and the widths of the three hydrogen spectral lines marked as H $\beta$ , H $\gamma$ , and H $\delta$ . (Sec. 14.3) The redshift indicates the quasar’s enormous distance. The width of the lines implies rapid internal motion within the quasar. (Adapted from Palomar/Caltech)



▲ **FIGURE 24.29 Typical Quasar** Although quasars are the most luminous objects in the universe, they are often unimpressive in appearance. In this optical image, a distant quasar (marked by an arrow) is seen close (in the sky) to nearby normal stars. The quasar's much greater distance makes it appear fainter than the stars, but intrinsically it is much, much brighter. Often starlike in appearance, quasars are generally identified via their unusual nonstellar colors or spectra. (SDSS)



▲ **FIGURE 24.30 Quasar Jets** This radio image of the quasar 3C 175, which is some 3000 Mpc away, shows radio jets feeding faint radio lobes. The lobes themselves span approximately a million light-years—comparable in size to the radio galaxies discussed earlier (see also the chapter-opening image on page 606.). (NRAO)

## 24.5 The Central Engine of an Active Galaxy

The present consensus among astronomers is that, despite their differences in appearance and luminosity, Seyferts, radio galaxies, quasars—as well as “normal” galactic nuclei—share a common energy-generation mechanism.


As a class, active galactic nuclei have some or all of the following properties:

1. They have *high luminosities*, generally greater than the  $10^{37}$  W characteristic of a bright normal galaxy.
2. Their energy emission is mostly *nonstellar*—it cannot be explained as the combined radiation of even trillions of stars.
3. Their energy output can be highly *variable*, implying that their energy is emitted from a small central nucleus much less than a parsec across.
4. They may exhibit *jets* and other signs of explosive activity.
5. Their optical spectra may show broad emission lines, indicating *rapid internal motion* within the energy-producing region.
6. Often the activity appears to be associated with *interactions* between galaxies.

The principal questions, then, are How can such vast quantities of energy arise from these relatively small regions of space? Why is the radiation nonstellar? and What is the origin of the jets and extended radio-emitting lobes? We first consider how the energy is *produced* and then turn to the question of how it is actually *emitted* into intergalactic space.

### Energy Production

As illustrated in Figure 24.31, the leading model for the central engine of active galaxies is a scaled-up version of the process powering X-ray binaries in our own Galaxy and the activity in our Galactic center—accretion of gas



**▲ FIGURE 24.31 Active Galactic Nucleus** The leading theory for the energy source in active galactic nuclei holds that these objects are powered by material accreting onto a supermassive black hole. As matter spirals toward the hole, it heats up, producing large amounts of energy. At the same time, high-speed jets of gas can be ejected perpendicular to the accretion disk, forming the jets and lobes observed in many active objects. Magnetic fields generated in the disk by charged matter in motion are carried by the jets out to the radio lobes, where they play a crucial role in producing the detected radiation.

onto a supermassive black hole, releasing huge amounts of energy as the matter sinks onto the central object. ∞ (Secs. 22.3, 22.8, 23.7) In order to power the brightest active galaxies, theory suggests that the black holes involved must be *billions* of times more massive than the Sun.

As with this model's smaller scale counterparts, the infalling gas forms an *accretion disk* and spirals down toward the black hole, heating up to high temperatures by friction within the disk and emitting large amounts of radiation as a result. In the case of an active galaxy, however, the origin of the accreted gas is not a binary companion, as it is in stellar X-ray sources, but entire stars and clouds of interstellar gas—most likely diverted into the galactic center by an encounter with another galaxy—that come too close to the hole and are torn apart by its strong gravity.


Accretion is extremely efficient at converting infalling mass (in the form of gas) into energy (in the form of electromagnetic radiation). Detailed calculations indicate that as much as 10 or 20 percent of the total mass–energy of the infalling matter can be radiated away before it crosses the hole's event horizon and is lost forever. ∞ (Sec. 22.5) Since the total mass–energy of a star like the Sun—the mass times the speed of light squared—is about  $2 \times 10^{47}$  J, it follows that the  $10^{38}$ -W luminosity of a bright active galaxy can be accounted for by the consumption of “only” 1 solar mass of gas per decade by a billion-solar-mass black hole. More or less luminous active galaxies would require correspondingly more or less fuel. Thus, to power a  $10^{40}$ -W quasar, which is 100 times brighter, the black hole simply consumes 100 times more fuel, or 10 stars per year. The central black hole of a  $10^{36}$ -W Seyfert galaxy would devour only one Sun's worth of material every thousand years.

The small size of the emitting region is a direct consequence of the compact central black hole. Even a billion-solar-mass black hole has a radius of only  $3 \times 10^9$  km, or  $10^{-4}$  pc—about 20 AU—and theory suggests that the part of the accretion disk responsible for most of the emission would be much less than 1 pc across. ∞ (Sec. 22.5) Instabilities in the accretion disk can cause fluctuations in the energy released, leading to the variability observed in many objects.

The broadening of the spectral lines seen in the nuclei of many active galaxies results from the rapid orbital motion of the gas in the black hole's intense gravity.

Jets appear to be a common feature of accretion flows, large and small. The jets shown in Figure 24.31 consist of material (mainly electrons and protons) blasted out into space—and completely out of the visible portion of the galaxy—from the inner regions of the disk. They are most likely formed by strong magnetic fields produced within the accretion disk itself. These fields accelerate charged particles to nearly the speed of light and eject them parallel to the disk's rotation axis. Figure 24.32 shows a *Hubble Space Telescope* image of a disk of gas and dust at the core of the radio galaxy NGC 4261 in the Virgo Cluster. Consistent with the model just described, the disk is perpendicular to the huge jets emanating from the galaxy's center.






**◀ FIGURE 24.32 Giant Elliptical Galaxy** (a) A combined optical/radio image of the giant elliptical galaxy NGC 4261, in the Virgo Cluster, shows a white visible galaxy at center, from which blue-orange (false-color) radio lobes extend for about 60 kpc. (b) A more detailed view of the galaxy's nucleus reveals a 100-pc-diameter disk surrounding a bright hub thought to harbor a black hole. (NRAO; NASA)

Figure 24.33 shows further evidence in favor of this model, in the form of imaging and spectroscopic data from the center of M87, suggesting a rapidly rotating disk of matter orbiting the galaxy's center, again perpendicular to the jet. Measurements of the gas velocity on opposite sides of the disk indicate that the mass within a few parsecs of the center is approximately  $3 \times 10^9$  solar

masses; we assume that this is the mass of the central black hole. At M87's distance, *HST*'s resolution of 0.05 arc second corresponds to a scale of about 5 pc, so we are still far from seeing the (solar system-sized) central black hole itself, but the improved "circumstantial" evidence has convinced many astronomers of the basic correctness of the theory.



**▲ FIGURE 24.33 M87 Disk** Both images and spectra of M87 support the idea of a rapidly whirling accretion disk at this galaxy's heart. (a) The central region of M87, similar to that shown in Figure 24.26(c), shows the galaxy's bright nucleus and jet. (b) A magnified view of the nucleus suggests a spiral swarm of stars, gas, and dust. (c) Spectral-line features observed on opposite sides of the nucleus show contrasting red and blue Doppler shifts, implying that matter on one side is coming toward us and matter on the other side is moving away. Apparently, an accretion disk spins perpendicular to the jet, and at its center is a black hole having some 3 billion times the mass of the Sun. (NASA)


## Energy Emission

Theory suggests that the radiation emitted by the hot accretion disk surrounding a supermassive black hole should span a broad range of wavelengths, from infrared through X-rays, corresponding to the broad range of temperatures in the disk as the gas heats up. This accounts for the spectra of some active galactic nuclei, but it also appears that in many cases the high-energy radiation emitted from the accretion disk is “reprocessed”—that is, absorbed and reemitted at longer wavelengths—by material beyond the nucleus before eventually reaching our detectors.

Researchers think that the most likely site of this reprocessing is a rather fat, donut-shaped ring of gas and dust surrounding the inner accretion disk where the energy is actually produced. As illustrated in Figure 24.34, if our line of sight to the black hole does not intersect this dusty donut, then we see the “bare” energy source emitting large amounts of high-energy radiation (with broad emission lines, since we can see the rapidly moving gas near the black hole). ∞ (Sec. 4.5) If the donut intervenes, we see instead large amounts of infrared radiation reradiated from the dust (and only narrow emission lines, from gas farther from the center). The structure of the donut itself is quite uncertain and may in reality bear little resemblance to the rather regular-looking ring shown in the figure.


A different reprocessing mechanism operates in many jets and radio lobes. This mechanism involves the *magnetic fields* possibly produced within the accretion disk and transported by the jets into intergalactic space (Figure 24.31). As sketched in Figure 24.35(a), whenever a charged particle (here an electron) encounters a magnetic field, the particle tends to spiral around the magnetic field lines. We have already encountered this idea in the discussions of Earth’s magnetosphere and solar activity. ∞ (Secs. 7.5, 16.5)

As the particles whirl around, they emit electromagnetic radiation. ∞ (Sec. 3.2) The radiation produced in



▲ **FIGURE 24.34 Dusty Donut** The accretion disk surrounding a massive black hole, drawn here with some artistic license, consists of hot gas at many different temperatures (hottest nearest the center). When viewed from above or below, the disk radiates a broad spectrum of electromagnetic energy extending into the X-ray band. However, the dusty infalling gas that ultimately powers the system is thought to form a fat, donut-shaped region outside the accretion disk (shown here in red), which effectively absorbs much of the high-energy radiation reaching it, reemitting it mainly in the form of cooler, infrared radiation. When the accretion disk is viewed from the side, strong infrared emission is observed. (Compare with Figure 24.25.) (Adapted from D. Berry)

this way—called **synchrotron radiation**, after the type of particle accelerator in which it was first observed—is *nonthermal* in nature, meaning that there is no link between the emission and the temperature of the radiating object. Hence, the radiation is not described by a blackbody curve. Instead, its intensity decreases with increasing frequency, as shown in Figure 24.35(b). This is just what is needed to explain the overall spectrum of radiation from radio galaxies and radio-loud quasars.



**▲ FIGURE 24.35 Nonthermal Radiation** (a) Charged particles, especially fast-moving electrons (red), emit synchrotron radiation (blue) while spiraling in a magnetic field (black). This process is not confined to active galaxies. It occurs on smaller scales as well, when charged particles interact with magnetism in Earth’s Van Allen belts (Sec. 7.5), when charged matter arches above sunspots (Sec. 16.5), in the vicinity of neutron stars (Sec. 22.2), and at the center of our own Galaxy (Sec. 23.7). (b) Thermal and synchrotron (nonthermal) radiation vary differently with frequency. Thermal radiation, described by a blackbody curve, peaks at a frequency that depends on the temperature of the source. By contrast, nonthermal synchrotron radiation is more intense at low frequencies and is independent of the temperature of the emitting object. (Compare with Figure 24.19.)

Observations of the radiation received from the jets and radio lobes of active galaxies are completely consistent with synchrotron radiation.

Eventually, the jet is slowed and stopped by the intergalactic medium, the flow becomes turbulent, and the magnetic field grows tangled. The result is a gigantic radio lobe emitting virtually all of its energy in the form of synchrotron radiation. Thus, even though the radio *emission* comes from an enormously extended volume of space that dwarfs the visible galaxy, the *source* of the energy is still the accretion disk—a billion billion times smaller in volume than the radio lobe—lying at the galactic center. The jets transport energy from the nucleus, where it is generated, into the lobes, where it is finally radiated into space.

The existence of the inner lobes of Centaurus A and the blobs in M87’s jet imply that the formation of a jet may be an intermittent process (or, as in the case of the Seyferts discussed earlier, may not occur at all), and, as we have seen, there is also evidence to indicate that much, if not all, of the activity observed in nearby active galaxies has been sparked

by recent interaction with a neighbor. Many nearby active galaxies (e.g., Centaurus A) appear to have been “caught in the act” of interacting with another galaxy, suggesting that the fuel supply can be “turned on” by a companion. The tidal forces involved divert gas and stars into the galactic nucleus, triggering an outburst that may last for many millions of years.

What do active galaxies look like between active outbursts? What other connections exist between them and the normal galaxies we see? To answer these important questions, we must delve more deeply into the subject of *galaxy evolution*, to which we turn in Chapter 25.

### CONCEPT Check

- ✓ How does accretion onto a supermassive black hole power the energy emission from the extended radio lobes of a radio galaxy?

**The Big Question** Galactic research lags stellar research by about 50 years. That's because galaxies were discovered only in the 20th century, and we are still learning about them. How did they form, and how do they evolve? Those are the biggest questions regarding galaxies, and they will not be answered until more and better data accumulate, especially regarding the most distant systems. With much larger galaxy surveys on the horizon, breakthroughs that will help solve these important issues may be imminent.


# Chapter Review

## SUMMARY

**1** The **Hubble classification scheme** (p. 608) divides galaxies into several classes, depending on their appearance. **Spiral galaxies** (p. 608) have flattened disks, central bulges, and spiral arms. Their halos consist of old stars, whereas the gas-rich disks are the sites of ongoing star formation. **Barred-spiral galaxies** (p. 609) contain an extended “bar” of material projecting beyond the central bulge. **Elliptical galaxies** (p. 611) have no disk and contain little or no cool gas or dust, although very hot interstellar gas is observed. In most cases, they consist entirely of old stars. They range in size from dwarf ellipticals, which are much less massive than the Milky Way Galaxy, to giant ellipticals, which may contain trillions of stars. **S0 and SB0 galaxies** (p. 612) are intermediate in their properties between ellipticals and spirals. **Irregular galaxies** (p. 612) are galaxies that do not fit into any of the other categories. Many are rich in gas and dust and are the sites of vigorous star formation.



**2** Astronomers often use **standard candles** (p. 615) as distance-measuring tools. These are objects that are easily identifiable and whose luminosities lie within some reasonably well-defined range. Comparing luminosity and apparent brightness, astronomers determine distance with the use of the inverse-square law. An alternative approach is the **Tully-Fisher relation** (p. 616), an empirical correlation between rotational velocity and luminosity in spiral galaxies.




**3** The Milky Way, Andromeda, and several other smaller galaxies form a small gravitationally bound collection of galaxies called the **Local Group** (p. 618). **Galaxy clusters** (p. 618) consist of many galaxies orbiting one another, bound together by their own gravity. The nearest large galaxy cluster to the Local Group is the Virgo Cluster.



**4** Distant galaxies are observed to be receding from the Milky Way at speeds proportional to their distances from us. This relationship is called **Hubble's law**


(p. 620). The constant of proportionality in the law is **Hubble's constant** (p. 620). Its value is thought to be around 70 km/s/Mpc. Astronomers use Hubble's law to determine distances to the most remote objects in the universe. The redshift associated with the Hubble expansion is called the **cosmological redshift** (p. 620).



**5** **Active galaxies** (p. 622) can be much more luminous than normal galaxies and have nonstellar spectra, emitting most of their energy outside the visible part of the electromagnetic spectrum. Often, the nonstellar activity suggests rapid internal motion and is associated with a bright **active galactic nucleus** (p. 624). Many active galaxies have high-speed, narrow jets of matter shooting out from their central nuclei. The jets transport energy from the nucleus, where it is generated, to enormous **radio lobes** (p. 625) lying far beyond the visible portion of the galaxy, where it is radiated into space. The jets often appear to be made up of distinct “blobs” of gas, suggesting that the process that generates the energy is intermittent.




**6** A **Seyfert galaxy** (p. 624) looks like a normal spiral, but has an extremely bright central galactic nucleus. Spectral lines from Seyfert nuclei are very broad, indicating rapid internal motion, and the rapid variability in the luminosity of Seyferts implies that the source of the radiation is much less than 1 light-year across. **Radio galaxies** (p. 625) emit large amounts of energy in the radio part of the spectrum. The corresponding visible galaxy is usually elliptical. **Quasars** (p. 629), or quasi-stellar objects, are the most luminous objects known. In visible light they appear starlike, and their spectra are usually substantially redshifted. All quasars are very distant, indicating that we see them as they were in the remote past.



**7** The generally accepted explanation for the observed properties of all active galaxies is that their energy is generated by the accretion of galactic gas onto a supermassive (million- to billion-solar-mass) black hole lying in the galactic center. The small size of the accretion disk explains the compact extent of the emitting region, and the high-speed orbit of gas in the black hole's intense gravity accounts for the rapid motion that is observed. Typical luminosities of active galaxies require the consumption of about 1 solar mass of material every few years. Some of the infalling matter may be blasted out into space, producing magnetized jets that create and feed the galaxy's radio lobes. The accretion

disk emits at a broad range of temperatures, producing a nonstellar spectrum. In addition, much of the radiation may be reprocessed into the infrared by a ring of dust surrounding the disk. On larger scales, charged particles spiraling around magnetic field lines produce **synchrotron radiation** (p. 633), whose spectrum is consistent with the radio emission from radio galaxies and jets.



**MasteringAstronomy**® For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- LO1** What distinguishes one type of spiral galaxy from another?
- Describe some similarities and differences between elliptical galaxies and the halo of our own Galaxy.
- LO2** Describe the four rungs in the distance-measurement ladder used to determine the distance to a galaxy lying 5 Mpc away.
- LO3** Describe the contents of the Local Group. How much space does it occupy compared with the volume of the Milky Way?
- What is the Virgo Cluster?
- What are standard candles, and why are they important to astronomy?
- How is the Tully-Fisher relation used to measure distances to galaxies?
- LO4 POS** What is Hubble's law? How is it used by astronomers to measure distances to galaxies?
- What is the most likely range of values for Hubble's constant? What are the uncertainties in its value?
- LO5** Name two basic differences between normal galaxies and active galaxies.
- POS** What is the evidence that the radio lobes of some active galaxies consist of material ejected from the galaxy's center?
- LO6 POS** How do we know that the energy-emitting regions of many active galaxies must be very small?
- What was it about the spectra of quasars that was so unexpected and surprising?
- How do we know that quasars are extremely luminous?
- LO7** Briefly describe the leading model for the central engine of an active galaxy.

## Conceptual Self-Test: Multiple Choice

- Young stars in a galactic disk are (a) evenly distributed within and between spiral arms; (b) mostly found in the space between spiral arms; (c) mostly found in the spiral arms; (d) older than stars in the halo.
- Astronomers classify elliptical galaxies by (a) the number of stars they contain; (b) their colors; (c) how flattened they appear; (d) their diameters.
- Using the method of standard candles, we can, in principle, find the distance to a campfire if we know (a) the number of logs used; (b) the fire's temperature; (c) the length of time the fire has been burning; (d) the type of wood used in the fire.
- VIS** If the galaxy in Figure 24.11 ("Galaxy Rotation") were smaller and spinning more slowly, then, in order to represent it correctly, the figure should be redrawn to show (a) a greater blueshift; (b) a greater redshift; (c) a narrower combined line; (d) a larger combined amplitude.
- Within 30 Mpc of the Sun, there are about (a) 3 galaxies; (b) 30 galaxies; (c) a few thousand galaxies; (d) a few million galaxies.
- VIS** According to Figure 24.17 ("Hubble's Law"), a galaxy 500 million parsecs away has a velocity of roughly (a) 25,000 km/s away from us; (b) 35,000 km/s toward us; (c) 35,000 km/s away from us; (d) 75,000 km/s toward us.

7. **VIS** According to Figure 24.19 (“Galaxy Energy Spectra”), active galaxies (a) emit most of their energy at long wavelengths; (b) emit very little energy at high frequencies; (c) emit large amounts of energy at all wavelengths; (d) emit most of their energy in the visible part of the spectrum.
8. If the light from a galaxy fluctuates in brightness very rapidly, the region producing the radiation must be (a) very large; (b) very small; (c) very hot; (d) rotating very rapidly.
9. Quasar spectra (a) are strongly redshifted; (b) show no spectral lines; (c) look like the spectra of stars; (d) contain emission lines from unknown elements.
10. Active galaxies are very luminous because they (a) are hot; (b) contain black holes in their cores; (c) are surrounded by hot gas; (d) emit jets.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

1. ● A supernova of luminosity 1 billion times the luminosity of the Sun is used as a standard candle to measure the distance to a faraway galaxy. From Earth, the supernova appears as bright as the Sun would appear from a distance of 10 kpc. What is the distance to the galaxy?
2. ●● A Cepheid variable star in the Virgo Cluster has an absolute magnitude of  $-5$  and is observed to have an apparent magnitude of 26.3. Use these figures to calculate the distance to the Virgo Cluster.
3. ● According to Hubble’s law, with  $H_0 = 70$  km/s/Mpc, what is the recessional velocity of a galaxy at a distance of 200 Mpc? How far away is a galaxy whose recessional velocity is 4000 km/s? How do these answers change if  $H_0 = 60$  km/s/Mpc? If  $H_0 = 80$  km/s/Mpc?
4. ●● According to Hubble’s law, with  $H_0 = 70$  km/s/Mpc, how long will it take for the distance from the Milky Way Galaxy to the Virgo Cluster to double?
5. ● Use the data in Table 24.2 to estimate the absolute magnitude and luminosity of a quasar with a redshift of 5 and an apparent magnitude of 22.
6. ●● A certain quasar has a redshift of 0.25 and an apparent magnitude of 13. Using the data in Table 24.2, calculate the quasar’s absolute magnitude and hence its luminosity. Compare the apparent brightness of the quasar, viewed from a distance of 10 pc, with that of the Sun as seen from Earth.
7. ●● Spectral lines from a Seyfert galaxy are observed to be redshifted by 0.5 percent and to have broadened emission lines indicating an orbital speed of 250 km/s at an angular distance of  $0.1''$  from its center. Assuming circular orbits, use Kepler’s laws to estimate the mass within this  $0.1''$  radius. ∞ (Sec. 23.6)
8. ● A quasar consumes 1 solar mass of material per year, converting 15 percent of it directly into energy. What is the quasar’s luminosity, in solar units?

## Activities

### Collaborative

1. Observe the Virgo Cluster of galaxies. An 8-inch telescope is the optimal size for this project. The constellation Virgo is visible from the United States during much of fall through spring. To locate the cluster, first find the constellation Leo. The eastern part of Leo is composed of a distinct triangle of stars, Denebola ( $\beta$ ), Chort ( $\theta$ ), and Zosma ( $\delta$ ). Go from Chort to Denebola in a straight line east; continue on the same distance as between the two stars, and you will be approximately at the center of the Virgo Cluster. Look for the following Messier objects, the brightest galaxies in the cluster: M49, M58, M59, M60, M84, M86, M87, M89, and M90. Examine each galaxy for unusual features; some have very bright nuclei. Sketch or photograph what you see, and construct your own pictorial catalog of the brightest galaxies in Virgo.

### Individual

1. Quasar 3C 273 is the nearest and brightest one, but that does not mean it is easy to find! Its coordinates are  $RA = 12^h 29.2^m$ ,  $dec = +2^\circ 03'$ . It is located in the southern part of the Virgo Cluster, but it is not associated with it. At magnitude 12–13 (it is variable), it may require a 10- or 12-inch telescope to see, but try it first with an 8-inch. It should appear as a very faint star. The significance of seeing this object is that it is 640 Mpc distant. The light you are seeing left this object over 2 billion years ago! It is the most distant object observable with a small telescope.



# Galaxies and Dark Matter

## THE LARGE-SCALE STRUCTURE OF THE COSMOS

On scales much larger than even the largest galaxy clusters, the dynamics of the universe itself becomes apparent, new levels of structure are revealed, and a humbling new reality emerges. We may be star stuff, the product of countless cycles of stellar evolution, but we are not the stuff of the cosmos. The universe in the large is composed of matter fundamentally different from the familiar atoms and molecules that make up our bodies, our planet, our star and Galaxy, and all the luminous matter we observe in the heavens.

By comparing and classifying the properties of galaxies near and far, astronomers have begun to understand their formation, dynamics, and evolution. By mapping out the distribution of those galaxies in space, we can trace out the immense realms of the universe. Points of light in the uncharted darkness, they remind us that our position in the universe is no more special than that of a boat adrift at sea.

**The Big Picture** Galaxies are among the grandest, most beautiful objects in the universe—each one a colossal collection of hundreds of billions of stars held together loosely by gravity. Galaxies dominate our view of deep space—they seem to be everywhere—yet they represent just a tiny fraction of all matter in the cosmos. Vast quantities of unseen cosmic material—dark matter—actually account for most of the mass in the universe.

**LEFT:** Some galaxies are bright and splendid, like the two big ones in this image. Others are dim and distant, like several that appear smaller in the background. This pair of galaxies, nearly 300 million light-years away and known collectively as Arp 273, is in the process of colliding over millions of years. Notice the roselike shape of the top galaxy, caused by the gravitational pull of the bottom one, and the swath of clusters of young blue stars glowing like jewels. Mergers and acquisitions are common among galaxies, but astronomers still don't fully understand how galaxies formed long ago. (*STSci*)

### Learning Outcomes

*Studying this chapter will enable you to*


- 1 Describe some methods used to determine the masses of galaxies and galaxy clusters.
- 2 Explain why astronomers think that most of the matter in the universe is dark.
- 3 Describe how galaxies form and evolve, and outline the role of collisions in the process.
- 4 Present the evidence for supermassive black holes in the centers of galaxies, and explain how active galaxies fit into current theories of galactic evolution.
- 5 Summarize what is known about the large-scale distribution of galaxies in the universe.
- 6 Outline some techniques used by astronomers to probe the universe on very large scales.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.






## 25.1 Dark Matter in the Universe


In Chapter 23, we saw how measurements of the orbital velocities of stars and gas in our own Galaxy reveal the presence of an extensive *dark-matter halo* surrounding the galaxy we see.  (Sec. 23.6) Do other galaxies have similar dark halos? And what evidence do we have for dark matter on larger scales? To answer these questions, we need a way to calculate the masses of galaxies and galaxy clusters, then compare those masses with the luminous matter we actually observe.

How can we measure the masses of such large systems? Surely, we can neither count all their stars nor estimate their interstellar content very well: Galaxies are just too complex for us to take a direct inventory of their material makeup. Instead, we must rely on indirect techniques. Despite their enormous sizes, galaxies and galaxy clusters obey the same physical laws as do the planets in our own solar system. To calculate galaxy masses, we turn as usual to Newton's law of gravity.

### Masses of Galaxies and Galaxy Clusters

Astronomers can calculate the masses of some spiral galaxies by determining their *rotation curves*, which plot rotation speed versus distance from the galactic center, as illustrated in Figure 25.1. Rotation curves for a few nearby spirals are shown in Figure 25.1(b). The mass within any given radius then follows directly from Newton's laws.  (Sec. 2.7) The rotation curves shown imply masses ranging from about  $10^{11}$  to  $5 \times 10^{11}$  solar masses within about 25 kpc of the center—comparable to the results obtained for our own Galaxy using the same technique.  (Sec. 23.6)


Distant galaxies are generally too far away for such detailed curves to be drawn. Nevertheless, by observing the broadening of their spectral lines—as discussed in Chapter 24 in the context of the Tully-Fisher relation—we can still measure the overall rotation speed of these galaxies.  (Sec. 24.2) Estimating a galaxy's size then leads to an estimate of its mass. Similar techniques have been applied to ellipticals and irregulars. In general, the approach is useful for measuring the mass lying within about 50 kpc of a galaxy's center—the extent of the electromagnetic emission from stellar and interstellar material.

To probe farther from the centers of galaxies, astronomers turn to *binary* galaxy systems (Figure 25.2a), whose components may lie hundreds of kiloparsecs apart. The orbital period of such a system is typically billions of years, far too long for the orbit to be accurately measured. However, by estimating the period and semimajor axis from the available information—the line-of-sight velocities and the angular separation of the components—an approximate total mass can be derived.  (More Precisely 2-2)

Galaxy masses obtained in this way are uncertain, but by combining many such measurements, astronomers can obtain quite reliable *statistical* information about




(a)




(b)

### Interactive FIGURE 25.1 Galaxy Rotation Curves


 (a) Orbital velocities can be measured at different distances from the center of a disk galaxy, as illustrated here for M64, the “Evil Eye” galaxy, some 5 Mpc distant. (b) Rotation curves for some nearby spiral galaxies indicate masses of a few hundred billion times the mass of the Sun. (NASA)

galaxy masses. Most normal spirals (the Milky Way Galaxy included) and large ellipticals contain between  $10^{11}$  and  $10^{12}$  solar masses of material. Irregular galaxies often contain less mass, about  $10^8$  to  $10^{10}$  times that of the Sun. Dwarf ellipticals and dwarf irregulars can contain as little as  $10^6$  or  $10^7$  solar masses of material.

We can use another statistical technique to derive the combined mass of all the galaxies within a galaxy cluster.



(a)



(b)

**▲ FIGURE 25.2 Galaxy Masses** (a) In a binary galaxy system, galaxy masses can be estimated by observing the orbit of one galaxy about the other. (b) The mass of a galaxy cluster can be estimated by observing the motion of many galaxies in the cluster and then estimating how much mass is needed to prevent the cluster from flying apart.

As depicted in Figure 25.2(b), each galaxy within a cluster moves relative to all other members of the cluster, and we can estimate the cluster’s mass simply by asking how massive it must be in order to bind its galaxies gravitationally. For example, if we find that galaxies in a cluster are moving with an average speed of 1000 km/s and the cluster radius is 3 Mpc (both typical values), it follows from Newton’s laws—*assuming* that the cluster is gravitationally bound—that the mass of the cluster must be around  $(3 \text{ Mpc}) \times (1000 \text{ km/s})^2 / G \approx 7 \times 10^{14}$  solar masses. [∞ \(More Precisely 2-2\)](#) Cluster masses obtained in this way generally lie in the range of  $10^{14}$ – $10^{15}$  solar masses. Notice

that this calculation gives us no information whatsoever about the masses of individual galaxies. It tells us only about the *total* mass of the cluster.

## Visible Matter and Dark Halos

The rotation curves of the spiral galaxies shown in Figure 25.1 remain flat (that is, do not decline, and even rise slightly) far beyond the galaxies’ visible images, implying that those galaxies—and perhaps all spiral galaxies—contain large amounts of *dark matter*, in the form of invisible dark halos similar to that surrounding the Milky Way. [∞ \(Sec. 23.6\)](#) Overall, spiral galaxies seem to contain from 3 to 10 times more mass than can be accounted for in the form of luminous matter. Studies of elliptical galaxies suggest similarly large dark halos surrounding these galaxies, too.


Astronomers find even greater discrepancies between visible light and total mass when they study galaxy clusters. Calculated cluster masses range from 10 to nearly 100 times the mass suggested by the light emitted by individual cluster galaxies. Put another way, a lot more mass is needed to bind galaxy clusters than we can see. Thus, the problem of dark matter exists not just in our own Galaxy, but also in other galaxies and, to an even greater degree, in galaxy clusters as well. In that case, we are compelled to accept the fact that *upwards of 90 percent of the matter in the universe is dark*—and not just in the visible portion of the spectrum. The mass goes undetected at *any* electromagnetic wavelength.

As discussed in Chapter 23, many possible explanations for the dark matter have been suggested, ranging from stellar remnants of various sorts to exotic subatomic particles. [∞ \(Sec. 23.6\)](#) Whatever its nature, the dark matter in clusters cannot simply be the accumulation of dark matter within individual galaxies. Even including the galaxies’ dark halos, we still cannot account for all the dark matter in galaxy clusters. As we look on larger and larger scales, we find that a larger and larger fraction of the matter in the universe is dark.

## Intracluster Gas

In addition to the luminous matter observed within the cluster galaxies themselves, astronomers also have evidence for large amounts of *intracluster gas*—superhot (more than 10 million K), diffuse intergalactic matter filling the space among the galaxies. Satellites orbiting above Earth’s atmosphere have detected substantial amounts of X-ray radiation from many clusters. Figure 25.3 shows false-color X-ray images of one such system. The X-ray-emitting region is centered on, and comparable in size to, the visible cluster image.

Further evidence for intracluster gas can be found in the appearance of the radio lobes of some active galaxies. [∞ \(Sec. 24.4\)](#) In some systems, known as *head-tail* radio galaxies, the lobes seem to form a “tail” behind the main part of the galaxy. For example, the lobes of radio galaxy



Images like these show that the space between the galaxies within galaxy clusters is filled with superheated gas.

**◀ FIGURE 25.3 Galaxy Cluster X-Ray Emission** (a) Superposition of infrared and X-ray radiation from the distant galaxy cluster Abell 1835. The infrared (in red) hugs the galaxies, but the X-rays display a fuzzy, bluish cloud of hot gas filling the intracluster spaces among the galaxies. (b) A longer infrared exposure of the central region, showing the richness of this cluster, which spans about a million parsecs. (NASA; ESA)

solve the dark-matter problem. To account for the total masses of galaxy clusters implied by dynamical studies, we would have to find from 10 to 100 times more mass in gas than exists in stars.


Why is the intracluster gas so hot? Simply because its particles are bound by gravity and hence are moving at speeds comparable to

NGC 1265, shown in Figure 25.4, appear to be “swept back” by some onrushing wind, and, indeed, this is the most likely explanation for the galaxy’s appearance. If NGC 1265 were at rest, it would be just another double-lobe source, perhaps quite similar to Centaurus A (Figure 24.22). However, the galaxy is traveling through the intergalactic medium of its parent galaxy cluster (known as the Perseus Cluster), and the outflowing matter forming the lobes tends to be left behind as NGC 1265 moves.

How much gas do these observations reveal? At least as much matter—and, in most cases, significantly *more*—exists within clusters in the form of hot gas as is visible in the form of stars. This is a lot of material, but it still doesn’t

those of the galaxies in the cluster—1000 km/s or so. Since temperature is just a measure of the speed at which the gas particles move, this speed translates (for protons) to a temperature of 40 million K. ∞ (More Precisely 8-1)

Where did the gas come from? There is so much of it that it could not have been expelled from the galaxies themselves. Instead, astronomers think that it is mainly *primordial*—gas that has been around since the universe formed and that never became part of a galaxy. However, the intracluster gas does contain some heavy elements—carbon, nitrogen, and so on—implying that at least some of it is material ejected from galaxies after enrichment by stellar evolution. ∞ (Sec. 21.5) Just how this occurred remains a mystery.



**PROCESS OF SCIENCE Check**

- ✓ What assumptions are we making when we infer the mass of a galaxy cluster from observations of the spectra of its constituent galaxies?


**◀ FIGURE 25.4 Head-Tail Radio Galaxy** (a) Radiograph, in false color, of the head-tail radio galaxy NGC 1265. (b) The same radio data, in contour form, superposed on the optical image of the galaxy. Astronomers reason that this object is moving rapidly through space, trailing a “tail” behind as it goes. (NRAO; Palomar/Caltech)

## 25.2 Galaxy Collisions

Contemplating the congested confines of a rich galaxy cluster (such as Virgo or Coma), with thousands of member galaxies orbiting within a few megaparsecs, we might expect that collisions among galaxies would be common. ∞ (Sec. 24.2) Gas particles collide in our atmosphere, and hockey players collide in the rink. So, do galaxies in clusters collide, too? The answer is yes, and this simple fact plays a pivotal role in our understanding of how galaxies evolve.


Figure 25.5 apparently shows the aftermath of a bull’s-eye collision between a small galaxy (perhaps one of the two at the right, although that is by no means certain) and the larger galaxy at the left. The result is the “Cartwheel” galaxy, about 150 Mpc from Earth, its halo of young stars resembling a vast ripple in a pond. The ripple is most likely a density wave created by the passage of the smaller galaxy through the disk of the larger one. ∞ (Sec. 23.5) The disturbance is now spreading outward from the region of impact, creating new stars as it goes.

Figure 25.6 shows an example of a close encounter that hasn’t (yet) led to an actual collision. Two spiral galaxies are apparently passing each other like majestic ships in the night. The larger and more massive galaxy on the left is called NGC 2207; the smaller one on the right is IC 2163. Analysis of this image suggests that IC 2163 is now swinging past NGC 2207 in a counterclockwise direction, having made a close approach some 40 million years ago. The two galaxies seem destined to undergo further close encounters, as IC 2163 apparently does not have enough energy to escape the gravitational pull of NGC 2207. Each time the two galaxies experience a close encounter, bursts of star formation erupt in both as their interstellar clouds of gas and dust are pushed, shoved, and shocked. In roughly a billion years, these two galaxies will probably merge into a single, massive galaxy.



▲ **FIGURE 25.5 Cosmic Cartwheel** The “Cartwheel” galaxy (left) may have resulted from a collision (possibly with one of the smaller galaxies at right) that has led to an expanding ring of star formation moving outward through the galactic disk. This is a false-color composite image combining four spectral bands: infrared in red (from *Spitzer*), optical in green (from *Hubble*), ultraviolet in blue (from *Galex*), and X-ray in purple (from *Chandra*). (NASA)

These examples illustrate how an interaction with another galaxy—a close encounter or an actual collision—can have dramatic consequences for a galaxy, especially its interstellar gas. The rapidly varying gravitational forces during the interaction compress the gas, often resulting in a galaxy-wide episode of star formation. The result is a




◀ **FIGURE 25.6 Galaxy Encounter** This encounter between two spirals, NGC 2207 (left) and IC 2163, has already led to bursts of star formation in each. Eventually the two will merge, but not for a billion years or so. (NASA)

MA

ANIMATION/VIDEO Starburst Galaxy

ANIMATION/VIDEO Galaxy Collision



**Interactive FIGURE 25.7 Starburst Galaxy** This interacting galaxy pair (IC 694, at the left, and NGC 3690) shows starbursts now under way in both galaxies—hence the bluish tint. Such intense, short-lived bursts probably last for no more than a few tens of millions of years—a small fraction of a typical galaxy’s lifetime. (W. Keel)

allow astronomers to better understand the effects of a collision on the galaxies involved and even estimate the eventual outcome of the interaction.


The particular calculation shown in Figure 25.8(b) began with two colliding spiral galaxies, not so different from those shown in Figure 25.6, but the details of the original structure have been largely obliterated by the collision. Notice the similarity to the real image of NGC 4038/4039 (Figure 25.8a), the so-called Antennae galaxies, which show extended tails, as well as double galactic centers only a few hundred parsecs across. Star formation induced by the collision is clearly traced by the blue light from thousands of young, hot stars. The simulations indicate that, as with the galaxies in Figure 25.6, ultimately the two galaxies will merge into one.

Galaxies in clusters apparently collide quite often. Many collisions and near misses similar to those shown in the previous figures have been observed (see also Section 24.4), and a straightforward calculation reveals that, given the crowded conditions in even a modest cluster, close


**starburst galaxy**, a spectacular example of which is shown in Figure 25.7.

No human will ever witness an entire galaxy collision, for it lasts many millions of years. However, computers can follow the event in a matter of hours. Simulations modeling in detail the gravitational interactions among stars and gas, and incorporating the best available models of gas dynamics,

*Collisions seen in these real images at left can be studied in computer simulations like that at right.*



*Such simulations demonstrate the crucial role played by dark-matter halos during galaxy interactions.*



**▲ FIGURE 25.8 Galaxy Collision** (a) The long tidal “tails” (black-and-white image at left) mark the final plunge of the “Antennae” galaxies a few tens of millions of years ago. Strings of young, bright “super star clusters” (magnified color image at center) were caused by violent shock waves produced in the gas disks of the two colliding galaxies. (b) A computer simulation of the encounter shows many of the same features as the real object at left. (AURA; NASA; J. Barnes)

encounters are the norm rather than the exception. The reason is simple: The distance between adjacent galaxies in a cluster averages a few hundred thousand parsecs, which is not much greater (certainly less than five times more) than the size of a typical galaxy, including its extended dark halo. Galaxies simply do not have that much room to roam around without bumping into one another. Many researchers think that most galaxies in most clusters have been strongly influenced by collisions, in some cases in the relatively recent past.

Computer simulations clearly show that the extensive dark-matter halos surrounding most, if not all, galaxies are crucial to galaxy collisions. The dark halos make the galaxies much larger than their optical appearance would suggest, making interactions and mergers all the more likely. Consider two galaxies approaching one another. As they orbit, the galaxies interact with each other's dark halos, slowing the galaxies' motion and stripping halo material by tidal forces. The halo matter is redistributed between the galaxies or is entirely lost from the system. In either case, the result is a much stronger interaction that can greatly change the orbits of both galaxies.

In the smaller groups, the galaxies' speeds are low enough that interacting galaxies tend to “stick together,” and mergers, as shown in the computer simulation, are the most common outcome. In larger groups, galaxies move faster and tend to pass through one another without sticking. Either way, the encounters have substantial effects on the galaxies involved (see Section 25.3). If we wait long enough, we will have an opportunity to see for ourselves what a galaxy collision is really like: Our nearest large neighbor, the Andromeda Galaxy (Fig. 23.2), is currently approaching the Milky Way at a velocity of 120 km/s. In a few billion years it will collide with our Galaxy, and we will then be able to test astronomers' theories firsthand!

Curiously, although a collision may wreak havoc on the large-scale structure of the galaxies involved, it has essentially no effect on the individual stars they contain. The stars within each galaxy just glide past one another. In contrast to galaxies in the cluster, the stars in a galaxy are so small compared with the distances between them that when two galaxies collide, the star population merely doubles for a time, and the stars continue to have so much space that they do not run into each other. Collisions can rearrange the stellar and interstellar contents of each galaxy, often producing a spectacular burst of star formation that may be visible to enormous distances, but from the point of view of the stars, it's clear sailing.

### CONCEPT Check

✓ What role do collisions play in the evolution of galaxies?

## 25.3 Galaxy Formation and Evolution

With Hubble's law as our guide to distances in the universe, and armed now with knowledge of the distribution of dark matter on galactic and larger scales, let's turn to the question of how galaxies came to be the way they are. Can we explain the different types of galaxy we see? Astronomers know of no simple evolutionary connections among the various categories in the Hubble classification scheme. ∞ (Sec. 24.1) To answer the question, we must understand how galaxies formed.


Unfortunately, compared with the theories of star formation and stellar evolution, the theory of galaxy formation and evolution is still very much in its infancy. Galaxies are far more complex than stars, they are harder to observe, and the observations are harder to interpret. In addition, we have only partial information on conditions in the universe during the formation process, quite unlike the corresponding situation for stars. ∞ (Sec. 18.3) Finally, and most important, stars almost never *collide* with one another, with the result that most single stars and binaries evolve in isolation. Galaxies, however, may suffer numerous collisions during their lives, making it much harder to decipher their pasts. Indeed, collisions like those described in the previous section blur the distinction between formation and evolution to the point where it can be hard to separate one from the other.

Nevertheless, some general ideas have gained widespread acceptance, and we can offer some insights into the processes responsible for the galaxies we see. We first describe a general scenario for how small galaxies merge to form larger ones, then discuss how galaxies change in time due to both internal stellar evolution and external influences. Finally, we consider how the galaxy types in Hubble's classification fit in to this broad picture.

### Mergers and Acquisitions

The seeds of galaxy formation were sown in the very early universe, when small density fluctuations in the primordial matter began to grow (see Section 27.5). Our discussion here begins with these “pregalactic” blobs of gas already formed. The masses of the various fragments were quite small—only a few million solar masses, comparable to the masses of the smallest present-day dwarf galaxies, which may in fact be remnants of that early time. Most astronomers think that galaxies grew by repeated *merging* of smaller objects, as illustrated in Figure 25.9(a). Contrast this with the process of star formation, in which a large cloud fragments into smaller pieces that eventually become stars. ∞ (Sec. 19.2)

Theoretical evidence for this picture of **hierarchical merging** is provided by computer simulations of the early



**▲ FIGURE 25.9 Galaxy Formation** (a) The best current theory of galaxy formation holds that large systems were built up from smaller ones through collisions and mergers, as shown schematically in the drawing at left. (b) This photograph, one of the deepest ever taken of the universe, provides “fossil evidence” for hundreds of galaxy shards and fragments, up to 5000 Mpc distant. (c) Enlargements of selected portions of (b) reveal rich (billion-star) “star clusters,” all lying within a relatively small volume of space (about 1 Mpc across). Such pregalactic fragments might be about to merge to form a galaxy. The events pictured took place about 10 billion years ago. (NASA)

universe, which clearly show the process taking place. Further strong support comes from observations that galaxies at large redshifts (meaning that they are very distant and the light we see was emitted long ago) are distinctly smaller and more irregular than those found nearby. Figure 25.9(b) (see also Figure 25.10) shows some of these images, which include objects up to 5 billion parsecs away. The vague bluish patches are separate small galaxies, each containing only a few percent of the mass of the Milky Way Galaxy. Their irregular shape is thought to be the result of galaxy mergers; the bluish coloration comes from young stars that formed during the merging process.

Figure 25.9(c) shows more detailed views of some of the objects in Figure 25.9(b), all lying in the same region of space, about 1 Mpc across and almost 5000 Mpc from Earth. Each blob seems to contain several billion stars spread throughout a distorted spheroid about a kiloparsec across. Their decidedly bluish tint suggests that active star formation is already underway. We see them as they were nearly 10 billion years ago, a group of young galaxies possibly poised to merge into one or more larger objects.


Hierarchical merging provides the conceptual framework for all modern studies of galaxy evolution. It describes a process that began billions of years ago and

continues (albeit at a greatly reduced rate) to the present day, as galaxies continue to collide and merge. By studying how galaxy properties vary with distance, and hence look-back time, astronomers try to piece together the merger history of the universe. [∞ \(More Precisely 24-1\)](#)

Figure 25.10 is a remarkable image from the *Hubble Space Telescope* showing billions of years of galaxy evolution in a single tiny patch of the sky. The large, bright galaxies with easily discernible Hubble types are mostly (according to their redshifts) relatively nearby objects. They are seen here against a backdrop of small, faint, irregular galaxies lying much farther away. The size and appearance of these distant galaxies compared with those in the foreground strongly support the basic idea that galaxies were smaller and less regular in the past.

## Evolution and Interaction

Left alone, a galaxy will evolve slowly and fairly steadily as interstellar clouds of gas and dust are turned into new generations of stars and main-sequence stars evolve into giants and, ultimately, into compact remnants—white dwarfs, neutron stars, and black holes. The galaxy’s overall color, composition, and appearance change in a more or



◀ **FIGURE 25.10 Hubble Deep Field** Numerous small, irregularly shaped young galaxies can be seen in this very deep optical image. Known as the Hubble Deep Field–North, this image, made with an exposure of approximately 100 hours, captured objects as faint as 30th magnitude. ∞ (Sec. 17.2) (As in Figure 25.9, “deep” in this context implies “faint,” meaning that we are looking at objects far away and as they were long ago.) Redshift measurements (as denoted by the superposed values observed at the Keck Observatory in Hawaii) indicate that some of these galaxies lie well over 1000 Mpc from Earth. ∞ (More Precisely 24-1) The field of view is about 2 arc minutes across, less than one-tenth the angular diameter of the full Moon. (NASA; Keck)

correspondingly more frequent. ∞ (More Precisely 24-1) We see the majority of these violent events as they unfolded roughly 10 billion years ago. The galaxy interactions observed locally are extensions of this same basic process into the present day. Figure 25.11 presents a graphical and artistic summary of these (mostly) ancient events.

Although the hierarchical merging scenario accounts

less predictable way as the cycle of stellar evolution recycles and enriches the galaxy’s interstellar matter. ∞ (Sec. 21.5) If the galaxy is an elliptical, lacking interstellar gas, it will tend to become fainter and redder in time as the more massive stars burn out and are not replaced. ∞ (Sec. 20.5) For a gas-rich galaxy, such as a spiral or an irregular, hot, bright stars will lend a bluish coloration to the overall light for as long as gas remains available to form them.


But many—perhaps most—galaxies are *not* alone. They reside in small groups and clusters, and, as we have just seen, may *interact with other galaxies* repeatedly over extended periods of time. As described in the previous section, these interactions can rearrange a galaxy’s internal structure, compressing interstellar gas, and triggering sudden, intense bursts of star formation. Encounters may also divert fuel to a central black hole, powering violent activity in some galactic nuclei. ∞ (Sec. 24.4) Thus, starbursts and nuclear activity are key indicators of interactions and mergers between galaxies.

Careful studies of starburst galaxies and active galactic nuclei indicate that most galactic encounters probably took place long ago—at redshifts greater than about 1, when the clusters were more compact and galaxy collisions were

well for the numbers and total masses (including dark matter) of galaxies over the history of the universe, in recent years astronomers have come to realize that it is not the whole story. Specifically, this scenario has difficulty explaining both the distribution of gas in galaxies and the *rates* at which stars are observed to form (Figure 25.11a), up to and including the present day. Many disk galaxies have asymmetric or warped gas disks, often with more gas found above or below the disk plane than would be expected from theory. In addition, the observed star formation rate in the universe is *larger* than we would expect if galaxies were simply consuming the gas they formed with long ago. Instead, as in our own Milky Way Galaxy, star formation and galaxy growth appear to be both enhanced and prolonged by the continual infall of fresh gas from intergalactic space. ∞ (Sec. 23.4) Supporting evidence for this picture comes from radio observations of some “small” distant galaxies, like those shown in Figure 25.9, which reveal that the visible galaxies are in fact surrounded by large, cool disks of mostly hydrogen gas.

Thus we have ample evidence that galaxies evolved, and are still evolving, in response to external factors long after the first pregalactic fragments formed and merged.





**▲ FIGURE 25.11 Galaxies Build and Stars Form** (a) This graph, derived from observations of the luminosities of many different galaxies at various distances from us, implies that the star formation rate peaked a few billion years after the Big Bang, when accreting material contained within dwarf galaxies helped to grow the bigger galaxies, as suggested by the artist's sketch in (b).

## Types of Merger

The different types and masses of galaxies can lead to an almost bewildering variety of possible interactions. Here we consider just a few of the many possibilities.




**▲ FIGURE 25.12 Galactic Cannibalism** This is a dramatic glimpse of a large and massive galaxy under assembly by the merging of smaller, lighter galaxies. Most galaxies probably developed in this way in the earlier universe—by means of a “bottom up” scenario that hierarchically built really big objects by merging star-rich building blocks. This image captures a formative process that occurred about 10 billion years ago, only a few billion years after the Big Bang. The bigger image highlights a region at upper left (in the white box) catalogued MRC 1138-262 and nicknamed the “Spiderweb” Galaxy. The inset shows more clearly dozens of small galaxies about to merge into a single huge object. (NASA/ESA)

If one galaxy of an interacting pair happens to have a much lower mass than the other, its interaction with the other's halo causes it to spiral inward, ultimately to be disrupted near the center of the larger system. This process is colloquially termed *galactic cannibalism* and may explain why supermassive galaxies are often found at the cores of rich galaxy clusters. Having “dined” on their companions, they now lie at the center of the cluster, waiting for more “food” to arrive. Figure 25.12 is a remarkable combination of images that has apparently captured this process at work in a distant cluster.

We also have examples of galactic cannibalism closer to home. The small Sagittarius dwarf galaxy (Figure 24.13) is already well on its way to suffering a similar fate at the center of the Milky Way, and theory indicates that the Magellanic Clouds (Figure 24.7) will eventually meet the same end. Figure 25.13(a) illustrates how the




disruption of a dwarf galaxy by the Milky Way leaves behind a *tidal stream* of stripped stars, all with similar orbits and composition, still following the orbital path of their parent galaxy.


Astronomers have discovered numerous such streams in the halo of the Milky Way that are thought to be the result of precisely this process.  (Sec. 23.3) Figure 25.13(b) is a wide-angle *Sloan Digital Sky Survey* mosaic (see *Discovery 25-1*) of roughly half of the northern sky, looking in the direction away from the Galactic center, showing several streams of stars crossing the field of view. The most prominent streams (marked) represent two orbits of the Sagittarius dwarf over the past 500 million years. Their locations in the sky are consistent with the measured properties of the Sagittarius galaxy, which currently lies in the opposite direction, as seen from Earth.

Now consider two interacting disk galaxies, one a little smaller than the other, but each having a mass comparable to the Milky Way Galaxy. As shown in the computer-generated frames of Figure 25.14, the smaller galaxy can distort the larger one, substantially, causing spiral arms to appear where none existed before, triggering an extended episode of star formation. The entire event requires several hundred million


years—a span of evolution that a supercomputer can model in minutes. The final frame of the figure looks remarkably similar to the double galaxy shown in the opening photo for Chapter 23, and in fact, the simulation was constructed to mimic the sizes, shapes, and velocities in that binary galaxy system. The magnificent spiral galaxy is M51, popularly known as the Whirlpool Galaxy, about 10 Mpc from Earth. Its smaller companion is an irregular galaxy that may have drifted past M51 millions of years ago.

What if the colliding galaxies are comparable in size and mass? Computer simulations reveal that such a merger can destroy a spiral galaxy's disk, creating a galaxywide starburst episode. The violence of the merger and the effects of subsequent supernovae eject much of the remaining gas into intergalactic space, creating the hot intracluster gas noted in Section 24.1.  (Sec. 24.1) Once the burst of star formation has subsided, the resulting object looks very much like an elliptical galaxy. The elliptical's hot X-ray halo is the last vestige of the original spiral's disk. The merging galaxies in Figures 25.7 and 25.8 may be examples of this phenomenon in progress.


## Making the Hubble Sequence

If galaxies form and evolve by repeated mergers, can we account for the Hubble sequence and, specifically, differences between spirals and ellipticals?  (Sec. 24.1) The details are still far from certain, but, remarkably, the answer now seems to be a qualified yes. Collisions and close encounters are random events and do *not* represent a “genuine” evolutionary connection between galaxies. Nevertheless, observations and computer simulations do suggest some plausible ways in which the observed Hubble types might have arisen, starting from a universe populated only by irregular, gas-rich galaxy fragments.

As we have just seen, the simulations reveal that “major” mergers—collisions between large galaxies of comparable size—tend to destroy galactic disks, effectively turning spirals into ellipticals (Figure 25.15a). On the other hand, “minor” mergers, in which a small galaxy interacts with, and ultimately




(a)



(b)

**◀ FIGURE 25.13 Tidal Streams in the Milky Way** (a) This illustration depicts the breakup and dispersion of an incoming star-rich galaxy companion captured by our Milky Way. Eventually, the smaller galaxy dissolves within the larger one, a case of the littler one being “digested,” much as other dwarf companion galaxies were consumed by our Galaxy long ago. (b) This outer region of the Milky Way shows innumerable stars that have been torn from our Galaxy’s disrupted satellite galaxies (colors indicate distance, with blue being the closest). Several tidal streams are evident, the biggest one at center showing two orbits of the enormous, arching death spiral of the Sagittarius dwarf galaxy. (V. Belokurov; SDSS)

Environmental influences can seriously affect how galaxies evolve.




**▲ FIGURE 25.14 Galaxy Interaction** Galaxies can change their shapes long after their formation. In this computer-generated sequence, two galaxies closely interact over several hundred million years. The smaller galaxy, in red, has gravitationally disrupted the larger galaxy, in blue, changing it into a spiral galaxy. Compare the result of this supercomputer simulation with Figure 24.2(b), a photograph of the Whirlpool Galaxy and its small companion. (*J. Barnes & L. Hernquist*)

ANIMATION/VIDEO Galaxy Merger

is absorbed by, a larger one, generally leave the larger galaxy intact, with more or less the same Hubble type as it had before the merger (Figure 25.15b). This is the most likely way for large spirals to grow—in particular, our own Galaxy probably formed in such a manner.

Supporting evidence for this general picture comes from observations that spiral galaxies are relatively rare in regions of high galaxy density, such as the central regions of rich galaxy clusters. These observations are consistent with the view that the fragile disks of spiral galaxies are easily destroyed by collisions, which are more common in dense galactic environments. Spirals also seem to be more common at larger redshifts (that is, in the past), implying that their numbers are decreasing with time, presumably also as the result of collisions. However, nothing in this area of astronomy is clear-cut, and astronomers know of numerous isolated elliptical galaxies in low-density regions of the universe that are hard to explain as the result of mergers. In addition, the competition between infall, which acts to sustain galactic disks, and collisions, which tend to destroy them, remains poorly understood, as is the effect of activity in galactic nuclei, to be discussed in Section 25.4.

In principle, the starbursts associated with galaxy mergers leave their imprint on the star-formation history of the universe in a way that can be correlated with the properties of galaxies. As a result, studies of star formation in distant galaxies have become a very important way of testing and quantifying the details of the entire hierarchical merger scenario.



**▲ FIGURE 25.15 Galaxy Mergers** (a) When comparably sized galaxies come together, the result is probably an elliptically shaped galaxy, as their original arms and disks do not likely survive the encounter. (b) By contrast, if a large spiral absorbs a smaller companion, the probable result is merely a larger spiral, with much of its original geometry unchanged.

### CONCEPT Check

- ✓ Other than scale, in what important ways does galaxy evolution differ from that of stars?

## DISCOVERY 25-1

## The Sloan Digital Sky Survey

Many of the photographs used in this book—not to mention most of the headline-grabbing imagery found in the popular media—come from large, high-profile, and usually very expensive instruments such as NASA’s *Hubble Space Telescope* and the European Southern Observatory’s Very Large Telescope in Chile. ∞ (Secs. 5.3, 5.4) Their spectacular views of deep space have revolutionized our view of the universe. Yet a less well-known, considerably cheaper, but no less ambitious, project may, in the long run, have every bit as great an impact on astronomy and our understanding of the cosmos.

The Sloan Digital Sky Survey (SDSS), a 5-year project that began scientific operation in 2000 and has since been extended until 2014, was designed to systematically map out a quarter of the entire sky on a scale and at a level of precision never before attempted. It has cataloged almost 1 billion celestial objects, recording their apparent brightnesses at five different colors (wavelength ranges) spread across the optical and near-infrared part of the spectrum. In addition, spectroscopic follow-up observations have determined redshifts and hence distances to 1.5 million galaxies and 230,000 quasars. These data have been used to construct detailed redshift surveys (see Section 26.1), and to probe the structure of the universe on very large scales. The sensitivity of the survey is such that it can detect bright galaxies like our own out to distances of more than 1 billion parsecs. Very bright objects, such as quasars and young starburst galaxies, are detectable almost throughout the entire observable universe.

The first figure shows the Sloan Survey telescope, a special-purpose 2.5-m instrument sited in Apache Point Observatory, near Sunspot, New Mexico. This reflecting telescope is not space-based, does not employ active or adaptive optics, and cannot probe as deeply (that is, far) into space as larger instruments. How can it possibly compete with these other systems? The answer is that, unlike most other large telescopes in current use, where hundreds or even thousands of observers share the instrument and compete for its time, the SDSS telescope was designed specifically for the purpose of the survey. It has a wide field of view and is dedicated to the task, carrying out observations of the sky on *every* clear night during the duration of the project.

The use of a single instrument night after night, combined with tight quality controls on which nights’ data are actually incorporated into the survey (nights with poor seeing or other problematic conditions are discarded and the observations repeated) mean that the end product is a database of exceptionally high quality and uniformity spanning an enormous volume of space—a monumental achievement and an indispensable tool for the study of the universe. The survey field of view covers much of the sky away from the Galactic plane in the north, together with a large swath of the sky around the Galactic south pole.

Archiving images and spectra on millions of galaxies produces a lot of data. The full survey consists of roughly 60 *trillion* bytes of information—comparable to the entire Library of Congress! All of it has been released to the public. The second figure shows an image of the Perseus galaxy cluster, just one of hundreds of thousands of images that make up the full dataset. Among



recent highlights, SDSS has detected the largest known structure in the universe, observed the most distant known galaxies and quasars, and has been instrumental in pinning down the key observational parameters describing our universe (see Chapter 26).

SDSS impacts astronomy in areas as diverse as the large-scale structure of the universe, the origin and evolution of galaxies, the nature of dark matter, the structure of the Milky Way, the properties and distribution of interstellar matter, and the properties of exoplanetary systems. Its uniform, accurate, and detailed database is likely to be used by generations of scientists for decades to come. Its success has spawned several even more ambitious follow-up surveys; the first is due to become operational around 2015.



(SDSS; R. Lupton)



## 25.4 Black Holes in Galaxies


Now let's ask how quasars and active galaxies fit into the framework of galaxy evolution just described. The fact that quasars are more common at great distances from us demonstrates that they were much more prevalent in the past than they are today. ∞ (Sec. 24.4) Quasars have been observed with redshifts of up to 7.1 (the current record, as of early 2013), so the process must have started at least 13 billion years ago (see Table 24.2). However, most quasars have redshifts between 2 and 3, corresponding to an epoch some 2 billion years later. Most astronomers agree that quasars represent an early stage of galaxy evolution—an “adolescent” phase of development, prone to frequent flare-ups and “rebellion” before settling into more steady “adulthood.” This view is reinforced by the fact that the same black hole energy-generation mechanism can account for the luminosities of quasars, active galaxies, and the central regions of normal galaxies like our own.

### Black Hole Masses

In Chapter 24 we saw the standard model of active galactic nuclei accepted by most astronomers—accretion of gas onto a supermassive black hole. ∞ (Sec. 24.5) We also saw that a large fraction of all “bright” galaxies exhibit activity of some sort, even though in many cases it represents only a small fraction of the galaxy's total energy output. This suggests that these galaxies may also harbor central black holes, with the potential of far greater activity under the right circumstances. Our own Galaxy is a case in point. ∞ (Sec. 23.7) The 4-million-solar-mass black hole at the center of the Milky Way is not currently active, but if fresh fuel were supplied (say, by a star or molecular cloud coming too close to the hole's intense gravitational field), it might well become a (relatively weak) active galactic nucleus.

In recent years, astronomers have found that many bright normal galaxies contain supermassive black holes at their centers. Figure 25.16 presents perhaps the most compelling evidence for such a black hole in a normal galaxy. It comes from a radio study of NGC 4258, a spiral galaxy about 6 Mpc away. Using the Very Long Baseline Array, a continent-wide interferometer comprising 10 radio telescopes, a U.S.-Japanese team has achieved an angular resolution hundreds of times better than that attainable with *HST*. ∞ (Sec. 5.6) The observations reveal a group of molecular clouds swirling in an organized fashion about the galaxy's center. Doppler measurements indicate a slightly warped, spinning disk centered precisely on the galaxy's heart. The rotation speeds imply the presence of more than 40 million solar masses packed into a region less than 0.2 pc across.

Similar evidence exists for supermassive black holes in the nuclei of several dozen bright galaxies—some normal, some active—within a few tens of megaparsecs of the Milky Way. Some observers would go so far as to say that in every




▲ **FIGURE 25.16 Galactic Black Hole** A network of radio telescopes has probed the core of the spiral galaxy NGC 4258, shown here in the light of mostly hydrogen emission. Within the innermost region (inset), a disk of Doppler-shifted molecular clouds (designated by red, green, and blue dots) obey Kepler's third law perfectly, apparently revealing a huge black hole at the center of the disk. (J. Moran)

case where a galaxy has been surveyed and a black hole *could* have been detected, given the resolution and the sensitivity of the observations, a black hole *has* in fact been found. It is a small step to the remarkable conclusion that *every bright galaxy—active or not—contains a central supermassive black hole*. This unifying principle connects our theories of normal and active galaxies in a fundamental way.

Astronomers have also found that there is a correlation between the masses of the central black holes and the properties of the galaxy in which they reside. As illustrated in Figure 25.17, the largest black holes tend to be found in the most massive galaxies (as measured by the mass of the bulge). The reason for this correlation is not fully understood, but most astronomers take it to mean that, at the very least, the evolution of normal and active galaxies must be very closely connected, as we now discuss.

### The Quasar Epoch

Where did the supermassive black holes in galaxies come from? To be honest, the processes whereby the first



▲ **FIGURE 25.17 Black-Hole Masses** Observations of nearby normal and active galaxies reveal that the mass of the central black hole is closely correlated with the mass of the galactic bulge. In this diagram, each point represents a different galaxy. The straight line is the best fit to the data points for many galaxies, implying a black-hole mass of 1/200 the mass of the bulge.

billion-solar-mass black holes formed early in the history of the universe are not fully understood. However, the accretion responsible for quasar energy emission also naturally accounts for the mass of the black hole—only a few percent of the infalling mass is converted to energy; the rest is trapped forever in the black hole once it crosses the event horizon. ∞ (Sec. 22.5) Simple estimates suggest that the accretion rates needed to power the quasars are generally consistent with black hole masses inferred by other means.

Since the brightest known quasars devour about a thousand solar masses of material every year, it is unlikely that they could maintain their luminosity for very long—even a million years would require a billion solar masses, enough to account for the most massive black holes known. ∞ (Sec. 24.4) This suggests that a typical quasar spends only a fairly short amount of time in its highly luminous phase—perhaps only a few million years in some cases—before running out of fuel. Thus, most quasars were relatively brief events that occurred long ago.


To make a quasar, we need a black hole and enough fuel to power it. Although fuel was abundant at early times in the universe’s history in the form of gas and newly formed stars, black holes were not. They had yet to form, most probably by the same basic stellar evolutionary processes we saw in Chapter 21, although the details are again not well-known. ∞ (Sec. 21.2) The building blocks of the supermassive black holes that would ultimately power the quasars may well have


been relatively small black holes having masses tens or perhaps a few hundreds of times the mass of the Sun, formed by the first generations of stars. These small black holes sank to the center of their still-forming parent galaxy and merged to form a single, more massive black hole.


As galaxies merged, so, too, did their central black holes, and eventually supermassive (1-million-solar-mass to 1-billion-solar-mass) black holes existed in the centers of many young galaxies. Some supermassive black holes may have formed directly by the gravitational collapse of the dense central regions of a protogalactic fragment or perhaps by accretion or a rapid series of mergers in a particularly dense region of the universe. These events resulted in the earliest (redshift 6–7) quasars known, shining brightly 13 billion years ago. However, in most cases, the mergers took longer—roughly another 2 billion years. By then (at redshifts between 2 and 3, roughly 11 billion years ago), many supermassive black holes had formed, and there was still plenty of merger-driven fuel available to power them. This was the height of the “quasar epoch” in the universe.

Until recently, astronomers were confident that black holes would merge when their parent galaxies collided, but they had no direct evidence of the process—no image of two black holes “caught in the act.” In 2002, the *Chandra X-ray Observatory* discovered a binary black hole—two supermassive objects, each having a mass a few tens of millions of times that of the Sun—in the center of the ultraluminous starburst galaxy NGC 6240, itself the product of a galaxy merger some 30 million years ago. Figure 25.18 shows optical and X-ray views of the system. The black holes are the two blue-white objects near the center of the (false-color) X-ray image. Orbiting just 1000 pc apart, they are losing energy through interactions with stars and gas and are predicted to merge in about 400 million years. Astronomers now know of several binary black holes in relatively nearby galaxies, caught in the act of spiraling together on their way to merging. NGC 6240 lies just 120 Mpc from Earth, so we are far from seeing a quasar merger in the early universe. Nevertheless, astronomers think that events similar to this must have occurred countless times billions of years ago, as galaxies collided and quasars blazed.

Distant galaxies are generally much fainter than their bright quasar cores. As a result, until quite recently, astronomers were hard-pressed to discern any galactic structure in quasar images. Since the mid-1990s, several groups of astronomers have used the *Hubble Space Telescope* to search for the “host” galaxies of moderately distant quasars. After removing the bright quasar core from the *HST* images and carefully analyzing the remnant light, the researchers have reported that, in every case studied—several dozen quasars so far—a host galaxy can be seen enveloping the quasar. Figure 25.19 shows some of the longest quasar exposures ever taken. Even without sophisticated computer processing, the hosts are clearly visible.




**▲ FIGURE 25.18 Binary Black Hole** These (a) optical (*Hubble*) and (b) X-ray (*Chandra*) images of the starburst galaxy NGC 6240 show two supermassive black holes (the blue-white objects near the center of the X-ray image) orbiting about 1 kpc apart. Theoretical estimates imply that they will merge in about 400 million years, releasing an intense burst of gravitational radiation.  (Discovery 22-2) The colors of the optical image are real; the false colors in the X-ray image indicate a range of energies. (NASA)

As we saw in Chapter 24, the connection between active galaxies and galaxy clusters is well established, and many relatively nearby quasars are also known to be members of clusters.  (Sec. 24.4) The link is less clear-cut for the most distant quasars, however, simply because they are so far away that other cluster members are very faint and extremely hard to see. However, as the number of known quasars continues to increase, evidence for quasar clustering (and presumably, therefore, for quasar membership in young galaxy clusters) mounts. Thus, as best we can tell,

quasar activity is intimately related to interactions and collisions in young galaxy clusters.

This connection also suggests a possible way in which the growth of black holes might be tied to the growth of their parent galaxies. Many astronomers think that a process called **quasar feedback**, in which some fraction of the quasar’s enormous energy output is absorbed by the surrounding galactic gas, might explain the correlation of black hole and bulge masses shown in Figure 25.17. In this picture, which is appealing but by no means certain, the absorbed energy expels the gas from the galaxy, simultaneously shutting down both galactic star formation and the quasar’s own fuel supply, thus tying the growth of the central black hole to the formation of new stars in the bulge.



**◀ FIGURE 25.19 Quasar Host Galaxies** These long-exposure images of distant quasars show the young host galaxies in which the quasars reside, lending support to the idea that quasars represent an early, highly luminous phase of galactic evolution. The quasar at the top left is the best example, having the catalog name PG0052 + 251 and residing roughly 700 Mpc from Earth. (NASA)

## Active and Normal Galaxies


Early on, frequent mergers may have replenished the quasar's fuel supply, extending its luminous lifetime. However, as the merger rate declined, these systems spent less and less of their time in the "quasar" phase. The rapid decline in the number of bright quasars roughly 10 billion years ago marks the end of the quasar epoch. Today, the number of quasars has dropped virtually to zero (recall that the nearest lies hundreds of megaparsecs away). ∞ (Sec. 24.4)

Large black holes do not simply vanish. If a galaxy contained a bright quasar 10 billion years ago, the black hole responsible for all that youthful activity must still be present in the center of the galaxy today. We see some of these black holes as active galaxies. The remainder reside dormant in normal galaxies all around us. In this view, *the difference between an active galaxy and a normal one is mainly a matter of fuel supply*. When the fuel runs out and a quasar shuts down, its central black hole remains behind, its energy output reduced to a relative trickle. The black holes at the hearts

of normal galaxies are simply quiescent, awaiting another interaction to trigger a new active outburst. Occasionally, two nearby galaxies may interact with each other, causing a flood of new fuel to be directed toward the central black hole of one or both. The engine starts up for a while, giving rise to the nearby active galaxies—radio galaxies, Seyferts, and others—we observe.

Should this general picture be correct, it follows that many relatively nearby galaxies (but probably *not* our own Milky Way, whose central black hole is even now only a paltry 3–4 million solar masses) must once have been brilliant quasars. ∞ (Sec. 23.7) Perhaps some alien astronomer, thousands of megaparsecs away, is at this very moment observing the progenitor of M87 in the Virgo cluster—seeing it as it was billions of years ago—and is commenting on its enormous luminosity, nonstellar spectrum, and possibly its high-speed jets, and wondering what exotic physical process can possibly account for its violent activity! ∞ (Sec. 24.4)

Finally, Figure 25.20 suggests some possible (but unproven!) evolutionary connections among quasars, active



**Narrated FIGURE 25.20 Galaxy Evolution** Most evolutionary sequences for galaxies begin with galaxy



mergers that lead to highly luminous quasars, after which they decrease in violence through the radio and Seyfert galaxies, eventually resulting in normal ellipticals and spirals. The central black holes that powered the early activity remain at later times, but many of them have run out of fuel.



galaxies, and normal galaxies. If the largest black holes reside in the most massive galaxies, and also tend to power the brightest active galactic nuclei, then we would expect that the most luminous nuclei should reside in the largest galaxies, which probably came into being via “major” mergers of other large galaxies. Since the products of such mergers are elliptical galaxies, we have a plausible explanation of why the brightest active galaxies—the radio galaxies—should be associated with large ellipticals. ∞ (Sec. 24.4) Furthermore, the path to spiral galaxies would necessarily have entailed a series of mergers involving smaller galaxies, resulting in the less-violent Seyferts along the way.

## Active Galaxies and the Scientific Method

When active galactic nuclei—especially quasars—were first discovered, their extreme properties defied conventional explanation. Initially, the idea of supermassive (million- to billion-solar-mass) black holes in galaxies was just one of several competing, and very different, hypotheses advanced to account for the enormous luminosities and small sizes of those baffling objects. However, as observational evidence mounted, the other hypotheses were abandoned one by one, and massive black holes in galactic nuclei became first the leading, and eventually the standard, theory of active galaxies.

As often happens in science, a theory once itself considered extreme is now the accepted explanation for these phenomena. Far from threatening the laws of physics, as some

astronomers once feared, active galaxies are now an integral part of our understanding of how galaxies form and evolve. The synthesis of studies of normal and active galaxies, galaxy formation, and large-scale structure is one of the great triumphs of extragalactic astronomy.

### CONCEPT Check

✓ Does every galaxy have the potential for activity?

## 25.5 The Universe on Large Scales


Many galaxies, including our own, are members of galaxy clusters—megaparsec-sized structures held together by their own gravity. ∞ (Sec. 24.2) Our own small cluster is called the Local Group. Figure 25.21 shows the locations of the Virgo Cluster, the closest “large” large cluster, and several other well-defined clusters in our cosmic neighborhood. The region displayed is about 70 Mpc across. Each point in the figure represents an entire galaxy whose distance has been determined by one of the methods described in Chapter 24.

### Clusters of Clusters

Do galaxy clusters top the cosmic hierarchy, or does the universe have even larger groupings of matter? Most astronomers have concluded that the galaxy clusters are themselves clustered, forming titanic agglomerations of matter known as **superclusters**.

Together, the galaxies and clusters in the vicinity of the Milky Way form the *Local Supercluster*, also known as the Virgo Supercluster. Aside from the Virgo Cluster itself, it contains the Local Group and numerous other clusters lying within about 20–30 Mpc of Virgo. Figure 25.21 shows a three-dimensional rendering of our extended cosmic neighborhood, illustrating the Virgo Supercluster (near the center) relative to other “nearby” galaxy superclusters within a vast imaginary rectangle roughly 100 Mpc on its short side.

All told, the Local Supercluster is about 40–50 Mpc across, contains some  $10^{15}$  solar masses of material (several tens of thousands of galaxies), and is very irregular in shape. The Local Supercluster is significantly elongated



◀ **FIGURE 25.21 Virgo Supercluster in 3-D** The elongated structure of the Virgo Supercluster (left center) is mapped relative to other neighboring galaxy superclusters within about a billion light-years of the Milky Way (which resides within the small dot labeled Local Group near the center of this vast map). Individual galaxies are not shown; rather, smoothed contour plots outline galaxy clusters, each named or numbered by its most prominent member.

perpendicular to the line joining the Milky Way to Virgo, with its center lying near the Virgo Cluster. By now it should come as no surprise that the Local Group is *not* found at the heart of the Local Supercluster—we live far off in the periphery, about 18 Mpc from the center.

## Redshift Surveys

The farther we peer into deep space, the more galaxies, clusters of galaxies, and superclusters we see. Is there structure on scales even larger than superclusters? To answer these questions, astronomers use Hubble's law to map out the distribution of galaxies in the universe.

Figure 25.22 shows part of an early survey of the universe performed by astronomers at Harvard University in the 1980s. Using Hubble's law as a distance indicator, the team systematically mapped out the locations of galaxies within about 200 Mpc of the Milky Way in a series of wedge-shaped "slices," each  $6^\circ$  thick, starting in the northern sky. The first slice (shown in the figure) covered a region of the sky containing the Coma Cluster (see Figure 24.1), which happens to lie in a direction almost perpendicular to our Galaxy's plane. Because redshift is used as the primary distance indicator, these studies are known as *redshift surveys*.


The most striking feature of maps such as that of Figure 25.22 is that the distribution of galaxies on very large scales is decidedly nonrandom. The galaxies appear to be arranged in a network of strings, or filaments, surrounding large, relatively unpopulated regions of space known as **voids**. Voids account for some 50 percent of the total volume of the nearby universe, but only 5–10 percent of the mass. The largest measure some 100 Mpc across. The most likely

explanation for the voids and the filamentary structure shown in the figure is that the galaxies and galaxy clusters are spread across the surfaces of vast "bubbles" in space. The voids are the interiors of these gigantic bubbles. The galaxies seem to be distributed like beads on strings only because of the way our slice of the universe cuts through the bubbles. Like suds on soapy water, these bubbles fill the entire universe. The densest clusters and superclusters lie in regions where several bubbles meet. The elongated shape of the Virgo Supercluster (Figure 25.21) is a local example of this same filamentary structure.


Most theorists think that this "frothy" distribution of galaxies, and in fact all structure on scales larger than a few megaparsecs, traces its origin directly to conditions in the very earliest stages of the universe (Chapter 27). Consequently, studies of large-scale structure are vital to our efforts to understand the origin and nature of the cosmos itself.

The idea that the filaments are the intersection of a survey slice with much larger structures (the bubble surfaces) was confirmed when the next three slices of the survey, lying above and below the first, were completed. The region of Figure 25.22 indicated by the red outline was found to continue through both the other slices. This extended sheet of galaxies, which has come to be known as the *Great Wall*, measures at least 70 Mpc (out of the plane of the page) by 200 Mpc (across the page). It is one of the largest known structures in the universe.

Figure 25.23 shows a more recent redshift survey, considerably larger than the one presented in Figure 25.22. This survey includes nearly 24,000 galaxies within about 750 Mpc of the Milky Way. Numerous voids and "Great Wall-like" filaments can be seen (some are marked), but



◀ **FIGURE 25.22 Galaxy Survey** A "slice" of the universe, covering 1732 galaxies out to a distance of about 200 Mpc, shows that galaxies and clusters are not randomly distributed on large scales. Instead, they appear to have a filamentary structure, surrounding vast, nearly empty voids. (*Harvard-Smithsonian Center for Astrophysics*)



◀ **FIGURE 25.23 The Universe on Larger Scales** This large-scale galaxy survey, carried out at the Las Campanas Observatory in Chile, consists of 23,697 galaxies within about 1000 Mpc, in two  $80^\circ$  (wide)  $\times$   $4.5^\circ$  (thick) wedges of the sky. Many voids and “walls” on scales of up to 100–200 Mpc are evident, but nothing much larger. For scale, the extent of the survey shown in Figure 25.22 is marked as a thin blue arc in the northern sky.

apart from the general falloff in numbers of galaxies at large distances—basically because the more distant galaxies are harder to see due to the inverse-square law—there is no obvious evidence for any structures on scales *larger* than about 200 Mpc. Careful statistical analysis confirms this impression. Apparently, voids and walls represent the largest structures in the universe. We will return to the far-reaching implications of this fact in Chapter 26.

## Quasar Absorption Lines


How can we probe the structure of the universe on very large scales? As we have seen, much of the matter is dark, and even the “luminous” component is so faint that it is hard to detect at large distances. One way to study large-scale structure is to take advantage of the great distances, pointlike appearance, and large luminosities of quasars. Since quasars are so far away, light traveling from a quasar to Earth has a pretty good chance of passing through or near “something interesting” en route. By analyzing quasar images and spectra, it is possible to piece together a partial picture of the intervening space.

The quasar approach is reminiscent of the use of bright stars to probe the interstellar medium near the

Sun, and it suffers from the same basic drawback: We can study only regions of the sky where quasars happen to be found. ∞ (Sec. 18.1) However, this problem will diminish in time as ongoing and planned large-scale surveys scan the sky for fainter and fainter objects. Foremost among these surveys is the Sloan Digital Sky Survey (*Discovery 25-1*), which has constructed a map of much of the northern sky, including several million galaxies and more than 100,000 quasars.

In addition to exhibiting their own strongly redshifted spectra, many quasars show additional absorption features that are redshifted by substantially *less* than the lines from the quasar itself. For example, the quasar PHL 938 has an emission-line redshift of 1.954, placing the quasar at a distance of some 5200 Mpc, but it also shows absorption lines having redshifts of just 0.613. These lines with lesser redshifts are interpreted as arising from intervening gas that is much closer to us (only about 2300 Mpc away) than the quasar itself. Most probably, this gas is part of an otherwise invisible galaxy lying along the line of sight. Quasar spectra, then, afford astronomers a means of probing previously undetected parts of the universe.

The absorption lines of atomic hydrogen are of particular interest, since hydrogen makes up so much of all matter in the cosmos. Specifically, hydrogen’s ultraviolet (122-nm) “Lyman-alpha” line, associated with transitions between the ground and first excited states, is often used in this context. ∞ (Sec. 4.3) As illustrated in Figure 25.24, when astronomers observe the spectrum of a high-redshift quasar, they typically see a “forest” of absorption lines, starting at the (redshifted) wavelength of the quasar’s own Lyman-alpha emission line and extending to shorter wavelengths. These lines are interpreted as Lyman-alpha absorption features produced by gas clouds in foreground structures—galaxies, clusters, and so on—giving astronomers crucial information about the distribution of matter along the line of sight.



**Interactive FIGURE 25.24**  
**Absorption Line “Forest”** The huge number of absorption lines in the spectrum of quasar QSO 1422 + 2309 are the ultraviolet Lyman-alpha lines from hundreds of clouds of foreground hydrogen gas, each redshifted by a slightly different amount (but less than the quasar itself). The peak at left marks the Lyman-alpha emission line from the quasar, emitted at 122 nm but redshifted here to 564 nm, in the visible range.

The quasar light thus explores an otherwise invisible component of cosmic gas. In principle, every intervening cloud of atomic hydrogen leaves its own characteristic imprint on the quasar’s spectrum in a form that lets us explore the distribution of matter in the universe. By comparing these *Lyman-alpha forests* with the results of simulations, astronomers hope to refine many key elements of the theories of galaxy formation and the evolution of large-scale structure.

## Quasar “Mirages”


In 1979, astronomers were surprised to discover what appeared to be a binary quasar—two quasars with exactly the same redshift and similar spectra, separated by only a few arc seconds on the sky. Remarkable as the discovery of such a binary would have been, the truth about this pair of quasars turned out to be even more amazing: Closer study of the quasars’ radio emission revealed that they were *not* two distinct objects; instead, they were two separate images of the *same* quasar! Optical views of such a *twin quasar* are shown in Figure 25.25.

What could produce such a “doubling” of a quasar image? The answer is gravitational lensing—the deflection and focusing of light from a background object by the gravity of some foreground body (Figure 25.26). In Chapter 23, we saw how lensing by compact objects in the halo of the Milky Way Galaxy may amplify the light from a distant star, allowing astronomers to detect otherwise invisible stellar dark matter. (Sec. 23.6) In the case of quasars, the idea is the same, except that the foreground lensing object is an entire galaxy or galaxy cluster, and the deflection of the light is so great (a few arc seconds) that several separate images of the quasar may be formed, as shown in Figure 25.27.\* About two dozen such gravitational lenses are


\*In fact, much of the theory of gravitational lensing was worked out after the first lensed quasar observations and subsequently applied to dark-matter searches in our Galaxy.

known. As telescopes probe the universe with greater and greater sensitivity, astronomers are beginning to realize that gravitational lenses are relatively common features of the cosmos.

The existence of these multiple images provides astronomers with a number of useful observational tools. First, lensing by a foreground galaxy tends to amplify the light of the quasar, as just mentioned, making it easier to observe. At the same time, **microlensing** by individual stars within the galaxy may cause large fluctuations in the quasar’s brightness, allowing astronomers to study both the quasar and the galaxy’s stellar content. (Sec. 23.6) The amount of brightening due to microlensing depends on the size of



**▲ FIGURE 25.25 Twin Quasar** This “double” quasar (designated AC114 and located about 2 billion parsecs away) is not two separate objects at all. Instead, the two large “blobs” (at upper left and lower right) are images of the same object, created by a gravitational lens. The lensing galaxy itself is probably not visible in this image—the two objects near the center of the frame are thought to be unrelated galaxies in a foreground cluster. (NASA)





**Interactive FIGURE 25.26 Gravitational Lens** When light from a distant object passes close to a galaxy or a cluster of galaxies along the line of sight, the image of the background object (here, the quasar) can sometimes be split into two or more separate images (A and B). The foreground object is a gravitational lens.

the emitting region, and this in turn depends on the wavelength of the radiation observed—for example, the X-rays are emitted from a smaller region closer to the central black hole than is the quasar’s visible light (see Figure 24.34). By carefully comparing the amounts of brightening at different wavelengths, astronomers can probe the structure of the quasar’s accretion disk on scales inaccessible by any other means.


Second, because the light rays forming the images usually follow paths of different lengths, there is often a *time delay*,

ranging from days to years, between them. This delay provides advance notice of interesting events, such as sudden changes in the quasar’s brightness. Thus, if one image flares up, in time the other(s) will, too, giving astronomers a second chance to study the event. The time delay also allows astronomers to determine the *distance* to the lensing galaxy. This method provides

**▼ FIGURE 25.27 Einstein Cross** (a) The “Einstein Cross,” a multiply imaged quasar that spans only a couple of arc seconds, shows four separate images of the same quasar produced by the galaxy at the center. (b) A simplified artist’s conception of what might be occurring here, with Earth at right and the distant quasar at left. (NASA; D. Berry)



(a) (b)



**▲ FIGURE 25.28 Galaxy Cluster Lensing** (a) This spectacular example of gravitational lensing shows more than a hundred faint arcs from very distant galaxies. The wispy pattern spread across the foreground galaxy cluster (A 2218, about a billion parsecs distant) resembles a spider’s web, but it is really an illusion caused by A 2218’s gravitational field, which deflects the light from background galaxies and distorts their appearance. By measuring the extent of the distortion, astronomers can estimate the mass of the intervening cluster. (b) Another galaxy cluster, known only by its catalog name 0024 + 1654 and residing some 1.5 billion pc away, shows reddish-yellow blobs that are mostly normal elliptical galaxies and bluish looplike features that are images of a single background galaxy. (NASA)

an alternative means of measuring Hubble’s constant that is independent of any of the techniques discussed previously. The average value of  $H_0$  reported by workers using this approach is 65 km/s/Mpc, a little less than the value we have assumed throughout the text.

Finally, by studying the lensing of background quasars and galaxies by foreground galaxy clusters, astronomers can obtain a better understanding of the distribution of dark matter in those clusters, an issue that has great bearing on the large-scale structure of the cosmos.

## Mapping Dark Matter

Astronomers have extended the ideas first learned from studies of quasars to the lensing of any distant object in order to better probe the universe. Distant, faint irregular galaxies—the raw material of the universe if current theories are correct (see Section 25.3)—are of particular interest here; because they are far more common than quasars, they provide much better coverage of the sky. By studying the lensing of background quasars and galaxies by foreground galaxy clusters, astronomers can obtain a better understanding of the distribution of dark matter on large scales.


Figures 25.28(a) and (b) show how the images of faint background galaxies are bent into arcs by the gravity of a foreground galaxy cluster. The degree of bending allows the total mass of the cluster (*including* the mass of the dark matter) to be measured. The (mostly blue) loop- and arc-shaped features visible in Figure 25.28(b) are multiple images of a

single distant (unseen) spiral or ring-shaped galaxy, lensed by the foreground galaxy cluster (the yellow-red blobs in the image).

It is even possible to reconstruct the foreground dark-matter distribution by carefully analyzing the distortions of the background objects, thereby providing a means of tracing out the distribution of mass on scales far larger than have previously been possible. Figure 25.29 is a reconstructed map showing the presence of dark matter many megaparsecs from the center of a small galaxy cluster (the brightest blob near the center of the map). Notice the elongated structure of the dark-matter distribution, reminiscent of the Virgo Supercluster and filamentary structure seen in large-scale galaxy surveys.

In 2006, astronomers used these techniques to obtain what may be the first direct observational evidence for dark matter. Figure 25.30 shows combined optical and X-ray images of a distant galaxy cluster called 1E 0657-56. The fuzzy red region shows the location of the hot X-ray-emitting gas in the system, the dominant *luminous* component of the mass. The blue regions indicate where most of the mass actually lies, as determined from lensing studies of background galaxies. Note that the bulk of the mass is *not* found in the form of hot gas, implying that the dark matter is distributed differently from the “normal” matter in the cluster.

The explanation for this odd state of affairs is that we are witnessing a collision between two clusters. Each initially contained hot gas and dark matter distributed



**▲ FIGURE 25.29 Dark-Matter Map** Measured distortions in the images of background objects can be displayed as maps of dark matter in the universe. Analysis of an optical view (a) of a region of the sky containing a small galaxy cluster (the clump of yellowish galaxies near the center of the frame indicated by the arrow) reveals the distribution of dark matter (b) in and near the visible cluster and on the same scale as (a). (*J. A. Tyson, Alcatel-Lucent; NOAO*)

throughout the cluster, but when the two collided, the pressure of each gas cloud effectively stopped the other, leaving the gas behind in the middle as the galaxies and dark matter moved on. This separation between the gas and the dark matter directly contradicts some alternative theories of gravity that have been invoked to explain away the “dark matter problem” in galaxies and clusters and may prove to



be a crucial piece of evidence in our understanding of large-scale structure in the universe.

**CONCEPT Check**

- ✓ How do observations of distant quasars tell us about the structure of the universe closer to home?

*The arrows indicate the directions in which the two clusters are now moving, subsequent to what might have been the most energetic collision in the universe since the Big Bang.*

**▶ Interactive FIGURE 25.30 Cluster Collision** Clusters of galaxies must also occasionally collide, as is apparently the case here for this combined cluster with the innocuous catalog name 1E 0657-56 and the nickname “bullet cluster.” This is a composite image of a region about 1 billion parsecs away, showing optical light from the galaxies themselves in white and X-ray-emitting gas from the hot intracluster gas in red. By contrast, the blue color represents the inferred dark matter within the two large clusters that is distinctly displaced from their normal matter. (*NOAO/NASA*)



ANIMATION/VIDEO Bullet Cluster Collision  
ANIMATION/VIDEO Dark Matter Collision




**The Big Question** What is this dark matter that everyone frets over? Is it actual matter that exerts a gravitational pull but is impossible to detect by electromagnetic means, or does it imply that something is badly wrong with our theoretical understanding of the way gravity works on very large scales? Dark matter—and now dark energy, too (see Chapter 26)—represent the foremost scientific conundrums in astronomy today, and whoever solves them will become immediately famous. Lots of scientists are trying very hard, but no one has yet succeeded.


# Chapter Review

## SUMMARY


**1** The masses of nearby spiral galaxies can be determined by studying their rotation curves. Astronomers also use studies of binary galaxies and galaxy clusters to obtain statistical estimates of the masses of the galaxies involved.



**2** Measurements of galaxy and cluster masses reveal the presence of large amounts of dark matter. The fraction of dark matter grows as the scale under consideration increases. More than 90 percent of the mass in the universe is dark. Large amounts of hot X-ray-emitting gas have been detected among the galaxies in many clusters, but not enough to account for the dark matter inferred from dynamical studies.




**3** Researchers know of no simple evolutionary sequence that links spiral, elliptical, and irregular galaxies. Most astronomers think that large galaxies formed by the merger of smaller ones and that collisions and mergers among galaxies, as well as accretion of intergalactic gas, play very important roles in galactic evolution. A **starburst galaxy** (p. 644) may result when a galaxy has a close encounter or a collision with a neighbor. The strong tidal distortions caused by the encounter compress galactic gas, resulting in a widespread burst of star formation. Mergers between spirals most likely result in elliptical galaxies.




**4** Quasars, active galaxies, and normal galaxies may represent an evolutionary sequence. When galaxies began to form and merge, conditions may have been suitable for the formation of large black holes at their centers, and a highly luminous quasar could have been the result. The brightest quasars consume so much fuel that their energy-emitting lifetimes must be quite short. As the fuel supply diminished, the quasar dimmed, and the galaxy in which it was embedded became intermittently visible as an active galaxy. At


even later times, the nucleus became virtually inactive, and a normal galaxy was all that remained. Many normal galaxies have been found to contain massive central black holes, suggesting that most galaxies have the capacity for activity should they interact with a neighbor. **Quasar feedback** (p. 654) may provide a partial explanation of why the masses of black holes are correlated with the masses of their parent galaxies.



**5** Galaxy clusters themselves tend to clump together into **superclusters** (p. 656). The Virgo Cluster, the Local Group, and several other nearby clusters form the Local Supercluster. On even larger scales, galaxies and galaxy clusters are arranged on the surfaces of enormous “bubbles” of matter surrounding vast low-density regions called **voids** (p. 657). The origin of this structure is thought to be closely related to conditions in the very earliest epochs of the universe.



**6** Quasar spectra can be used as probes of the universe along the observer’s line of sight. Some quasars have been observed to have double or multiple images, caused by gravitational lensing, in which the gravitational field of a foreground galaxy or galaxy cluster bends and focuses the light from the more distant quasar. Analysis of the images of distant galaxies, distorted by the gravitational effect of a foreground cluster, provides a means of determining the masses of galaxy clusters—including the dark matter within them—far beyond the information that the optical images of the galaxies themselves afford.





**MasteringAstronomy**® For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- LO1** Describe two techniques for measuring the mass of a galaxy.
- LO2** Why do astronomers think that galaxy clusters contain a lot more mass than we can see?
- POS** What evidence do we have that galaxies collide with one another?
- Describe the role of collisions in the formation and evolution of galaxies. How can mergers transform one type of galaxy into another?
- LO3** Do you think that collisions between galaxies constitute “evolution” in the same sense as the evolution of stars?
- Do we have any evidence that our own Galaxy has collided with other galaxies in the past?
- What are starburst galaxies, and what do they have to do with galaxy evolution?
- Why do astronomers think that quasars represent an early, relatively short-lived stage of galaxy evolution?
- What happened to the energy source at the center of a quasar?
- LO4 POS** Why does the theory of galaxy evolution suggest that there should be supermassive black holes at the centers of many normal galaxies?
- What evidence do astronomers have for supermassive black holes in galactic nuclei?
- What is a redshift survey? What are voids?
- LO5** Describe the distribution of galactic matter on very large (more than 100 Mpc) scales.
- LO6 POS** How can observations of distant quasars be used to probe the space between them and Earth?
- POS** How do astronomers “see” dark matter?

## Conceptual Self-Test: Multiple Choice

- The more massive a galaxy is, **(a)** the more distant it is; **(b)** the faster star formation in it occurs; **(c)** the larger the proportion of old stars it contains; **(d)** the faster it rotates.
- A galaxy containing substantial amounts of dark matter will **(a)** appear darker; **(b)** spin faster; **(c)** repel other galaxies; **(d)** have more tightly wound arms.
- According to X-ray observations, the space between galaxies in a galactic cluster is **(a)** completely devoid of matter; **(b)** very cold; **(c)** very hot; **(d)** filled with faint stars.
- Relative to luminous stellar matter, the fraction of dark matter in clusters is **(a)** greater than the fraction in galaxies; **(b)** less than the fraction in galaxies; **(c)** the same as the fraction in galaxies; **(d)** unknown.
- VIS** The Hubble Deep Field (Figure 25.10) shows a patch of sky that has the same angular size as **(a)** the thickness of a piece of string; **(b)** a dime; **(c)** a clenched fist; **(d)** a basketball held at arm's length.
- Galaxies evolve by **(a)** fragmenting into smaller galaxies; **(b)** merging to form larger galaxies; **(c)** ejecting their gas and dust into intergalactic space; **(d)** using up all their gas and eventually becoming ellipticals.
- According to current theories of galactic evolution, quasars occur **(a)** early in the evolutionary sequence; **(b)** near the Milky Way; **(c)** when elliptical galaxies merge; **(d)** late in the evolutionary sequence.
- Many nearby galaxies **(a)** will become black holes; **(b)** contain quasars; **(c)** have radio lobes; **(d)** were more active in the past.
- VIS** If light from a distant quasar did not pass through any intervening atomic hydrogen clouds, then Figure 25.24 (“Absorption Line”) would have to be redrawn to show **(a)** more absorption features; **(b)** few absorption features; **(c)** a single large absorption feature; **(d)** more features at short wavelengths, but fewer at long wavelengths.
- VIS** If Figure 25.26 (“Gravitational Lens”) showed a more massive lensing galaxy, the quasar images would be **(a)** farther apart; **(b)** closer together; **(c)** fainter; **(d)** redder.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

- The Andromeda Galaxy is approaching our Galaxy with a radial velocity of 120 km/s. Given the galaxies' present separation of 800 kpc, and neglecting *both* the transverse component of the velocity *and* the effect of gravity in accelerating the motion, estimate when the two galaxies will collide.
- Based on the data in Figure 25.1, estimate the mass of the galaxy NGC 4984 inside 20 kpc.
- Use Kepler's third law (Section 23.6) to estimate the mass required to keep a galaxy moving at 750 km/s in a circular orbit of radius 2 Mpc around the center of a galaxy cluster. Given the approximations involved in calculating this mass, do you think it is a good estimate of the cluster's true mass?
- Calculate the average speed of hydrogen nuclei (protons) in a gas of temperature 20 million K. Compare your answer with the speed of a galaxy moving in a circular orbit of radius 1 Mpc around a galaxy cluster of mass  $10^{14}$  solar masses.
- A small satellite galaxy moves in a circular orbit around a much more massive parent and happens to be moving exactly parallel to the line of sight as seen from Earth. The recession velocities of the satellite and the parent galaxy are measured to be 6450 km/s and 6500 km/s, respectively, and the two galaxies are separated by an angle of  $0.1^\circ$  in the sky. Assuming that  $H_0 = 70$  km/s/Mpc, calculate the mass of the parent galaxy.
- In a galaxy collision, two similar-sized galaxies pass through each other with a combined relative velocity of 1500 km/s. If each galaxy is 100 kpc across, how long does the event last?
- Assuming an energy-generation efficiency (i.e., the ratio of energy released to total mass-energy available) of 10 percent, calculate how much mass a  $10^{41}$ -W quasar would consume if it shone for 10 billion years.
- The spectrum of a quasar with a redshift of 0.20 contains two sets of absorption lines redshifted by 0.15 and 0.155, respectively. If  $H_0 = 70$  km/s/Mpc, estimate the distance between the intervening galaxies responsible for the two sets of lines.

## Activities

### Collaborative

- Figure 25.10 is called the "Hubble Deep Field." It contains too many galaxies for one person to easily count. Each group member should count the galaxies in a random area  $2\text{ cm} \times 2\text{ cm}$  and then determine a group average. Since the entire image is approximately  $500\text{ cm}^2$ , multiply your group's average number of galaxies in a  $2\text{ cm} \times 2\text{ cm}$  area by 125 to estimate the number of galaxies in the image. How does your value compare to that of another group?

### Individual

- Look for a copy of the *Atlas of Peculiar Galaxies* by Halton Arp. It is available in book form, but it will be more convenient to find a version online. Search for examples of interacting galaxies of various types: (1) tidal interactions, (2) starburst galaxies, (3) collisions between two spirals, and (4) collisions between a spiral and an elliptical. For (1) look for galactic material pulled away from a galaxy by a neighboring galaxy. Is the latter galaxy also tidally distorted? In (2) the surest signs of starburst activity are bright knots of star formation. In what type(s) of galaxies do you find starburst activity? For (3) and (4), how do collisions differ depending on the types of galaxies involved? What typically happens to a spiral galaxy after a near miss or collision? Do ellipticals suffer the same fate?



# Cosmology

## THE BIG BANG AND THE FATE OF THE UNIVERSE

Our field of view now extends for billions of parsecs into space and billions of years back in time. We have asked and answered many questions about the structure and evolution of planets, stars, and galaxies. At last we are in a position to address the central issues of the biggest puzzle of all: How big is the universe? How long has it been around, and how long will it last? What was its origin, and what will be its fate? Is the universe a one-time event, or does it recur and renew itself, in a grand cycle of birth, death, and rebirth? How and when did matter, atoms, and our Galaxy form? These are basic questions, but they are hard questions.

In this and the next chapter, we will see how modern scientific cosmology addresses these important issues and what it has to tell us about the universe we inhabit. After more than 10,000 years of civilization, science may be ready to provide some insight regarding the origin of all things.

**The Big Picture** The universe began in a fiery expansion some 14 billion years ago, and out of this maelstrom emerged all the energy that would later form galaxies, stars, and planets. That expansion continues today, yet to what end remains unknown. This is the science of cosmology—the study of the origin, structure, evolution, and fate of the cosmos on the largest scales.

**LEFT:** This image—called the Ultra Deep Field—was taken with the Advanced Camera for Surveys aboard the *Hubble* telescope. It is one of the finest photographs of deep space ever made. More than a thousand galaxies are crowded into this one image, displaying many different types, shapes, and colors. In all, astronomers estimate that the observable universe contains about 100 billion such galaxies. (NASA/ESA)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 State the cosmological principle, and explain its significance and observational underpinnings.
- 2 Explain what observations of the dark night sky tell us about the age of the universe.
- 3 Describe the Big Bang theory of the expanding universe.
- 4 List and discuss the possible outcomes of the present cosmic expansion.
- 5 Describe the relationship between the density of the universe and the overall geometry of space.
- 6 Say why astronomers think the expansion of the universe is accelerating, and discuss the cause.
- 7 Explain what dark energy implies for the composition and age of the universe.
- 8 Describe the cosmic microwave background and explain its importance to the science of cosmology.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.

## 26.1 The Universe on the Largest Scales

The universe shows structure on every scale we have examined so far. Subatomic particles form nuclei and atoms. Atoms form planets and stars. Stars form star clusters and galaxies. Galaxies form galaxy clusters, superclusters, and even larger structures—voids, filaments, and sheets that stretch across the sky. ∞ (Sec. 25.5) From the protons in a nucleus to the galaxies in the Great Wall, we can trace a hierarchy of “clustering” of matter from the very smallest to the very largest scales. It is natural to ask, “Does the clustering ever end? Is there some scale on which the universe can be regarded as more or less smooth and featureless?” Perhaps surprisingly, given the trend we have just described, most astronomers think the answer is yes. This turns out to be a crucial assumption in the science of **cosmology**—the study of the structure and evolution of the entire universe.


### The End of Structure

We saw in Chapter 25 how astronomers use redshift surveys to construct three-dimensional maps of the universe on truly “cosmic” scales. ∞ (Sec. 25.5) Figure 26.1 is a map similar to those shown previously, but based on data from the most extensive redshift survey to date—the Sloan Digital Sky Survey. ∞ (Discovery 25-1) It extends out to a

distance of almost 1000 Mpc, comparable to Figure 25.23, but because it includes much fainter galaxies, the Sloan map contains many more galaxies than the earlier figure, making structure easier to discern, particularly at large distances. The extended “filament” of galaxies near the center of the wedge, some 300 Mpc from Earth, is called the *Sloan Great Wall*. Measuring some 250 Mpc long by 50 Mpc thick, it is currently the largest known structure in the universe.

Plots such as this contain huge amounts of information about the structure and evolution of the universe. Yet, although they cover wide areas of the sky and enormous volumes of space, the studies on which they are based are still relatively “local,” in the sense that they span only about 10 percent of the distance to the farthest quasars (which lie over 9000 Mpc from Earth). ∞ (Sec. 24.4) The main obstacle to extending these wide-angle surveys to much greater distances is the sheer observational effort involved in measuring the redshifts of all the galaxies within larger and larger volumes of space.

An alternative approach is to narrow the field of view to only a few small patches of the sky, but to study extremely faint (and hence very distant) galaxies within those patches. The volume surveyed then becomes a long, thin “pencil beam” extending deep into space rather than a wide swath through the local universe. As illustrated in Figure 26.2, along the line of sight clusters and walls show up as “spikes” in the distribution—groups of galaxies with similar redshifts, separated by broad empty regions of space (the voids).



▲ **FIGURE 26.1 Galaxy Survey** This map of the universe is drawn using data from the Sloan Digital Sky Survey. ∞ (Discovery 25-1) It shows the locations of 66,976 galaxies lying within  $12^\circ$  of the celestial equator and extending to a distance of almost 1000 Mpc. The largest known structure in the universe, the Sloan Great Wall, is marked, stretching nearly 300 Mpc across the center of the frame. There is no evidence for any structure on larger scales. (SDSS)

The data from both kinds of survey seem to agree that the largest known structures in the local universe are “only” 200–300 Mpc across. *No larger voids, superclusters, or walls of galaxies are seen.* Rich superclusters measure tens of megaparsecs across, whereas the largest voids are perhaps 100 Mpc in diameter. Most walls and filaments are less than 100 Mpc in length, and even the largest structures—the Great Walls mentioned previously—can be explained statistically as chance superpositions of smaller structures. Studies of Lyman-alpha forests in quasar spectra lead to generally similar conclusions. ∞ (Sec. 25.5) In short, there is *no* evidence for structure in the universe on scales greater than about 300 Mpc.

We will turn to the *origin* of large-scale cosmic structure in Chapter 27. In the current chapter, however, we focus on the *absence* of structure on the very largest scales to frame our discussion of the future of the universe.


## The Cosmological Principle

The results of the large-scale studies just mentioned strongly suggest that the universe is **homogeneous** (the same everywhere) on scales greater than a few hundred megaparsecs. In other words, if we took a huge cube—300 Mpc on a side, say—and placed it anywhere in the universe, its overall contents would look much the same no matter where it was centered. Some of the galaxies it contained would be clustered and clumped into fairly large structures and some would not; we would see numerous walls and voids, but the total numbers of these objects would not vary much as the cube was moved from place to place. In this sense, the universe appears *smooth* on the largest scales.


The universe also appears to be **isotropic** (the same in all directions) on these large scales. Excluding directions that are obscured by our Galaxy, we count roughly the same number of galaxies per square degree in any patch of the sky we choose to observe, provided that we look deep (far) enough that local inhomogeneities don’t distort our sample. In other words, any deep pencil-beam survey of the sky should count about the same number of galaxies, regardless of which patch of the sky is chosen.

Cosmologists generally assume that the universe is *homogeneous* and *isotropic* on sufficiently large scales. These twin assumptions are known as the **cosmological principle**. No one knows whether these assumptions are precisely correct, but we can at least say that they are consistent with current observations, and they provide helpful guidance to our studies of the cosmos. Note that the cosmological principle also includes the important assumption made throughout this book (and indeed throughout astronomy) that the laws of physics are the same everywhere. In this chapter, we simply assume that it holds.

The cosmological principle has far-reaching implications. For example, it implies that there can be no edge to the universe, because that would violate the assumption of



(a)



(b)

▲ **FIGURE 26.2 Pencil-Beam Survey** The results of a deep “pencil-beam” survey of two small portions of the sky in opposite directions from Earth, perpendicular to the Galactic plane (a), are plotted in (b). The graph shows the number of galaxies found at different distances from us, out to about 2000 Mpc. Wherever we look on the sky, the distinctive “picket fence” pattern highlights voids and sheets of galaxies on scales of 100–200 Mpc, but gives no indication of any larger structures.

homogeneity. Furthermore, it implies that there is *no center*, because that would mean that the universe would not be the same in all directions from any noncentral point, a violation of the assumption of isotropy. This is the familiar Copernican principle expanded to truly cosmic proportions—not only that we are not central to the universe, but that *no one* can be central, because *the universe has no center!* ∞ (Sec. 2.3)

### CONCEPT Check

✓ In what sense, and on what scale, is the universe homogeneous and isotropic?

## 26.2 The Expanding Universe

Every time you go outside at night and notice that the sky is dark, you are making a profound cosmological observation. Here's why.

### Olbers's Paradox

Let's assume that, in addition to being homogeneous and isotropic, the universe is infinite in spatial extent and unchanging in time—precisely the view of the universe that prevailed until the early part of the 20th century. On average, then, the universe is uniformly populated with galaxies filled with stars. In that case, when you look up at the night sky, your line of sight must *eventually* encounter a star, as illustrated in Figure 26.3. The star may lie at an enormous distance in some remote galaxy, but the laws of probability dictate that, in an infinite universe, sooner or later any line drawn outward from Earth will run into a bright stellar surface.

Of course, faraway stars appear fainter than those nearby because of the inverse-square law.  $\infty$  (Sec. 17.2) However, they are also much more numerous because the number of stars we see at any given distance in fact increases as the *square* of the distance. (Just consider the area of a sphere of increasing radius.) Thus, the diminishing brightnesses of distant stars are exactly balanced by their increasing numbers, and stars at all distances contribute equally to the total amount of light received on Earth. This fact has a dramatic implication: No matter where you look, the sky should be as bright as the surface of a star; in other words, the entire night sky should be as brilliant as the surface of the Sun! The obvious difference between this prediction and the actual appearance of the night sky is known as **Olbers's paradox**, after the 19th-century German astronomer Heinrich Olbers, who popularized the idea.

So why is it dark at night? Given that the universe appears to be homogeneous and isotropic, one (or both) of the other two assumptions must be false: Either the universe is finite in extent, or it evolves over time. In fact, the answer involves aspects of each and is intimately tied to the behavior of the universe on the largest scales.


### The Birth of the Universe

In Chapter 24, we saw that all the galaxies in the universe are rushing away from us in a manner described by Hubble's law,

$$\text{recession velocity} = H_0 \times \text{distance},$$

where we take Hubble's constant  $H_0$  to be 70 km/s/Mpc.  $\infty$  (Sec. 24.3) Up to now, we have used this relation as a convenient means of determining the distances to galaxies and quasars, but it is much more than that.

A good analogy to Olbers's paradox is a thick forest where every line of sight eventually intersects a tree.



▲ **FIGURE 26.3 Olbers's Paradox** If the universe were homogeneous, isotropic, infinite in extent, and unchanging, then any line of sight from Earth should eventually meet a star and the entire night sky should be bright. Since the night sky is obviously dark, this contradiction is known as Olbers's paradox.

Assuming for the moment that all velocities have remained constant in time, we can ask a simple question: How long has it taken for any given galaxy to reach its present distance from us? The answer follows from Hubble's law. The time taken is simply the distance traveled divided by the velocity, so

$$\begin{aligned} \text{time} &= \frac{\text{distance}}{\text{velocity}} \\ &= \frac{\text{distance}}{H_0 \times \text{distance}} \quad (\text{using Hubble's law} \\ &= \frac{1}{H_0}. \quad \quad \quad \text{for the velocity}) \end{aligned}$$

For  $H_0 = 70$  km/s/Mpc, this time is about 14 billion years. Notice that it is *independent* of the distance: Galaxies twice as far away are moving twice as fast, so the time they took to cross the intervening distance is the same.

Hubble's law therefore implies that, at some time in the past—14 billion years ago, according to the foregoing simple calculation—all the galaxies in the universe lay right on top of one another. In fact, astronomers think that *everything* in the universe—matter and radiation alike—was confined at that instant to a single point of enormously high temperature and density, often referred to as the **primeval fireball**. Then the universe began to expand at a furious rate, its density and temperature

falling rapidly as the volume increased. This stupendous, unimaginably violent event, involving literally everything in the cosmos, is known as the **Big Bang**. It marked the beginning of the universe.

Thus, by measuring Hubble's constant, we can estimate the age of the universe to be  $1/H_0 \approx 14$  billion years. The range of possible error in this age is considerable, both because Hubble's constant is not known precisely and because the assumption that galaxies moved at constant speed in the past is not a good one. We will refine our estimate in a moment, but regardless of the details, the critical fact here is that the age of the universe is *finite*.

The Big Bang provides the resolution of Olbers's paradox. Whether the universe is actually finite or infinite in extent is irrelevant, at least as far as the appearance of the night sky is concerned. The finite *age* of the universe implied by Hubble's law is the key. We see only a finite part of the cosmos—the region lying within roughly 14 billion light-years of us. What lies beyond is unknown—its light has not yet had time to reach us.

Note that, even though it appears to place us at the center of the expansion, Hubble's law does *not* violate the cosmological principle in any way. To see this, consider Figure 26.4, which shows how observers in five hypothetical galaxies perceive the motion of their neighbors. For simplicity, the galaxies are taken to be equally spaced, 100 Mpc apart, and they are separating in accordance with Hubble's law with  $H_0 = 70$  km/s/Mpc, as seen by an observer in the middle galaxy, number 3. The first pair of numbers beneath each galaxy represents its distance and recessional velocity as measured by that observer. For definiteness, let's take


galaxy number 3 to be the Milky Way and the observer to be an astronomer on Earth.

Now consider how the expansion looks from the point of view of the observer in galaxy 2. Galaxy 4, for example, is moving with velocity 7000 km/s to the right relative to galaxy 3, and galaxy 3 in turn is moving at 7000 km/s to the right as seen by observer 2. Therefore, galaxy 4 is moving at a velocity of 14,000 km/s to the right as seen by the observer in galaxy 2. But the distance between the two galaxies is 200 Mpc, so the Hubble constant measured by the observer on galaxy 2 is  $14,000 \text{ km/s}/200 \text{ Mpc} = 70 \text{ km/s/Mpc}$ , the same as the Hubble constant measured by the observer on galaxy 3. The distances and velocities that would be measured by observer 2 are noted in the second row. You can verify for yourself that the ratio of recession velocity to distance is the same for all galaxies.

Similarly, the measurements made by an observer on galaxy 1 are noted in the third row. Again, the ratio of velocity to distance is the same. The conclusion is clear: *Each* observer sees an overall expansion described by Hubble's law, and the constant of proportionality—Hubble's constant—is the same in all cases. Far from singling out any one observer as central, Hubble's law is in fact the *only* expansion law possible if the cosmological principle holds.

## Where Was the Big Bang?

Now we know *when* the Big Bang occurred. Is there any way of telling *where*? We think that the universe is the same everywhere, yet we have just seen that the observed recession of the galaxies described by Hubble's law suggests that all



**▲ FIGURE 26.4 Hubble Expansion** Hubble's law is the same, regardless of who makes the measurements. The top numbers are the distances and recessional velocities as seen by an observer on the middle of five galaxies, galaxy 3. The bottom two sets of numbers are from the points of view of observers on galaxies 2 and 1, respectively. In all cases, Hubble's law holds: The ratio of the observed recession velocity to the distance is the same.




the galaxies expanded from a point at some time in the past. Wasn't that point, then, different from the rest of the universe, violating the assumption of homogeneity expressed in the cosmological principle? The answer is a definite *no*!

To understand why there is no “center” to the expansion, we must make a great leap in our perception of the universe. If we were to imagine the Big Bang as simply an enormous explosion that spewed matter out into space, ultimately to form the galaxies we see, then the foregoing reasoning would be quite correct—there would be a center and an edge, and the cosmological principle would not apply. But the Big Bang was *not* an explosion in an otherwise featureless, empty universe. The only way that we can have Hubble's law hold *and* retain the cosmological principle is to realize that the Big Bang involved the entire universe—not just the matter and radiation within it, but the universe *itself*.

In other words, the galaxies are not flying apart into the rest of the universe. *The universe itself is expanding.* Like raisins in a loaf of raisin bread that move apart as the bread expands in an oven, the galaxies are just along for the ride.

Let's consider again some of our earlier statements in light of this new perspective. We now recognize that Hubble's law describes the expansion of the universe itself. Although galaxies have some small-scale, individual random motions, on average they are *not* moving with respect to the fabric of space—any such overall motion would pick out a “special” direction in space and violate the assumption of isotropy. On the contrary, the portion of the galaxies' motion that makes up the Hubble flow is really an expansion of space itself. The expanding universe remains homogeneous at all times. There is no “empty space” beyond the galaxies into which they rush. At the time of the Big Bang, the galaxies did not reside at a point located at some well-defined place within the universe. Rather, the *entire universe* was a point. That point was in no way different from the rest of the universe; that point *was* the universe. Therefore, there was no one point where the Big Bang “happened”—because the Big Bang involved the entire universe, it happened *everywhere* at once.

To illustrate these ideas, imagine an ordinary balloon with coins taped to its surface, as shown in Figure 26.5.



(Better yet, do the experiment yourself!) The coins represent galaxies, and the two-dimensional surface of the balloon represents the “fabric” of our three-dimensional universe. The cosmological principle applies to the balloon because every point on the balloon looks pretty much the same as every other. Imagine yourself as a resident of one of the three dark-colored coin “galaxies” in the leftmost frame, and note your position relative to your neighbors. As the balloon inflates (i.e., as the universe expands), the other galaxies recede from you; more distant galaxies recede more rapidly. Notice, incidentally, that the coins themselves do *not* expand along with the balloon, any more than people, planets, stars, or galaxies—all of which are held together by their own internal forces—expand along with the universe.

∞ (Sec. 24.3)

Regardless of which galaxy you chose to consider, you would see all the other galaxies receding from you. Nothing is special or peculiar about the fact that all the galaxies are receding from you. Such is the cosmological principle: No observer anywhere in the universe has a privileged position. There is no center to the expansion and no position that can be identified as the location from which the universal expansion began. Everyone sees an overall expansion described by Hubble's law, with the same value of Hubble's constant in all cases.

Now imagine letting the balloon deflate. This corresponds to running the universe backward from the present time to the Big Bang. *All* the galaxies (coins) would arrive at the same place at the same time—at the instant the balloon reached zero size. But there is no one point on the balloon that could be said to be *the* place where that occurred. The entire balloon expanded from a point, just as the Big Bang encompassed the entire universe and expanded from a point.

This analogy has its shortcomings. The main difficulty with it is that we see the balloon, which, in our illustration, we imagined as two dimensional, expanding into the third dimension of space. This might suggest that the three-dimensional universe is expanding “into” some fourth spatial dimension. It is not, so far as we know. At the very least, if higher spatial dimensions are involved, they are not relevant to our theory of the universe.

### Interactive FIGURE 26.5 Receding Galaxies

Coins taped to the surface of a spherical balloon recede from each other as the balloon inflates (left to right).

Similarly, galaxies recede from each other as the universe expands. As the coins separate, the distance between any two of them increases, and the rate of increase of this distance is proportional to the distance between them.

## The Cosmological Redshift

This view of the expanding universe requires us to reinterpret the cosmological redshift. ∞ (Sec. 24.3) Previously, we discussed the redshift of galaxies as though it were a Doppler shift—a consequence of their motion relative to us. However, we have just argued that the galaxies are *not* in fact moving with respect to the universe, in which case the Doppler interpretation is incorrect. The true explanation is that, as a photon moves through space, its wavelength is influenced by the expansion of the universe. In a sense, we can think of the photon as being attached to the expanding fabric of space, so its wavelength expands along with the universe, as illustrated in Figure 26.6. Although it is common practice in astronomy to refer to the cosmological redshift in terms of recessional velocity, bear in mind that, strictly speaking, that is not the right thing to do. The cosmological redshift is a consequence of the changing size of the universe—it is *not* related to velocity at all.


The redshift of a photon measures the amount by which the universe has expanded since that photon was emitted. For example, when we measure the light from a quasar and find that it has a redshift of 5, it means that the observed wavelength is 6 times (1 plus the redshift) greater than the wavelength at the time of emission, and this in turn means that the light was emitted at a time when the universe was just one-sixth its present size (and we are observing the quasar as it was at that time). ∞ (More Precisely 24-1) In general, the larger a photon's redshift, the smaller the universe was at the time the photon was emitted, so the longer ago that emission occurred. Because the universe expands with time and redshift is related to that expansion, cosmologists routinely use redshift as a convenient means of expressing time.

These concepts are difficult to grasp. The notion of the entire universe expanding from a hot, dense fireball—with *nothing*, not even space and time, outside—takes some getting used to. Nevertheless, this description of the universe lies at the heart of modern cosmology.

### PROCESS OF SCIENCE Check

✓ Why does Hubble's law imply a Big Bang?

Radiation also shifts with cosmic expansion.



**Interactive FIGURE 26.6**  
**Cosmological Redshift** As the universe expands, photons of radiation are stretched in wavelength, giving rise to the cosmological redshift. In this case, as the baseline in the diagram stretches, the radiation shifts from the short-wavelength blue region of the spectrum to the longer wavelength red region. ∞ (Sec. 3.1)

## 26.3 The Fate of the Cosmos


Will the universe expand forever? This fundamental question about the fate of the universe has been at the heart of cosmology since Hubble's law was first discovered. Until the late 1990s, the prevailing view among cosmologists was that the answer would most likely be found by determining the extent to which gravity would slow, and perhaps ultimately reverse, the current expansion. However, it now appears that the answer is more subtle—and perhaps a lot more profound in its implications—than was hitherto thought.

### Critical Density

Let's begin with another analogy. Assume for the moment that gravity is the only force affecting large-scale motion in the universe, and consider a rocket ship launched from the surface of a planet. Until relatively recently, this scenario was much more than just an analogy—this basic picture and its implications represented the conventional wisdom among cosmologists. However, as we will see in Section 26.5, new observations have forced fundamental changes in astronomers' view of the universe. Nevertheless, the simplified view we now present is a convenient starting point, as it allows us to define some basic ideas and terminology.

What are the likely outcomes of the rocket ship's motion? According to Newtonian mechanics, there are just two basic possibilities, depending on the launch speed of the ship relative to the escape speed of the planet. ∞ (Sec. 2.8) If the launch speed is high enough, it will exceed the planet's escape speed, and the ship will never return to the surface. The speed will diminish because of the planet's gravitational pull, but it will never reach zero. The spacecraft leaves the planet on an unbound trajectory, as illustrated in Figure 26.7(a). Alternatively, if the launch speed is lower than the escape speed, the ship will reach a maximum distance from the planet and then fall back to the surface. Its bound trajectory is shown in Figure 26.7(b).

Similar reasoning applies to the expansion of the universe. Imagine two galaxies at some known distance from each other, moving apart with their current relative velocity given by Hubble's law. The same two basic possibilities exist




▲ **FIGURE 26.7 Escape Speed** (a) A spacecraft (arrows) leaving a planet (blue ball) with a speed greater than the planet's escape speed follows an unbound trajectory and escapes. (b) If the launch speed is less than the escape speed, the ship eventually drops back to the planet. Its distance, as graphed, from the planet first rises and then falls.

for these galaxies as for our rocket ship: The distance between them can increase forever, or it can increase for a while and then start to decrease. What's more, the cosmological principle says that, whatever the outcome for galaxy A and galaxy B, it must be the same for *any* two galaxies—in other words, the same statement applies to the universe *as a whole*. Thus, as illustrated in Figure 26.8, the universe has only two options: It can continue to expand forever, or the present expansion will someday stop and turn around into a contraction. The two curves in the figure are drawn so that they pass through the same point at the present time. Both are possible descriptions of the universe, given its current size and expansion rate.

What determines which of the two possibilities will actually occur? In the case of a rocket ship of fixed launch speed (analogous to a universe with a given expansion rate), the *mass* of the planet (for given radius) determines whether or not escape will occur—a more massive planet has a higher escape speed. For the universe, the corresponding factor is the *density* of the cosmos. A high-density universe contains enough mass to stop the expansion and eventually cause a collapse. A low-density universe, conversely, will expand forever.

The dividing line between these outcomes—the density corresponding to a universe in which *gravity acting alone* would be just sufficient to halt the present expansion—is called the universe's **critical density**. For  $H_0 = 70 \text{ km/s/Mpc}$ , the critical density is about  $9 \times 10^{-27} \text{ kg/m}^3$ . That's an extraordinarily




▲ **FIGURE 26.8 Model Universes** Distance between two galaxies as a function of time in each of the two basic universes discussed in the text: a low-density universe that expands forever and a high-density cosmos that collapses. The point where the two curves touch represents the present time.

low density—just five hydrogen atoms per cubic meter, a volume the size of a small household closet. In more “cosmological” terms, it corresponds to about 0.1 Milky Way Galaxy (including the dark matter) per cubic megaparsec.

## Two Futures

The two possibilities just presented represent radically different futures for our universe. If the cosmos emerged from the Big Bang with sufficiently high density, then it contains enough matter to halt its own expansion, and the recession of the galaxies will eventually stop. At some time in the future, astronomers everywhere—on any planet within any galaxy—will announce that the radiation received from nearby galaxies is no longer redshifted. (The light from *distant* galaxies will still be redshifted, however, because we will see them as they were in the past, at a time when the universe was still expanding.) The bulk motion of the universe will be stilled—at least momentarily.

The expansion may stop, but the pull of gravity will not. The universe will begin to contract. Nearby galaxies will begin to show blueshifts, and both the density and the temperature of the universe will start to rise as matter collapses back onto itself. As illustrated in Figure 26.9(a), the universe will collapse to a point, requiring just as much time to fall back as it took to rise. First galaxies and then stars will collide with increasing frequency and violence as the available space diminishes and the entire universe shrinks toward a superdense, superhot singularity much like the one from which it originated. The cosmos will ultimately—billions of years from now—experience a “heat death,” in which all matter and life are destined to be incinerated—a “Big Crunch.” Cosmologists do not know what would happen to the universe if it ever reached this point. The laws of physics



◀ **FIGURE 26.9 Two Futures** If gravity is the only force influencing the cosmic expansion, then the mass density of the universe determines its fate. (a) A high-density universe has a beginning, an end, and a finite lifetime. The lower frames illustrate its evolution, from initial expansion, to maximum size, to collapse. (b) A low-density universe expands forever, with galaxies getting farther and farther apart as time passes.

as we presently understand them are simply inadequate to describe those extreme conditions.

A quite different fate awaits a low-density universe whose gravity is too weak to halt the present expansion. As illustrated in Figure 26.9(b), such a universe will expand forever, the galaxies continually receding, their radiation steadily weakening with increasing distance. In time, an observer on Earth will see no galaxies in the sky beyond the Local Group (which is not itself expanding). Even with the most powerful telescope, the rest of the observable universe will appear dark, the distant galaxies too faint to be seen. Eventually, the Milky Way and the Local Group, too, will peter out as their fuel supply is consumed. This universe will ultimately experience a “cold death”: All radiation, matter, and life are eventually destined to freeze.

How long might the “cold death” of the universe take? Astronomers estimate that our Galaxy probably contains enough gas to keep forming stars for several tens of billions of years, and the majority of stars (the low-mass red dwarfs) can shine for hundreds of billions of years or more. ∞ (Sec. 17.8) Thus, we can expect our Galaxy (and our neighbor Andromeda) to shine on—albeit feebly—for another trillion years or so.

We will see in a moment that the separation between never-ending expansion and cosmic collapse is not quite as straightforward as the foregoing simple reasoning would suggest. Several independent lines of evidence now indicate that gravity is *not* the only influence on the dynamics of the universe on large scales (Section 26.5). As a result, while the “futures” just described are still the only two possibilities for the long-term evolution of the universe, the distinction between them turns out to be more than just a matter of density alone. Nevertheless, the density of the universe—or, more precisely, the *ratio of the total density to the critical value*—is a vitally important quantity in cosmology.

### CONCEPT Check

- ✓ What are the two basic possibilities for the future expansion of the universe?

## 26.4 The Geometry of Space

Our discussion in the previous section used the familiar notions of Newtonian mechanics and gravity because speaking in Newtonian terms made the evolution of the universe easier to understand. But in reality, the proper description of the universe as a dynamic, evolving object is far beyond the capabilities of Newtonian mechanics, which up to now has been our indispensable tool for understanding the cosmos. ∞ (Sec. 2.8) Instead, the more powerful techniques of Einstein’s *general relativity*, with its built-in notions of warped space and dynamic spacetime, are needed. ∞ (Sec. 22.6)

### Relativity and the Universe

We encountered general relativity in Chapter 22 when we discussed the strange properties of black holes. ∞ (Sec. 22.5) We can loosely summarize its description of the universe by saying that the presence of matter or energy causes a warping, or curvature, of *spacetime* and that the curved trajectories of freely falling particles within warped spacetime are what Newton thought of as orbits under the influence of gravity. The amount of warping depends on the amount of matter present.

When applied to the orbits of planets, stars, even of galaxies, the predictions of general relativity are, for the most part, in accord with those of Newtonian mechanics. But on the scale of the entire universe, relativity has some implications that simply have no counterpart within Newton’s theory. Foremost among these non-Newtonian predictions are the facts that the space around us is *curved* and that the degree of curvature is determined by the total *density* of the cosmos. ∞ (Sec. 22.6)

Furthermore, general relativity is very clear on precisely what “density” means here. Both matter *and* energy must be taken into account, with energy ( $E$ ) properly “converted” into mass ( $m$ ) units by Einstein’s famous relation  $E = mc^2$ . ∞ (More Precisely 22-1) [That is, an energy of 1 joule is counted as its mass equivalent of  $1 \text{ J}/(3 \times 10^8 \text{ m/s})^2$ , or  $1.1 \times 10^{-17} \text{ kg}$ —not much, but it adds up!] The total density of the universe includes not just the atoms and molecules

that make up the familiar “normal” matter around us, but also the invisible dark matter that dominates the masses of galaxies and galaxy clusters, as well as *everything* that carries energy—photons, relativistic neutrinos, gravity waves, and anything else we can think of.


## Cosmic Curvature

In a homogeneous universe, the curvature (on sufficiently large scales) must be the same everywhere, so there are really only three distinct possibilities for the large-scale geometry of space. (For more information on the different types of geometry involved, see *More Precisely 26-1*.) General relativity tells us that the geometry of the universe depends only on the ratio of the density of the universe to the critical density (defined in the previous section). As just noted, for  $H_0 = 70 \text{ km/s/Mpc}$ , the critical density is  $9 \times 10^{-27} \text{ kg/m}^3$ . Cosmologists call the ratio of the universe’s actual density to the critical value the *cosmic density parameter* and denote it by the symbol  $\Omega_0$  (“omega nought”). In terms of this quantity, then, a universe with density equal to the critical value has  $\Omega_0 = 1$ , a “low-density” cosmos has  $\Omega_0$  less than 1, and a “high-density” universe has  $\Omega_0$  greater than 1.

In a high-density universe ( $\Omega_0$  greater than 1), space is curved so much that it bends back on itself and “closes off,” making this universe *finite* in size. Such a universe is known as a **closed universe**. It is difficult to visualize a three-dimensional volume uniformly arching back on itself in this way, but the two-dimensional version is well known: It is just the surface of a sphere, like that of the balloon we discussed earlier. Figure 26.5, then, is the two-dimensional likeness of a three-dimensional closed universe. Like the surface of a sphere, a closed universe has no boundary, yet is finite in extent.\* One remarkable property of a closed universe is illustrated in Figure 26.10: Just as a traveler on the surface of a sphere can keep moving forward in a straight line and eventually return to her starting point, a flashlight beam shone in some direction in space might eventually traverse the entire universe and return from the opposite direction!

The surface of a sphere curves, loosely speaking, “in the same direction,” no matter which way we move from a given point. A sphere is said to have *positive curvature*. However, if the average density of the universe is below the critical value, the surface curves like a saddle, in which case it has *negative curvature*. Most people have a good idea of what a saddle looks like—it curves “up” in one direction and “down” in another—but no one has ever seen a uniformly negatively

\*Notice that for the sphere analogy to work we must imagine ourselves as two-dimensional “flatlanders” who cannot visualize or experience in any way the third dimension perpendicular to the sphere’s surface. Flatlanders and their light rays are confined to the sphere’s surface, just as we are confined to the three-dimensional volume of our universe.



*In a curved universe, light might return from the opposite direction!*

▲ **FIGURE 26.10 Einstein’s Curve Ball** In a closed universe, a beam of light launched in one direction might return someday from the opposite direction after circling the universe, just as motion in a “straight line” on Earth’s surface will eventually encircle the globe.

curved surface, for the simple reason that it cannot be constructed in three-dimensional Euclidean space! It is just “too big” to fit. A low-density, saddle-curved universe is infinite in extent and is usually called an **open universe**.

The intermediate case, in which the density is precisely equal to the critical density (i.e.,  $\Omega_0 = 1$ ), is the easiest to visualize. This universe, called a **critical universe**, has no curvature. It is said to be “flat” and is infinite in extent. In this case, and *only* in this case, the geometry of space on large scales is precisely the familiar Euclidean geometry taught in high schools. Apart from its overall expansion, this is basically the universe that Newton knew.

Euclidean geometry—the geometry of flat space—is familiar to most of us because it is a good description of space in the vicinity of Earth. It is the geometry of everyday experience. Does this mean that the universe is flat, which would in turn mean that it has exactly the critical density? Not necessarily: Just as a flat street map is a good representation of a city, even though we know Earth is really a sphere, Euclidean geometry is a good description of space within the solar system, or even the Galaxy, because the curvature of the universe is negligible on scales smaller than about 1000 Mpc. Only on the very largest scales would the geometric effects we have just discussed become evident.

### CONCEPT Check

- ✓ How is the curvature of space related to the density of the universe?

## MORE PRECISELY 26-1

### Curved Space

*Euclidean geometry* is the geometry of flat space—the geometry taught in high school. Set forth by one of the most famous of the ancient Greek mathematicians, Euclid, who lived around 300 B.C., it is the geometry of everyday experience. Houses are usually built with flat floors. Writing tablets and blackboards are also flat. We work easily with flat, straight objects, because the straight line is the shortest distance between any two points.

When we construct houses or any other straight-walled buildings on the surface of Earth, the other basic axioms of Euclid's geometry also apply: Parallel lines never meet, even when extended to infinity; the angles of any triangle always sum to  $180^\circ$ ; the circumference of a circle equals  $\pi$  times the diameter of the circle. (See the accompanying figure.) If these axioms did not hold, walls and roof would never meet to form a house!

In reality, though, the geometry of Earth's surface is not really flat; it is curved. We live on the surface of a sphere, and on that *surface* Euclidean geometry breaks down. Instead, the rules for the surface of a sphere are those of *Riemannian geometry*, named after the 19th-century German mathematician Georg Friedrich Riemann. There are no parallel “straight” lines on a sphere. The analog of a straight line on a sphere's surface is a “great circle”—the arc formed when a plane passing through the center of the sphere intersects the surface. Any two such lines must eventually intersect. The sum of a triangle's angles, when drawn on the surface of a sphere, exceeds  $180^\circ$ —in the  $90^\circ$ – $90^\circ$ – $90^\circ$  triangle shown in the accompanying figure, the sum is actually  $270^\circ$ —and the circumference of a circle is less than  $\pi$  times the circle's diameter.


We see that the curved surface of a sphere, governed by the spherical geometry of Riemann, differs greatly from the flat-space geometry of Euclid. The two are approximately the same only if we confine ourselves to a small patch on the surface. If the patch is small enough compared with the sphere's radius, the surface looks “flat” nearby, and Euclidean geometry is approximately valid. This is why we can draw a usable map of our home, our city, and even our state, on a flat sheet of paper, but an accurate map of the entire Earth must be drawn on a globe.

When we work with larger parts of Earth, we must abandon Euclidean geometry. World navigators are fully aware of this. Aircraft do not fly along what might appear on most maps as a straight-line path from one point to another. Instead, they follow a great circle on Earth's surface. On the curved surface of a sphere, such a path is always the shortest distance between two points.

For example, as illustrated in the figure, a flight from Los Angeles to London does not proceed directly across the United States and the Atlantic Ocean, as you might expect from looking at a flat map. Instead, it goes far to the north, over Canada and Greenland, above the Arctic Circle, finally coming in over Scotland for a landing at London. This is the great-circle route—the shortest path between the two cities, as you can easily see if you inspect a globe.

The “positively curved” space of Riemann is not the only possible departure from flat space. Another is the “negatively curved” space first studied by Nikolai Ivanovich Lobachevsky, a 19th-century Russian mathematician. In this geometry, there are an *infinite* number of lines through any given point that are parallel to another line, the sum of the angles of a triangle is *less* than  $180^\circ$  (see the first figure), and the circumference of a circle is *greater* than  $\pi$  times its diameter. This type of space is described by the surface of a curved saddle, rather than a flat plane or a curved sphere. It is a hard geometry to visualize!

Most of the local realm of the *three-dimensional* universe (including the solar system, the neighboring stars, and even our Milky Way Galaxy) is correctly described by Euclidean geometry. If the currently favored cosmology described in the text turns out to be correct, then the whole universe is, too!



## 26.5 Will the Universe Expand Forever?

Is there any way for us to determine which of the futures we have described actually applies to our universe (that is, apart from simply waiting to find out)? Will the universe end as a dense fireball much like that from which it began? Or will it expand forever? And can we hope to measure the geometry of the vast cosmos we inhabit? Finding answers to these questions has been the dream of astronomers for decades. We are fortunate to live at a time when astronomers can subject these questions to intensive observational tests and come up with definite answers—even though they aren't what most cosmologists expected! Let's begin by looking at the density of the universe (or, equivalently, the cosmic density parameter  $\Omega_0$ ).

### The Density of the Universe

How might we determine the density of the universe? On the face of it, it would seem simple: Just measure the total mass of the galaxies residing within some large parcel of space, calculate the volume of that space, and then divide the mass by the volume to compute the average density. When astronomers do this, they usually find a little less than  $10^{-28}$  kg/m<sup>3</sup> in the form of luminous matter. Largely independent of whether the chosen region contains many scattered galaxies or only a few rich galaxy clusters, the resulting density is about the same, within a factor of two or three. Galaxy counts thus yield a value of  $\Omega_0$  of only a few percent. If that measure were correct, and galaxies were all that existed, then we would live in a low-density open universe destined to expand forever.

But there is a catch. We have noted (Chapters 23 and 25) that most of the matter in the universe is *dark*—it exists in the form of invisible material that has been detected only through its gravitational effect in galaxies and galaxy clusters. [∞ \(Secs. 23.6, 25.1\)](#) Currently, we do not know what the dark matter is, but we *do* know that it is there. Galaxies may contain as much as 10 times more dark matter than luminous material, and the figure for galaxy clusters is even higher—perhaps as much as 95 percent of the total mass in clusters is invisible. Even though we cannot see it, dark matter contributes to the density of the universe and plays its part in opposing the cosmic expansion. Including all the dark matter that is known to exist in galaxies and galaxy clusters increases the value of  $\Omega_0$  to about 0.25.

Unfortunately, although we can detect and quantify the effects of dark matter in galaxies and galaxy clusters, its distribution on larger scales is harder to measure and is not very well-known. Astronomers have developed techniques to study matter on supercluster and larger scales, using gravitational lensing of distant objects and the large-scale motions of galaxies and clusters to probe the gravitational fields of cosmic concentrations of invisible matter. [∞ \(Sec. 25.5\)](#) Yet the results of all these studies add little to

the overall density. As best we can tell, there doesn't seem to be much additional dark matter “tucked away” on very large scales. Most cosmologists agree that the overall density of matter (luminous plus dark) in the universe is between 25 and 30 percent of the critical value—not enough to halt the universe's current expansion.


### Cosmic Acceleration

Determining the mass density of the universe is an example of a *local* measurement that provides an estimate of  $\Omega_0$ . But the result we obtain depends on just how local our measurement is and there are many uncertainties in the result, especially on large scales. In an attempt to get around this problem, astronomers have devised alternative methods that rely instead on *global* measurements, covering much larger regions of the observable universe. In principle, such global tests should indicate the universe's overall density, not just its value in our cosmic neighborhood.


One such global method is based on observations of Type I (carbon-detonation) supernovae. [∞ \(Sec. 21.3\)](#) Recall that these objects are very bright and have a remarkably narrow spread in luminosities, making them particularly useful as standard candles. [∞ \(Sec. 24.2\)](#) They can be used as probes of the universe because, by measuring their distances (*without* using Hubble's law) and their redshifts, we can determine the rate of cosmic expansion in the distant past. Here's how the method works.

Suppose the universe is decelerating, as we would expect if gravity were slowing its expansion. Then, because the expansion rate is decreasing, objects at great distances—that is, objects that emitted their radiation long ago—should appear to be receding *faster* than Hubble's law predicts. Figure 26.11(a) illustrates this concept. If the universal expansion were constant in time, recessional velocity and distance would be related by the black line. (The line is not quite straight, because it takes the expansion of the universe properly into account in computing the distance.) [∞ \(More Precisely 24-1\)](#) In a decelerating universe, the velocities of distant objects should lie *above* the black curve, and the deviation from that curve should be greater for a denser universe in which gravity has been more effective at slowing the expansion.

How does theory compare with reality? In the late 1990s, two groups of astronomers announced the results of independent, systematic surveys of distant supernovae. Some of these supernovae are shown in Figure 26.11(b); the data are marked on Figure 26.11(a). Far from clarifying the picture of cosmic deceleration, these findings indicated that the expansion of the universe is not slowing, but actually accelerating! According to the supernova data, galaxies at large distance are receding *less* rapidly than Hubble's law would predict. The deviations from the decelerating curves appear small in the figure, but they are statistically very significant, and both groups report similar findings. Subsequent supernova



(a)



(b)

▲ **FIGURE 26.11 Accelerating Universe** (a) In a decelerating universe (purple and red curves), redshifts of distant objects are greater than would be predicted from Hubble's law (black curve). The reverse is true for an accelerating universe. The data points show observations of some 50 supernovae that strongly suggest cosmic expansion is accelerating. (b) The bottom frames show three Type-I supernovae (marked by arrows) that exploded in distant galaxies when the universe was nearly half its current age. The top frames show the same areas prior to the explosions. (CfA; NASA)

observations, most recently from the Sloan Digital Sky Survey in 2009, are generally in agreement with the initial findings. ∞ (*Discovery 25-1*)

These observations are *inconsistent* with the “gravity only” Big Bang model just described and have sparked a major revision of our view of the cosmos. The measurements

are difficult, and the results depend quite sensitively on just how “standard” the supernovae luminosities really are; some astronomers initially questioned the accuracy of the method. In particular, if supernovae at great distances (i.e., long ago) were for some reason slightly less luminous than those nearby, then we would think that these distant supernovae were farther away than they actually are, and the error would appear as a deviation to the right of the black curve in Figure 26.11(a)—in other words, as an acceleration in the cosmic expansion rate.

Not surprisingly, since so much hangs on this measurement, the reliability of the supernova measurement technique has been the subject of intense scrutiny by cosmologists. However, no convincing argument against the method has yet been put forward, so there is no reason to think that we are somehow being “fooled” by nature. As far as we can tell, the measurements are good, and the acceleration is real.

## Dark Energy


What could cause an overall acceleration of the universe? Frankly, cosmologists don't know, although several possibilities have been suggested. Whatever it is, the mysterious cosmic field causing the universe to accelerate is neither matter nor radiation. Although it carries energy, it exerts an overall *repulsive* effect on the universe, speeding up the expansion of empty space. It has come to be called **dark energy**, and it is perhaps *the* leading puzzle in astronomy today.

As illustrated in Figure 26.12, the repulsive effect of dark energy is proportional to the size of the universe, so it increases as the universe expands. Thus, it was negligible at early times, but today, given the magnitude of the observed acceleration, it is the major factor controlling the cosmic expansion. Furthermore, since the effect of gravity *weakens* as the expansion proceeds, it follows that once dark energy begins to dominate, gravity can never catch up, and the universe will continue to accelerate at an ever-increasing pace. Thus, although there is considerable uncertainty as to the nature of dark energy, we can at least say that, by opposing the attractive force of gravity, dark energy's repulsion strengthens our earlier conclusion that the universe will expand forever.

One leading dark-energy candidate is an additional “vacuum energy” force associated with empty space and effective only on very large scales. Known simply as the **cosmological constant**, it has a long and checkered history. It was first proposed by Einstein as a way to force his new theory of general relativity into “predicting” a static universe, but was subsequently dropped from Einstein's equations following Hubble's discovery that the universe is not static, but instead is expanding (see *Discovery 26-1*). Since the 1990s, the cosmological constant has arisen again, to become a staple of astronomers' models of the universe. Note, however, that although models that take this force into account can fit the observational data, as described in



► **FIGURE 26.12 Dark Energy** The expansion of the universe is opposed by the attractive force of gravity and sped up by the repulsion due to dark energy. As the universe expands, the gravitational force weakens, whereas the force due to dark energy increases. A few billion years ago, dark energy began to dominate, and the expansion of the cosmos has been accelerating ever since.



the next section, astronomers have *no* clear physical interpretation of what the force actually is. It is neither required nor explained by any known law of physics.

An additional problem for cosmologists is the fact that the present value of the repulsive force is *comparable* to the attractive force of gravity opposing further expansion. Why is that a problem? Because, when we calculate the evolution of a universe containing a cosmological constant consistent with current observations (see Figure 26.14), we find that this state of affairs was not true in the early universe (when galaxies were forming, say), nor will it be true in 10 or 20 billion years' time. In other words, the observations seem to suggest that we live at a special time in the history of the universe—a conclusion viewed with great suspicion by astronomers who grew up with the Copernican principle as their guide.

A promising alternative dark-energy candidate, called *quintessence*,\* might offer a means of avoiding this problem. Whereas the cosmological constant is a property of empty space and is independent of any matter or “normal” energy that space contains, quintessence evolves in time in a way that depends on the matter and radiation in the universe. By coupling the behavior of dark energy to the other contents of the cosmos, quintessence may provide a natural mechanism for dark energy to emerge as the dominant force as the universe expands and cools, and galaxies begin to form and grow.

With little hard data to constrain them, theorists have considerable freedom in constructing models of the dark-matter content of the universe. Cosmologists are searching for experimental and observational tests to refine their models and to distinguish between competing theories.

### CONCEPT Check

✓ Why do astronomers think the universe will expand forever?

\*In ancient alchemy, *quintessence* was the “fifth element,” after earth, air, fire, and water. It was believed to be the perfect substance composing the heavens and all heavenly bodies.

## 26.6 Dark Energy and Cosmology

As we proceed through the remainder of this chapter and the next, it is worth bearing in mind that the Big Bang *is* a scientific theory and, like any other, must continually be challenged and scrutinized. ∞ (Sec. 1.2) The Big Bang theory makes detailed, testable predictions about the state and history of the cosmos and must change—or be replaced—if these predictions are found to be at odds with observations. The supernova observations just described are a case in point.

Even though the supernova observations and their interpretation have so far withstood intense scrutiny, the idea of an accelerating universe driven by some completely unknown field called dark energy probably would not have gained such rapid and widespread acceptance among cosmologists were it not strongly supported by several other pieces of evidence. In this section, we discuss how the existence of dark matter fits in with observations of the universe and even helps resolve some long-standing riddles. Every independent piece of evidence, and every old puzzle solved, provides further support not just for the idea of dark energy, but also for the entire Big Bang theory of the universe.

### Composition of the Universe

In addition to measuring density and acceleration, astronomers have several other means of estimating the “cosmological parameters” that describe the large-scale properties of our universe.

Theoretical studies of the early universe (to be discussed in more detail in Chapter 27) strongly suggest that the geometry of the universe should be precisely flat—that is, that the total density of the cosmos should exactly equal the critical value. This idea first became widespread in the 1980s, and for many years there seemed to be a major discrepancy between it and observations that clearly showed a cosmic matter density of less than 30 percent of the critical

## DISCOVERY 26-1

## Einstein and the Cosmological Constant

Even the greatest minds are fallible. The first scientist to apply general relativity to the universe was, not surprisingly, the theory's inventor, Albert Einstein. When he derived and solved the equations describing the behavior of the universe, Einstein discovered that they predicted a universe that evolved in time. But in 1917, neither he nor anyone else knew about the expansion of the universe, as described by Hubble's law, which would not be discovered for another 10 years. [☞ \(Sec. 24.3\)](#) At the time, Einstein, like most scientists, believed that the universe was static—that is, unchanging and everlasting. The discovery that there was no static solution of his equations seemed to Einstein to be a near-fatal flaw in his new theory.

To bring the theory into line with his beliefs, Einstein tinkered with the equations, introducing a “fudge factor” describing a hypothetical repulsive force operating on large scales in the universe. This factor is now known as the cosmological constant. One possible solution to Einstein's modified equations described a universe in which the repulsive cosmological constant just balanced the attractive force of gravity, allowing the size of the cosmos to remain constant for an indefinite period of time. Einstein took this to be the static universe he expected.

Instead of predicting an evolving cosmos, which would have been one of general relativity's greatest triumphs, Einstein yielded to a preconceived notion of the way the universe “should be,” unsupported by observational evidence. Later, when the expansion of the universe was discovered and Einstein's equations—without the fudge factor—were found to describe it perfectly, he declared that the cosmological constant was the biggest blunder of his scientific career.

Scientists are reluctant to introduce unknown quantities into their equations purely to make the results “come out right.”

Einstein introduced the cosmological constant to fix what he thought was a serious problem with his equations, but he discarded it immediately once he realized that no problem actually existed. As a result, the cosmological constant fell out of favor among astronomers for many years.

In the 1980s, the concept made something of a comeback with the realization by physicists that the very early universe may have gone through a phase when its evolution was determined by a “cosmological constant” of sorts (see Section 27.4), and this idea is now firmly entrenched in many cosmologists' models of the universe. Today, as discussed in the text, the cosmological constant has apparently been completely rehabilitated and identified as a leading candidate for the “dark energy” whose existence is inferred from studies of the universe on very large scales. As shown in Figure 26.15, inclusion of a suitable dark-energy term in Einstein's equations can cause the expansion of the universe to accelerate, instead of slowing down as it would if only gravity were involved.

For many researchers—Einstein included—the main problem with the cosmological constant was (and still is) the fact that we have no clear explanation for either its existence or its present value. The leading theories of the structure of matter do in fact predict repulsive forces of this sort, but these forces generally operate only under extreme conditions, and, in any case, their “natural” energy scale is vastly greater (by something like a factor of  $10^{120}$ !) than anything consistent with cosmological observations.


As discussed in the text, theoretical efforts are underway to combine aspects of the cosmological constant with a more reasonable scale on which such a cosmic repulsive force might act. But before we make too many sweeping statements about the role of the cosmological constant in cosmology, we should probably remember the experience of its inventor and bear in mind that—at least for now—its physical meaning remains completely unknown.

value, even taking the dark matter into account. Dark energy resolves that conflict by providing another form in which the “extra” density can exist, although not all astronomers are happy at the price of this resolution, which has introduced yet another unknown component into the cosmic mix!

Detailed measurements of the radiation field known to fill the entire cosmos (see Section 26.7 and Chapter 27) strongly support the theoretical prediction that  $\Omega_0 = 1$  and are also consistent with the dark energy inferred from the supernova studies. Further independent corroboration comes from careful analyses of galaxy surveys such as those discussed in Section 26.1, which allow astronomers to measure the growth of large-scale structure in the universe. Simply put, the more mass there is in the universe, the easier it is for clusters, superclusters, walls, and voids to grow as gravity gathers matter into larger and larger clumps. Higher density implies more rapid formation of structure—or, equivalently, *less* structure in the past, given the structure we see around us today. Thus, structure measurements constrain the value of  $\Omega_0$ .

Remarkably, all the approaches just described yield consistent results! The current consensus among cosmologists is that the universe is of precisely critical density,  $\Omega_0 = 1$ , but that this density is made up of both matter (mostly dark) *and* dark energy (converted into mass units as discussed earlier in Section 26.4). Radiation contributes negligibly to the total (see Section 27.1). Based on all available data—and noting that some significant discrepancies exist between recent measurements, as discussed in Section 27.6—our best current estimate is that normal “luminous” matter accounts for 5 percent of the total, dark matter for roughly 25 percent, and dark energy for the remaining 70 percent of the cosmic density. This composition is illustrated schematically in Figure 26.13. This is the assumption underlying Table 24.2 and used consistently throughout this book.


Note that such a universe will expand forever and, the heavy machinery of general relativity and curved spacetime notwithstanding, is perfectly flat (Figure 26.14)—an irony that would no doubt have amused Newton!



▲ **FIGURE 26.13 Composition of the Universe** The universe today is made mostly of mysterious dark energy, accounting for more than two-thirds of the total. Dark matter comprises about a quarter. Normal matter accounts for only a few percent—and of that most is galactic and intergalactic gas. Only a minuscule fraction—about half of 1 percent—consists of stars, planets, and life-forms.


## Cosmic Age Estimates

We have at least one other independent, noncosmological way of testing the preceding important conclusion. In Section 26.2, when we estimated the age of the universe from the accepted value of Hubble’s constant, we made the assumption that the expansion speed of the cosmos was constant in the past. However, as we have just seen, this is a considerable oversimplification. Gravity tends to slow the universe’s expansion, whereas dark energy acts to accelerate it, and the actual expansion of the universe is the result of the competition between the two. In the absence of a cosmological constant, the universe would have expanded faster in



◀ **FIGURE 26.15 Cosmic Age** The age of a universe without dark energy (represented by all three lower curves colored red, purple, and blue) is always less than  $1/H_0$  and decreases for larger values of the present-day density. The existence of a repulsive cosmological constant increases the age of the cosmos, as shown by the green curve that is drawn using the best available cosmic parameters, as described in the text.

*Geometry is simple when all is flat.*



▲ **FIGURE 26.14 Geometry of the Universe** The universe on the largest scales is geometrically flat—governed by the same familiar Euclidean geometry taught in high schools.

the past than it does today, so the assumption of a constant expansion rate leads to an overestimate of the universe’s age—such a universe is younger than the 14 billion years calculated earlier. Conversely, the repulsive effect of dark energy tends to increase the age of the cosmos.

Figure 26.15 illustrates these points. It is similar to Figure 26.8, except that we have added two extra lines, one corresponding to a constant expansion rate at the present value—a completely *empty* 14-billion-year-old universe—the other to the best-fit accelerating universe with the parameters just described. The age of a critical-density universe with no cosmological constant is about 9 billion years. A low-density open universe (again with no cosmological constant) is older than 9 billion years, but still less than 14 billion years old. The age corresponding to the accelerating universe is 13.8 billion years, coincidentally very close to the value for constant expansion.

How does this kind of calculation compare with an age estimated by other means? On the basis of the theory of stellar evolution, the oldest globular clusters formed about 12 billion years ago, and most are estimated to be between 10 and 12 billion years old. ∞ (Secs. 19.6, 20.5) This range is indicated in Figure 26.15. These ancient star clusters are thought to have formed at around the same time as our Galaxy, so they date the epoch of galaxy formation. More important, they can’t be older than the universe! The figure shows that globular cluster

ages are consistent with a 14-billion-year-old cosmos and even allow a couple of billion years for galaxies to form and grow, as discussed in Chapter 25. ∞ (Sec. 25.3) Note also that the cluster ages are *not* consistent with a critical-density universe without dark energy. This independent check of a key prediction is an important piece of evidence supporting the modern version of the Big Bang theory.

Thus, for  $H_0 = 70$  km/s/Mpc, our current best guess of the history of the universe places the Big Bang at roughly 14 billion years ago. The first quasars appeared about 13 billion years ago (at a redshift of 7), the peak quasar epoch (redshifts 2–3) occurred during the next 1 billion years, and the oldest known stars in our Galaxy formed during the 2 billion years after that. Even though astronomers do not currently understand the nature of dark energy, the good agreement between so many separate lines of reasoning has convinced many that the dark-matter, dark-energy Big Bang theory just described is the correct description of the universe. But astronomers aren't ready to relax just yet: The history of this subject suggests that there may be a few more unexpected twists and turns in the road before the details are finally resolved.

### CONCEPT Check

✓ Why have astronomers concluded that dark energy is the major constituent of the universe?

## 26.7 The Cosmic Microwave Background

Looking out into space is equivalent to looking back into time. ∞ (More Precisely 24-1) But how far back in time can we probe? Is there any way to study the universe beyond the most distant quasar? How close can we come to perceiving the edge of time—the very origin of the universe—directly?

A partial answer to these questions was discovered by accident in 1964, during an experiment designed to improve the U.S. telephone system. As part of a project to identify and eliminate interference in planned satellite communications, Arno Penzias and Robert Wilson, two scientists at Bell Telephone Laboratories in New Jersey, were studying the Milky Way's emission at microwave (radio) wavelengths,




▲ **FIGURE 26.16 Microwave Background Discoverers** This “sugarscoop” antenna was built to communicate with Earth-orbiting satellites, but was used by Robert Wilson (right) and Arno Penzias to discover the 2.7-K cosmic background radiation. (Alcatel-Lucent)


using the horn-shaped antenna shown in Figure 26.16. In their data, they noticed a bothersome background “hiss” that just would not go away—a little like the background static on an AM radio station. Regardless of where and when they pointed their antenna, the hiss persisted. Never diminishing or intensifying, the weak signal was detectable at any time of the day, any day of the year, apparently filling all space.

What was the source of this radio noise? And why did it appear to come uniformly from all directions, unchanging in time? Unaware that they had detected a signal of great cosmological significance, Penzias and Wilson sought many different origins for the excess emission, including atmospheric storms, interference from the ground, short circuits of equipment—even pigeon droppings inside the antenna! Eventually, after conversations with colleagues at Bell Labs and theorists at nearby Princeton University, the two experimentalists realized that the origin of the mysterious static was nothing less than the fiery creation of the universe itself. The radio hiss that Penzias and Wilson detected is now known as the **cosmic microwave background**. Their discovery won them the 1978 Nobel Prize in physics.

In fact, researchers had predicted the existence and general properties of the microwave background well before its discovery. As early as the 1940s, physicists had realized that, in addition to being extremely dense, the early universe must also have been very hot, and shortly after the Big Bang the universe must have been filled with extremely high-energy thermal radiation—gamma rays of very short wavelength. Researchers at Princeton had extended these ideas, reasoning that the frequency of this primordial radiation would have been redshifted (simply by cosmic expansion) from gamma ray, to X-ray, to ultraviolet, and eventually all the way into the radio range of the electromagnetic spectrum as the universe expanded and cooled (Figure 26.17). ∞ (Sec. 3.4) By




▲ **FIGURE 26.17 Cosmic Blackbody Curves** Theoretically derived blackbody curves for the entire universe (a) 1 second after the Big Bang, (b) 100,000 years later, (c) 10 million years after that, and (d) today, approximately 14 billion years after the Big Bang.



**▲ FIGURE 26.18 Microwave Background Spectrum** The intensity of the cosmic background radiation, as measured by the *COBE* satellite, agrees very well with theory. The curve is the best fit to the data, corresponding to a temperature of 2.725 K. Experimental errors in this remarkably accurate observation are smaller than the dots representing the data points.

the present time, they argued, this redshifted “fossil remnant” of the primeval fireball should have a temperature of no more than a few tens of kelvins, peaking in the microwave part of the spectrum. The Princeton group was in the process of constructing a microwave antenna to search for this radiation when Penzias and Wilson announced their discovery.


The Princeton researchers confirmed the existence of the microwave background and estimated its temperature at about 3 K. However, because of atmospheric absorption, this part of the electromagnetic spectrum happens to be difficult to observe from the ground, and it was 25 years until astronomers could demonstrate conclusively that the radiation was described by a blackbody curve. In 1989, the *Cosmic Background Explorer (COBE)* satellite measured the intensity of the microwave background at wavelengths straddling the peak of the curve, from a half millimeter up to about 10 cm. The results are shown in Figure 26.18. The solid line is the black



**▲ FIGURE 26.19 Microwave Sky** A *COBE* map of the entire sky reveals that the microwave background appears a little hotter in the direction of the constellation Leo and a little cooler in the opposite direction. (NASA)

body curve that best fits the *COBE* data. The near-perfect fit corresponds to a universal temperature of about 2.7 K.

Figure 26.19 shows a *COBE* map of the microwave background temperature over the entire sky. The blue regions are hotter than average, by about 0.0034 K, the red regions cooler by the same amount. This temperature range is not an inherent property of the microwave background, however. Rather, it is a consequence of Earth’s *motion* through space. If we were precisely at rest with respect to the universal expansion (like the coin taped to the surface of the expanding balloon in Figure 26.5), then we would see the microwave background as almost perfectly isotropic, as illustrated in Figure 26.20(a). However, if we are moving with respect to that frame of reference, as in Figure 26.20(b), then the radiation from in front of us should be slightly blueshifted by our motion, whereas that from behind should be redshifted. Thus, to a moving observer, the microwave background should appear a little hotter than average in front and slightly cooler behind.



**◀ FIGURE 26.20 Earth’s Motion Through the Cosmos** (a) To an observer at rest with respect to the expanding universe, the microwave background radiation appears isotropic. (b) A moving observer measures “hot” blueshifted radiation in one direction (the direction of motion) and “cool” redshifted radiation in the opposite direction.

The data indicate that Earth's velocity is about 380 km/s in the approximate direction of the constellation Leo. Once the effects of this motion are corrected for, the cosmic microwave background is found to be strikingly isotropic. Its intensity is virtually constant (in fact, to about one part in  $10^5$ ) from one direction on the sky to another, lending strong support to one of the key assumptions underlying the cosmological principle.

When we observe the microwave background, we are looking almost all the way to the very beginning of the universe. The photons that we receive as these radio waves today have not interacted with matter since the universe was a mere

400,000 years old, when, according to our models, it was less than a thousandth of its present size. To probe further, back to the Big Bang itself, requires that we enter the world of nuclear and particle physics. The Big Bang was the biggest and the most powerful particle accelerator of all! In the next chapter, we will see how studies of conditions in the primeval fireball aid us in understanding the present-day structure and future evolution of the universe in which we live.

### CONCEPT Check


✓ When was the cosmic microwave background formed?

**The Big Question** Alpha and omega; beginning and end. What is the origin of the universe, and what will be its ultimate fate? Dare humanity ask such really big questions, and can astronomers hope to answer them? Scientists today are actively sorting through a plethora of ideas, supported by steadily improving data, trying to address questions so fundamental they were once posed only by philosophers and theologians. But this is science today. We may be on the cusp of answering some of the deepest queries any human has ever asked.


# Chapter Review

## SUMMARY


1 Redshift surveys reveal that, on scales larger than a few hundred megaparsecs, the universe appears roughly **homogeneous** (the same everywhere, p. 669) and **isotropic** (the same in all directions, p. 669). In **cosmology** (p. 668)—the study of the universe as a whole—researchers usually assume that the universe is homogeneous and isotropic. This assumption is known as the **cosmological principle** (p. 669) and implies that the universe cannot have a center or an edge.



2 If the universe were homogeneous, isotropic, infinite, and unchanging, the night sky would be bright because any line of sight would eventually intercept a star. The fact that the night sky is dark is called **Olbers's paradox** (p. 670). Its resolution lies in the fact that, regardless of whether the universe is finite or infinite, we see only a finite part of it from Earth—the region from which light has had time to reach us since the universe began.




3 Tracing the observed motions of galaxies back in time implies that some 14 billion years ago the universe consisted of a hot, dense **primeval fireball** (p. 670) that expanded rapidly in the **Big Bang** (p. 671). However, the galaxies are not flying apart into the rest of an otherwise empty universe; rather, space itself is expanding. The Big Bang did not happen at any particular location in space, because space itself




was compressed to a point at that instant—the Big Bang happened everywhere at once. The cosmological redshift occurs as a photon's wavelength is “stretched” by cosmic expansion. The extent of the observed redshift is a direct measure of the expansion of the universe since the photon was emitted.

4 There are only two possible outcomes to the current expansion: Either the universe will expand forever, or it will eventually recollapse. The **critical density** (p. 674) is the density of matter needed for gravity alone to overcome the present expansion and cause the universe to collapse. Most astronomers think that the total mass density of the universe today is no more than about 30 percent of the critical value.




5 General relativity provides a description of the geometry of the universe on the largest scales. The curvature of spacetime is determined by the total density of the universe, including that of matter, radiation, and dark energy. The curvature in a high-density (greater than critical) universe is sufficiently large that the universe “bends back” on itself and is finite in extent, somewhat like the surface of a sphere. Such a universe is said to be a **closed universe** (p. 676). A low-density **open universe** (p. 676) is infinite in extent and has a “saddle-shaped” geometry.




The **critical universe** (p. 676) has a density precisely equal to the critical value and is spatially flat.

6 Observations of distant supernovae indicate that the expansion of the universe is accelerating, apparently driven by the effects of **dark energy** (p. 679), a mysterious repulsive force that exists throughout all space. The physical nature of dark energy is unknown. Possible candidates include the **cosmological constant** (p. 679) and quintessence.




7 The best available observational data are consistent with the idea that the universe is flat—that is, of exactly critical density—with matter (mostly dark) making up 27 percent of the total and dark energy making up the rest. Such a universe is spatially flat and will expand forever. For  $H_0 = 70$  km/s/Mpc, the age of a critical-density universe without dark energy would be about 9 billion years.

This age estimate conflicts with the 10- to 12-billion-year ages of globular clusters derived from studies of stellar evolution. The inclusion of dark energy increases the age of the universe to 14 billion years, consistent with the cluster ages.



8 The **cosmic microwave background** (p. 683) is an isotropic black-body radiation field that fills the entire universe. Its present temperature is about 3 K. The existence of the microwave background is direct evidence that the universe expanded from a hot, dense state. As the universe has expanded, the initially high energy radiation has been redshifted to lower and lower temperatures.



**MasteringAstronomy**® For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- POS** What evidence do we have that there is no structure in the universe on very large scales? How large is “very large”?
- LO1** What is the cosmological principle?
- LO2** What is Olbers’s paradox? How is it resolved?
- Explain how an accurate measure of Hubble’s constant leads to an estimate of the age of the universe.
- LO3** Why isn’t it correct to say that the expansion of the universe involves galaxies flying outward into empty space?
- Where did the Big Bang occur?
- How does the cosmological redshift relate to the expansion of the universe?
- LO4** What properties of the universe determine whether it will or will not expand forever?
- Is there enough matter to halt the current cosmic expansion?
- LO5** Do we live in a “flat” universe?
- LO6** What do observations of distant supernovae tell us about the expansion of the universe?
- LO7** What is dark energy, and what does it have to do with the future of the universe?
- POS** Why are measurements of globular cluster ages important to cosmology?
- LO8** What is the significance of the cosmic microwave background?
- POS** Do you think it is good science to explain the universe mainly in terms of dark matter and dark energy, neither of which is known or understood?

## Conceptual Self-Test: Multiple Choice

- If observations made from the middle of a large city are isotropic, then (a) there are tall buildings in every direction; (b) all buildings are exactly the same height; (c) all buildings are the same color; (d) some buildings are taller than others.
- The cosmological principle would be disproved if we found (a) the universe is not expanding; (b) galaxies are older than currently estimated; (c) the number of galaxies per square degree is the same in every direction; (d) the observed structure of the universe depends on the direction in which we look.
- When we use Hubble’s law to estimate the age of the universe, the result (a) depends on which galaxies we choose; (b) is the same for all galaxies; (c) depends on the direction in which we look; (d) proves that we are at the center of the universe.

4. Olbers's paradox is resolved by (a) the finite size of the universe; (b) the finite age of the universe; (c) light from distant galaxies being redshifted so we can't see it; (d) the fact that there is an edge to the universe.
5. **VIS** The data points in Figure 26.11 ("Accelerating Universe") (a) prove that the universe is not expanding; (b) imply that the expansion is decelerating faster than expected; (c) allow a measurement of Hubble's constant; (d) indicate that the redshifts of distant galaxies are greater than would be expected if gravity alone were acting.
6. The galactic distances used to measure the acceleration of the universe are determined by observations of (a) trigonometric parallax; (b) line broadening; (c) Cepheid variable stars; (d) exploding white dwarfs.
7. The observed acceleration of the universe means that (a) we understand the nature of dark energy; (b) the amount of dark energy is small compared with the luminous mass in galaxies; (c) the amount of dark energy exceeds the total mass-energy of matter in the universe; (d) dark energy has a higher temperature than expected.
8. On the basis of our current best estimate of the present mass density of the universe, astronomers think that (a) the universe is finite in extent and will expand forever; (b) the universe is finite in extent and will eventually collapse; (c) the universe is infinite in extent and will expand forever; (d) the universe is infinite in extent and will eventually collapse.
9. The age of the universe is estimated to be (a) less than Earth's age; (b) the same as the age of the Sun; (c) the same as the age of the Milky Way Galaxy; (d) greater than the age of the Milky Way Galaxy.
10. The cosmic background radiation is observed to come from (a) the center of our Galaxy; (b) the center of the universe; (c) radio antennae in New Jersey; (d) all directions equally.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

1. • What is the greatest distance at which a galaxy survey sensitive to objects as faint as 20th magnitude could detect a galaxy as bright as the Milky Way (absolute magnitude  $-20$ )?
2. •• Assuming  $H_0 = 70$  km/s/Mpc, estimate the redshift of the Milky Way at the distance calculated in the previous question.
3. • If the entire universe were filled with Milky Way-like galaxies, with an average density of 0.1 galaxy per cubic megaparsec, calculate the total number of galaxies observable by the survey in Problem 1 if it covered the entire sky.
4. • According to the Big Bang theory described in this chapter, *without a cosmological constant*, what is the maximum possible age of the universe if  $H_0 = 60$  km/s/Mpc? 70 km/s/Mpc? 80 km/s/Mpc?
5. •• Eight galaxies are located at the corners of a cube. The present distance from each galaxy to its nearest neighbor is 10 Mpc, and the entire cube is expanding according to Hubble's law, with  $H_0 = 70$  km/s/Mpc. Calculate the recession velocity of one corner of the cube relative to the opposite corner.
6. •• The Virgo Cluster is observed to have a recession velocity of 1200 km/s. For  $H_0 = 70$  km/s/Mpc and a critical-density universe, calculate the total mass contained within a sphere centered on Virgo and just enclosing the Milky Way. What is the escape speed from the surface of this sphere?
7. • For a Hubble constant of 70 km/s/Mpc, the critical density is  $9 \times 10^{-23}$  kg/m<sup>3</sup>. (a) How much mass does this correspond to within a volume of 1 cubic astronomical unit? (b) How large a cube would be required to enclose 1 Earth mass of material?
8. •• (a) What is the present peak wavelength of the cosmic microwave background? Calculate the size of the universe relative to its present size when the radiation background peaked in (b) the infrared, at 10  $\mu$ m, (c) in the ultraviolet, at 100 nm, and (d) in the gamma-ray region of the spectrum, at 1 nm.

## Activities

### Collaborative


1. Make a model of a two-dimensional universe and examine Hubble's law on it. Find a balloon that will expand into a nice large sphere. Blow it up about halfway and mark dots all over its surface, representing galaxies. Each group member should choose one dot as his/her home galaxy. Measure the distances to various other galaxies, numbering the dots so you will not confuse them later. Now blow the balloon up to full size and measure the distances again. Calculate the change in the distances for each galaxy; this is a measure

of recession velocity. Plot the velocities versus the new distances as in Figure 24.17 or 26.11. Do you find a "Hubble" law? Does it matter which dot you choose as home?

### Individual

1. Isotropy is the extent to which things look the same in every direction. Considering buildings, geographic features, and similar objects within a few miles of your current location, is your local universe isotropic? If not, is there any scale on which isotropy applies, even approximately?





# The Early Universe

## TOWARD THE BEGINNING OF TIME

What were the conditions during the first few seconds of the universe, and how did those conditions change to give rise to the universe we see today? In studying the earliest moments of our universe, we enter a truly alien domain. As we move backward in time toward the Big Bang, our customary landmarks slip away one by one. Atoms vanish, then nuclei, and then even the elementary particles themselves.

In the beginning, the universe consisted of pure energy at unimaginably high temperatures. As it expanded and cooled, the ancient energy gave rise to the particles that make up everything we see around us today. Modern physics has now arrived at the point where it can reach back almost to the instant of the Big Bang itself, allowing scientists to unravel some of the mysteries of our beginnings in time.

**The Big Picture** Modern cosmology maintains that the entire universe can be traced back to an extraordinarily hot and dense energy state billions of years ago. As mind-boggling as it may seem, all that we see around us apparently arose from microscopic “quantum” fluctuations that occurred a fraction of a second after the Big Bang. Ironically, the largest-scale structures observed in astronomy today are inextricably tied to the smallest scales known in physics.

**LEFT:** Underground on the Swiss-French border near Geneva, scientists are using the world’s biggest physics experiment to explore matter more deeply than ever before. The Large Hadron Collider simulates events that occurred within the first second after the origin of the universe by violently smashing together subatomic protons. Here, inside a blue-shaded detector, two protons (red streaks) collide, producing a spray of particles (yellow) that enable scientists to test their best theoretical ideas about how the universe began. (CERN)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Describe the characteristics of the universe immediately after its birth.
- 2 Explain how matter emerged from the primeval fireball.
- 3 Describe how radiation and matter evolved as the universe expanded and cooled.
- 4 State how and when the simplest nuclei formed.
- 5 Explain the consequences of the formation of the first atoms.
- 6 Summarize the horizon and flatness problems, and describe how the theory of cosmic inflation solves them.
- 7 Describe the formation of large-scale structure in the cosmos.
- 8 Explain how studies of the microwave background allow astronomers to test and quantify their models of the universe.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.

## 27.1 Back to the Big Bang

On the very largest scales, the universe is a roughly homogeneous mixture of matter (mostly dark), radiation, and dark energy. ∞ (Sec. 26.5) As we have seen, “matter” includes both *normal matter*, made up of protons, neutrons, and electrons, and *dark matter*, whose composition is still being debated by astronomers. *Dark energy* is the mysterious repulsive force that permeates even the apparent vacuum of intergalactic space. As best we can tell, we live in a geometrically “flat” universe in which the total mass—energy density of *all* the constituents of the cosmos exactly equals the critical value. ∞ (Secs. 26.3, 26.4, 26.6) According to theoretical models, there is not enough matter in the cosmos for the attractive force of gravity to overcome the repulsion of dark energy and reverse the current expansion.

Thus, the future of the cosmos seems clear: The universe is destined to expand forever. In this chapter, we turn our attention to the past. To understand the early universe, just after the Big Bang, we must look more closely at the roles played by matter, radiation, and dark energy in the cosmos. We begin by taking stock of their contributions to the total energy density of the universe.

### Cosmic Composition

On the basis of the best available observational data, cosmologists have concluded that, today, just over 70 percent of the total mass-energy of the universe exists in the form of dark energy. ∞ (Sec. 26.6) Virtually all of the remaining 30 percent is accounted for by matter. Thus, at the present moment, *dark energy dominates the density of the universe*, with matter a rather distant second. We can quantify this statement using the results of Chapter 26. For a Hubble constant  $H_0 = 70 \text{ km/s/Mpc}$ , the critical density is  $9 \times 10^{-27} \text{ kg/m}^3$ . ∞ (Sec. 26.3) Thus, in round numbers, the density of dark energy in the universe today is just over  $6 \times 10^{-27} \text{ kg/m}^3$ ; the current density of matter is slightly less than  $3 \times 10^{-27} \text{ kg/m}^3$ .

Most of the *radiation* in the universe is in the form of the cosmic microwave background—the low-temperature (3 K) radiation field that fills all space. ∞ (Sec. 26.7) Surprisingly, although the microwave background radiation is very weak, it still contains more energy than has been emitted by all the stars and galaxies that have ever existed! The reason is that stars and galaxies, though very intense sources of radiation, occupy only a tiny fraction of space. Averaged out over the volume of the entire universe, their energy falls short of the energy of the microwave background by at least a factor of 10. For our current purposes, then, we can ignore most of the first 26 chapters of this book and regard the cosmic microwave background as the only significant form of radiation in the universe!

Does radiation play an important role in the evolution of the universe on large scales? In order to compare matter

and radiation, we must first as usual convert them to a “common currency”—either mass or energy. We will compare their masses. We can express the energy in the microwave background as an equivalent density by first calculating the number of photons in any meter of space and then converting the total energy of these photons into a mass using the relation  $E = mc^2$ . ∞ (Sec. 16.6) When we do this, we arrive at an equivalent density for the microwave background of about  $5 \times 10^{-31} \text{ kg/m}^3$ . Thus, *at the present moment*, the densities of both dark energy and matter in the universe far exceed the density of radiation.


### Radiation in the Universe

Was the universe always dominated by dark energy? To answer this question, we must ask how the densities of dark energy, matter, and radiation changed as the universe expanded. To this end, cosmologists construct *theoretical models* of the universe, taking into account the effects of Einstein’s general relativity and incorporating both the known properties of matter and radiation and the assumed properties of dark energy. ∞ (Sec. 22.6) These models describe how cosmic quantities (such as the densities of the various components) change as the universe evolves. They also make detailed *predictions*, which can be compared directly with observations. ∞ (Sec. 1.2) The outstanding agreement between models and reality (see Section 27.5) is the main reason that astronomers attach so much weight to the measurements of cosmic density, composition, and evolution described in the previous chapter.

As illustrated in Figure 27.1, the models indicate that, as the scale of the universe increases, the densities of matter and radiation both decrease, with the expansion diluting the numbers of atoms and photons alike. But the radiation is also diminished in energy by the cosmological redshift, so its density falls *faster* than that of matter as the universe grows. The dark energy behaves in a very different way. According to theory, it is a large-scale phenomenon, increasing in importance as the universe expands (see Figure 26.12). In fact (at least, if it behaves like Einstein’s cosmological constant), the density associated with the dark energy remains *constant* as the universe expands. ∞ (Discovery 26-1)

Hence, as we look *back* in time, closer and closer to the Big Bang, the density of the radiation increases faster than that of matter, and both increase faster than that of dark energy. These facts allow us to draw two important conclusions about the composition of the universe in the past.

1. Even though dark energy dominates the density of the universe today, it was *unimportant* at early times, and we can neglect it in our discussion of conditions in the very early universe. Astronomers estimate that the densities of matter and dark energy were equal about 4 billion



**▲ FIGURE 27.1 Radiation–Matter Dominance** As the universe expanded, the number of both matter particles and photons per unit volume decreased. The photons were additionally reduced in energy by the cosmological redshift. As a result, the density of radiation (blue curve) fell faster than the density of matter (red curve) as the universe grew, and radiation dominated matter at early times, before the crossover point. Today, dark energy (green line) dominates both matter and radiation.

years ago. Before then, in cosmological parlance, the universe was **matter-dominated**.

2. Although the radiation density is currently much less than that of matter, there must have been a time even farther in the past when they, too, were equal. Before that time, radiation was the main constituent of the cosmos, which is said to have been **radiation-dominated**. The crossover point—the time at which the densities of matter and radiation were equal—occurred about 50,000 years after the Big Bang, when the universe was about 6000 times smaller than it is today. The temperature of the background radiation at that time was about 16,000 K, so it peaked in the near-ultraviolet portion of the spectrum. [∞ \(Sec. 26.7\)](#)

Throughout this book, we have been concerned with the history of the universe long after it became dominated by matter and/or dark energy—the formation and evolution of galaxies, stars, and planets as the universe thinned and cooled toward the state we see today. In this chapter, we consider some important events in the early, hot, radiation-dominated universe, long before any star or galaxy existed, that played no less a role in determining the present condition of the cosmos.

## Particle Production

The existence of the microwave background implies that the early universe was dominated by an intense radiation field



whose temperature fell steadily as the cosmos expanded. The temperatures and densities prevailing at these times were far greater than anything we have encountered thus far, even in the hearts of supernovae. To understand conditions in the universe shortly after the Big Bang, we must delve more deeply into the behavior of matter and radiation at very high temperatures.

The key to understanding events at very early times lies in a process called **pair production**, in which two photons give rise to a *particle–antiparticle* pair, as shown in Figure 27.2(a) for the particular case of electrons and positrons. Through pair production, matter is created directly from energy in the form of electromagnetic radiation. The reverse process can also occur: A particle and its antiparticle can *annihilate* each other to produce radiation, as depicted in Figure 27.2(b). Energy in the form of radiation can be converted into matter in the form of particles and antiparticles, and particles and antiparticles can be converted back into radiation, subject only to the law of conservation of mass and energy.


The higher the temperature of a radiation field, the greater the energy of the typical constituent photons, and the greater the masses of the particles that can be created by pair production. [∞ \(Secs. 3.4, 4.2, 16.6\)](#) For any given particle, the critical temperature above which pair production is possible and below which it is not is called the particle's *threshold temperature*. The threshold temperature increases as the mass of the particle increases. For electrons, it is about  $6 \times 10^9$  K. For protons, which are nearly 2000 times more massive, it is just over  $10^{13}$  K.

As an example of how pair production affected the composition of the early universe, consider the production of electrons and positrons as the universe expanded and cooled. At high temperatures—above about  $10^{10}$  K—most photons had enough energy to form an electron or a positron, and pair production was commonplace. Space seethed with electrons and positrons, constantly created from the radiation field and annihilating one another to form photons again. Particles and radiation are said to have been in *thermal equilibrium*: New particle–antiparticle pairs were created by pair production at the same rate as they annihilated one another. As the universe expanded and the temperature decreased, so did the average photon energy. By the time the temperature had fallen below 1 billion kelvins, photons no longer had enough energy for pair production to occur, and only radiation remained. Figure 27.3 illustrates how this change took place.

Pair production in the very early universe was directly responsible for all the matter that exists in the universe today. *Everything we see around us was created out of radiation as the cosmos expanded and cooled.* Because we are here to ponder the subject, we know that some matter must have survived those early violent moments. For some



**▲ FIGURE 27.2 Pair Production** (a) Two photons can produce a particle–antiparticle pair—in this case an electron and a positron—if their total energy exceeds the mass energy of the particles produced. (b) The reverse process is particle–antiparticle annihilation, in which an electron and a positron destroy each other, vanishing in a flash of gamma rays. (c) This actual observation, on submicroscopic scales, shows two gamma rays (whose paths at left are invisible because they are electrically neutral) dislodging an atomic electron and sending it flying (the longest track). At the same time, the gamma rays provide enough energy to produce an electron–positron pair (the spiral paths, which curve in opposite directions in the detector’s magnetic field because of their opposite electric charges). (*Fermi Laboratory*)



**▲ FIGURE 27.3 Thermal Equilibrium** (a) At 10 billion K, most photons have enough energy to create particle–antiparticle (electron–positron) pairs, so these particles exist in great numbers in equilibrium with the radiation. The label  $e^-$  refers to the electrons,  $e^+$  to positrons. (b) Below about  $10^9$  K, photons have too little energy for pair production to occur, so electrons and positrons are no longer in thermal equilibrium with the background radiation field.

reason, there was a slight excess of matter over antimatter at early times—about one extra proton for every billion proton–antiproton pairs. That small residue of particles that outnumbered their antiparticles was left behind as the temperature dropped below the threshold for creating them. With no antiparticles left to annihilate them, the number of particles has remained constant ever since. These survivors are said to have *frozen out* of the radiation field as the universe expanded and cooled.

According to the models, the first hundred or so seconds of the universe’s existence saw the creation of all of the basic “building blocks” of matter we know today. Protons and neutrons froze out when the temperature dropped below  $10^{13}$  K, when the universe was only 0.0001 seconds old; the lighter electrons froze out somewhat later, about a minute or so after the Big Bang, when the temperature fell below  $10^9$  K. This “matter-creation” phase of the universe’s evolution ended when the electrons—the lightest known elementary particles—appeared out of the cooling primordial fireball. From then on, matter has continued to evolve, clumping into more and more complex structures, eventually forming the atoms, planets, stars, galaxies, and large-scale structure we see today, but for all practical purposes no new matter has been created since that early time.

**CONCEPT Check**

✓ What does it mean to say that the early universe was radiation dominated?

**27.2 Evolution of the Universe**

For the first few thousand years after the Big Bang, the universe was small, dense, and dominated by radiation. We will refer to this period as the *radiation era*. Some matter existed during this time, but it was a mere contaminant in the blinding gamma-ray light of the primeval Big Bang fireball. Afterward, in the *matter era*, matter came to dominate. Atoms, molecules, and galaxies formed as the universe cooled and thinned toward the state we see today. Today we live in the *dark-energy era*, in which dark energy is becoming an increasingly important component of the cosmos.

Let's begin our study of the early universe by summarizing in broad terms the history of the cosmos, starting at the Big Bang. Figure 27.4 illustrates how the cosmic temperature and density dropped rapidly during the radiation and matter eras and identifies eight significant epochs in the development of the universe. Notice how the time scale on the horizontal axis increases from tiny fractions of a second to thousands of years as we move from left to

right—the rate of change of the cosmos slowed dramatically as the universe expanded. We will focus on these epochs in greater detail in the next few sections, but let's not lose sight of the big picture and the place of each epoch in it.

**Before the Big Bang?**

The Big Bang was a *singularity* in space and time—an instant when the present laws of physics imply that the universe had zero size and infinite temperature and density. As we saw in Chapter 22, where we discussed the singularities at the center of black holes, these predictions should not be taken too literally. ∞ (Sec. 22.7) The presence of singularities signals that, under extreme conditions, the theory making the predictions—in this case, general relativity—has broken down.


At present, no theory exists to let us penetrate the singularity at the start of the universe. We have no means of describing these earliest of times, so we have no way of answering the question “What came *before* the Big Bang?” Indeed, given the laws of physics as we currently know them, the question itself may be meaningless. The Big Bang represented the beginning of the entire universe—mass, energy, space, *and* time came into being at that instant. Without time, the notion of “before” does not exist. Consequently, some cosmologists maintain that asking what

happened before the Big Bang is a little like asking what lies north of the North Pole! Others disagree, however, arguing that when the correct theory of quantum gravity—the “Theory of Everything” that unifies gravity and quantum mechanics—is constructed it will remove the singularity and allow us to address the question of what came before.

**The Birth of the Cosmos**

Although ignorant of the moment of creation itself, theorists nevertheless think that the physical conditions in the universe can be understood in terms of present-day physics back to an extraordinarily short time—a mere  $10^{-43}$  s, in fact—after the Big Bang.

Why can't theorists push our knowledge back to the Big Bang itself? The answer is that we presently have no theory capable of describing the universe at these earliest of times. Under the extreme conditions of density and temperature within  $10^{-43}$  s of the Big Bang, gravity and the other fundamental forces (electromagnetism, the strong force, and the weak force, as described in *More Precisely 27-1*) were



▲ **FIGURE 27.4 Epochs in Cosmic History** The average temperature and average density throughout the history of the universe. At the earliest times, the universe was a sea of radiation with a single unified force of nature. Some subsequent key events in the history of the cosmos as it expanded and cooled are marked; they are discussed later in the chapter.

## MORE PRECISELY 27-1

### More on Fundamental Forces

In *More Precisely 16-1*, we noted that the behavior of all matter in the universe is ruled by just three *fundamental forces*: gravity, the electroweak force (the unification of the electromagnetic and weak forces), and the strong (nuclear) force. In terrestrial laboratories, these forces display properties that are very different from one another (see Table 27.1). Gravity and electromagnetism are long-range, inverse-square forces, whereas the strong and weak forces have very short ranges— $10^{-15}$  and  $10^{-17}$  m, respectively. Furthermore, the forces do not all affect the same particles. Gravity affects everything. The electromagnetic force affects only charged particles. The strong force operates between nuclear particles, such as protons and neutrons, but it does not affect electrons and neutrinos. The weak force shows up in certain nuclear reactions and radioactive decays. The strong force is 137 times stronger than the electromagnetic force, 100,000 times stronger than the weak force, and  $10^{39}$  times stronger than gravity.

In fact, there is more structure below the level of the nucleus. Protons and neutrons are not truly “elementary” in nature, but are actually made of subparticles called **quarks**. (The name derives from a meaningless word coined by novelist James Joyce in his book *Finnegans Wake*.) According to current theory, there are six distinct types of quark in the universe (with the obscure names *up*, *down*, *charm*, *strange*, *top*, and *bottom*). What we call the strong nuclear force is actually a manifestation of the interactions that bind quarks to one another.

On the face of it, one might not imagine that there could be any deep underlying connection between forces as dissimilar as those just described, yet there is strong evidence that they are really just different aspects of a single basic phenomenon. In the 1960s, theoretical physicists succeeded in explaining the electromagnetic and weak forces in terms of the electroweak force. Shortly thereafter, the first attempts were made at combining the strong and electroweak forces into a single all-encompassing “superforce.” A central idea in the modern version of this superforce is that there is a one-to-one correspondence between the quarks, which interact via the strong force, and particles called **leptons**, which are affected only by the electroweak force. The six known types of quark are paired with six distinct types of lepton: the electron, two related “electron-like” particles (called muons and taus), and three types of neutrino.

Theories that combine the strong and electroweak forces into one are generically known as **Grand Unified Theories**, or GUTs for short. (Note that the term is plural—no one GUT has yet been proven to be “the” correct description of nature.) One general prediction of GUTs is that the three nongravitational forces are indistinguishable from one another only at enormously high energies, corresponding to temperatures in excess of  $10^{28}$  K. Below that temperature, the superforce splits into two, displaying its separate strong and electroweak aspects. In particle physics parlance, we say that there is a *symmetry* between the strong and the electroweak forces that is broken at

temperatures below  $10^{28}$  K, allowing the separate characters of the two forces to become apparent. At “low” temperatures—less than about  $10^{15}$  K, a range that includes almost everything we know on Earth and in the stars—there is a second symmetry breaking, and the electroweak force splits to reveal its more familiar electromagnetic and weak natures.

The key predictions of the electroweak theory were experimentally verified in the 1970s, winning the theory’s originators (Sheldon Glashow, Steven Weinberg, and Abdus Salam) the 1979 Nobel Prize in physics. The GUTs have not yet been experimentally verified (or refuted), in large part because of the extremely high energies that must be reached in order to observe their predictions.

An important idea that has arisen from the realization that the strong and the electroweak forces can be unified is the notion of **supersymmetry**, which extends the idea of symmetry between fundamental forces to place all particles—those that are acted on by forces (such as protons and electrons) and those that transmit those forces (such as photons and gluons; see Section 27.4)—on an equal footing. One particularly important prediction of supersymmetry is that all particles should have so-called *supersymmetric partners*—extra particles that must exist in order for the theory to remain self-consistent. None of these new particles has yet been detected, yet many physicists are convinced of the theory’s essential correctness.

These new particles, if they exist, would have been produced in abundance in the Big Bang and should still be around today. They are also expected to be very massive—at least a thousand times heavier than a proton. So-called supersymmetric relics, the new particles, are among the current leading candidates for the dark matter in the universe (see Section 27.5).

Efforts to include gravity within this picture have so far been unsuccessful. Gravitation has not yet been incorporated into a single “Super GUT,” in which all the fundamental forces are united. Some theoretical efforts to merge gravity with the other forces have tried to fit gravity into the quantum world by postulating extra particles—called *gravitons*—that transmit the gravitational force. However, this is a different view of gravity from the geometric picture embodied in Einstein’s general relativity, and combining the two into a consistent theory of quantum gravity has proved very difficult.

One promising theory that is currently under active investigation seeks to interpret all particles and forces in terms of particular modes of vibration of submicroscopic objects known as *strings*. **String theory** is complex, but it solves a number of intractable technical problems that plagued previous efforts, and many theorists feel that it currently offers the greatest promise of unifying the forces of nature. Promise aside, though, realize that at present, *no* theory has yet succeeded in making any definite statement about conditions in the very early universe. A complete theory of quantum gravity continues to elude researchers.

**TABLE 27.1** Fundamental Forces and Particles

Force	Range (m)	Particles Affected	Unification (temperature)	
strong	$10^{-15}$	matter composed of quarks (protons, neutrons, etc.)	electroweak ( $10^{15}$ K)	GUT/superforce ( $10^{28}$ K)
electromagnetic	infinite	charged particles (protons, electrons, etc.)		
weak	$10^{-17}$	leptons (electrons, muons, taus, neutrinos)		
gravity	infinite	everything		quantum gravity ( $10^{32}$ K)

indistinguishable from one another—a far cry from the radically different characteristics we see today, listed in Table 27.1. The four forces are said to have been *unified* at that early time—there was, in effect, only one force of nature.

The theory that combines quantum mechanics (the proper description of microscopic phenomena) with general relativity (which describes the universe on the largest scales) is generically known as quantum gravity. ∞ (Sec. 22.7) The period from the beginning to  $10^{-43}$  s is often referred to as the *Planck epoch*, after Max Planck, one of the creators of quantum mechanics. Unfortunately, for now at least, there is no working theory of quantum gravity, so we simply cannot talk meaningfully about the universe during the Planck epoch.

By the end of the Planck epoch, the temperature was around  $10^{32}$  K, and the universe was filled with radiation and a vast array of subatomic particles created by the mechanism of pair production. At around that time, gravity parted company with the other forces of nature—it became distinguishable from them and has remained so ever since. The strong, weak, and electromagnetic forces were still unified. The present-day theories that describe this epoch are collectively known as Grand Unified Theories, or GUTs for short (see *More Precisely 27-1*). Accordingly, we refer to this period as the *GUT epoch*.


## Freeze-Out

Grand Unified Theories predict that three of the four basic forces of nature—electromagnetism and the strong and weak nuclear forces—are in reality aspects of a single, all-encompassing “superforce.” However, this unification is evident only at enormously high energies, corresponding to temperatures in excess of  $10^{28}$  K. At lower temperatures, the superforce reveals its separate electromagnetic, strong, and weak characters.

A fundamental concept in quantum physics is the idea that forces between elementary particles are exerted, or *mediated*, by the exchange of another type of particle,

generically called a **boson**. We might imagine the two particles as playing a rapid game of catch, using a boson as a ball, as illustrated in Figure 27.5. As the ball is thrown back and forth, the force is transmitted. For example, in ordinary electromagnetism, the boson involved is the *photon*—a bundle of electromagnetic energy that always travels at the speed of light. The strong force is mediated by particles known as *gluons*. The electroweak theory includes a total of four bosons—the massless photon and three other massive particles, called (for historical reasons)  $W^+$ ,  $W^-$ , and  $Z^0$ , all of which have been observed in laboratory experiments. Gravity is (theoretically) mediated by *gravitons*, and so on. All of the particles we have encountered so far in this book—electrons, protons, neutrons, neutrinos—play “catch” with at least some of these “balls.”

In Section 27.1 we saw how particles “froze out” of the universe as its temperature dropped below the threshold temperature for their creation by pair production. Now that we know that the basic forces of nature are mediated by particles, we can understand—in general terms, at least—how the fundamental forces froze out, too, as the universe cooled. According to the GUTs, the particle that unifies the strong and electroweak forces is extremely massive—at least  $10^{15}$  times the mass of the



▲ **FIGURE 27.5 Fundamental Forces** Forces between elementary particles are transmitted through the exchange of other particles called bosons. As two particles interact, they exchange bosons, a little like playing catch with a submicroscopic ball.



proton (and possibly much more). It is because this particle is so massive that the unification of the strong and electroweak forces becomes evident only at extremely high temperatures.

At temperatures below  $10^{28}$  K, the strong nuclear force becomes distinguishable from the electroweak force (the unified weak and electromagnetic forces). Once the universe had cooled to that temperature, about  $10^{-35}$  s after the Big Bang, the GUT epoch ended. According to many GUTs, one important legacy of that epoch may have been the appearance and subsequent freeze-out of a veritable “zoo” of very massive (and as yet unobserved) elementary particles that interact only very weakly with normal matter. These “exotic” particles are prime candidates for the dark matter of unknown composition thought to exist in abundance both within galaxies and in the unseen depths of intergalactic space.

∞ (Secs. 23.6, 25.1)

## Quarks and Leptons

Our next major subdivision of the radiation era covers the period when all “heavy” elementary particles—that is, all the way down in mass to protons, neutrons, and their constituent quarks—were in thermal equilibrium with the radiation. We refer to this period as the *quark epoch*, since quarks are the fundamental components of all particles that interact via the strong force.

The universe continued to expand and cool. At a temperature of about  $10^{15}$  K ( $10^{-10}$  s after the Big Bang), the weak and the electromagnetic components of the electroweak force began to display their separate characters. The W and Z particles responsible for the electroweak force have masses about 100 times the mass of a proton. The threshold temperature for their production—roughly  $10^{15}$  K—marks the point at which the weak and electromagnetic forces parted company.

By about 0.1 millisecond ( $10^{-4}$  s) after the Big Bang, the temperature had dropped well below the  $10^{13}$  K threshold for the creation of protons and neutrons (the lightest stable particles composed of quarks), and the quark epoch ended. The main constituents of the universe were now lightweight particles—muons (see *More Precisely 27-1*), electrons, neutrinos, and their antiparticles—all still in thermal equilibrium with the radiation. Compared with the numbers of these lighter particles, only very few protons and neutrons remained at this stage, because most had been annihilated.

Electrons, muons, and neutrinos are collectively known as *leptons*, after the Greek word meaning “light” (i.e., not heavy). Accordingly, we refer to this period in the history of the universe as the *lepton epoch*.

During that epoch, at a temperature of about  $3 \times 10^{10}$  K—approximately 1 second after the Big Bang—the rapidly thinning universe became transparent to neutrinos, and these ghostly particles have been streaming freely through space ever since. (Most neutrinos have not interacted with any other particle since the universe was a few seconds old!) The lepton epoch ended when the universe was about 100 seconds old and the temperature fell to about  $10^9$  K—too low for electron–positron pair production to occur. The density of the universe by this time was about 10 times the density of water.

The final significant event in the radiation era occurred when protons and neutrons began to fuse into heavier nuclei. At the start of this period, which we will call the *nuclear epoch*, the temperature was a few hundred million kelvins, and fusion occurred very rapidly, forming deuterium (“heavy” hydrogen—see Section 16.6) and helium in quick succession before conditions became too cool for further reactions to occur. By the time the universe was about 15 minutes old, much of the helium we observe today had been formed.

## The Matter and Dark-Energy Eras

Time passed, the universe continued to expand and cool, and radiation gave way to matter as the dominant constituent of the universe. Our next major epoch extends in time from 50,000 years (the end of the radiation era) to about 100 million years after the Big Bang. As the primeval fireball diminished in intensity, a crucial change occurred—perhaps the most important change in the history of the universe. At the end of the nuclear epoch, radiation still overwhelmed matter. As fast as protons and electrons combined, radiation broke them apart again, preventing the formation of even simple atoms or molecules. However, as the universe expanded and cooled, the early dominance of radiation eventually ended. Once formed, atoms remained intact. We will call this period the *atomic epoch*. It ended about 200 million years after the Big Bang, when the first stars formed and their intense radiation *reionized* the universe.

The last two epochs together bring us to the current age of the universe. During these late stages, change happened at a much more sedate pace. By the time the universe was about 3 billion years old, large-scale structure and most galaxies had formed. For the first time, the universe departed from homogeneity on macroscopic scales. The largely uniform universe of the radiation era became a universe containing large agglomerations of matter. We call the period from 200 million to 3 billion years after the Big Bang the *galactic epoch*, given that the main events at the time concerned galaxy construction. At its end,

large-scale structure and the bulk of most galaxies had formed, quasars were shining brightly, and early generations of stars were burning and exploding, helping to determine the future shape of their parent galaxies.

Since then, galaxies have continued to merge and evolve, stars peaked their formation rate, and planets and life appeared in the universe. These last two epochs, including the current *stellar epoch*—so named for the myriad stars that are still forming within galaxies—have been the subject of the first 25 chapters of this book.

### CONCEPT Check


✓ Why did lighter and lighter particles “freeze out” of the universe as the cosmos expanded?

## 27.3 Formation of Nuclei and Atoms

We now have all the ingredients needed to complete our story of the creation of the elements, begun in Chapter 21, but never quite finished. ∞ (Sec. 21.4) The theory of stellar nucleosynthesis accounts very well for the observed abundances of heavy elements in the universe, but there are discrepancies between theory and observation when it comes to the abundances of the light elements, especially helium. Simply put, the total amount of helium in the universe today—about 25 percent by mass—is far too large to be explained by nuclear fusion in stars. The accepted explanation is that this base level of helium is *primordial*—that is, it was created during the early, hot epochs of the universe, before any stars had formed. The production of elements heavier than hydrogen by nuclear fusion shortly after the Big Bang is called **primordial nucleosynthesis**.

### Helium Formation in the Early Universe




By about 100 s after the Big Bang, the temperature had fallen to about 1 billion K, and apart from “exotic” dark-matter particles, matter in the universe consisted of electrons, protons, and neutrons, with the protons outnumbering the neutrons by about five to one. The stage was set for nuclear fusion to occur. Protons and neutrons combined to produce deuterium nuclei (also called *deuterons*), containing 1 proton and 1 neutron:



Although this reaction must have occurred frequently during the lepton epoch, the temperature then was still so high

that the deuterium nuclei were broken apart by high-energy gamma rays as soon as they formed. The universe had to wait until it became cool enough for the deuterium to survive. This waiting period is sometimes called the *deuterium bottleneck*.

Only when the temperature of the universe fell below about 900 million K, roughly 2 minutes after the Big Bang, was deuterium at last able to form and endure. Once that occurred, the deuterium was quickly converted into heavier elements by numerous reactions, including:




The result was that, once the universe passed the deuterium bottleneck, fusion proceeded rapidly and large amounts of helium were formed. In just a few minutes most of the free neutrons were consumed, leaving a universe whose matter content was primarily hydrogen and helium. Figure 27.6 illustrates some of the reactions responsible for helium formation. Contrast it with Figure 16.27, which depicts how helium is formed today in the cores of main-sequence stars such as the Sun.\* ∞ (Sec. 16.6)

We might imagine that fusion could have continued to create heavier and heavier elements, just as it does in the cores of stars, but that did not occur. In stars, the density and the temperature both *increase* slowly with time, allowing more and more massive nuclei to form, but in the early universe exactly the opposite was true. The temperature and density were both *decreasing* rapidly, making conditions less and less favorable for fusion as time went on. Even before the supply of neutrons was completely used up, the nuclear reactions had effectively ceased. Reactions between helium nuclei and protons may also have formed trace amounts of lithium (the next element beyond helium) by this time, but for all practical purposes, the expansion of the universe caused fusion to stop at helium. The brief epoch of primordial nucleosynthesis was over about 15 minutes after it began.

By the end of the period of nucleosynthesis, some 1000 seconds after the Big Bang, the temperature of the universe was about 300 million K and the cosmic elemental abundances were set. Careful calculations indicate that about one helium nucleus had formed for every 12 protons remaining. Because a helium nucleus is four times more massive than a

\*The proton–proton chain that powers the Sun played no significant role in primordial helium formation. The proton–proton reaction that starts the chain is very slow compared with the proton–neutron discussed here and is important in the Sun only because the solar interior contains no free neutrons to make the latter reaction possible.



**FIGURE 27.6 Helium Formation** Some of the reaction sequences that led to the formation of helium in the early universe. (Recall that a deuteron is a nucleus of deuterium, the heavy form of hydrogen.) Compare this figure with Figure 16.27, which depicts the proton–proton chain in the Sun.

proton, helium accounted for about one-quarter of the total mass of matter in the universe:

$$\begin{aligned} \frac{1 \text{ helium nucleus}}{12 \text{ protons} + 1 \text{ helium nucleus}} &= \frac{4 \text{ mass units}}{12 \text{ mass units} + 4 \text{ mass units}} \\ &= \frac{4}{16} = \frac{1}{4}. \end{aligned}$$

The remaining 75 percent of the matter in the universe was hydrogen. It would be almost a billion years before nucleosynthesis in stars would change these numbers.

∞ (Sec. 21.4)

The foregoing calculation implies that all stars and galaxies should contain *at least 25 percent* helium by mass. The figure for the Sun, for example, is about 28 percent. However, it is difficult to disentangle the contributions to the present-day helium abundance from primordial nucleosynthesis and later hydrogen burning in stars. Our best hope of determining the amount of primordial helium is to study the oldest stars known, since they formed early on, before stellar nucleosynthesis had had time to change the helium content of the universe significantly. Unfortunately, stars surviving from that early time are of low mass and hence quite cool, making the helium lines in their spectra very weak and hard to measure accurately. ∞ (Secs. 17.5, 17.8) Nevertheless, despite this uncertainty, the observations are generally consistent with the theory just described.


Bear in mind that while all this was going on, matter was just an insignificant “contaminant” in the radiation-dominated universe. Radiation outmassed matter by about

a factor of 5000 at the time helium formed. The existence of helium is very important in determining the structure and appearance of stars today, but its creation was completely irrelevant to the evolution of the universe at the time.

## Deuterium and the Density of the Cosmos

During the nuclear epoch, although most deuterium was quickly fused into helium as soon as it formed, a small amount was left over when the primordial nuclear reactions ceased. Observations of deuterium—especially those made by orbiting satellites able to capture deuterium’s strongest spectral feature, which happens to be emitted in the ultraviolet part of the spectrum—indicate a present-day abundance of about two deuterium nuclei for every 100,000 protons. However, unlike helium, deuterium is not produced to any significant degree in stars (in fact, deuterium tends to be destroyed in stars), so any deuterium we see today *must* be primordial.

This observation is of great importance to astronomers because it provides them with a sensitive method—and one that is completely independent of the techniques discussed in previous chapters—of probing the present-day density of matter in the universe. According to theory, as illustrated in Figure 27.7, the denser the universe is today, the more particles there were at early times to react with deuterium as it formed and the less deuterium was left over when nucleosynthesis ended. A comparison of the observed deuterium abundance (marked on the figure) with the theoretical results implies a present-day



**▲ FIGURE 27.7 Deuterium Abundance** Today's abundance of deuterium depends strongly on the amount of matter present at early times, and this, in turn, determines the present-day density of the universe. Thus, measuring the amount of deuterium in the universe gives us an estimate of the overall density of matter. The best deuterium measurements reside within the blue band, and imply that the density of matter in the universe is at most a few percent of the critical value.

density of *at most*  $5 \times 10^{-28} \text{ kg/m}^3$ —only a few percent of the critical density.

But before we jump to any far-reaching cosmic conclusions based on this number, we must make a very important qualification. As just described, primordial nucleosynthesis depends *only* on the presence of protons and neutrons in the early universe. Thus, measurements of the abundance of helium and deuterium tell us only about the density of “normal” matter—matter made up of protons and neutrons—in the cosmos. This finding has a momentous implication for the overall composition of the universe. As we saw earlier, astronomers have concluded, for a variety of reasons, that the total density of matter is about the critical value.  $\infty$  (Sec. 26.5) In that case, if the density of normal matter is only a few percent of the critical value, then we are forced to admit that not only is most of the matter in the universe dark, but most of the dark matter is *not* composed of protons and neutrons.

We will see in Section 27.5 that, because normal matter and dark matter interact differently with the background radiation field, studies of the cosmic microwave background allow us to distinguish between these two types of matter. Observations made by the *WMAP* and *Planck* spacecraft have found the density of normal matter to be just 5 percent of the critical density,

in excellent agreement with estimates based on the abundance of deuterium.


Thus, the bulk (about 90 percent) of the matter in the universe apparently exists in the form of elusive subatomic particles (for example, the WIMPs discussed as dark-matter candidates in Chapter 23) whose nature we do not fully understand and whose very existence has yet to be conclusively demonstrated in laboratory experiments.  $\infty$  (Sec. 23.6) For the sake of brevity, from here on we will adopt the convention that the term “dark matter” refers only to these unknown particles and not to “stellar” dark matter, such as black holes and brown and white dwarfs (also discussed in Chapter 23), which are made of relatively well-understood normal matter.

## The First Atoms

A few tens of thousands of years after the Big Bang, radiation ceased to be the dominant component of the universe. The matter era had begun. At the start of the atomic epoch, matter consisted of electrons, protons, helium nuclei (formed by primordial nucleosynthesis), and dark matter. The temperature was several tens of thousands of kelvins—far too hot for atoms of hydrogen to exist (although some helium ions may already have formed). During the next few hundred thousand years, a major change occurred: The universe expanded by another factor of 10, the temperature dropped to a few thousand kelvins, and electrons and nuclei combined to form neutral atoms. By the time the temperature had fallen to about 3000 K, the universe consisted of atoms, photons, and dark matter.

The period during which nuclei and electrons combined to form atoms is called the epoch of **decoupling**, for it was during this period that the radiation background parted company with normal matter. Many astronomers also refer to this period as **recombination** (although, technically speaking, protons and electrons had never previously been combined in the form of atoms).

At early times, when matter was ionized, the universe was filled with large numbers of free electrons that interacted frequently with electromagnetic radiation of all wavelengths. As a result, a photon could not travel far before encountering an electron and scattering off it. In effect, the universe was opaque to radiation. Matter and radiation were strongly “tied,” or *coupled*, to one another by these interactions. After the electrons combined with nuclei to form atoms of hydrogen and helium, only certain wavelengths of radiation—the ones corresponding to the spectral lines of those atoms—could interact with matter.  $\infty$  (Sec. 4.2) Radiation of other wavelengths could travel virtually forever without being absorbed. Thus, the universe became nearly transparent. From that time on, most photons passed generally unhindered



▲ **FIGURE 27.8 Radiation–Matter Decoupling** When atoms formed, the universe became virtually transparent to radiation. Thus, observations of the cosmic background radiation reveal conditions in the universe around a time when the redshift was 1100 and the temperature was less than about 3000 K. For an explanation of how we can see a region of space 14,000 Mpc (46 billion light-years) away when the universe is just 14 billion years old, see *More Precisely 24-1*.

through space. As the universe expanded, the radiation simply cooled, eventually becoming the microwave background we see today.

The microwave photons now detected on Earth have been traveling through the universe ever since they decoupled. According to the models that best fit the observational data, the last interaction these photons had with matter (at the epoch of decoupling) occurred when the universe was about 400,000 years old and roughly 1100 times smaller (and hotter) than it is today—that is, at a redshift of 1100. As illustrated in Figure 27.8, the epoch of atom formation created a kind of “photosphere” in the universe, completely surrounding Earth at a distance of approximately 14,000 Mpc, the distance at which photons last interacted before they decoupled. [∞ \(More Precisely 24-1\)](#)

On our side of the photosphere—that is, since decoupling—the universe is transparent. On the far side—before decoupling—it was opaque. Thus, by observing the microwave background, we are probing conditions in the universe almost all the way back in time to the Big Bang, in much the same way as studying sunlight tells us about the surface layers of the Sun.

## PROCESS OF SCIENCE Check


- ✓ How do we know that most of the dark matter in the universe is not of “normal” composition?

## 27.4 The Inflationary Universe

In the late 1970s, cosmologists trying to piece together the evolution of the universe were confronted with two nagging problems that had no easy explanation within the standard Big Bang model. The resolution of these problems has caused cosmologists to completely rethink their views of the very early universe.

### The Horizon and Flatness Problems

The first problem is known as the **horizon problem**, and it concerns the remarkable *isotropy* of the cosmic microwave background. [∞ \(Sec. 26.7\)](#) Recall that the temperature of this radiation is virtually constant, at about 2.7 K, in all directions. Imagine observing the microwave background in two opposite directions of the sky, as illustrated in Figure 27.9. As we have just seen, that radiation last interacted with matter in the universe at around a redshift of 1100. Thus, in observing these two distant regions of the universe, marked A and B on the figure, we are studying



▲ **FIGURE 27.9 Horizon Problem** The isotropy of the microwave background indicates that regions A and B in the universe were very similar to each other when the radiation we now observe left them, but there has not been enough time since the Big Bang for them ever to have interacted with one another. Why then should they look the same?


regions that were separated by several million parsecs when they emitted this radiation. The fact that the background radiation is isotropic to high accuracy means that regions A and B had similar densities and temperatures at the time the radiation we see left them.

The problem is, according to the Big Bang theory as just described, there is *no* good reason why these regions should in fact be similar to each other. To take an everyday example, we all know that heat flows from regions of high temperature to regions of low temperature, but it takes time for this to occur. If we light a fire in one corner of a room, we have to wait a while for the other corners to warm up. Eventually, the room reaches a more or less uniform temperature, but only after the heat from the fire—or, more generally, the *information* that the fire is there—has had time to spread.

Similar reasoning applies to regions A and B in Figure 27.9. These regions are separated by many megaparsecs, and there has not been enough time for information, which can go no faster than the speed of light, to travel from one to the other. In cosmological parlance, the two regions are said to be outside each other's *horizon*. But if that is so, then how do they “know” that they are supposed to look the same? With no possibility of communication between them, the only alternative is that regions A and B simply started off looking alike—an assumption that cosmologists are reluctant to make.

The second problem with the standard Big Bang model is called the **flatness problem**. Whatever the exact value of  $\Omega_0$ , it appears to be very close to unity—the total density of the universe is fairly near the critical value. In terms of spacetime curvature, the universe is remarkably close to being flat.  $\infty$  (Sec. 26.4) We say “remarkably” here because, again, there is no particular reason that the universe should have formed with a density very close to the critical value. Why not a millionth of, or a million times, that value? Furthermore, as illustrated in Figure 27.10, a universe that starts off close to, but not exactly on, the critical curve soon deviates greatly from it, so if the universe is close to critical now, it must have been *extremely* close to critical in the past. (The acceleration due to dark energy *does* in fact tend to push the universe toward critical density, but dark energy has not dominated the expansion for long enough for this fact to change our basic conclusion.) For example, if  $\Omega_0 = 0.3$  today (approximately the density of “known” normal and dark matter), then the departure from critical density at the time of nucleosynthesis would have been only 1 part in  $10^{15}$  (a thousand trillion)!

These observations constitute “problems” because cosmologists want to be able to *explain* the present condition of the universe, not just accept it “as is.” They would prefer to resolve the horizon and flatness problems in terms of physical processes that could have taken a universe with no




▲ **FIGURE 27.10 Flatness Problem** If the universe deviates even slightly from critical density, that deviation grows rapidly in time. For the universe to be as close to critical as it is today, it must have differed from the critical density in the past by only a tiny amount.

special properties and caused it to evolve into the cosmos we now see. The resolution of both problems takes us back in time even earlier than nucleosynthesis or the formation of any of the elementary particles we know today—back, in fact, almost to the instant of the Big Bang itself.

## Cosmic Inflation

As we saw in Section 27.2, at very early times, during the GUT and Planck epochs, most or all of the fundamental forces of nature were unified—that is, indistinguishable from one another. The various theories that describe this unification (for example, those outlined in *More Precisely 27-1*) predict—and in fact, rely upon—the existence of certain quantum-mechanical fields, generically called **scalar fields** in particle physics jargon, whose interactions with particles in the theory determine those particles’ properties. For our purposes, we can think of these fields as cosmic forces permeating all space, separate from, but closely related to, physical particles in the universe. These fields define the differences between the various forces of nature, and ultimately set the scale on which unification occurs.

What does all this have to do with cosmology? In the early 1980s, physicists realized that it was possible for these scalar fields to become temporarily *increased* in energy above their normal equilibrium states. Due to



▲ **FIGURE 27.11 Cosmic Inflation** During the period of inflation, the universe expanded enormously in a very short time. Afterward, it resumed its earlier “normal” expansion rate, except that the size of the cosmos had become about  $10^{50}$  times bigger than it was before inflation.

random fluctuations at the quantum level, regions of the universe could find themselves in this “elevated” state for some period of time. Under these circumstances, theory indicates that these parts of the universe would find themselves in a very odd and unstable condition—empty space would have acquired **vacuum energy**. Abstract as this may seem, these regions are of direct interest to us—if theorists are correct, we live in one!

The temporary appearance of vacuum energy within such a region—ours, say—had dramatic consequences. For a short while, as illustrated in Figure 27.11, the extra energy caused the region to expand at an enormously accelerated rate. The vacuum energy density remained almost constant as the region grew, and the expansion *accelerated* with time while this condition persisted. In fact, the size of the region doubled many times over. For definiteness, Figure 27.11 shows the expansion occurring near the end of the GUT epoch, with a doubling time of roughly  $10^{-34}$  s. This period of unchecked cosmic expansion is known as the **epoch of inflation**.

Actually, bizarre though this may seem, we have already seen a universal expansion along these lines, albeit at a much more leisurely pace. The leading models for dark energy (the cosmological constant and quintessence) are both scalar fields. Their nonzero vacuum energy is responsible for the cosmic acceleration discussed in Chapter 26.

∞ (Sec. 26.5)

Eventually, the scalar field returned to its equilibrium state, the region recovered its normal vacuum, and inflation stopped. For the example shown in Figure 27.11, the whole episode lasted a mere  $10^{-32}$  s, but during that time the patch of the universe that had become unstable swelled in size by the incredible factor of about  $10^{50}$ . After the inflationary phase, the universe once again resumed its (relatively) leisurely expansion. However, a number of important changes had occurred that would have far-reaching ramifications for the evolution of the cosmos.


The original theory of inflation was developed in the early 1980s and associated the inflationary period (as in Figure 27.11) with the end of the GUT epoch. The scalar field in that case was the one responsible for distinguishing between the strong and electroweak forces. However, since that time, researchers have realized that conditions suitable for inflation could have occurred under many different circumstances—and possibly *many times*—during the evolution of the early universe. This generalization actually strengthens inflation as a theory by loosening the restrictions on when it might have happened, although it blurs the question of exactly when the inflationary epoch(s) leading to “our” universe actually occurred.

Nevertheless, the basic idea of a *quantum fluctuation* expanding to become the universe we know is now quite well established. Some theorists have gone so far as to suggest that a quantum fluctuation during the Planck epoch may have been the trigger that caused the Big Bang. Others even speculate that we might be living in a sort of “self-creating universe” that erupted into existence spontaneously from inflation in one such random fluctuation! This sort of “statistical” creation of the primal cosmic energy from absolutely nothing has been dubbed “the ultimate free lunch.”

Note that it is possible—even likely, according to many theorists—that not all of the universe underwent inflation. Only some regions became unstable, causing huge inflated “bubbles” to appear in the cosmos. We apparently live in one such bubble; the universe outside is probably unknowable to us. Henceforth, we will use the term “universe” to refer to just this bubble and its contents.

## Implications for the Universe

The inflationary epoch provides a natural solution to the horizon and flatness problems. The horizon problem is solved because inflation took regions of the universe that had already had time to communicate with one another—and so had established similar physical properties—and then dragged them far apart, well out of communication range of one another. For example (again following Figure 27.11), regions A and B in Figure 27.9 have been out of contact since  $10^{-32}$  s after creation, but they were in contact before then. As illustrated in Figure 27.12, their



**▲ FIGURE 27.12 Inflation and the Horizon Problem** Inflation solves the horizon problem by taking a small region of the very early universe—whose parts had already had time to interact with one another and that had thus already become homogeneous—and expanding it to enormous size. In (a), points A and B are well within the (shaded) homogeneous region of the universe centered on the eventual site of the Milky Way Galaxy. In (b), after inflation, A and B are far outside the horizon (indicated by the dashed line), so they are no longer visible from our location. Subsequently, the horizon expands faster than does the universe as a whole, so that today (c) A and B are just reentering our field of view. They have similar properties now because they had similar properties before the inflationary epoch.

properties are the same today because they were the same long ago, before inflation separated them.

Figure 27.12(a) shows a small piece of the universe just before the onset of inflation. The point that will one day become the site of the Milky Way Galaxy is at the center of the shaded region, which represents the portion of space “visible” to that point at that time—that is, there has been enough time since the Big Bang for light to have traveled from the edge of this region to its center. That entire region is more or less homogeneous, because different parts of it have been able to interact with one another, so any initial differences between the parts have largely been smoothed out. The points A and B of Figure 27.9 are also marked. They lie within the homogeneous patch, so they have very similar properties. The actual size of the shaded region is about  $10^{-26}$  m—only a trillionth the size of a proton.


Immediately after inflation, as shown in Figure 27.12(b), the homogeneous region has expanded by 50 orders of magnitude, to a diameter of about  $10^{24}$  m, or 30 Mpc—larger than the largest supercluster. By contrast, the visible portion of the universe, indicated by the dashed line, has grown only by a factor of a thousand and is still microscopic in size. In effect, the universe expanded much faster than the speed of light during the inflationary epoch, so what was once well within the horizon of the point that is to become the site of our Galaxy now lies far beyond it. In particular, points A and

B are no longer visible, either to us or to each other, at this time. (Note that, while the theory of relativity restricts matter and energy to speeds less than the speed of light, it imposes no such limit on the universe as a whole.) ∞ (Discovery 22-1)

Since the end of inflation, the universe has expanded by a further factor of  $10^{27}$ , so the size of the homogeneous region of space surrounding us is now about  $10^{51}$  m ( $10^{28}$  Mpc)—10 trillion trillion times greater than the distance to the most distant quasar. As shown in Figure 27.12(c), the horizon has expanded faster than the universe, so points A and B are just now becoming visible again. As the portion of the universe that is now observable from Earth grows in time, it remains homogeneous because our cosmic field of view is simply reexpanding into a region of the universe that was within our horizon long ago. We will have to wait a very long time—at least  $10^{35}$  years—before the edge of the homogeneous patch surrounding us comes back into view.

To see how inflation solves the flatness problem, let’s return to our earlier balloon analogy. ∞ (Sec. 26.2) Imagine that you are a 1-mm-long ant sitting on the surface of the balloon as it expands, as illustrated in Figure 27.13. When the balloon is just a few centimeters across, you can easily perceive the surface to be curved—its circumference is only a few times your own size. When the balloon expands to, say, a few meters in diameter, the curvature of the surface is less pronounced, but perhaps still perceptible. However, by the time the balloon has expanded to a few





◀ **FIGURE 27.13 Inflation and the Flatness Problem** Inflation solves the flatness problem by enormously expanding a curved surface, here represented by the surface of a balloon. To an ant on the surface, the balloon looks virtually flat when the expansion is complete.

kilometers across, an “ant-sized” patch of the surface will look quite flat, just as the surface of Earth looks flat to us.

Now imagine that the balloon expands 100 trillion trillion trillion times, as the universe did during the period of inflation. Your local patch of the surface is now completely indistinguishable from a perfectly flat plane, deviating from flatness by no more than one part in  $10^{50}$ . Exactly the same argument applies to the universe: Because it has expanded so much, for all practical purposes the universe is perfectly flat on all scales we can ever hope to observe.

Notice that this resolution of the flatness problem—the universe appears close to being flat because the universe *is* in fact precisely flat, to very high accuracy—has a very important consequence: Because the universe is geometrically flat, relativity tells us that the total density must be exactly equal to the critical value of  $\Omega_0 = 1$ . ∞ (Sec. 26.4) This is the key result that led us to conclude in Chapter 26 that dark energy—whatever it is—must dominate the density of the universe. ∞ (Sec. 26.5) Thus, the combined weight of theory and observation forces us to the conclusion that not only is most matter dark (Section 27.3), but also *most of the cosmic density isn’t made up of matter at all*.

## Inflation as a Theory

Even though inflation solves the horizon and flatness problems in a quite convincing way, for nearly two decades after it was first proposed the theory was resisted by many astronomers. The main reason was that its prediction of  $\Omega_0 = 1$  was clearly at odds with the growing evidence that the density of matter in the universe was no more than 30 percent or so of the critical value. Actually, many cosmologists *had* considered the possibility that a cosmological constant

offered a way to account for the remaining 70 percent of the cosmic density, but without independent corroboration a conclusive case could not be made. That is why the supernova observations were so important: By providing empirical evidence for acceleration in the cosmic expansion rate, they established independent evidence for the effects of dark energy and, in doing so, reconciled inflation with the otherwise discrepant observations. ∞ (Sec. 26.5)

Physicists will probably never create in terrestrial laboratories conditions even remotely similar to those that existed in the universe during the inflationary epoch. The creation of our own vacuum energy is (safely) beyond our reach. Nevertheless, cosmic inflation seems to be a natural consequence of many Grand Unified Theories. It explains two otherwise intractable problems within the Big Bang theory—and, following the empirical observations of cosmic acceleration, it is now reconciled with measurements of the matter density of the universe with the inclusion of dark energy into the cosmic mix.

For all these reasons, despite the absence of direct evidence for the process, inflation theory has become an integral part of modern cosmology. Inflation makes definite, testable predictions about the large-scale geometry and structure of the universe that are critically important to current theories of galaxy formation. As we will see in the next section, astronomers are now subjecting these predictions to rigorous scrutiny.

### CONCEPT Check

- ✓ Why does the theory of inflation imply that much of the energy density of the universe may be neither matter nor radiation?

## 27.5 Formation of Structure in the Universe

Just as stars formed from *inhomogeneities*—deviations from perfectly uniform density—in interstellar clouds, so too are galaxies, galaxy clusters, and larger structures thought to have grown from small density fluctuations in the matter of the expanding universe. ∞ (Sec. 19.1)

Where did these density fluctuations come from? According to current theories of structure formation, they were the result of microscopic “quantum” fluctuations in the very early universe, expanded to macroscopic scales by the effects of inflation! In a very real sense, the quantum universe just after the Big Bang is the progenitor of all the cosmic structure we see around us today.

Given the conditions in the universe during the atomic and galactic epochs (Table 27.1), cosmologists calculate that regions of higher-than-average density that contained more than about a million times the mass of the Sun would have begun to contract. There was thus a natural tendency for million-solar-mass “pregalactic” objects to form. In Chapter 25, we learned how these pregalactic fragments might have interacted and merged to form galaxies. ∞ (Sec. 25.3) In the rest of this chapter, we concern ourselves mostly with the formation of structure on much larger scales.

### Growth of Inhomogeneities

By the early 1980s, cosmologists had come to realize that galaxies could not have formed from the contraction of inhomogeneities involving only *normal* matter. The following lines of reasoning led to this conclusion:

1. Calculations show that, before decoupling (which occurred at a redshift of 1100), the intense background radiation would have prevented clumps of normal matter from contracting. Matter and radiation were just too strongly coupled for structure to form. Thus, any such clumps would have had to wait until *after* decoupling before their densities could start to increase.
2. Because radiation was “tied” to normal matter up until decoupling, any variations in the matter density at that time would have led to temperature variations in the cosmic background radiation—denser regions would have been a little hotter than less dense ones. The high degree of isotropy observed in the microwave background indicates that any density variations from one region of space to another at the time of decoupling must have been *small*—at most a few parts in  $10^5$ . ∞ (Sec. 26.7)
3. Galaxies—or, at least, quasars—are known to have formed by a redshift of 6. Furthermore, some theorists think that, in order to produce the densest galactic

nuclei we see today, the formation process must have already been well established as long ago as a redshift of 20. ∞ (Sec. 25.3) Thus, the initial fluctuations, which, as we have just seen, must have been very small at a redshift of 1100, had to grow to form the first stars and galaxies by a redshift of 20.

4. The contracting matter had to “fight” the general expansion of the universe. As a result, theory shows, these contracting pregalactic clumps could have increased in density by a factor of at most 50 to 100 in the time available. As a result, the small inhomogeneities permitted by observations of the microwave background *could not have grown into galaxies in the time available*—the universe would still have been almost perfectly homogeneous at a time when we know galaxies had already formed.

Put another way, if galaxies had grown from density fluctuations in the normal-matter component of the early universe, then the fluctuations would have had to be so large as to leave a clearly observable imprint on the cosmic microwave background. That imprint is not observed.







### Dark Matter

Normal matter, then, cannot account for the large-scale structure we see today. Fortunately for cosmology (and for life on Earth), much of the universe is made of *dark* matter, which has properties quite different from those of normal matter and which provides a natural explanation for the large-scale structure we see today. Whatever the nature of dark matter, its defining property is that it interacts only very weakly with normal matter and radiation, so its natural tendency to clump and contract under gravity was not hindered by the radiation background. Dark matter started clumping well before decoupling (redshift 1100)—in fact, density inhomogeneities in the dark-matter component of the universe probably began to grow as soon as matter first began to dominate the universe at a redshift of about 6000.

Because the dark matter was not directly tied to the radiation, these inhomogeneities could have been quite large at the time of decoupling, without having a correspondingly large effect on the microwave background. In short, dark matter could clump to form large-scale structure in the universe without running into any of the problems just described for normal matter.

Thus, as illustrated in Figure 27.14, dark matter determined the overall distribution of mass in the universe and clumped to form the observed large-scale structure without violating any observational constraints on the microwave background. Then, at later times, normal matter was drawn by gravity into the regions of highest density, eventually forming galaxies and galaxy clusters. This picture explains why so much dark matter is found outside the

The plots below are graphical representations of structure growth.



The maps above show what those structures might look like on the sky.

### Narrated FIGURE 27.14 Structure Formation

The formation of structure in the cosmos depended crucially on the existence of dark matter. (a) The very early universe was a mixture of (mostly) dark and normal matter. (b) A few thousand years after the Big Bang, the dark matter began to clump. (c) Eventually, the dark matter formed large structures (represented here by the two high-density peaks) into which normal matter flowed, ultimately to form the galaxies we see today.

**Cold dark matter** consists of very massive particles, possibly formed during the GUT epoch or even before. Computer simulations modeling the universe with these particles as the dark matter easily produce small-scale structure. With the understanding that galaxies form preferentially in the densest regions—as is predicted particularly by models that include a cosmological constant—these models also predict large-scale structure that is in excellent agreement with what is actually observed.

Figure 27.15 shows the results of a supercomputer simulation of a universe consisting of approximately 30 percent matter (most of it dark) and 70 percent dark energy (in the

form of a cosmological constant). (Sec. 26.6) Yellow dots represent regions in each frame where significant star formation is occurring—quasars at redshift 6 and bright interacting galaxies today. The similarities with real observations of cosmic structure, shown in Figures 25.21 and 26.1, are striking, and more detailed statistical analysis confirms that these models agree extremely well with reality. Notice the large-scale extended filamentary structure evident in the last two frames, which is comparable in both scale and appearance to the observed structure presented earlier in the text. Such filaments, which contain both dark and normal matter, are a generic feature of cold dark-matter cosmological simulations. The visible galaxies are also surrounded by extensive dark-matter halos.


Although calculations like this cannot prove that these models are the correct description of the universe, the agreement in detail between models and reality strongly favor the dark-energy/cold dark-matter model of the cosmos.

Given that the nature of the dark matter is still unknown, theorists have considerable freedom in choosing its properties when they attempt to simulate the formation of structure in the universe. Cosmologists conventionally classify dark matter as either “hot” or “cold,” on the basis of its temperature at the time when galaxies began to form. The two types predict quite different kinds of structure in the present-day universe.

**Hot dark matter** consists of lightweight particles—much less massive than the electron. Neutrinos, which appear to have small, but nonzero, masses, are leading candidates for hot dark-matter particles. (Sec. 16.6) However, simulations of a universe filled with hot dark matter indicate that, whereas large structures, such as superclusters and voids, form fairly naturally, structure on smaller scales does not. Small amounts of hot material tend to disperse, not clump together. As a result, most cosmologists have concluded that models based on hot dark matter are unable to explain the observed structure of the universe.

**CONCEPT Check**

✓ Why was dark matter necessary for structure formation in the universe?



▲ **FIGURE 27.15 Structure Simulated** The large-scale structure we see in the universe today is the direct descendent of quantum fluctuations in the early universe, inflated to macroscopic scales. These frames show three views of a (present-day)  $100 \times 100 \times 100$ -Mpc cube in a simulated cold dark-matter universe with  $\Omega_0 = 1$ . The cube expands with the universe, so that it encloses the same material at all times. The three frames show structures progressively growing from small density fluctuations in the early universe, displayed at three different times after the Big Bang. (*V. Springel*)

## 27.6 Cosmic Structure and the Microwave Background

Because dark matter does not interact directly with photons, its density variations do not cause large (and hence easily observable) temperature variations in the microwave background. However, theory suggests that, before decoupling, the cosmos must have been filled with *sound waves*—tiny fluctuations in the densities of normal matter and the background radiation field—moving through space at just over half the speed of light. Most astronomers think that these fluctuations originated at the end of the inflationary epoch, as the universe regained its “normal” vacuum state after the period of unchecked expansion.

Because radiation and matter were tied to one another at this time, the fluctuations in the matter density corresponded to temperature fluctuations in the radiation field, and these features were “imprinted” in the microwave background when matter and radiation finally parted company at a redshift of 1100. As a result, cosmological models predict that there should be tiny “ripples” in the microwave background—temperature variations of only a few tens of parts per million from place to place on the sky.



### Ripples in the Radiation Background

Until the late 1980s these ripples were too small to be measured accurately, although cosmologists were confident that

they would be found. In 1992, after almost 2 years of careful observation, the *COBE* team announced that the expected ripples had indeed been detected. ∞ (Sec. 26.7) The temperature variations are tiny—only 30–40 millionths of a kelvin from place to place in the sky—but they are there. The *COBE* results are displayed as a temperature map of the microwave sky in Figure 27.16. The temperature variation due to Earth’s motion (see Figure 26.18) has been subtracted out, as has the radio emission from the Milky Way, and temperature deviations from the average are displayed.

The ripples seen by *COBE*, combined with computer simulations such as that shown in Figure 27.15, predict present-day structure that is consistent with the superclusters, voids, filaments, and Great Walls we see around us. Although the *COBE* data were limited by relatively low (roughly  $7^\circ$ ) resolution, detailed analysis of the ripples also supports the key prediction of inflation theory—that the universe is of exactly critical density and hence spatially flat. For these reasons, the *COBE* observations rank alongside the discovery of the microwave background itself in terms of their importance to the field of cosmology. The lead investigators of the *COBE* program won the 2006 Nobel Prize in physics for their groundbreaking work.





Subsequent missions have radically improved our view of the microwave background, confirming and extending the *COBE* results. NASA’s *Wilkinson Microwave Anisotropy Probe (WMAP)* operated from 2001 until 2009. Its angular resolution was roughly  $20$ – $30'$ , some 20 times finer than that



◀ **FIGURE 27.16 Cosmic Microwave Background Map** This *COBE* map of temperature fluctuations in the cosmic microwave background over the entire sky shows hotter-than-average regions in yellow and cooler-than-average regions in blue. The total range of temperature fluctuations shown is extremely small, in fact only  $\pm 200$  millionths of a kelvin. The temperature variation due to Earth's motion has been subtracted, as has the radio emission from the Milky Way Galaxy. (NASA)

of *COBE*, allowing extraordinarily detailed measurements of many cosmological parameters to be made. More recently, the European Space Agency's *Planck* mission, launched in 2009, has refined *WMAP*'s observations by a further factor of 3 in resolution and 10 in sensitivity, making still more precise measurements of the microwave sky and largely confirming *WMAP*'s basic findings. Figure 27.17 shows an all-sky map of temperature fluctuations in the microwave background, based on the first year of *Planck* observations and released in 2013.

Unlike *COBE*, neither *WMAP* nor *Planck* orbited Earth. Rather, both were permanently stationed some 1.5 million km outside Earth's orbit along the Sun–Earth line, always pointing away from the Sun to keep their delicate heat-sensitive detectors in shadow and completing a scan of the entire sky every 6 months. The inset to Figure 27.17 shows a smaller scale (just  $2^\circ$  wide) but even higher resolution ( $7'$ ) image returned by *Cosmic Background Imager*, a ground-based microwave telescope located high in the Chilean Andes. (Recall from Chapter 3 that the microwave part of the



**Interactive FIGURE 27.17 Early Structure** The entire microwave sky, as seen by the *Planck* spacecraft at frequencies as high as 90 GHz (3-mm wavelength), can be compared directly with the lower-resolution *COBE* map shown in Figure 27.16. The inset at right shows an even higher resolution map of a small patch of sky, obtained by the ground-based *Cosmic Background Imager* instrument at 30 GHz (1-cm wavelength). The bright blobs are slightly denser-than-average regions of the universe at an age of roughly 400,000 years; they will eventually contract to form clusters of galaxies. (ESA; CBI)

spectrum is only partly transparent to electromagnetic radiation, so microwave detectors must be placed above as much of Earth's absorbing atmosphere as possible.)  $\infty$  (Sec. 3.3)


Both of these high-resolution maps show temperature fluctuations of a few hundred microkelvins, with a characteristic angular scale of about  $1^\circ$ . This temperature range is larger than the fluctuations seen by *COBE* because *COBE*'s low resolution effectively averaged the data over a large area of the sky, smearing out the peaks and troughs seen in the higher resolution observations.

The amount of structure evident in the data on different angular scales shows a peak at about  $1^\circ$ —pretty much what your eyes see in Figure 27.17. This scale is related to conditions in the early universe in a very important way: It corresponds to the maximum distance a sound wave can travel between the end of inflation and the time of decoupling. With reasonable assumptions, cosmologists can compute this distance, allowing direct comparison between observations and theory. In fact, although *WMAP* and *Planck* differ slightly in their detailed measurements of the cosmological parameters, the observed  $1^\circ$  fluctuations are in excellent agreement with the theoretical prediction for a universe with  $\Omega_0 = 1$ , having roughly 30 percent matter and 70 percent dark energy, as noted in Chapter 26 and earlier in this chapter. This strongly supports the implication of inflation that  $\Omega_0$  must be very close to 1, with a small margin of error.

More detailed analysis of the data provides a wealth of information on the history and composition of the universe. For example, before decoupling, although the radiation field was largely unaffected by dark matter, it was influenced slightly by the gravity of the growing dark clumps, resulting in a slight gravitational redshift that varied from place to place, depending on the dark-matter density. As a result, careful analysis of the signal allows astronomers to infer the dark-matter density at decoupling. The *WMAP* and *Planck* data are our primary sources for the cosmological parameters used in *More Precisely 24-1* and throughout this text.

## Matter Oscillations

Further analysis of the epoch of decoupling allows us to obtain additional important information about the universe. Consider in more detail what happens to a sound wave in the early (pre-decoupling) universe. Imagine a small region of space that is slightly denser than its surroundings. As we saw in Section 27.5, the dark matter in this overdense clump will one day grow to form a galaxy, but focus here on the normal matter and radiation in the same region. Because radiation and matter are strongly coupled, the radiation “pushes” on the normal matter, causing it to expand rapidly outward in a shell, as illustrated in Figure 27.18. The gravity of the dark matter clump




▲ **FIGURE 27.18 Acoustic Oscillation** This sketch depicts a two-dimensional rendering of a three-dimensional wave of normal matter pushed by radiation away from a clump of dark matter in the early universe. In reality, myriad such sound waves would have propagated like this all over the sky wherever there were dark-matter concentrations.

is far too weak to prevent the shell from escaping. Such interactions would have caused slightly overdense regions of space to oscillate, sending out waves much as when a pebble strikes the surface of a pond (see Figure 3.2)—like a bell ringing in space and time, its tone getting quieter and deeper as the universe expands. This is the origin of the cosmic sound waves discussed at the start of this section—the tongue-twisting technical name for which is **baryon acoustic oscillations**.

The shell continues to expand until the epoch of decoupling, at which time the push from the radiation stops and the shell stalls. Subsequently, the shell simply expands along with the rest of the universe. But because the shell itself represents a denser-than-average part of the universe, it too will tend to attract more matter, and it will eventually form galaxies of its own. The result is that *every* dark matter region that forms a galaxy or galaxy cluster is expected to have associated with it a secondary shell of galaxies. The radius of this shell is the distance sound can travel between the end of inflation and decoupling, which amounts to about 150 Mpc in present-day units (that is, adjusted for the expansion of the universe since decoupling).

The importance of this result is that this feature is imprinted on the galaxy distribution throughout the universe and at all redshifts. Figure 27.19 shows how such a ripple grows with the expansion of the universe; today, its radius would measure about 150 Mpc. If such features could be detected at different redshifts, they would constitute a new “standard yardstick,” telling us precisely the scale of the universe at different times in the past. As such,



◀ **FIGURE 27.19 Acoustic Remnants** The record of baryon acoustic oscillations (white circles) allow astronomers to retrace cosmic history. This simulation shows how the small density variations of the early universe (at left) grew to become the clusters, walls, and filaments seen in more recent times. (Z. Rostomian/SDSS)

it represents an alternative powerful means of probing the expansion of the universe, *independent* of supernova studies described in Chapter 26. ∞ (Sec. 26.5)

Of course, the universe is much more complicated than depicted artistically in Figures 27.18 and 27.19 (as can be seen by the observational mess in Figure 27.17). Every density fluctuation gave rise to a similar wave, and so every galaxy cluster should have a corresponding shell, with all the shells overlapping and mixed together on the sky. Nevertheless, astronomers can infer statistically the existence of these shells, and that has been a major goal of the *Sloan Digital Sky Survey*, which includes data from hundreds of thousands of galaxies. ∞ (Discovery 25-1) Preliminary results show that the separations of those

galaxies, again measured statistically, are very close to predictions of the acoustical process just described.

The first decade of the 21st century saw the basic parameters of the universe measured (even if not yet fully understood) to an accuracy only dreamed of just a few years ago. It now appears that the second decade is well on track to explore the nature of dark matter and dark energy with unprecedented precision.

### PROCESS OF SCIENCE Check


- ✓ What do observations of fluctuations in the microwave background tell us about the structure of the universe?

**The Big Question** How did the universe begin? Did it actually have an origin or has it existed forever? While no one knows the answers, for the first time in recorded history, human beings are using logic, rationality, and some very sophisticated (and expensive) experimental equipment to try to address these fundamental questions. It's hard to gauge when success might be achieved, but the very fact that scientists are now fully engaged in this quest illustrates the breathtaking scope of modern scientific inquiry.


# Chapter Review

## SUMMARY


1 At present, the universe is dominated by dark energy, which exceeds the matter density by more than a factor of 2. The densities of dark energy and matter both greatly exceed the equivalent mass density of radiation: The density of matter was much greater in the past, when the universe was smaller and the universe was **matter-dominated** (p. 691), a few billion years ago. However, because radiation is redshifted as the universe expands, the density of radiation at early times was greater still. Thus, the early universe was **radiation-dominated** (p. 691).




2 During the first few minutes after the Big Bang, matter was formed out of the primordial fireball by the process of **pair production** (p. 691). In the early universe, matter and radiation were linked by this process. Particles and forces “froze out” of the radiation background as the temperature fell below the threshold for creating them. The existence of matter today means that there must have been unequal amounts of matter and antimatter early on.




3 The physical state of the universe can be understood in terms of present-day physics back to about 10<sup>-43</sup> s after the Big Bang. Before that, the four fundamental forces of nature—gravity, electromagnetism, the strong force, and the weak force—were all indistinguishable. There is presently no theory that can describe these extreme conditions. As the universe expanded and its temperature dropped, the forces became distinct from one another. First gravity, then the strong force, and then the weak and electromagnetic forces separated out. Subsequently, nuclei, then atoms, and eventually large clumps of matter destined to become galaxies and stars formed as the universe continued to cool.




4 All of the hydrogen in the universe is primordial, formed from radiation as the universe expanded and cooled. Most of the helium observed in the universe today is also primordial, created by **primordial nucleosynthesis** (p. 697) in the early universe a few minutes after the Big Bang. Some deuterium was also formed at these early times, and it provides a sensitive indicator of the present density of the universe in the form of “normal” (as opposed to dark) matter. Studies of deuterium indicate that normal matter can account for at most 3 or 4 percent of the critical density. The remaining mass inferred from studies of clusters must then be made of dark matter, in the form of unknown particles formed at some very early epoch.




5 When the universe was about 1100 times smaller than it is today, the temperature became low enough for atoms to form. At that time, the (then-optical) background radiation **decoupled** (p. 699) from the matter. The universe became transparent. The photons that now make up the microwave background have been traveling freely through space ever since.




6 Very early on, the universe underwent a brief period of rapid expansion called the **epoch of inflation** (p. 702), during which the size of the cosmos increased by a huge factor—10<sup>50</sup> or more. The **horizon problem** (p. 700) is the fact that, according to the standard (that is, noninflationary) Big Bang model, there is no good reason for widely separated parts of the universe to be as similar as they are. Inflation solves the horizon problem by taking a small homogeneous patch of the early universe and expanding it enormously. The patch is still homogeneous, but it is now much larger than the portion of the universe we can see today. Inflation also solves the **flatness problem** (p. 701), which is the fact that there is no obvious reason why the present density of the universe is so close to the critical value. Inflation implies that the cosmic density is, for all practical purposes, exactly critical.



7 Large-scale structure in the universe formed when density fluctuations in the dark matter clumped and grew to form the “skeleton” of the structure we observe. Normal matter then flowed into the densest regions of space, eventually forming the galaxies we now see. Cosmologists distinguish between hot dark matter and **cold dark matter** (p. 706), depending on the temperature of dark matter at the end of the radiation era. In order to explain the observed large-scale structure in the universe, much of the dark matter must be cold.



8 “Ripples” in the microwave background are the imprint of early density inhomogeneities on the radiation field. These ripples were observed by the COBE satellite. Subsequent observations made by the WMAP spacecraft provided accurate measurements of many cosmological parameters and lend strong support to the inflationary prediction that we live in a flat, critical-density universe. Detailed observations of the microwave background, combined with studies of large-scale structure in the cosmos, provide precise information on the basic cosmological parameters of the universe.





**MasteringAstronomy**<sup>®</sup> For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- LO1** How long was the universe dominated by radiation? How hot was the universe when this period ended?
- What was the role of dark energy in the very early universe?
- LO2** How did matter “freeze out” of the early radiation field as the universe expanded?
- LO3** Describe how the relative importance of matter and radiation changed as the universe increased in size.
- LO4** When and how did the first helium nuclei form?
- Why do all stars, regardless of their abundance of heavy elements, seem to contain at least one-quarter helium by mass?
- Why didn't heavier and heavier elements form in the early universe, as they do in stars?
- POS** How do we know that most matter in the universe is not “normal”?
- When and how did the first atoms form?
- LO5** How can we observe the epoch at which the universe became transparent?
- LO6** What is the epoch of inflation, and what happened to the early universe during that time?
- POS** How does inflation solve the horizon and flatness problems?
- What does inflation tell us about the total density of the universe?
- LO7** What is the connection between dark matter and the formation of large- and small-scale structures?
- LOB POS** What key measurements were made by the *COBE* and *WMAP* experiments?

## Conceptual Self-Test: Multiple Choice

- Immediately after its birth, the universe (a) was dominated by photons; (b) was made mostly of protons; (c) had equal amounts of matter and antimatter; (d) formed stars and galaxies.
- Present-day Grand Unified Theories unite all of the fundamental forces except (a) the strong force; (b) the weak force; (c) the electromagnetic force; (d) the gravitational force.
- About half a million years after the Big Bang, the universe had cooled to the point that (a) protons and electrons could combine to form atoms; (b) particle–antiparticle annihilation ceased; (c) gas could condense to form stars; (d) carbon condensed to make dust.
- One of the problems with the standard Big Bang model is that (a) galaxies are redshifted; (b) the temperature is almost exactly the same everywhere; (c) the universe is hottest in the center; (d) the galaxy will expand forever.
- VIS** According to our best estimates, the line that best describes the universe in Figure 27.10 (“Flatness Problem”) is (a) accelerating; (b) open; (c) critical; (d) closed.
- It is likely that the density of the universe is made up mostly of (a) hydrogen; (b) electromagnetic radiation; (c) dark energy; (d) cold dark matter.
- The horizon problem in the standard Big Bang model is solved by having the universe (a) accelerate; (b) inflate rapidly early in its existence, (c) have tiny, but significant fluctuations in temperature; (d) be geometrically flat.
- The structure we observe in the universe is the result of (a) dark matter clumping long ago; (b) galaxies colliding; (c) the freezing out of electrons; (d) radiation dominance in the early universe.
- Elements more massive than lithium were not formed in the early universe because the temperature was (a) too high; (b) too low; (c) not related to density; (d) unstable.
- Matter and energy clumping in the early universe result in (a) the formation of atoms; (b) rapid inflation; (c) small but observable red shifts; (d) lower temperatures.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

- What was the distance between the points that would someday become, respectively, the center of the Milky Way Galaxy and the center of the Virgo Cluster at the time of decoupling? (The present separation is 18 Mpc.)
- What was the equivalent mass density of the cosmic radiation field when the universe was one-thousandth its present size? (Hint: Don't forget the cosmological redshift!)

3. ●● Of matter and radiation, which dominated the universe, and by what factor in density (assuming critical density today), at the start of (a) decoupling, (b) nucleosynthesis?
4. ●● Given that the threshold temperature for the production of electron–positron pairs is about  $6 \times 10^9$  K and that a proton is 1800 times more massive than an electron, calculate the threshold temperature for proton–antiproton pair production.
5. ● At what wavelength did the background radiation peak at the start of the epoch of nucleosynthesis? In what part of the electromagnetic spectrum does this wavelength lie?
6. ●● By what factor did the volume of the universe increase during the epoch of primordial nucleosynthesis, from the time when deuterium could first survive until the time at which all nuclear reactions ceased? By what factor did the matter density of the universe decrease during that period?
7. ● From Table 24.1, the “photosphere” of the universe corresponding to the epoch of decoupling presently lies some 14,000 Mpc from us. (See Figure 27.8.) How far away was a point on the photosphere when the background radiation we see today was emitted?
8. ●● The blobs evident in the inset to Figure 27.17 are about 20' across. If those blobs represent clumps of matter around the time of decoupling (redshift = 1100), estimate the diameter of the clumps at the time of decoupling, assuming Euclidean geometry.

## Activities

---

### Collaborative

1. Discuss the philosophical differences between living in an infinite universe that will expand forever and a closed, spatially finite universe that will some day recollapse. Are there aspects of any of these two possibilities that are hard to accept? It appears, given the current state of cosmological observations, that the former outcome is more likely. Our current models rest squarely on two quantities—dark matter and dark energy—whose nature remains largely unknown. How confident are you that our models of the universe are “set” and will not see further large shifts in the future?

### Individual

1. Go online and read about the *steady-state universe*, which enjoyed some measure of popularity in the 1950s and 1960s. How does it differ from the standard Big Bang model? Can you find any similarities between the steady-state model and our current view of the cosmos? Why do you think the steady-state model is not widely accepted today?



# Life in the Universe

## ARE WE ALONE?

Are we unique? Is life on our planet the only example of life in the universe? If so, then what might be the implications of such cosmic loneliness? If not, then how and where should we search for other intelligent beings? These are difficult questions since the subject of extraterrestrial life is one for which we have no data, yet they are important questions, with profound implications for the human species.

In this final chapter, we examine how humans evolved on Earth and then consider whether those evolutionary steps might have happened elsewhere. We then assess the likelihood of our having Galactic neighbors and consider how we might learn about them if they do exist.

**The Big Picture** Earth is the only place in the universe where we know for certain that life exists. Despite the likelihood of life elsewhere in the universe, we have no clear evidence for it. None of the hundreds of extrasolar planets discovered recently has yet shown any sign of life, intelligent or otherwise. Even so, astronomers keep watchful eyes on the sky, constantly aware that evidence for extraterrestrial intelligence (ETI) might emerge at any moment.

**LEFT:** This fanciful painting, entitled *Galaxyrise Over Alien Planet*, suggests a plurality of life—some perhaps extinct, some perhaps exotic—on alien worlds well beyond Earth. Despite blockbuster movies, science-fiction novels, and a host of claims for extraterrestrial contact, astronomers have so far found no unambiguous evidence for life of any kind anywhere else in the universe. (© Dana Berry)

### Learning Outcomes

*Studying this chapter will enable you to*

- 1 Summarize the process of cosmic evolution as it is currently understood.
- 2 Describe the basic ingredients of life on Earth.
- 3 Identify the most promising sites for life elsewhere in the solar system, and explain why they are promising.
- 4 Summarize the various probabilities used to estimate the number of advanced civilizations that might exist in the Galaxy.
- 5 Outline some techniques we might use to search for extraterrestrials and to communicate with them.

### MasteringAstronomy®

Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.

## 28.1 Cosmic Evolution

In our study of the universe, we have been very careful to avoid any inference or conclusion that places Earth in a special place in the cosmos. This Copernican principle, or principle of mediocrity, has been our invaluable guide in helping us define our place in the “big picture.” However, when discussing life in the universe, we face a problem: Ours is the only planet we know of on which life and intelligence have evolved, making it hard for any discussion of intelligent life *not* to treat humans as special cases.

Accordingly, in this final chapter we adopt a decidedly different approach. We first describe the chain of events leading to the only technologically proficient, intelligent civilization we know—us. Then we try to assess the likelihood of finding and communicating with intelligent life elsewhere in the cosmos.

### Life in the Universe

With this human-centered view clearly evident, Figure 28.1 identifies seven major evolutionary phases that have contributed to development of life on our planet: *particulate*, *galactic*, *stellar*, *planetary*, *chemical*, *biological*, and *cultural* evolution. Matter formed from energy in the early universe, then cooled and clumped to form galaxies and stars. Within galaxies, generation after generation of stars formed and died, seeding the interstellar medium with heavy elements so that, when our Sun formed billions of years after the first star blazed, the rocky planet Earth formed along with it. Eventually, on Earth, life appeared and slowly evolved into the diverse environment we see today.

Together, these evolutionary phases represent the grand sweep of **cosmic evolution**—the continuous transformation


of matter and energy that has led to the appearance of life and civilization on our planet. The first four represent, in reverse order, the contents of this book. We now expand our field of view beyond astronomy to include the other three.

From the Big Bang, to the formation of galaxies, to the birth of the solar system, to the emergence of life, to the evolution of intelligence and culture, the universe has evolved from simplicity to complexity. We are the result of an incredibly complex chain of events that spanned billions of years. Were those events random, making us unique, or are they in some sense *natural*, so that *technological civilization*—which, as a practical matter, we will take to mean “civilization capable of off-planet communication by electromagnetic or other means”—is inevitable? Put another way, are we alone in the universe, or are we just one among countless other intelligent life-forms in our Galaxy?

Before trying to answer these important questions, we need a working definition of *life*. Defining life, however, is not an easy task: The distinction between the living and the nonliving is not as obvious as we might at first think. Although most physicists would agree on the definitions of matter and energy, biologists have not arrived at a clear-cut definition of life. Generally speaking, scientists regard the following as characteristics of living organisms: (1) They can *react* to their environment and can often heal themselves when damaged; (2) they can *grow* by taking in nourishment from their surroundings and processing it into energy; (3) they can *reproduce*, passing along some of their own characteristics to their offspring; and (4) they have the capacity for genetic change and can therefore *evolve* from generation to generation and adapt to a changing environment.

These rules are not strict, and there is great leeway in interpreting them. Stars, for example, react to the gravity of their neighbors, grow by accretion, generate energy, and

▼ **FIGURE 28.1 Arrow of Time** Some highlights of cosmic history, as it relates to the emergence of life on Earth, are noted along this arrow of time, from the beginning of the universe to the present. At the bottom of the arrow are seven “windows” outlining the major phases of cosmic evolution: evolution of primal energy into elementary particles; of atoms into galaxies and stars; of stars into heavy elements; of elements into solid, rocky planets; of those elements into the molecular building blocks of life; of those molecules into life itself; and of advanced life forms into intelligence, culture, and technological civilization. (*D. Berry*)



## DISCOVERY 28-1

### The Virus


The central idea of chemical evolution is that life evolved from nonliving molecules. But aside from insight based on biochemical knowledge and laboratory simulations of some key events on primordial Earth, do we have any direct evidence that life could have developed from nonliving molecules? The answer is yes.

The smallest and simplest entity that sometimes appears to be alive is a virus. We say “sometimes” because viruses seem to have the attributes of both nonliving molecules and living cells. *Virus* is the Latin word for “poison,” an appropriate name, since viruses often cause disease. Although they come in many sizes and shapes—a representative example is the polio virus, shown here magnified 300,000 times—all viruses are smaller than the size of a typical modern cell. Some are made of only a few thousand atoms. In terms of size, then, viruses seem to bridge the gap between cells that are living and molecules that are not.


Viruses contain some proteins and genetic information (in the form of DNA or the closely related molecule RNA, the two molecules responsible for transmitting genetic characteristics from one generation to the next), but not much else—none of the material by which living organisms normally grow and reproduce. How, then, can a virus be considered alive? Indeed, alone, it cannot; a virus is absolutely lifeless when it is isolated from living organisms. But when it is inside a living system, a virus has all the properties of life.

Viruses come alive by transferring their genetic material into living cells. The genes of a virus seize control of a cell and establish themselves as the new master of chemical activity.

Viruses grow and reproduce copies of themselves by using the genetic machinery of the invaded cell, often robbing the cell of its usual function. Some viruses multiply rapidly and wildly, spreading disease and—if unchecked—eventually killing the invaded organism. In a sense, then, viruses exist within the gray area between the living and the nonliving.



(R. Williams)



“reproduce” by triggering the formation of new stars, but no one would suggest that they are alive. By contrast, a virus (see *Discovery 28-1*) is inert when isolated from living organisms, but once inside a living system, it exhibits all the properties of life, seizing control of a living cell and using the cell’s own genetic machinery to grow and reproduce. Most researchers now think that the distinction between living and nonliving matter is more one of structure and complexity than a simple checklist of rules.

The general case in favor of extraterrestrial life is summed up in what are sometimes called the *assumptions of mediocrity*: (1) Because life on Earth depends on just a few basic molecules, and (2) because the elements that make up these molecules are (to a greater or lesser extent) common to all stars, and (3) if the laws of science we know apply to the entire universe, as we have supposed throughout this book, then—given sufficient time—life must have originated elsewhere in the cosmos. The opposing view maintains that intelligent life on Earth is the product of a series of extremely fortunate accidents—astronomical, geological, chemical, and biological events unlikely to


have occurred anywhere else in the universe. The purpose of this chapter is to examine some of the arguments for each of these viewpoints.

### Chemical Evolution


What information do we have about the earliest stages of planet Earth? Unfortunately, not very much. Geological hints about the first billion years or so were largely erased by violent surface activity, as volcanoes erupted and meteorites bombarded our planet; subsequent erosion by wind and water has seen to it that little evidence of that era has survived to the present. Scientists think that the early Earth was barren, with shallow, lifeless seas washing upon grassless, treeless continents. Gases emanating from our planet’s interior through volcanoes, fissures, and geysers produced an atmosphere rich in hydrogen, nitrogen, and carbon compounds and poor in free oxygen. As Earth cooled, ammonia, methane, carbon dioxide, and water formed. The stage was set for the appearance of life.

The surface of the young Earth was a very violent place. Natural radioactivity, lightning, volcanism, solar ultraviolet radiation, and meteoritic impacts all provided large amounts of energy that eventually shaped the ammonia, methane, carbon dioxide, and water on our planet into more complex molecules known as **amino acids** and **nucleotide bases**—organic (carbon-based) molecules that are the building blocks of life as we know it. Amino acids build *proteins*, and proteins control metabolism, the daily utilization of food and energy by means of which organisms stay alive and carry out their vital activities. Sequences of nucleotide bases form *genes*—parts of the DNA molecule—which direct the synthesis of proteins and thus determine the characteristics of the organism (Figure 28.2). These same genes, via the DNA contained within every cell in the organism, transfer hereditary characteristics from one generation to the next through reproduction. In all living creatures on Earth—from bacteria to amoebas to humans—genes mastermind life and proteins maintain it.

The idea that complex molecules could have evolved naturally from simpler ingredients found on the primitive Earth has been around since the 1920s. The first experimental verification was provided in 1953, when scientists Harold Urey and Stanley Miller, using laboratory equipment somewhat similar to that shown in Figure 28.3. The **Urey-Miller experiment** took a mixture of the materials thought to be present on Earth long ago—a “primordial soup” of water,



▲ **FIGURE 28.2 DNA Molecule** DNA (deoxyribonucleic acid) is the molecule containing all the genetic information needed for a living organism to reproduce and survive. Often consisting of literally tens of billions of individual atoms, its double-helix structure allows it to “unzip,” exposing its internal structure to control the creation of proteins needed for a cell to function. The ordering of its constituent parts is unique to each individual organism.




▲ **FIGURE 28.3 Urey-Miller Experiment** This chemical apparatus is designed to synthesize complex biochemical molecules by energizing a mixture of simple chemicals. Gases (ammonia, methane, carbon dioxide, and water vapor) are placed in the upper bulb to simulate Earth’s primordial atmosphere and then zapped by spark-discharge electrodes akin to lightning. After about a week, amino acids and other complex molecules emerge in the trap at the bottom, which simulates the primordial oceans into which heavy molecules produced in the overlying atmosphere would have fallen.

methane, carbon dioxide, and ammonia—and energized it by passing an electrical discharge (“lightning”) through the gas. After a few days, they analyzed their mixture and found that it contained many of the same amino acids found in all living things on Earth. About a decade later, scientists succeeded in constructing nucleotide bases in a similar manner. These experiments have been repeated in many different forms, with more realistic mixtures of gases and a variety of energy sources, but always with the same basic outcomes.

Although none of these experiments has ever produced a living organism, or even a single strand of DNA, they do demonstrate conclusively that “biological” molecules—the molecules involved in the functioning of living organisms—can be synthesized by strictly *nonbiological* means, using raw materials available on the early Earth. More advanced experiments, in which amino acids are united under the influence of heat, have fashioned protein-like blobs that behave to some extent like true biological cells. Such near-protein material resists dissolution in water (so it would remain intact when it fell from the primitive atmosphere into the ocean) and tends to cluster into small droplets called microspheres—a little like oil globules floating on the surface of water. Figure 28.4 shows some of these laboratory-made proteinlike microspheres, whose walls permit the inward

These three photographs, taken through a microscope, show structures on the scale of 1 micrometer, which equals 1/10,000 of a centimeter.



**▲ FIGURE 28.4 Chemical Evolution** These carbon-rich, protein-like droplets contain as many as a billion amino acid molecules in a liquid sphere. The droplets can “grow,” and parts of them can separate from their “parent” droplet to become new individual droplets (as at A, B, and C). (S. Fox)

passage of small molecules, which then combine within the droplet to construct more complex molecules that are too large to pass back out through the walls. As the droplets “grow,” they tend to “reproduce,” forming smaller droplets.

Can we consider these protein-like microspheres to be alive? Almost certainly not. Most biochemists would say that the microspheres are not life itself, but they contain many of the basic ingredients needed to form life. The microspheres lack the hereditary DNA molecule. However, as illustrated in Figure 28.5, they do have similarities to ancient cells found in the fossil record, which in turn have many similarities to modern organisms (such as blue-green algae). Thus, while no actual living cells have yet been created “from scratch” in any laboratory, many biochemists feel that the chain of events leading from simple nonbiological molecules almost to the point of life itself has been amply demonstrated.


## An Interstellar Origin?

Recently, a dissenting view has emerged. Some scientists have argued that Earth’s primitive atmosphere might *not* in fact have been a particularly suitable environment for the production of complex molecules. These scientists say that there may not have been sufficient energy available to power the necessary chemical reactions and that the early atmosphere may

not have contained enough raw material for the reactions to have become important in any case. They suggest instead that much, if not all, of the organic (carbon-based) material that combined to form the first living cells was produced in *interstellar space* and subsequently arrived on Earth in the form of comets, interplanetary dust, and meteors that did not burn up during their descent through the atmosphere.


Several pieces of evidence support this idea. Interstellar molecular clouds are known to contain complex molecules—indeed, there have even been reports (still unconfirmed) of at least one amino acid (glycine) in interstellar space. [∞ \(Sec. 18.5\)](#)

To test the interstellar space hypothesis, NASA researchers have carried out their own version of the Urey-Miller experiment in which they exposed an icy mixture of water, methanol, ammonia, and carbon monoxide—representative of many interstellar grains—to ultraviolet radiation to simulate the energy from a nearby newborn star. As shown in Figure 28.6, when they later placed the irradiated ice in water and examined the results, they found that the ice had formed droplets surrounded by membranes and containing complex organic molecules. As with the droplets found in earlier experiments, no amino acids, proteins, or DNA were observed in the mix, but the results, repeated numerous times, clearly show that even the harsh, cold vacuum of interstellar space can be a suitable medium in which complex molecules and primitive cellular structures can form.



**▲ FIGURE 28.5 Primitive Cells** (a) This photograph shows primitive fossils that display concentric spheres or walls connected by smaller spheroids. They were found in sediments radioactively dated to be about 2 billion years old. (b) For comparison, modern blue-green algae found in a backyard stream are shown on approximately the same scale. (E. Barghoorn)






▲ **FIGURE 28.6 Interstellar Globules** These oily, hollow droplets rich in organic molecules were made by exposing a mixture of primordial matter to harsh ultraviolet radiation. When immersed in water, the larger ones display cell-like membrane structure. Although they are not alive, they bolster the idea that life on Earth could have come from space. (*NASA*)

As described in Chapter 15, these icy interstellar grains are thought to have formed the comets in our own solar system. ∞ (Sec. 15.3) Large amounts of organic material were detected on comet Halley by space probes when Halley last visited the inner solar system, and similarly complex molecules have been observed on many other well-studied comets, such as Hale-Bopp. ∞ (Sec. 14.2) Also as discussed in Chapter 15, there is reason to suspect that cometary impacts were responsible for most of Earth's water, and it is perhaps a small step to imagining that this water already contained the building blocks for life. ∞ (Sec. 15.3)

In addition, a small fraction of the meteorites that survive the plunge to Earth's surface—including perhaps the controversial “Martian meteorite” discussed in Chapter 10—contain organic compounds. ∞ (*Discovery 10-1*) The Murchison meteorite (Figure 28.7), which fell near



▲ **FIGURE 28.7 Murchison Meteorite** The Murchison meteorite contains relatively large amounts of amino acids and other organic material, indicating that chemical evolution of some sort occurred beyond our own planet. In this magnified view of a fragment from the meteorite, an arrow points to a microscopic sphere of organic matter. (*Harvard-Smithsonian CfA*)

Murchison, Australia, in 1969, is a particularly well-studied example. Located soon after crashing to the ground, this meteorite has been shown to contain 12 of the amino acids normally found in living cells, although the detailed structures of these molecules indicate potentially important differences between those found in space and those found on Earth. At the very least, though, these discoveries argue that complex molecules can form in an interplanetary or interstellar environment and that they could have reached Earth's surface unscathed after their fiery descent.

Thus, the hypothesis that organic matter is constantly raining down on Earth from space in the form of interplanetary debris is quite plausible. However, whether this was the *primary* means by which complex molecules first appeared in Earth's oceans remains unclear.

## Diversity and Culture

However the basic materials appeared on Earth, we know that life *did* appear. The fossil record chronicles how life on Earth became widespread and diversified over the course of



time. The study of fossil remains shows the initial appearance of simple one-celled organisms such as blue-green algae more than 3.5 billion years ago. These were followed by more complex one-celled creatures, such as the amoeba, about 2 billion years ago. Multicellular organisms such as sponges did not appear until about 1 billion years ago, after which there flourished a wide variety of increasingly complex organisms—insects, reptiles, and mammals. Figure 28.8 illustrates some of the key developments in the evolution of life on our planet.

The fossil record leaves no doubt that biological organisms have changed over time—all scientists accept the reality of *biological evolution*. As conditions on Earth shifted and Earth's surface evolved, those organisms that could best take advantage of their new surroundings succeeded and thrived—often at the expense of organisms that could not make the necessary adjustments and consequently became extinct.

What led to these changes? Chance. An organism that happened to have a certain useful genetically determined trait—for example, the ability to run faster, climb higher, or even hide more easily—would find itself with the upper hand in a particular environment. That organism was therefore more likely to reproduce successfully, and its advantageous characteristic would then be more likely to be passed on to the next generation. The evolution of the rich variety of life on our planet, including human beings, occurred as chance mutations—changes in genetic structure—led to changes in organisms over millions of years.


What about the development of intelligence? Many anthropologists think that, like any other highly advantageous trait, intelligence *is* strongly favored by natural selection. As humans learned about fire, tools, and agriculture, the brain became more and more elaborate. The social cooperation that went with coordinated hunting efforts was

another important competitive advantage that developed as brain size increased.

Perhaps most important of all was the development of language. Indeed, some anthropologists have gone so far as to suggest that human intelligence *is* human language. Through language, individuals could signal one another while hunting for food or seeking protection. Even more importantly, now our ancestors could share ideas as well as food and shelter. Experience, stored in the brain as memory, could be passed down from generation to generation. A new kind of evolution had begun, namely, *cultural evolution*, the changes in the ideas and behavior of society. Within only the past 10,000 years or so, our more recent ancestors have created the entirety of human civilization.

To put all this into historical perspective, let's imagine the entire lifetime of Earth to be 46 years rather than 4.6 billion years. On this scale, we have no reliable record of the first decade of our planet's existence. Life originated at least 35 years ago, when Earth was about 10 years old. Our planet's middle age is largely a mystery, although we can be sure that life continued to evolve and that generations of mountain chains and oceanic trenches came and went. Not until about 6 years ago did abundant life flourish throughout Earth's oceans. Life came ashore about 4 years ago, and plants and animals mastered the land only about 2 years ago. Dinosaurs reached their peak about 1 year ago, only to die suddenly about 4 months later. [\(Discovery 11-1\)](#) Humanlike apes changed into apelike humans only last week, and the latest ice ages occurred only a few days ago. *Homo sapiens*—our species—did not emerge until about 4 hours ago. Agriculture was invented within the last hour, and the Renaissance—along with all of modern science—is just 3 minutes old!

▼ **FIGURE 28.8 Life on Earth** This simplified timeline of the origin and evolution of life on our planet begins at the far left with the origin of Earth about 4.6 billion years ago and extends linearly to the present day at right. Notice how life forms most familiar to us—fish, reptiles, mammals—emerged relatively recently in the history of our planet. Technological civilization has existed on Earth for just a few millionths of 1 percent of our planet's lifetime.



**CONCEPT Check**

✓ Has chemical evolution been verified in the laboratory?

**28.2 Life in the Solar System**

Simple one-celled life-forms reigned supreme on Earth for most of our planet's history. It took time—a great deal of time—for life to emerge from the oceans, to evolve into simple plants, to continue to evolve into complex animals, and to develop intelligence, culture, and technology. Have those (or similar) events occurred elsewhere in the universe? Let's try to assess what little evidence we have on the subject.

**Life as We Know It**

“Life as we know it” is generally taken to mean carbon-based life that originated in a liquid water environment—in other words, life on Earth. Might such life exist elsewhere in our solar system?

The Moon and Mercury lack liquid water, protective atmospheres, and magnetic fields, so these two bodies are subjected to fierce bombardment by solar ultraviolet radiation, the solar wind, meteoroids, and cosmic rays. Simple molecules could not survive in such hostile environments. Venus, by contrast, has *far too much* protective atmosphere! Its dense, dry, scorchingly hot atmospheric blanket effectively rules it out as an abode for life, at least like us.

The jovian planets have no solid surfaces (although some researchers have suggested that life might have evolved in their atmospheres), and most of their moons (apart from volcanic Io) have frozen surfaces far too cold to support Earth-like life. One possible exception is Saturn's moon Titan. With its thick atmosphere of methane, ammonia, and nitrogen gases; liquid hydrocarbon lakes; and apparent geological activity, Titan might conceivably be a place where surface life could have arisen, although the most recent results from the *Cassini-Huygens* mission suggest that the environment there would be hostile to anything remotely familiar to us. ∞ (Sec. 12.5)

A more promising scenario comes from the discovery that four jovian moons—Jupiter's Europa and Ganymede and Saturn's Titan and Enceladus—may contain significant amounts of liquid water in their interiors. ∞ (Sec. 11.5) This possibility has fueled speculation about the development of life within these bodies, making them prime candidates for future exploration. Europa in particular is high on the priority lists of both NASA and the European Space Agency. Again, conditions in or on these moons are far from ideal by Earthly standards, but, as we discuss below, scientists are finding more and more examples of terrestrial organisms that thrive in extreme environments once regarded as uninhabitable.


The planet most likely to harbor life (or, more likely, to have harbored it in the past) still seems to be Mars. The Red

Planet is harsh by Earth standards: Liquid water is scarce, the atmosphere is thin, and the absence of magnetism and an ozone layer allows solar high-energy particles and ultraviolet radiation to reach the surface unabated. But the Martian atmosphere was thicker, and the surface probably warmer and much wetter, in the past. ∞ (Sec. 10.5) Indeed, there is strong photographic evidence from orbiters such as *Viking* and *Mars Global Surveyor* for flowing and standing water on Mars in the distant (and perhaps even relatively recent) past. In 2004, the European *Mars Express* orbiter confirmed the long-hypothesized presence of water ice at the Martian poles, and NASA's *Opportunity* rover reported strong geological evidence that the region around its landing site was once “drenched” with water for an extended period.


All of these lines of reasoning strongly suggest that Mars—at least at some time in its past—harbored liquid water. However, none of the Mars landers has detected anything that might be interpreted as the remains (fossilized or otherwise) of large plants or animals, and only the *Viking* and *Curiosity* landers have carried equipment capable of performing the detailed biological analysis needed to detect bacterial life (or its fossil remnants). The *Viking* robots scooped up Martian soil and tested it for the presence of life by conducting chemical experiments designed to detect the waste gases and other products of metabolic activity, but no unambiguous evidence of Martian life has emerged. ∞ (Discovery 10-1) The *Curiosity* rover that landed in 2012 (Figure 28.9) is now at work near what resembles ancient, dry lake beds, but so far no Martian life, dead or alive, has been detected.

Some scientists have suggested that a different type of biology may be operating on the Martian surface. They propose that Martian microbes capable of eating and digesting oxygen-rich compounds in the Martian soil could also explain the *Viking* results. This speculation would be greatly strengthened if recent announcements of fossilized bacteria in meteorites originating on Mars were confirmed (although it seems that the weight of scientific opinion is currently running against that interpretation of the data). The consensus among biologists and chemists today is that Mars does not house any life similar to that on Earth, but a solid verdict regarding past life on Mars will not be reached until we have thoroughly explored our intriguing neighbor.

In considering the emergence of life under adversity, we should perhaps not be too quick to rule out an environment based solely on its extreme properties. Figure 28.10 shows a very hostile environment on a deep-ocean floor, where hydrothermal vents spew forth boiling hot water from vertical tubes a few meters tall. The conditions are quite unlike anything on our planet's surface, yet life thrives in an environment rich in sulfur, poor in oxygen, and completely dark. Such underground hot springs might conceivably exist on alien worlds, raising the possibility of



▲ **FIGURE 28.9 Search for Martian Life** Samples of Martian dirt were collected and examined by the *Curiosity* robot, seen partially at left while experimenting in a shallow depression called “Yellowknife Bay” near the equator. No signs of life have yet been found. (NASA)



life-forms with much greater diversity over a much wider range of conditions than those known to us on Earth.

In recent years scientists have discovered many instances of so-called **extremophiles**—life-forms that have adapted to live in extreme environments. The superheated hydrothermal vents in Figure 28.10 are one example, but extremophiles have also been found in frigid lakes buried deep under the Antarctic glaciers, in the dark, oxygen-poor and salt-rich floor of the Mediterranean Sea, in the mineral-rich superalkaline environment of California’s Mono lake, and even in the hydrogen-rich volcanic darkness far below Earth’s crust. In many cases these

◀ **FIGURE 28.10 Hydrothermal Vents** A two-person submarine (the *Alvin*, partly seen at bottom) took this picture of a hot spring, or “black smoker”—one of many along the midocean ridge in the eastern Pacific Ocean. As hot water rich in sulfur pours out of the top of the vent’s tube (near center), black clouds billow forth, providing a strange environment for many life-forms thriving near the vent. The inset shows a close-up of the vent base, where extremophilic life thrives, including, as seen here, giant red tube worms and huge crabs. (WHOI)

organisms have evolved to create the energy they need by purely chemical means, using *chemosynthesis* instead of *photosynthesis*, the process whereby plants turn sunlight into energy. These environments may present conditions not so different from those found on Mars, Europa, or Titan, suggesting that even “life as we know it” might well be able to thrive in these hostile, alien worlds.

## Alternative Biochemistries

Conceivably, some types of biology might be so different from life on Earth that we would not recognize them and would not know how to test for them. What might these other biologies be?

Some scientists have pointed out that the abundant element silicon has chemical properties somewhat similar to those of carbon and have suggested silicon as a possible alternative to carbon as the basis for living organisms. Ammonia (made of the common elements hydrogen and nitrogen) is sometimes put forward as a possible liquid medium in which life might develop, at least on a planet cold enough for ammonia to exist in the liquid state. Together or separately, these alternatives would surely give rise to organisms with biochemistries (the basic biological and chemical processes responsible for life) radically different from those we know on Earth. Conceivably, we might have difficulty even identifying these organisms as alive.

Although the possibility of such alien life-forms is a fascinating scientific problem, most biologists would argue that chemistry based on carbon and water is the one most likely to give rise to life. Carbon’s flexible chemistry and water’s wide liquid temperature range are just what are needed for life to develop and thrive. Silicon and ammonia seem unlikely to fare as well as bases for advanced life-forms. Silicon’s chemical bonds are weaker than those of carbon and may not be able to form complex molecules—an apparently essential aspect of carbon-based life. Also, the colder the environment, the less energy there is to drive biological processes. The low temperatures necessary for ammonia to remain liquid might inhibit or even completely prevent the chemical reactions leading to the equivalent of amino acids and nucleotide bases.

Still, we must admit that we know next to nothing about noncarbon, nonwater biochemistries, for the very good

reason that there are no examples of them to study experimentally. We can speculate about alien life-forms and try to make general statements about their characteristics, but we can say little of substance about them.

### CONCEPT Check

✓ Which solar system bodies (other than Earth) are the leading candidates in the search for extraterrestrial life?

## 28.3 Intelligent Life in the Galaxy


With humans apparently the only intelligent life in the solar system, we must broaden our search for extraterrestrial intelligence to other stars and perhaps even other galaxies. At such distances, though, we have little hope of actually detecting life with current equipment. Instead, we must ask, “How likely is it that life in any form—carbon based, silicon based, water based, ammonia based, or something we cannot even dream of—exists?” Let’s look at some numbers to develop estimates of the probability of life elsewhere in the universe.

### The Drake Equation

An early approach to this problem is known as the **Drake equation**, after the U.S. astronomer who pioneered the analysis (below). It attempts to express the probability of life in our Galaxy in terms of specific factors with roots in astronomy, biology, and anthropology.

Of course, several of the factors in this formula are largely a matter of opinion. We do not have nearly enough information to determine—even approximately—every factor in the equation, so the Drake equation cannot give us a hard-and-fast answer. Its real value is that it subdivides a large and difficult question into smaller pieces that we can attempt to answer separately. The equation provides the framework within which the problem can be addressed and parcels out the responsibility for the final solution among many different scientific disciplines. Figure 28.11 illustrates how, as our requirements become more and more stringent, only a small fraction of star systems in the Milky Way is likely to generate

$$\begin{array}{cccccccc}
 \text{number of} & \text{rate of star} & & \text{average} & & \text{fraction} & & \text{fraction} \\
 \text{technological,} & \text{formation,} & \text{fraction} & \text{number of} & \text{fraction} & \text{of those} & \text{fraction} & \text{of those} \\
 \text{intelligent} & \text{averaged} & \times & \text{habitable} & \times & \text{of those} & \times & \text{of those} \\
 \text{civilizations} & \text{over the} & \times & \text{planets} & \times & \text{habitable} & \times & \text{intelligent-} \\
 \text{now present in} & \text{lifetime} & \times & \text{within} & \times & \text{planets on} & \times & \text{life planets} \\
 \text{the Galaxy} & \text{of the} & \times & \text{those} & \times & \text{which life} & \times & \text{that develop} \\
 & \text{Galaxy} & \times & \text{planetary} & \times & \text{arises} & \times & \text{technological} \\
 & & \times & \text{systems} & \times & & \times & \text{society} \\
 & & \times & & \times & & \times & \text{average} \\
 & & \times & & \times & & \times & \text{lifetime of a} \\
 & & \times & & \times & & \times & \text{technologically} \\
 & & \times & & \times & & \times & \text{competent} \\
 & & \times & & \times & & \times & \text{civilization.}
 \end{array}$$



**Narrated** **FIGURE 28.11 Drake Equation** Of all the star systems in our Milky Way Galaxy, progressively fewer and fewer have each of the qualities typical of a long-lasting technological society.

the advanced qualities specified by the combination of factors on the right-hand side of the equation.

Let's examine the factors in the equation one by one and make some educated guesses about their values. Bear in mind, though, that if you ask two scientists for their best estimates of any given factor, you will likely get two very different answers!

## Rate of Star Formation

We can estimate the average number of stars forming each year in the Galaxy simply by noting that at least 100 billion stars now shine in the Milky Way. Dividing this number by the 10-billion-year lifetime of the Galaxy, we obtain a formation rate of 10 stars per year. This rate may be an overestimate, because we think that fewer stars are forming now than formed at earlier epochs of the Galaxy, when more interstellar gas was available. However, we do know that stars are forming today, and our estimate does not include stars that formed in the past and have since died, so our value of 10 stars per year is probably reasonable when averaged over the lifetime of the Milky Way.

## Fraction of Stars Having Planetary Systems

Many astronomers regard planet formation as a natural result of the star-formation process. If the condensation theory (Chapter 16) or some variant of it is correct, and if there is nothing special about our Sun, as we have argued throughout this book, then we would expect many stars to have at least one planet. (Secs. 6.7, 15.5) Indeed, as we

have seen, increasingly sophisticated observations indicate the presence of disks around young stars. Could these disks be protosolar systems? The condensation theory suggests that they are, and the short (theoretical) lifetimes of disks imply the existence of many planet-forming systems in the neighborhood of the Sun.

As observational techniques have improved over the past two decades, these expectations have been borne out, and there is now overwhelming evidence for planets orbiting hundreds of other stars. The first planets discovered were much larger than Earth, and mostly moved on eccentric or “hot” orbits, but as we saw in Chapter 15, these were the only planets that could have been detected with the instruments available at the time. (Sec. 15.4) However, as detection technology has advanced, more and more planets with masses comparable to Earth have been discovered, to the point that, today, several dozen Earth-sized planets, many on roughly Earth-like orbits, have been confirmed. (Sec. 15.5) These observations are at the very edge of current capabilities, and many astronomers expect the numbers of “Earth-like” planets to grow rapidly as new detectors come on line.


Only about 10 percent of the nearby stars surveyed to date have been found to have planets. However, most researchers think that this is a significant underestimate of the true fraction because of observational limitations and selection biases. (Sec. 15.4) Thus, accepting the condensation theory and its consequences, and without being either too conservative or naively optimistic, we assign a value near unity to this factor—that is, we think that essentially all stars form with planetary systems of some sort.

## Number of Habitable Planets per Planetary System

What determines the feasibility of life on a given planet? *Temperature* is perhaps the single most important factor, although the possibility of catastrophic external events, such as cometary impacts or even distant supernovae, must also be considered. (Discovery 11-1, Sec. 21.3)

The surface temperature of a planet depends on two things: the planet's distance from its parent star and the thickness of the planet's atmosphere. Planets with a nearby parent star (but not too close) and some atmosphere (though not too thick) should be reasonably warm, like Earth or Mars. Planets far from the star and with no atmosphere, like Pluto, will surely be cold by our standards. And planets too close to the star and with a thick atmosphere, like Venus, will be very hot indeed.

As discussed in Chapter 15, a three-dimensional *stellar habitable zone* of “comfortable” temperatures surrounds every star. (Sec. 15.7) (The zones are indicated as rings in our two-dimensional figure, Figure 28.12.) The habitable zone represents the range of distances within which a planet



**◀ FIGURE 28.12 Stellar Habitable Zones** Hot stars have bigger habitable zones than do cool ones. For a G-type star like the Sun, the zone extends from about 0.8 AU to 2 AU. For a hotter F-type star, the range is 1.2 to 2.8 AU. For a cool M-type star, only Earth-like planets orbiting between about 0.02 and 0.06 AU would be habitable.

of mass and composition similar to Earth’s would have a surface temperature between the freezing and boiling points of water. (Our Earth-based bias is again clear here!) The hotter the star, the larger is this zone (Figure 28.12). A- and F-type stars have rather large habitable zones, but the size of the zone diminishes rapidly as we proceed through G-, K-, and M-type stars (although, as we saw in Chapter 15, numerous Earths and super-Earths do in fact lie within the habitable zones of their low-mass parent stars). ∞ (Sec. 15.7)


In addition to their small habitable zones, lightweight M-type stars are thought to be prone to such violent surface activity that they are generally not considered likely hosts of life-bearing planets, despite their large numbers. ∞ (Sec. 17.8) At the other extreme, massive O- and B-type stars are also considered unlikely candidates, both because they are rare and because they are not expected to last long enough for life to develop, even if they do have planets.

Three planets—Venus, Earth, and Mars—reside in or near the habitable zone surrounding our Sun. Venus is too hot because of its thick atmosphere and proximity to the Sun. Mars is a little too cold because its atmosphere is too thin and it is too far from the Sun. But if the orbits of Venus and Mars were swapped—not inconceivable, since chance played such a large role in the formation of the terrestrial planets—then both of these nearby planets might conceivably have evolved surface conditions resembling those on Earth. ∞ (Secs. 10.5, 15.4) In that case our solar system would have had three habitable planets instead of one. Perhaps just as important, proximity to a giant planet may also render the interior of a moon (such as Europa or Titan) habitable, the planet’s tidal heating making up for the lack of sunlight. ∞ (Sec. 11.5) Sheltered by its parent planet’s gravity, such a moon might be largely immune to the habitable limitations just described for planets.

A planet moving on a “habitable” orbit may still be rendered uninhabitable by external events. Many scientists think that the outer planets in our own solar system are critical to the habitability of the inner worlds, both by stabilizing their orbits and by protecting them from cometary impacts, deflecting would-be impactors away from the inner part of the solar system. The theory presented in Chapter 15 suggests that a star with inner terrestrial planets on stable orbits would probably also have the jovian worlds needed to safeguard their survival. ∞ (Sec. 15.2) However, observations of extrasolar planets are not yet sufficiently refined to determine the fraction of stars having “outer planet” systems like our own. ∞ (Sec. 15.7)

Other external forces may also influence a planet’s survival. Some researchers have suggested that there is a *galactic habitable zone* for stars in general, outside of which conditions are unfavorable for life (see Figure 28.13). Far from the Galactic center, the star formation rate is low and few cycles of star formation have occurred, so there are insufficient heavy elements to form terrestrial planets or populate them with technological civilizations if any should form. ∞ (Sec. 21.5) Too close, and the radiation from bright stars and supernovae in the crowded inner part of the Galaxy might be detrimental to life. More importantly, the gravitational effects of nearby stars may send frequent showers of comets from the counterpart of the Oort cloud into the inner regions of a planetary system, striking the terrestrial planets and terminating any chain of evolution that might lead to intelligent life.

Thus, to estimate the number of habitable planets per planetary system, we must first take inventory of how many stars of each type shine in the Galactic habitable zone, then calculate the sizes of their stellar habitable zones and estimate the number of planets likely to be found there. In doing so,



◀ **FIGURE 28.13 Galactic Habitable Zone** Some regions of the Galaxy may be more conducive to life than others. Too far from the Galactic center, there may not be enough heavy elements for terrestrial planets to form or technological society to evolve. Too close, the radiative or gravitational effects of nearby stars may render life impossible. The result is a ring-shaped habitable zone, colored here in green, although its full extent is uncertain.

fact is that we still have insufficient data about most stars to make any strong statement about habitable worlds in their planetary systems.

Taking the many uncertainties into account as best we can, we assign a value of 1/10 to this factor in our equation. In other words, we think that, on average, there is 1 potentially habitable planet for every 10 planetary systems that might exist in our Galaxy. Single F-, G-, and K-type stars are the best candidates.


### Fraction of Habitable Planets on Which Life Actually Arises

we eliminate almost all of the stars around which planets have so far been observed, and presumably a similar fraction of stars in general. The large jovian planets seen in most cases have eccentric orbits that would destabilize the motion of any inner terrestrial world, either ejecting it completely from the system or making conditions so extreme that the chances for the development of life are severely reduced. ∞ (Sec. 15.6) We also exclude the majority of binary-star systems: Given the observed properties of binaries in our Galaxy, “habitable” planetary orbits in binary systems would be unstable in many cases, as illustrated in Figure 28.14, so there would not be time for life to develop.

The scant observational evidence currently available on Earth-like planets in habitable orbits suggests that only a few percent of known planetary systems contain a habitable planet. ∞ (Sec. 5.5) However, because these planets are so close to the limits of detectability with current equipment, many astronomers think the true fraction will turn out to be much higher. Potentially habitable jovian moons could increase the fraction still further. However, many uncertainties remain. The inner and outer radii of the Galactic habitable zone are not known with any certainty, and the simple

The number of possible combinations of atoms is incredibly large. If the chemical reactions that led to the complex molecules that make up living organisms occurred completely at random, then it is extremely unlikely that those molecules could have formed at all. In that case, life is extraordinarily rare, this factor is close to zero, and we are probably alone in the Galaxy, perhaps even in the entire universe.

▶ **FIGURE 28.14 Binary-Star Planets** In binary-star systems, planets are restricted to only a few kinds of orbits that are gravitationally stable. (a) This orbit is stable only if the planet lies very close to its parent star, so that the gravity of the other star is negligible. (b) A planet circulating at a great distance about both stars in an elliptical orbit is stable only if it lies far from both stars. (c) Another possible path interweaves between the two stars in a figure-eight pattern.





However, laboratory experiments (such as the Urey-Miller experiment described earlier) seem to suggest that certain chemical combinations are strongly favored over others—that is, the reactions are *not* random. Of the billions upon billions of basic organic groupings that could occur on Earth from the random combination of all sorts of simple atoms and molecules, only about 1500 actually do occur. Furthermore, these 1500 organic groups of terrestrial biology are made from only about 50 simple “building blocks” (including the amino acids and nucleotide bases mentioned earlier). This suggests that molecules critical to life are not assembled by chance alone; apparently, additional factors are at work on the microscopic level. If a relatively small number of chemical “evolutionary tracks” are likely to exist, then the formation of complex molecules—and hence, we assume, life—becomes much more likely, given sufficient time.

To assign a very low value to this factor in the equation is to think that life arises randomly and rarely. To assign a value close to unity is to think that life is inevitable, given the proper ingredients, a suitable environment, and a long enough time. No simple experiment can distinguish between these extreme alternatives, and there is little or no middle ground. To many researchers, the discovery of life (past or present) on Mars, Europa, Titan, or some other body in our solar system would convert the appearance of life from an unlikely miracle to a virtual certainty throughout the Galaxy. Working on the principle that a universe with other life forms is far more interesting than one without, we will take the optimistic view and adopt a value of unity.

## Fraction of Life-Bearing Planets on Which Intelligence Arises

As with the evolution of life, the appearance of a well-developed brain is a highly unlikely event if only chance is involved. However, biological evolution through natural selection is a mechanism that generates apparently highly improbable results by singling out and refining useful characteristics. Organisms that profitably use adaptations can develop more complex behavior, and complex behavior provides organisms with the *variety* of choices needed for more advanced development.

One school of thought maintains that, given enough time, intelligence is inevitable. In this view, assuming that natural selection is a universal phenomenon, at least one organism on a planet will always rise to the level of “intelligent life.” If this is correct, then the fifth factor in the Drake equation equals or nearly equals unity.

Others argue that there is only one known case of intelligence: human beings on Earth. For 2.5 billion years—from the start of life about 3.5 billion years ago to the first appearance of multicellular organisms about 1 billion years ago—life did not advance beyond the one-celled stage. Life remained simple and dumb, but it survived. If this latter

view is correct, then the fifth factor in our equation is very small, and we are faced with the depressing prospect that humans may be the smartest form of life anywhere in the Galaxy. As with the previous factor, we will be optimistic and simply adopt a value of unity here.

## Fraction of Planets on Which Intelligent Life Develops and Uses Technology

To evaluate the sixth factor of our equation, we need to estimate the probability that intelligent life eventually develops technological competence. Should the rise of technology be inevitable, this factor is close to unity, given a long enough time. If it is not inevitable—if intelligent life can somehow “avoid” developing technology—then this factor could be much less than unity. The latter scenario envisions a universe possibly teeming with intelligent civilizations, but very few among them ever becoming technologically competent. Perhaps only one managed it—ours.

Again, it is difficult to decide conclusively between these two views. We don’t know how many prehistoric Earth cultures failed to develop simple technology or rejected its use. We do know that the roots of our present civilization arose independently at several different places on Earth, including Mesopotamia, India, China, Egypt, Mexico, and Peru. Because so many of these ancient cultures originated at about the same time, it is tempting to conclude that the chances are good that some sort of technological society will inevitably develop, given some basic intelligence and enough time.

If technology is inevitable, then why haven’t other life-forms on Earth also found it useful? Possibly the competitive edge given to humans, by intellectual and technological skills the first species to develop them, allowed us to dominate so rapidly that other species—gorillas and chimpanzees, for example—simply haven’t had time to catch up. The fact that only one technological society exists on Earth does not imply that the sixth factor in our Drake equation must be very much less than unity. On the contrary, it is precisely because *some* species will probably always fill the niche of technological intelligence that we will take this factor to be close to unity.

## Average Lifetime of a Technological Civilization

The reliability of the estimate of each factor in the Drake equation declines markedly from left to right. For example, our knowledge of astronomy enables us to make a reasonably good stab at the first factor, namely, the rate of star formation in our Galaxy, but it is much harder to evaluate some of the later factors, such as the fraction of life-bearing planets that eventually develop intelligence. The last factor on the right-hand side of the equation, the longevity of

technological civilizations, is totally unknown. There is only one known example of such a civilization: humans on planet Earth. Our own civilization has survived in its “technological” state for only about 100 years, and how long we will be around before a natural or human-made catastrophe ends it all is impossible to tell. [∞ \(Discovery 11-1\)](#)

## Number of Technological Civilizations in the Galaxy

One thing is certain: If the correct value for *any one factor* in the equation is very small—and we have just seen at least two for which this could well be the case, our optimistic choices notwithstanding—then few technological civilizations now exist in the Galaxy. In other words, if the pessimistic view of the development of life or of intelligence is correct, then we are unique, and that is the end of our story. However, if both life and intelligence are inevitable consequences of chemical and biological evolution, as many scientists think, and if intelligent life always becomes technological, then we can plug the higher, more optimistic values into the Drake equation. In that case, combining our estimates for the other six factors (and noting that  $10 \times 1 \times 1/10 \times 1 \times 1 \times 1 = 1$ ), we have

$$\begin{array}{l} \text{number of technological} \\ \text{intelligent civilizations} \\ \text{now present in the} \\ \text{Milky Way Galaxy} \end{array} = \begin{array}{l} \text{average lifetime of a} \\ \text{technologically} \\ \text{competent civilization} \\ \text{in years.} \end{array}$$

Thus, if civilizations typically survive for 1000 years, there should be 1000 of them currently in existence scattered throughout the Galaxy. If they live for a million years, on average, we would expect there to be a million advanced civilizations in the Milky Way, and so on.

Note that, even setting aside language and cultural issues, the sheer size of the Galaxy presents a significant hurdle to communication between technological civilizations. The minimum requirement for a two-way conversation is that we can send a signal and receive a reply in a time shorter than our own lifetime. If the lifetime is short, then civilizations are literally few and far between—small in number, according to the Drake equation, and scattered over the vastness of the Milky Way—and the distances between them (in light-years) are much greater than their lifetimes (in years). In that case, two-way communication, even at the speed of light, will be impossible. However, as the lifetime increases, the distances get smaller as the Galaxy becomes more crowded, and the prospects improve.

Taking into account the size, shape, and distribution of stars in the Galactic disk (why do we exclude the halo?), and under the optimistic assumptions just made, we find that, unless the life expectancy of a civilization is *at least* a few thousand years, it is unlikely to have time to communicate with even its nearest neighbor.

## PROCESS OF SCIENCE Check

- ✓ If most of the factors are largely a matter of opinion, how does the Drake equation assist astronomers in refining their search for extraterrestrial life?

## 28.4 The Search for Extraterrestrial Intelligence


Let us continue our optimistic assessment of the prospects for life and assume that civilizations enjoy a long stay on their parent planet once their initial technological “teething problems” are past. In that case, intelligent, technological, and perhaps also communicative cultures are likely to be plentiful in the Galaxy. How might we become aware of their existence? The ongoing search for extraterrestrial intelligence (known to many by its acronym, SETI) is the topic of this final section.

### Meeting Our Neighbors

For definiteness, let’s assume that the average lifetime of a technological civilization is 1 million years—only 1 percent of the reign of the dinosaurs, but 100 times longer than human civilization has survived thus far. Given the size and shape of our Galaxy and the known distribution of stars in the Galactic disk, we can then estimate the average *distance* between these civilizations to be some 30 pc, or about 100 light-years. Thus, any two-way communication with our neighbors—using signals traveling at or below the speed of light—will take at least 200 years (100 years for the message to reach the planet and another 100 years for the reply to travel back to us).

One obvious way to search for extraterrestrial life would be to develop the capability to travel far outside our solar system. However, that may never be a practical possibility. At a speed of 50 km/s, the speed of the fastest space probes operating today, the round-trip to even the nearest Sun-like star, Alpha Centauri, would take about 50,000 years. The journey to the nearest technological neighbor (assuming a distance of 30 pc) and back would take 600,000 years—almost the entire lifetime of our species! Interstellar travel at these speeds is clearly not feasible. Speeding up our ships to near the speed of light would reduce the travel time, but doing that is far beyond our present technology.

Actually, our civilization has already launched some interstellar probes, although they have no specific stellar destination. Figure 28.15 is a reproduction of a plaque mounted on board the *Pioneer 10* spacecraft launched in the mid-1970s and now well beyond the orbit of Pluto, on its way out of the solar system. Similar information was included aboard the *Voyager* probes launched in 1978. Although these spacecraft would be incapable of reporting back to Earth the



▲ **FIGURE 28.15 Interstellar Message** This replica of a plaque mounted on board the *Pioneer 10* spacecraft shows a scale drawing of the craft, a man, and a woman; a diagram of the hydrogen atom undergoing a change in energy (top left); a starburst pattern representing various pulsars and the frequencies of their radio waves that can be used to estimate when the craft was launched (middle left); and a depiction of the solar system, showing that the spacecraft departed the third planet from the Sun and passed the fifth planet on its way toward interstellar space (bottom). (NASA)

news that they had encountered an alien culture, scientists hope that the civilization on the other end would be able to unravel most of its contents using the universal language of mathematics. The caption to Figure 28.15 notes how the aliens might discover from where and when the *Pioneer* and *Voyager* probes were launched.

Setting aside the many practical problems that arise in trying to establish direct contact with extraterrestrials, some scientists have argued that it might not even be a particularly good idea. Our recent emergence as a technological civilization implies that we must be one of the least advanced technological intelligences in the entire Galaxy. Any other civilization that discovers us will almost surely be more advanced than us. Consequently, a healthy degree of caution may be warranted. If extraterrestrials behave even remotely like human civilizations on Earth, then the most advanced aliens may naturally try to dominate all others. The behavior of the “advanced” European cultures toward the “primitive” races they encountered on their voyages of discovery in the seventeenth, eighteenth, and nineteenth centuries should serve as a clear warning of the possible undesirable consequences of contact. Of course, the aggressiveness of Earthlings may not apply to extraterrestrials, but given the history of the one intelligent species we know, the cautious approach may be in order.

## Radio Communication

A cheaper and much more practical alternative to direct contact is to try to communicate with extraterrestrials by


using only electromagnetic radiation, the fastest known means of transferring information from one place to another. Because light and other high-frequency radiation is heavily scattered while moving through dusty interstellar space, long-wavelength radio radiation seems to be the natural choice. We would not attempt to broadcast to *all* nearby candidate stars, however—that would be far too expensive and inefficient. Instead, radio telescopes on Earth would listen *passively* for radio signals emitted by other civilizations. Indeed, some preliminary searches of selected nearby stars are now underway, thus far without success.

In what direction should we aim our radio telescopes? The answer to this question, at least, is fairly easy: On the basis of our earlier reasoning, we should target all F-, G-, and K-type stars in our vicinity. But are extraterrestrials broadcasting radio signals? If they are not, this search technique will obviously fail. And even if they are, how do we distinguish their artificially generated radio signals from signals naturally emitted by interstellar gas clouds? To what frequency should we tune our receivers? The answer to this question depends on whether the signals are produced deliberately or are simply “waste radiation” escaping from a planet.

Consider how Earth would look at radio wavelengths to extraterrestrials. Figure 28.16 shows the pattern of radio signals we emit into space. From the viewpoint of a distant observer, the spinning Earth emits a bright flash of radio radiation every few hours. In fact, Earth is now a more intense radio emitter than the Sun. The flashes result from the periodic rising and setting of hundreds of FM radio stations and television transmitters. Each station broadcasts mostly parallel to Earth’s surface, sending a great “sheet” of electromagnetic radiation into interstellar space, as illustrated in Figure 28.16(a). (The more common AM broadcasts are trapped below our ionosphere, so those signals never leave Earth.)

Because the great majority of these transmitters are clustered in the eastern United States and western Europe, a distant observer would detect periodic blasts of radiation from Earth as our planet rotates each day (Figure 28.16b). This radiation races out into space and has been doing so since the invention of these technologies more than seven decades ago. Another civilization at least as advanced as ours might have constructed devices capable of detecting these blasts of radiation. If any sufficiently advanced (and sufficiently interested) civilization resides on a planet orbiting any of the thousand or so stars within roughly 70 light-years (20 pc) of Earth, then we have already broadcast our presence to them.

Of course, it may very well be that, having discovered cable and fiber-optics technology, most civilizations’ indiscriminate transmissions cease after a few decades. In that case, radio silence becomes the hallmark of intelligence, and we must find an alternative means of locating our neighbors.



▲ **FIGURE 28.16 Earth's Radio Leakage** Radio radiation leaks from Earth into space because of the daily activities of our technological civilization. (a) Most radio and television transmitters broadcast their energy parallel to Earth's surface (where people live), sending a great "sheet" of electromagnetic radiation into interstellar space. (b) Because most transmitters are clustered in the eastern United States and western Europe, a distant observer would detect blasts of radio radiation from Earth as our planet rotates each day.

## The Water Hole


Now let us suppose that a civilization has decided to assist searchers by actively broadcasting its presence to the rest of the Galaxy. At what frequency should we listen for such an extraterrestrial beacon? The electromagnetic spectrum is enormous; the radio domain alone is vast. To hope to detect a signal at some unknown radio frequency is like searching for a needle in a haystack. Are some frequencies more likely than others to carry alien transmissions?



Some basic arguments suggest that civilizations might communicate at a wavelength near 20 cm. As we saw in Chapter 18, the basic building blocks of the universe, namely, hydrogen atoms, radiate naturally at a wavelength of 21 cm. ∞ (Sec. 18.4) Also, one of the simplest molecules, hydroxyl (OH), radiates near 18 cm. Together, these two substances form water (H<sub>2</sub>O). Arguing that water is likely to be the interaction medium for life anywhere and that radio radiation travels through the disk of our Galaxy with the least absorption by interstellar gas and dust, some researchers have proposed that the interval between 18 and 21 cm is the best range of wavelengths for civilizations to transmit or monitor.

Called the **water hole**, this radio interval might serve as an "oasis" where all advanced galactic civilizations would gather to conduct their electromagnetic business. So, if ET wants to be found, the reasoning goes, this is where we should look!

The water-hole frequency interval is only a guess, of course, but it is supported by other arguments as well. Figure 28.17 shows the water hole's location in the electromagnetic spectrum and plots the amount of natural emission from our Galaxy and from Earth's atmosphere. The 18- to 21-cm range lies within the quietest part of the spectrum, where the galactic "static" from stars and interstellar clouds happens to be minimized. Furthermore, the atmospheres of typical planets—or, at least, planets comparable to Earth—are also expected to interfere least at these wavelengths. Thus, the water hole seems like a good choice for the frequency of an interstellar beacon, although we cannot be sure of this reasoning until contact is actually achieved.

A few radio searches are now in progress at frequencies in and around the water hole. One of the most sensitive and comprehensive projects in the search for extraterrestrial intelligent life (known to many by its acronym SETI) is now underway with the Allen Telescope Array (Figure 28.18a).



**◀ FIGURE 28.17 Water Hole** The “water hole” is bounded by the natural emission frequencies of the hydrogen (H) atom at 21-cm wavelength and the hydroxyl (OH) molecule at 18-cm wavelength.  (Secs. 18.4, 18.5) The topmost solid (blue) curve sums the natural emissions of our Galaxy (dashed line) and Earth’s atmosphere (dotted line), which in turn are superposed atop the cosmic background radiation.  (Sec. 26.7) This sum is minimized near the water-hole frequencies, and thus all intelligent civilizations might conduct their interstellar communications within this quiet “electromagnetic oasis.”

This collection of many small dishes is currently searching millions of channels simultaneously in the 1- to 3-GHz range. Actually, in these searches, computers do most of the “listening;” humans get involved only if the signals look intriguing. Figure 28.18(b) shows what a typical narrow-band, 1-Hz signal—a potential “signature” of an intelligent transmission—would look like on a computer monitor. However, this observation was merely a test to detect the weak, redshifted radio signal emitted by the *Pioneer 10* robot, now receding into the outer realm of our solar system—a sign of intelligence, but one that we put there. Nothing resembling an extraterrestrial signal has yet been detected.

The space surrounding all of us could be, right now, flooded with radio signals from extraterrestrial civilizations. If only we knew the proper direction and frequency, we might be able to make one of the most startling discoveries of all time. The result would likely provide whole new opportunities to study the cosmic evolution of energy, matter, and life throughout the universe.

### PROCESS OF SCIENCE Check


- ✓ Why do many researchers regard the “water hole” as a likely place to search for extraterrestrial signals?

### ► FIGURE 28.18 Project

**SETI** (a) This array of small radio dishes at the SETI Institute in California is designed to search for extraterrestrial intelligent signals. (b) A typical recording of an alien signal—here, as a test, the Doppler-shifted broadcast from the *Pioneer 10* spacecraft, now well beyond the Kuiper belt—shows a diagonal line across the computer monitor, in contrast to the random noise in the background. (SETI Institute)



(a)



(b)

**The Big Question** The “Big Question” of Chapter 1 asked if planets might orbit the innumerable stars observed in the universe. It further wondered if there might be intelligent beings on some of those planets. Perhaps the grandest of all unsolved questions in astronomy is whether alien beings reside beyond Earth. For that reason alone, the search for extraterrestrial life will continue. The quest will never end.

# Chapter Review


## SUMMARY

### 1 Cosmic evolution (p. 716)

is the continuous process that has led to the appearance of galaxies, stars, planets, and life on Earth. Living organisms may be characterized by their ability to react to their environment, to grow by taking in nutrition from their surroundings, and to reproduce, passing along some of their own characteristics to their offspring. Organisms that can best take advantage of their new surroundings succeed at the expense of those organisms that cannot make the necessary adjustments. Intelligence is strongly favored by natural selection.



2 Powered by natural energy sources, reactions between simple molecules in the oceans of the primitive Earth may have led to the formation of **amino acids** (p. 718) and **nucleotide bases** (p. 718), the basic molecules of life. Amino acids build proteins, which control metabolism, while sequences of nucleotide bases make up DNA, the genetic blueprint of a living organism. Alternatively, some of these complex molecules, or their progenitors, may have been formed in interstellar space and then delivered to Earth by meteors or comets.




3 The best hope for life beyond Earth in the solar system is the planet Mars, although no direct evidence for life—current or extinct—has yet been found. Some of the icy moons of the outer planets—Jupiter’s Europa and Ganymede, and Saturn’s Titan and Enceladus—may also be possibilities for life of some sort. Conditions on those frozen bodies




are harsh by terrestrial standards, although **extremophiles** (p. 723) on Earth have been found to thrive in hostile

environments in which life had previously been thought impossible.

4 The **Drake equation** (p. 724) provides a means of estimating the probability of intelligent life in the Galaxy. The astronomical factors in the equation are the galactic star-formation rate, the likelihood of planets, and the number of habitable planets. Chemical and biological factors are the probability that life appears and the probability that it subsequently develops intelligence. Cultural and political factors are the probability that intelligence leads to technology and the lifetime of a civilization in the technological state. Taking an optimistic view of the development of life and intelligence leads to the conclusion that the total number of technologically competent civilizations in the Galaxy is approximately equal to the lifetime of a typical civilization, expressed in years. Even with optimistic assumptions, the distance to our nearest intelligent neighbor is likely to be many hundreds of parsecs.



5 Currently, space travel is not a feasible means of searching for intelligent life. Existing programs to discover extraterrestrial intelligence involve scanning the electromagnetic spectrum for signals. So far, no intelligible broadcasts have been received. A technological civilization would probably “announce” itself to the universe by the radio and television signals it emits into space. Observed from afar, our planet would appear as a radio source with a 24-hour period, as different regions of the planet rise and set. The “**water hole**” (p. 731) is a region in the radio range of the electromagnetic spectrum, near the 21-cm line of hydrogen and the 18-cm line of hydroxyl, where natural emissions from the Galaxy happen to be minimized. Many researchers regard this region as the best part of the spectrum for communication purposes.



**MasteringAstronomy**® For instructor-assigned homework go to MasteringAstronomy.

Problems labeled **POS** explore the process of science. **VIS** problems focus on reading and interpreting visual information. **LO** connects to the introduction's numbered Learning Outcomes.

## Review and Discussion

- LO1** Outline the processes that led to life on Earth. Why is life difficult to define in general?
- LO2** What is chemical evolution? What are the basic ingredients from which biological molecules formed on Earth?
- What is the Urey-Miller experiment? What important organic molecules did it produce? What other experiments have attempted to produce organic molecules by inorganic means?
- Why do some scientists think life might have originated in space?
- POS** How do we know anything at all about the early episodes of life on Earth?
- What is the role of language in cultural evolution?
- LO3** Where—besides Earth and the planet Mars—might we hope to find signs of life in our solar system?
- POS** Do we know whether Mars ever had life at any time during its past? What argues in favor of the position that it may once have harbored life?
- POS** What is generally meant by “life as we know it”? What other forms of life might be possible?
- LO4** How many of the factors in the Drake equation are known with any degree of certainty? Which factor is least well known?
- What factors determine the suitability of a star as the parent of a planet on which life might arise?
- What is the relationship between the average lifetime of galactic civilizations and the possibility of our someday communicating with them?
- How would Earth appear at radio wavelengths to extraterrestrial astronomers? What are the advantages in using radio waves for communication over interstellar distances?
- LOS POS** What is the “water hole”? What advantages does it offer for interstellar communication?
- If you were designing a SETI experiment, what parts of the sky would you monitor?

## Conceptual Self-Test: Multiple Choice

- The “assumptions of mediocrity” suggest that (a) life should be common throughout the cosmos; (b) lower forms of life must evolve to higher forms; (c) lower forms of life have lower intelligence; (d) viruses are actually life-forms.
- The chemical elements that form the basic molecules needed for life are found (a) in the cores of Sun-like stars; (b) commonly throughout the cosmos; (c) only on planets that have liquid water; (d) only on Earth.
- Fossil records of early life-forms on Earth suggest that life began about (a) 6000 years ago; (b) 65 million years ago; (c) 3.5 billion years ago; (d) 14 billion years ago.
- The discovery of bacteria on another planet would be an important discovery because bacteria (a) can easily survive in high temperatures; (b) are the only life-form to exist on Earth for most of the planet’s history; (c) are the lowest form of life known to exist; (d) eventually evolve into intelligent beings.
- The least-well-known factor in the Drake equation is (a) the rate of star formation; (b) the average number of habitable planets within planetary systems; (c) the average lifetime of a technologically competent civilization; (d) the diameter of the Milky Way Galaxy.
- Although the habitable zone around a large B-class star is large, we don’t often look for life on planets there because the star (a) has too much gravity; (b) is too short lived for life to evolve; (c) is at too low a temperature to sustain life; (d) would have only gas giant planets.
- VIS** From the data shown in Figure 28.12 (“Stellar Habitable Zones”) and your knowledge of stellar properties (Chapter 18), the habitable zone surrounding a main sequence K-type star (a) cannot be determined; (b) extends more than 3 AU from the star; (c) is larger than that of a G-type star; (d) is larger than that of an M-type star.
- VIS** If Figure 28.16 (“Leakage”) were to be redrawn for a planet spinning twice as fast, the new jagged line would be (a) unchanged; (b) taller; (c) stretched out horizontally; (d) compressed horizontally.
- Radio telescopes cannot simply scan the skies looking for signals, because (a) astronomers don’t know what frequencies alien civilizations might use; (b) many nonliving objects emit radio signals naturally; (c) Earth’s radio communications drown out extraterrestrial signals; (d) inclement weather in the winter prevents the use of radio telescopes.
- The strongest radio-wavelength emitter in the solar system is (a) human-made signals from Earth; (b) the Sun; (c) the Moon; (d) Jupiter.

## Problems

The number of dots preceding each Problem indicates its approximate level of difficulty.

1. ● If Earth's 4.6-billion-year age were compressed to 46 years, as described in the text, what would be your age, in seconds? On that scale, how long ago was the end of World War II? The Declaration of Independence? Columbus's discovery of the New World? The extinction of the dinosaurs?
2. ●● Using the data in the previous problem, how would the inner and outer radii of the Sun's habitable zone change if the solar luminosity increased by a factor of four?
3. ●● A planet orbits one component of a binary-star system at a distance of 1 AU. (See Figure 28.14a.) If both stars have the same mass and their orbit is circular, estimate the minimum distance between the stars for the tidal force due to the companion not to exceed a "safe" 0.01 percent of the gravitational force between the planet and its parent star.
4. ● Based on the numbers presented in the text, and assuming an average lifetime of 5 billion years for suitable stars, estimate the total number of habitable planets in the Galaxy.
5. ● Suppose that each of the fractional factors in the Drake equation turns out to have a value of 1/10, that stars form at an average rate of 20 per year, and that each star has exactly one habitable planet orbiting it. Estimate the present number of technological civilizations in the Milky Way Galaxy if the average lifetime of a civilization is (a) 100 years; (b) 10,000 years; (c) 1 million years.
6. ● Assuming that there are 10,000 FM radio stations on Earth, each transmitting at a power level of 50 kW, calculate the total radio luminosity of Earth in the FM band. Compare this value with the roughly  $10^6$  W radiated by the Sun in the same frequency range.
7. ● Convert the water hole's wavelengths to frequencies. For practical reasons, any search of the water hole must be broken up into channels, much like those you find on a television, except that the water hole's channels are very narrow in radio frequency, about 100 Hz wide. How many channels must astronomers search in the water hole?
8. ● There are 20,000 stars within 100 light-years that are to be searched for radio communications. How long will the search take if 1 hour is spent looking at each star? What if 1 day is spent per star?

## Activities

### Collaborative

1. As a group, compose a paragraph everyone agrees with that defines life. It should clearly show that rocks are not alive and that plants are alive. According to your definition, are stars alive? What about viruses? Compare and contrast your group's definition with that from another group.
2. If your group was appointed to "speak for Earth" upon establishing communication with an extraterrestrial world, what would you say? What questions would you ask, and what aspects of our planet would you choose to present? Write your group's speech and annotate it with explanations of why you chose to say this.

### Individual

1. It has been suggested that if extraterrestrial life is discovered, it will have a profound effect on human culture. Interview as many people as you can and ask the following two questions: (1) Do you think that extraterrestrial life exists? (2) Why? From your results, try to decide whether the discovery of extraterrestrial life would indeed profoundly affect life on Earth.
2. The Drake equation should be able to "predict" at least one civilization in our Galaxy: us. Try changing the values of various factors so that you end up with at least one. What do these various combinations of factors imply about how life arises and develops? Are there some combinations that just don't make any sense?



*This page intentionally left blank*

# Appendix 1

## Scientific Notation

The objects studied by astronomers range in size from the smallest particles to the largest expanse of matter we know—the entire universe. Subatomic particles have sizes of about 0.000000000000001 meter, while galaxies (like that shown in Figure 1.3) typically measure some 1,000,000,000,000,000,000 meters across. The most distant known objects in the universe lie on the order of 100,000,000,000,000,000,000,000 meters from Earth.

Obviously, writing all those zeros is both cumbersome and inconvenient. More important, it is also very easy to make an error—write down one zero too many or too few and your calculations become hopelessly wrong! To avoid this, scientists always write large numbers using a shorthand notation in which the number of zeros following or preceding the decimal point is denoted by a superscript power, or *exponent*, of 10. The exponent is simply the number of places between the first significant (nonzero) digit in the number (reading from left to right) and the decimal point. Thus, 1 is  $10^0$ , 10 is  $10^1$ , 100 is  $10^2$ , 1000 is  $10^3$ , and so on. For numbers less than 1, with zeros between the decimal point and the first significant digit, the exponent is negative: 0.1 is  $10^{-1}$ , 0.01 is  $10^{-2}$ , 0.001 is  $10^{-3}$ , and so on. Using this notation we can shorten the number describing subatomic particles to  $10^{-15}$  meter, and write the number describing the size of a galaxy as  $10^{21}$  meters.

More complicated numbers are expressed as a combination of a power of 10 and a multiplying factor. This factor is conventionally chosen to be a number between 1 and 10, starting with the first significant digit in the original number. For example, 150,000,000,000 meters (the distance from Earth to the Sun, in round numbers) can be more concisely written at  $1.5 \times 10^{11}$  meters, 0.000000025 meters as  $2.5 \times 10^{-8}$  meter, and so on. The exponent is simply the number of places the decimal point must be moved *to the left* to obtain the multiplying factor.

Some other examples of scientific notation are:

- the approximate distance to the Andromeda Galaxy  
= 2,500,000 light-years =  $2.5 \times 10^6$  light-years
- the size of a hydrogen atom  
= 0.00000000005 meter =  $5 \times 10^{-11}$  meter
- the diameter of the Sun  
= 1,392,000 kilometers =  $1.392 \times 10^6$  kilometers

- the U.S. national debt (as of May 1, 2013)  
= \$16,819,254,000,000.00 = \$16.819254 trillion  
=  $1.6819254 \times 10^{13}$  dollars.

In addition to providing a simpler way of expressing very large or very small numbers, this notation also makes it easier to do basic arithmetic. The rule for multiplication of numbers expressed in this way is simple: Just multiply the factors and add the exponents. Similarly for division: Divide the factors and subtract the exponents. Thus,  $3.5 \times 10^{-2}$  multiplied by  $2.0 \times 10^3$  is simply  $(3.5 \times 2.0) \times 10^{-2+3} = 7.0 \times 10^1$ —that is, 70. Again,  $5 \times 10^6$  divided by  $2 \times 10^4$  is just  $(5/2) \times 10^{6-4}$ , or  $2.5 \times 10^2$  (= 250). Applying these rules to unit conversions, we find, for example, that 200,000 nanometers is  $200,000 \times 10^{-9}$  meter (since 1 nanometer =  $10^{-9}$  meter; see Appendix 2), or  $2 \times 10^5 \times 10^{-9}$  meter, or  $2 \times 10^{5-9} = 2 \times 10^{-4}$  meter = 0.2 mm. Verify these rules for yourself with a few examples of your own. The advantages of this notation when considering astronomical objects will soon become obvious.

Scientists often use “rounded-off” versions of numbers, both for simplicity and for ease of calculation. For example, we will usually write the diameter of the Sun as  $1.4 \times 10^6$  kilometers, instead of the more precise number given earlier. Similarly, Earth’s diameter is 12,756 kilometers, or  $1.2756 \times 10^4$  kilometers, but for “ballpark” estimates we really don’t need so many digits and the more approximate number  $1.3 \times 10^4$  kilometers will suffice. Very often, we perform rough calculations using only the first one or two significant digits in a number, and that may be all that is necessary to make a particular point. For example, to support the statement, “The Sun is much larger than Earth,” we need only say that the ratio of the two diameters is roughly  $1.4 \times 10^6$  divided by  $1.3 \times 10^4$ . Since  $1.4/1.3$  is close to 1, the ratio is approximately  $10^6/10^4 = 10^2$ , or 100. The essential fact here is that the ratio is much larger than 1; calculating it to greater accuracy (to get 109.13) would give us no additional *useful* information. This technique of stripping away the arithmetic details to get to the essence of a calculation is very common in astronomy, and we use it frequently throughout this text.

# Appendix 2

## Astronomical Measurement

Astronomers use many different kinds of units in their work, simply because no single system of units will do. Rather than the *Système Internationale* (SI), or meter-kilogram-second (MKS), metric system used in most high school and college science classes, many professional astronomers still prefer the older centimeter-gram-second (CGS) system. However, astronomers also commonly introduce new units when convenient. For example, when discussing stars, the mass and radius of the Sun are often used as reference points. The solar mass, written as  $M_{\odot}$ , is equal to  $2.0 \times 10^{33}$  g, or  $2.0 \times 10^{30}$  kg (since 1 kg = 1000 g). The solar radius,  $R_{\odot}$ , is equal to 700,000 km, or  $7.0 \times 10^8$  m (1 km = 1000 m). The subscript  $\odot$  always stands for the Sun. Similarly, the subscript  $\oplus$  always stands for Earth. In this book, we try to use the units that astronomers commonly use in any given context, but we also give the “standard” SI equivalents where appropriate.

Of particular importance are the units of length astronomers use. On small scales, the *angstrom* (1 Å =  $10^{-10}$  m =  $10^{-8}$  cm), the *nanometer* (1 nm =  $10^{-9}$  m =  $10^{-7}$  cm), and the *micron* (1 μm =  $10^{-6}$  m =  $10^{-4}$  cm) are used. Distances within the solar system are usually expressed in terms of the *astronomical unit* (AU), the

mean distance between Earth and the Sun. One AU is approximately equal to 150,000,000 km, or  $1.5 \times 10^{11}$  m. On larger scales, the *light-year* (1 ly =  $9.5 \times 10^{15}$  m =  $9.5 \times 10^{12}$  km) and the *parsec* (1 pc =  $3.1 \times 10^{16}$  m =  $3.1 \times 10^{13}$  km = 3.3 ly) are commonly used. Still larger distances use the regular prefixes of the metric system: *kilo* for one thousand and *mega* for one million. Thus 1 kiloparsec (kpc) =  $10^3$  pc =  $3.1 \times 10^{19}$  m, 10 megaparsecs (Mpc) =  $10^7$  pc =  $3.1 \times 10^{23}$  m, and so on.

Astronomers use units that make sense within a context, and as contexts change, so do the units. For example, we might measure densities in grams per cubic centimeter ( $\text{g}/\text{cm}^3$ ), in atoms per cubic meter ( $\text{atoms}/\text{m}^3$ ), or even in solar masses per cubic megaparsec ( $M_{\odot}/\text{Mpc}^3$ ), depending on the circumstances. The important thing to know is that once you understand the units, you can convert freely from one set to another. For example, the radius of the Sun could equally well be written as  $R_{\odot} = 6.96 \times 10^3$  m, or  $6.96 \times 10^{10}$  cm, or  $109 R_{\oplus}$ , or  $4.65 \times 10^{-3}$  AU, or even  $7.363 \times 10^{-5}$  ly—whichever happens to be most useful. Some of the more common units used in astronomy, and the contexts in which they are most likely to be encountered, are listed below.

Length		
1 angstrom (Å)	= $10^{-10}$ m	
1 nanometer (nm)	= $10^{-9}$ m	atomic physics, spectroscopy
1 micron (μm)	= $10^{-6}$ m	interstellar dust and gas
1 centimeter (cm)	= 0.01 m	
1 meter (m)	= 100 cm	in widespread use throughout all astronomy
1 kilometer (km)	= 1000 m = $10^5$ cm	
Earth radius ( $R_{\oplus}$ )	= 6378 km	planetary astronomy
Solar radius ( $R_{\odot}$ )	= $6.96 \times 10^8$ m	
1 astronomical unit (AU)	= $1.496 \times 10^{11}$ m	solar system, stellar evolution
1 light-year (ly)	= $9.46 \times 10^{15}$ m = 63,200 AU	
1 parsec (pc)	= $3.09 \times 10^{16}$ m = 206,000 AU = 3.26 ly	galactic astronomy, stars and star clusters
1 kiloparsec (kpc)	= 1000 pc	
1 megaparsec (Mpc)	= 1000 kpc	galaxies, galaxy clusters, cosmology
Mass		
1 gram (g)		
1 kilogram (kg)	= 1000 g	in widespread use in many different areas
Earth mass ( $M_{\oplus}$ )	= $5.98 \times 10^{24}$ kg	planetary astronomy
Solar mass ( $M_{\odot}$ )	= $1.99 \times 10^{30}$ kg	“standard” unit for all mass scales larger than Earth
Time		
1 second (s)		in widespread use throughout astronomy
1 hour (h)	= 3600 s	
1 day (d)	= 86,400 s	planetary and stellar scales
1 year (yr)	= $3.16 \times 10^7$ s	virtually all processes occurring on scales larger than a star

# Appendix 3

## Tables

**TABLE 1** Some Useful Constants and Physical Measurements\*

astronomical unit	$1 \text{ AU} = 1.496 \times 10^8 \text{ km}$ ( $1.5 \times 10^8 \text{ km}$ )
light-year	$1 \text{ ly} = 9.46 \times 10^{12} \text{ km}$ ( $10^{13} \text{ km}$ , about 6 trillion miles)
parsec	$1 \text{ pc} = 3.09 \times 10^{13} \text{ km} = 206,000 \text{ AU} = 3.3 \text{ ly}$
speed of light	$c = 299,792.458 \text{ km/s}$ ( $3 \times 10^5 \text{ km/s}$ )
Stefan-Boltzmann constant	$a = 5.67 \times 10^{-8} \text{ W/m}^2 \cdot \text{K}^4$
Planck's constant	$h = 6.63 \times 10^{-34} \text{ J s}$
gravitational constant	$G = 6.67 \times 10^{-11} \text{ Nm}^2 / \text{kg}^2$
mass of Earth	$M_{\oplus} = 5.98 \times 10^{24} \text{ kg}$ ( $6 \times 10^{24} \text{ kg}$ , about 6000 billion billion tons)
radius of Earth	$R_{\oplus} = 6378 \text{ km}$ (6500 km)
mass of the Sun	$M_{\odot} = 1.99 \times 10^{30} \text{ kg}$ ( $2 \times 10^{30} \text{ kg}$ )
radius of the Sun	$R_{\odot} = 6.96 \times 10^5 \text{ km}$ ( $7 \times 10^5 \text{ km}$ )
luminosity of the Sun	$L_{\odot} = 3.90 \times 10^{26} \text{ W}$ ( $4 \times 10^{26} \text{ W}$ )
effective temperature of the Sun	$T_{\odot} = 5778 \text{ K}$ (5800 K)
Hubble's constant	$H_0 = 70 \text{ km/s/Mpc}$
mass of an electron	$m_e = 9.11 \times 10^{-31} \text{ kg}$
mass of a proton	$m_p = 1.67 \times 10^{-27} \text{ kg}$

\* The rounded-off values used in the text are shown above in parentheses.

### Conversions Between Common English and Metric Units

English	Metric
1 inch	= 2.54 centimeters (cm)
1 foot (ft)	= 0.3048 meters (m)
1 mile	= 1.609 kilometers (km)
1 pound (lb)	= 453.6 grams (g) or 0.4536 kilograms (kg) [on Earth]

TABLE 2 Periodic Table of Elements

Group	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
Period																		
1	1 <b>H</b> 1.0080 Hydrogen																	2 <b>He</b> 4.003 Helium
2	3 <b>Li</b> 6.939 Lithium	4 <b>Be</b> 9.012 Beryllium											5 <b>B</b> 10.81 Boron	6 <b>C</b> 12.011 Carbon	7 <b>N</b> 14.007 Nitrogen	8 <b>O</b> 15.9994 Oxygen	9 <b>F</b> 18.998 Fluorine	10 <b>Ne</b> 20.183 Neon
3	11 <b>Na</b> 22.990 Sodium	12 <b>Mg</b> 24.31 Magnesium											13 <b>Al</b> 26.98 Aluminum	14 <b>Si</b> 28.09 Silicon	15 <b>P</b> 30.974 Phosphorus	16 <b>S</b> 32.064 Sulfur	17 <b>Cl</b> 35.453 Chlorine	18 <b>Ar</b> 39.948 Argon
4	19 <b>K</b> 39.10 Potassium	20 <b>Ca</b> 40.08 Calcium	21 <b>Sc</b> 44.96 Scandium	22 <b>Ti</b> 47.87 Titanium	23 <b>V</b> 50.94 Vanadium	24 <b>Cr</b> 52.00 Chromium	25 <b>Mn</b> 53.94 Manganese	26 <b>Fe</b> 55.85 Iron	27 <b>Co</b> 58.93 Cobalt	28 <b>Ni</b> 58.69 Nickel	29 <b>Cu</b> 63.55 Copper	30 <b>Zn</b> 65.39 Zinc	31 <b>Ga</b> 69.72 Gallium	32 <b>Ge</b> 72.61 Germanium	33 <b>As</b> 74.92 Arsenic	34 <b>Se</b> 78.96 Selenium	35 <b>Br</b> 79.904 Bromine	36 <b>Kr</b> 83.80 Krypton
5	37 <b>Rb</b> 85.47 Rubidium	38 <b>Sr</b> 87.62 Strontium	39 <b>Y</b> 88.91 Yttrium	40 <b>Zr</b> 91.22 Zirconium	41 <b>Nb</b> 92.91 Niobium	42 <b>Mo</b> 95.94 Molybdenum	43 <b>Tc</b> (99) Technetium	44 <b>Ru</b> 101.07 Ruthenium	45 <b>Rh</b> 102.91 Rhodium	46 <b>Pd</b> 106.42 Palladium	47 <b>Ag</b> 107.87 Silver	48 <b>Cd</b> 112.41 Cadmium	49 <b>In</b> 114.82 Indium	50 <b>Sn</b> 118.71 Tin	51 <b>Sb</b> 121.76 Antimony	52 <b>Te</b> 127.60 Tellurium	53 <b>I</b> 126.904 Iodine	54 <b>Xe</b> 131.29 Xenon
6	55 <b>Cs</b> 132.91 Cesium	56 <b>Ba</b> 137.33 Barium	71 <b>Lu</b> 174.97 Lutetium	72 <b>Hf</b> 178.49 Hafnium	73 <b>Ta</b> 180.95 Tantalum	74 <b>W</b> 183.84 Tungsten	75 <b>Re</b> 186.21 Rhenium	76 <b>Os</b> 190.23 Osmium	77 <b>Ir</b> 192.22 Iridium	78 <b>Pt</b> 195.09 Platinum	79 <b>Au</b> 196.97 Gold	80 <b>Hg</b> 200.59 Mercury	81 <b>Tl</b> 204.38 Thallium	82 <b>Pb</b> 207.20 Lead	83 <b>Bi</b> 208.98 Bismuth	84 <b>Po</b> (209) Polonium	85 <b>At</b> (210) Astatine	86 <b>Rn</b> (222) Radon
7	87 <b>Fr</b> (223) Francium	88 <b>Ra</b> (226) Radium	103 <b>Lw</b> (262) Lawrencium	104 <b>Rf</b> (263) Rutherfordium	105 <b>Db</b> (262) Dubnium	106 <b>Sg</b> (266) Seaborgium	107 <b>Bh</b> (264) Bohrium	108 <b>Hs</b> (269) Hassium	109 <b>Mt</b> (268) Meitnerium	110 <b>Ds</b> (272) Darmstadtium	111 <b>Rg</b> (272) Roentgenium	112 <b>Cn</b> (277) Copernicium	113 <b>Uut</b> (284) Ununtrium	114 <b>Uuq</b> (289) Ununquadium	115 <b>Uup</b> (288) Ununpentium	116 <b>Uuh</b> (292) Ununhexium	117 <b>Uus</b> (294) Ununseptium	118 <b>Uuo</b> (294) Ununoctium
			57 <b>La</b> 138.91 Lanthanum	58 <b>Ce</b> 140.12 Cerium	59 <b>Pr</b> 140.91 Praseodymium	60 <b>Nd</b> 144.24 Neodymium	61 <b>Pm</b> (145) Promethium	62 <b>Sm</b> 150.36 Samarium	63 <b>Eu</b> 151.96 Europium	64 <b>Gd</b> 157.25 Gadolinium	65 <b>Tb</b> 158.93 Terbium	66 <b>Dy</b> 162.50 Dysprosium	67 <b>Ho</b> 164.93 Holmium	68 <b>Er</b> 167.26 Erbium	69 <b>Tm</b> 168.93 Thulium	70 <b>Yb</b> 173.04 Ytterbium		
			89 <b>Ac</b> (227) Actinium	90 <b>Th</b> 232.04 Thorium	91 <b>Pa</b> 231.03 Protactinium	92 <b>U</b> 238.03 Uranium	93 <b>Np</b> (237) Neptunium	94 <b>Pu</b> (242) Plutonium	95 <b>Am</b> (243) Americium	96 <b>Cm</b> (247) Curium	97 <b>Bk</b> (247) Berkelium	98 <b>Cf</b> (249) Californium	99 <b>Es</b> (252) Einsteinium	100 <b>Fm</b> (257) Fermium	101 <b>Md</b> (258) Mendelevium	102 <b>No</b> (259) Nobelium		

2 ← Atomic number  
**He** ← Symbol of element  
 4.003 ← Atomic weight  
 Helium ← Name of element

Element 117 was discovered in 2010. Element 118 was "discovered" in 1999, retracted in 2002, and reported again in 2006.

**TABLE 3A Planetary Orbital Data**

Planet	Semi-Major Axis		Eccentricity (e)	Perihelion		Aphelion	
	(AU)	(10 <sup>6</sup> km)		(AU)	(10 <sup>6</sup> km)	(AU)	(10 <sup>6</sup> km)
Mercury	0.39	57.9	0.206	0.31	46.0	0.47	69.8
Venus	0.72	108.2	0.007	0.72	107.5	0.73	108.9
Earth	1.00	149.6	0.017	0.98	147.1	1.02	152.1
Mars	1.52	227.9	0.093	1.38	206.6	1.67	249.2
Jupiter	5.20	778.4	0.048	4.95	740.7	5.46	816
Saturn	9.54	1427	0.054	9.02	1349	10.1	1504
Uranus	19.19	2871	0.047	18.3	2736	20.1	3006
Neptune	30.07	4498	0.009	29.8	4460	30.3	4537

Planet	Mean Orbital Speed (km/s)	Sidereal Period (tropical years)	Synodic Period (days)	Inclination to the Ecliptic (degrees)	Greatest Angular Diameter as Seen from Earth (arc seconds)
Mercury	47.87	0.24	115.88	7.00	13
Venus	35.02	0.62	583.92	3.39	64
Earth	29.79	1.00	—	0.01	—
Mars	24.13	1.88	779.94	1.85	25
Jupiter	13.06	11.86	398.88	1.31	50
Saturn	9.65	29.42	378.09	2.49	21
Uranus	6.80	83.75	369.66	0.77	4.1
Neptune	5.43	163.7	367.49	1.77	2.4

**TABLE 3B Planetary Physical Data**

Planet	Equatorial Radius		Mass		Mean Density (kg/m <sup>3</sup> )	Surface Gravity (Earth = 1)	Escape Speed (km/s)
	(km)	(Earth = 1)	(kg)	(Earth = 1)			
Mercury	2440	0.38	$3.30 \times 10^{23}$	0.055	5430	0.38	4.2
Venus	6052	0.95	$4.87 \times 10^{24}$	0.82	5240	0.91	10.4
Earth	6378	1.00	$5.97 \times 10^{24}$	1.00	5520	1.00	11.2
Mars	3394	0.53	$6.42 \times 10^{23}$	0.11	3930	0.38	5.0
Jupiter	71,492	11.21	$1.90 \times 10^{27}$	317.8	1330	2.53	60
Saturn	60,268	9.45	$5.68 \times 10^{26}$	95.16	690	1.07	36
Uranus	25,559	4.01	$8.68 \times 10^{25}$	14.54	1270	0.91	21
Neptune	24,766	3.88	$1.02 \times 10^{26}$	17.15	1640	1.14	24

Planet	Sidereal Rotation Period (solar days)*	Axial Tilt (degrees)	Surface Magnetic Field (Earth = 1)	Magnetic Axis Tilt (degrees relative to rotation axis)	Albedo <sup>†</sup>	Surface Temperature <sup>‡</sup> (K)	Number of Moons**
Mercury	58.6	0.0	0.011	<10	0.11	100–700	0
Venus	−243.0	177.4	<0.001		0.65	730	0
Earth	0.9973	23.45	1.0	11.5	0.37	290	1
Mars	1.026	23.98	0.001		0.15	180–270	2
Jupiter	0.41	3.08	13.89	9.6	0.52	124	16
Saturn	0.44	26.73	0.67	0.8	0.47	97	18
Uranus	−0.72	97.92	0.74	58.6	0.50	58	27
Neptune	0.67	29.6	0.43	46.0	0.5	59	13

\*A negative sign indicates retrograde rotation; <sup>†</sup>Fraction of sunlight reflected from surface; <sup>‡</sup>Temperature is effective temperature for jovian planets;

\*\*Moons more than 10 km in diameter.

**TABLE 4** The Twenty Brightest Stars in Earth's Night Sky

Name	Star	Spectral Type*		Parallax (arc seconds)	Distance (pc)	Apparent Visual Magnitude*	
		A	B			A	B
Sirius	$\alpha$ CMa	A1V	wd <sup>†</sup>	0.379	2.6	-1.44	+8.4
Canopus	$\alpha$ Car	F0Ib-II		0.010	96	-0.62	
Arcturus	$\alpha$ Boo	K2III		0.089	11	-0.05	
Rigel Kentaurus (Alpha Centauri)	$\alpha$ Gen	G2V	K0V	0.742	1.3	-0.01	+1.4
Vega	$\alpha$ Lyr	A0V		0.129	7.8	+0.03	
Capella	$\alpha$ Aur	GIII	M1V	0.077	13	+0.08	+10.2
Rigel	$\beta$ Ori	B8Ia	B9	0.0042	240	+0.18	+6.6
Procyon	$\alpha$ CMi	F5IV-V	wd <sup>†</sup>	0.286	3.5	+0.40	+10.7
Betelgeuse	$\alpha$ Ori	M2Iab		0.0076	130	+0.45	
Achernar	$\alpha$ Eri	B5V		0.023	44	+0.45	
Hadar	$\beta$ Cen	B1III	?	0.0062	160	+0.61	+4
Altair	$\alpha$ Aql	A7IV-V		0.194	5.1	+0.76	
Acrux	$\alpha$ Cru	B1IV	B3	0.010	98	+0.77	+1.9
Aldebaran	$\alpha$ Tau	K5III	M2V	0.050	20	+0.87	+13
Spica	$\alpha$ Vir	B1V	B2V	0.012	80	+0.98	2.1
Antares	$\alpha$ Sco	M1Ib	B4V	0.005	190	+1.06	+5.1
Pollux	$\beta$ Gem	K0III		0.097	10	+1.16	
Formalhaut	$\alpha$ PsA	A3V	?	0.130	7.7	+1.17	+6.5
Deneb	$\alpha$ Cyg	A2Ia		0.0010	990	+1.25	
Mimosa	$\beta$ Cru	B1IV		0.0093	110	+1.25	

Name	Visual Luminosity* (Sun = 1)		Absolute Visual Magnitude		Proper Motion (arc seconds/yr)	Transverse Velocity (km/s)	Radial Velocity (km/s)
	A	B	A	B			
Sirius	22	0.0025	+1.5	+11.3	1.33	16.7	-7.6 <sup>‡</sup>
Canopus	$1.4 \times 10^4$		-5.5		0.02	9.1	20.5
Arcturus	110		-0.3		2.28	119	-5.2
Rigel Kentaurus	1.6	0.45	+4.3	+5.7	3.68	22.7	-24.6
Vega	50		+0.6		0.34	12.6	-13.9
Capella	130	0.01	-0.5	+9.6	0.44	27.1	30.2 <sup>‡</sup>
Rigel	$4.1 \times 10^4$	110	-6.7	-0.3	0.00	1.2	20.7 <sup>‡</sup>
Procyon	7.2	0.0006	+2.7	+13.0	1.25	20.7	-3.2 <sup>‡</sup>
Betelgeuse	9700		-5.1		0.03	18.5	21.0 <sup>‡</sup>
Achernar	1100		-2.8		0.10	20.9	19
Hadar	$1.3 \times 10^4$	560	-5.4	-2.0	0.04	30.3	-12 <sup>‡</sup>
Altair	11		+2.2		0.66	16.3	-26.3
Acrux	4100	2200	-4.2	-3.5	0.04	22.8	-11.2
Aldebaran	150	0.002	-0.6	+11.5	0.20	19.0	54.1
Spica	2200	780	-3.5	-2.4	0.05	19.0	1.0 <sup>‡</sup>
Antares	$1.1 \times 10^4$	290	-5.3	-1.3	0.03	27.0	-3.2
Pollux	31		+1.1		0.62	29.4	3.3
Formalhaut	17	0.13	+1.7	+7.1	0.37	13.5	6.5
Deneb	$2.6 \times 10^5$		-8.7		0.003	14.1	-4.6 <sup>‡</sup>
Mimosa	3200		-3.9		0.05	26.1	—

\*Energy output in the visible part of the spectrum; A and B columns identify individual components of binary-star systems.

<sup>†</sup>“wd” stands for “white dwarf.”

<sup>‡</sup>Average value of variable velocity.

**TABLE 5** The Twenty Nearest Stars

Name	Spectral Type		Parallax (arc seconds)	Distance (pc)	Apparent Visual Magnitude*	
	A	B			A	B
Sun	G2V				-26.74	
Proxima Centauri	M5		0.772	1.30	+11.01	
Alpha Centauri	G2V	K1V	0.742	1.35	-0.01	+1.35
Barnard's Star	M5V		0.549	1.82	+9.54	
Wolf 359	M8V		0.421	2.38	+13.53	
Lalande 21185	M2V		0.397	2.52	+7.50	
UV Ceti	M6V	M6V	0.387	2.58	+12.52	+13.02
Sirius	A1V	wd <sup>†</sup>	0.379	2.64	-1.44	+8.4
Ross 154	M5V		0.345	2.90	+10.45	
Ross 248	M6V		0.314	3.18	+12.29	
ε Eridani	K2V		0.311	3.22	+3.72	
Ross 128	M5V		0.298	3.36	+11.10	
61 Cygni	K5V	K7V	0.294	3.40	+5.22	+6.03
ε Indi	K5V		0.291	3.44	+4.68	
Grm 34	M1V	M6V	0.290	3.45	+8.08	+11.06
Luyten 789-6	M6V		0.290	3.45	+12.18	
Procyon	F5IV-V	wd <sup>†</sup>	0.286	3.50	+0.40	+10.7
Σ 2398	M4V	M5V	0.285	3.55	+8.90	+9.69
Lacaille 9352	M2V		0.279	3.58	+7.35	
G51-15	MV		0.278	3.60	+14.81	

Name	Visual Luminosity* (Sun = 1)		Absolute Visual Magnitude*		Proper Motion (arc seconds/yr)	Transverse Velocity (km/s)	Radial Velocity (km/s)
	A	B	A	B			
Sun	1.0		+4.83				
Proxima Centauri	5.6 × 10 <sup>-5</sup>		+15.4		3.86	23.8	-16
Alpha Centauri	1.6	0.45	+4.3	+5.7	3.68	23.2	-22
Barnard's Star	4.3 × 10 <sup>-4</sup>		+13.2		10.34	89.7	-108
Wolf 359	1.8 × 10 <sup>-5</sup>		+16.7		4.70	53.0	+13
Lalande 21185	0.0055		+10.5		4.78	57.1	-84
UV Ceti	5.4 × 10 <sup>-5</sup>	0.00004	+15.5	+16.0	3.36	41.1	+30
Sirius	22	0.0025	+1.5	+11.3	1.33	16.7	-8
Ross 154	4.8 × 10 <sup>-4</sup>		+13.3		0.72	9.9	-4
Ross 248	1.1 × 10 <sup>-4</sup>		+14.8		1.58	23.8	-81
ε Eridani	0.29		+6.2		0.98	15.3	+16
Ross 128	3.6 × 10 <sup>-4</sup>		+13.5		1.37	21.8	-13
61 Cygni	0.082	0.039	+7.6	+8.4	5.22	84.1	-64
ε Indi	0.14		+7.0		4.69	76.5	-40
Grm 34	0.0061	0.00039	+10.4	+13.4	2.89	47.3	+17
Luyten 789-6	1.4 × 10 <sup>-4</sup>		+14.6		3.26	53.3	-60
Procyon	7.2	0.00055	+2.7	+13.0	1.25	2.8	-3
Σ 2398	0.0030	0.0015	+11.2	+11.9	2.28	38.4	+5
Lacaille 9352	0.013		+9.6		6.90	117	+10
G51-15	1.1 × 10 <sup>-5</sup>		+17.0		1.26	21.5	—

\*A and B columns identify individual components of binary-star systems.

<sup>†</sup>“wd” stands for “white dwarf.”



*This page intentionally left blank*

# Glossary

Key terms that are boldface in the text are followed by a page reference in the glossary.

## A

**A ring** One of three Saturnian rings visible from Earth. The A ring is farthest from the planet and is separated from the B ring by the Cassini division. (p. 298)

**aberration of starlight** Small shift in the observed direction to a star, caused by Earth's motion perpendicular to the line of sight. (p. 42)

**absolute brightness** The apparent brightness a star would have if it were placed at a standard distance of 10 parsecs from Earth.

**absolute magnitude** The apparent magnitude a star would have if it were placed at a standard distance of 10 parsecs from Earth. (p. 427)

**absolute zero** The lowest possible temperature that can be obtained; all thermal motion ceases at this temperature.

**absorption line** Dark line in an otherwise continuous bright spectrum, where light within one narrow frequency range has been removed. (p. 82)

**abundance** Relative amount of different elements in a gas.

**acceleration** The rate of change of velocity of a moving object. (p. 50)

**accretion** Gradual growth of bodies, such as planets, by the accumulation of other smaller bodies. (p. 150)

**accretion disk** Flat disk of matter spiraling down onto the surface of a neutron star or black hole. Often, the matter originated on the surface of a companion star in a binary-star system. (p. 522)

**active galactic nucleus** Region of intense emission at the center of an active galaxy, responsible for virtually all of the galaxy's nonstellar luminosity. (p. 624)

**active galaxies** The most energetic galaxies, which can emit hundreds or thousands of times more energy per second than the Milky Way, mostly in the form of long-wavelength nonthermal radiation. (p. 622)

**active optics** Collection of techniques used to increase the resolution of ground-based telescopes. Minute modifications are made to the overall configuration of an instrument as its temperature and orientation change; used to maintain the best possible focus at all times. (p. 112)

**active region** Region of the photosphere of the Sun surrounding a sunspot group, which can erupt violently and unpredictably. During sunspot maximum, the number of active regions is also at maximum. (p. 405)

**active Sun** The unpredictable aspects of the Sun's behavior, such as sudden explosive outbursts of radiation in the form of prominences and flares.

**adaptive optics** Technique used to increase the resolution of a telescope by deforming the shape of the mirror's surface under computer control while a measurement is being taken; used to undo the effects of atmospheric turbulence. (p. 112)

**aerosol** Suspension of liquid or solid particles in air.

**alpha particle** A helium-4 nucleus.

**alpha process** Process occurring at high temperatures, in which high-energy photons split heavy nuclei to form helium nuclei.

**ALSEP** Acronym for Apollo Lunar Surface Experiments Package.

**altimeter** Instrument used to determine altitude.

**amino acids** Organic molecules that form the basis for building the proteins that direct metabolism in living creatures. (p. 718)

**Amor asteroid** Asteroid that crosses only the orbit of Mars.

**amplitude** The maximum deviation of a wave above or below the zero point. (p. 61)

**angstrom** Distance unit equal to 0.1 nanometer, or one ten-billionth of a meter.

**angular diameter** Angle made between the top (or one edge) of an object, the observer, and the bottom (or opposite edge) of the object.

**angular distance** Angular separation between two objects as seen by some observer.

**angular momentum** Tendency of an object to keep rotating; proportional to the mass, radius, and rotation speed of the body. (p. 149)

**angular resolution** The ability of a telescope to distinguish between adjacent objects in the sky. (p. 106)

**annular eclipse** Solar eclipse occurring at a time when the Moon is far enough away from Earth that it fails to cover the disk of the Sun completely, leaving a ring of sunlight visible around its edge. (p. 21)

**antiparallel** Configuration of the electron and proton in a hydrogen (or other) atom when their spin axes are parallel but the two rotate in opposite directions.

**antiparticle** A particle of the same mass but opposite in all other respects (e.g., charge) to a given particle; when a particle and its antiparticle come into contact, they annihilate and release energy in the form of gamma rays.

**aphelion** The point on the elliptical path of an object in orbit about the Sun that is most distant from the Sun.

**Apollo asteroid** See Earth-crossing asteroid.

**apparent brightness** The brightness that a star appears to have, as measured by an observer on Earth. (p. 425)

**apparent magnitude** The apparent brightness of a star, expressed using the magnitude scale. (p. 426)

**association** Small grouping of (typically 100 or less) bright stars, spanning up to a few tens of parsecs across, usually rich in very young stars. (p. 486)

**assumption of mediocrity** Statements suggesting that the development of life on Earth did not require any unusual circumstances, suggesting that extraterrestrial life may be common.

**asteroid** One of thousands of very small members of the solar system orbiting the Sun between the orbits of Mars and Jupiter. Often referred to as “minor planets.” (p. 340)

**asteroid belt** Region of the solar system, between the orbits of Mars and Jupiter, in which most asteroids are found. (p. 340)

**asthenosphere** Layer of Earth’s interior, just below the lithosphere, over which the surface plates slide. (p. 174)

**astrology** Pseudoscience that purports to use the positions of the planets, Sun, and Moon to predict daily events and human destiny.

**astronomical unit (AU)** The average distance of Earth from the Sun. Precise radar measurements yield a value for the AU of 149,603,500 km. (p. 47)

**astronomy** Branch of science dedicated to the study of everything in the universe that lies above Earth’s atmosphere. (p. 6)

**asymptotic giant branch** Path on the Hertzsprung–Russell diagram corresponding to the changes that a star undergoes after helium burning ceases in the core. At this stage, the carbon core shrinks and drives the expansion of the envelope, and the star becomes a swollen red giant for a second time. (p. 502)

**Aten asteroid** Earth-crossing asteroid with semimajor axis less than 1 AU.

**atmosphere** Layer of gas confined close to a planet’s surface by the force of gravity. (p. 162)

**atom** Building block of matter, composed of positively charged protons and neutral neutrons in the nucleus surrounded by negatively charged electrons. (p. 84)

**atomic epoch** Period after decoupling when the first simple atoms and molecules formed.

**aurora** Event that occurs when atmospheric molecules are excited by incoming charged particles from the solar wind, then emit energy as they fall back to their ground states. Aurorae generally occur at high latitudes, near the north and south magnetic poles. (p. 180)

**autumnal equinox** Date on which the Sun crosses the celestial equator moving southward, occurring on or near September 21. (p. 17)

## B

**B ring** One of three Saturnian rings visible from Earth. The B ring is the brightest of the three, and lies just past the Cassini division, closer to the planet than the A ring. (p. 298)

**background noise** Unwanted light in an image, from unresolved sources in the telescope’s field of view, scattered light from the atmosphere, or instrumental “hiss” in the detector itself.

**barred-spiral galaxy** Spiral galaxy in which a bar of material passes through the center of the galaxy, with the spiral arms beginning near the ends of the bar. (p. 609)

**basalt** Solidified lava; an iron-magnesium-silicate mixture.

**baseline** The distance between two observing locations used for the purposes of triangulation measurements. The larger the baseline, the better the resolution attainable. (p. 24)

**belt** Dark, low-pressure region in the atmosphere of a jovian planet, where gas flows downward. (p. 268)

**Big Bang** Event that cosmologists consider the beginning of the universe, in which all matter and radiation in the entire universe came into being. (p. 671)

**Big Crunch** Point of final collapse of a bound universe.

**binary asteroid** Asteroid with a partner in orbit around it.

**binary pulsar** Binary system in which both components are pulsars.

**binary-star system** A system that consists of two stars in orbit about their common center of mass, held together by their mutual gravitational attraction. Most stars are found in binary-star systems. (p. 440)

**biological evolution** Change in a population of biological organisms over time.

**bipolar flow** Jets of material expelled from a protostar perpendicular to the surrounding protostellar disk. (p. 483)

**blackbody curve** The characteristic way in which the intensity of radiation emitted by a hot object depends on frequency. The frequency at which the emitted intensity is highest is an indication of the temperature of the radiating object. Also referred to as the Planck curve. (p. 68)

**black dwarf** The end point of the evolution of an isolated, low-mass star. After the white-dwarf stage, the star cools to the point where it is a dark “clinker” in interstellar space. (p. 507)

**black hole** A region of space where the pull of gravity is so great that nothing—not even light—can escape. A possible outcome of the evolution of a very massive star. (p. 556)

**blazar** Particularly intense active galactic nucleus in which the observer’s line of sight happens to lie directly along the axis of a high-speed jet of particles emitted from the active region. (p. 627)

**blue giant** Large, hot, bright star at the upper-left end of the main sequence on the Hertzsprung–Russell diagram. Its name comes from its color and size. (p. 436)

**blueshift** Motion-induced changes in the observed wavelength from a source that is moving toward us. Relative approaching motion between the object and the observer causes the wavelength to appear shorter (and hence bluer) than if there were no motion at all.

**blue straggler** Star found on the main sequence of the Hertzsprung–Russell diagram, but which should already have evolved off the main sequence, given its location on the diagram; thought to have formed from mergers of lower mass stars. (p. 508)

**blue supergiant** The very largest of the large, hot, bright stars at the uppermost-left end of the main sequence on the Hertzsprung–Russell diagram. (p. 436)

**Bohr model** First theory of the hydrogen atom to explain the observed spectral lines. This model rests on three ideas: that there is a state of lowest energy for the electron, that there is a maximum energy beyond which the electron is no longer bound to the nucleus, and that within these two energies the electron can only exist in certain energy levels. (p. 84)

**Bok globule** Dense, compact cloud of interstellar dust and gas on its way to forming one or more stars.

**boson** Particle that exerts or mediates forces between elementary particles in quantum physics. (p. 695)

**bound trajectory** Path of an object with launch speed too low to escape the gravitational pull of a planet.

**brown dwarf** Fragments of collapsing gas and dust that did not contain enough mass to initiate core nuclear fusion. Such an object is then frozen somewhere along its pre-main-sequence contraction phase, continually cooling into a compact dark object. Because of their small size and low temperature, brown dwarfs are extremely difficult to detect observationally. (pp. 378, 478)

**brown oval** Feature of Jupiter's atmosphere that appears only at latitudes near 20° N, this structure is a long-lived hole in the clouds that allows us to look down into Jupiter's lower atmosphere. (p. 272)

## C

**C ring** One of three Saturnian rings visible from Earth. The C ring lies closest to the planet and is relatively thin compared to the A and B rings. (p. 298)

**caldera** Crater that forms at the summit of a volcano.

**capture theory (Moon)** Theory suggesting that the Moon formed far from Earth but was later captured by it.

**carbonaceous asteroid** The darkest, or least reflective, type of asteroid, containing large amounts of carbon.

**carbon-based molecule** Molecule containing atoms of carbon.

**carbon-detonation supernova** See Type I supernova. (p. 527)

**cascade** Process of deexcitation in which an excited electron moves down through energy states one at a time.

**Cassegrain telescope** A type of reflecting telescope in which incoming light hits the primary mirror and is then reflected upward toward the prime focus, where a secondary mirror reflects the light back down through a small hole in the main mirror into a detector or eyepiece. (p. 103)

**Cassini division** A relatively empty gap in Saturn's ring system, discovered in 1675 by Giovanni Cassini. It is now known to contain a number of thin ringlets. (p. 298)

**cataclysmic variable** Collective name for novae and supernovae.

**catalyst** Something that causes or helps a reaction to occur, but is not itself consumed as part of the reaction.

**catastrophic theory** A theory that invokes statistically unlikely accidental events to account for observations.

**celestial equator** The projection of Earth's equator onto the celestial sphere. (p. 13)

**celestial mechanics** Study of the motions of bodies, such as planets and stars, that interact via gravity.

**celestial pole** Projection of Earth's North or South pole onto the celestial sphere. (p. 12)

**celestial sphere** Imaginary sphere surrounding Earth to which all objects in the sky were once considered to be attached. (p. 12)

**Celsius** Temperature scale in which the freezing point of water is 0 degrees and the boiling point of water is 100 degrees.

**center of mass** The "average" position in space of a collection of massive bodies, weighted by their masses. For an isolated system this point moves with constant velocity, according to Newtonian mechanics. (p. 53)

**centigrade** See Celsius.

**centripetal force** (literally "center seeking") Force directed toward the center of a body's orbit.

**centroid** Average position of the material in an object; in spectroscopy, the center of a spectral line.

**Cepheid variable** Star whose luminosity varies in a characteristic way, with a rapid rise in brightness followed by a slower decline. The period of a Cepheid variable star is related to its luminosity, so a determination of this period can be used to obtain an estimate of the star's distance. (p. 581)

**Chandrasekhar mass** Maximum possible mass of a white dwarf.

**chaotic rotation** Unpredictable tumbling motion that nonspherical bodies in eccentric orbits, such as Saturn's satellite Hyperion, can exhibit. No amount of observation of an object rotating chaotically will ever show a well-defined period.

**charge-coupled device (CCD)** An electronic device used for data acquisition; composed of many tiny pixels, each of which records a buildup of charge to measure the amount of light striking it. (p. 109)

**chemical bond** Force holding atoms together to form a molecule.

**chemosynthesis** Analog of photosynthesis that operates in total darkness.

**chromatic aberration** The tendency for a lens to focus red and blue light differently, causing images to become blurred.

**chromosphere** The Sun's lower atmosphere, lying just above the visible atmosphere. (p. 390)

**circumnavigation** Traveling all the way around an object.

**cirrus** High-level clouds composed of ice or methane crystals.

**closed universe** Geometry that the universe as a whole would have if the density of matter is above the critical value. A closed universe is finite in extent and has no edge, like the surface of a sphere. It has enough mass to stop the present expansion and will eventually collapse. (p. 676)

**CNO cycle** Chain of reactions that converts hydrogen into helium using carbon, nitrogen, and oxygen as catalysts.

**cocoon nebula** Bright infrared source in which a surrounding cloud of gas and dust absorb ultraviolet radiation from a hot star and reemits it in the infrared.

**cold dark matter** Class of dark-matter candidates made up of very heavy particles, possibly formed in the very early universe. (p. 706)

**collecting area** The total area of a telescope capable of capturing incoming radiation. The larger the telescope, the greater its collecting area, and the fainter the objects it can detect. (p. 105)

**collisional broadening** Broadening of spectral lines due to collisions between atoms, most often seen in dense gases.

**color index** A convenient method of quantifying a star's color by comparing its apparent brightness as measured through different filters. If the star's radiation is well described by a blackbody spectrum, the ratio of its blue intensity (B) to its visual intensity (V) is a measure of the object's surface temperature.

**color-magnitude diagram** A way of plotting stellar properties, in which absolute magnitude is plotted against color index. (p. 434)

**coma** An effect occurring during the formation of an off-axis image in a telescope. Stars whose light enters the telescope at a large angle acquire comet-like tails on their images. The brightest part of a comet, often referred to as the "head." (p. 346)

**comet** A small body, composed mainly of ice and dust, in an elliptical orbit about the Sun. As it comes close to the Sun, some of its material is vaporized to form a gaseous head and extended tail. (p. 345)

**common envelope** Outer layer of gas in a contact binary.

**comparative planetology** The systematic study of the similarities and differences among the planets, with the goal of obtaining deeper insight into how the solar system formed and has evolved in time. (p. 137)

**composition** The mixture of atoms and molecules that make up an object.

**condensation nuclei** Dust grains in the interstellar medium which act as seeds around which other material can cluster. The presence of dust was very important in causing matter to clump during the formation of the solar system. (p. 148)

**condensation theory** Currently favored model of solar system formation which combines features of the old nebular theory with new information about interstellar dust grains, which acted as condensation nuclei. (p. 148)

**conjunction** Orbital configuration in which a planet lies in the same direction as the Sun, as seen from Earth.

**conservation of angular momentum** See law of conservation of angular momentum. (p. 149)

**conservation of mass and energy** See law of conservation of mass and energy.

**constellation** A human grouping of stars in the night sky into a recognizable pattern. (p. 10)

**constituents** See composition.

**continental drift** The movement of the continents around Earth's surface.

**continuous spectrum** Spectrum in which the radiation is distributed over all frequencies, not just a few specific frequency ranges. A prime example is the blackbody radiation emitted by a hot, dense body. (p. 80)

**convection** Churning motion resulting from the constant upwelling of warm fluid and the concurrent downward flow of cooler material to take its place. (p. 163)

**convection zone** Region of the Sun's interior, lying just below the surface, where the material of the Sun is in constant convection motion. This region extends into the solar interior to a depth of about 20,000 km. (p. 390)

**co-orbital satellites** Satellites sharing the same orbit around a planet.

**Copernican principle** The removal of Earth from any position of cosmic significance.

**Copernican revolution** The realization, toward the end of the sixteenth century, that Earth is not at the center of the universe. (p. 39)

**core** The central region of Earth, surrounded by the mantle. (p. 162); The central region of any planet or star. (p. 390)

**core-accretion theory** Theory that the jovian planets formed when icy protoplanetary cores became massive enough to capture gas directly from the solar nebula. See gravitational instability theory. (p. 152)

**core-collapse supernova** See Type II supernova. (p. 526)

**core hydrogen burning** The energy burning stage for main-sequence stars, in which the helium is produced by hydrogen fusion in

the central region of the star. A typical star spends up to 90 percent of its lifetime in hydrostatic equilibrium brought about by the balance between gravity and the energy generated by core hydrogen burning. (p. 496)

**cornea** (eye) The curved transparent layer covering the front part of the eye. (p. 390)

**corona** One of numerous large, roughly circular regions on the surface of Venus, thought to have been caused by upwelling mantle material causing the planet's crust to bulge outward (plural, *coronae*) (p. 226); The tenuous outer atmosphere of the Sun, which lies just above the chromosphere and, at great distances, turns into the solar wind. (p. 390)

**coronal hole** Vast region of the Sun's atmosphere where the density of matter is about 10 times lower than average. The gas there streams freely into space at high speeds, escaping the Sun completely. (p. 407)

**coronal mass ejection** Giant magnetic "bubble" of ionized gas that separates from the rest of the solar atmosphere and escapes into interplanetary space. (p. 406)

**corpuscular theory** Early particle theory of light.

**cosmic density parameter** Ratio of the universe's actual density to the critical value corresponding to zero curvature.

**cosmic distance scale** Collection of indirect distance-measurement techniques that astronomers use to measure distances in the universe. (p. 24)

**cosmic evolution** The collection of the seven major phases of the history of the universe—namely particulate, galactic, stellar, planetary, chemical, biological, and cultural evolution. (p. 716)

**cosmic microwave background** The almost perfectly isotropic radio signal that is the electromagnetic remnant of the Big Bang. (p. 683)

**cosmic ray** Very energetic subatomic particle arriving at Earth from elsewhere in the Galaxy.

**cosmological constant** Quantity originally introduced by Einstein into general relativity to make his equations describe a static universe. Now one of several candidates for the repulsive "dark energy" force responsible for the observed cosmic acceleration. (p. 679)

**cosmological distance** Distance comparable to the scale of the universe.

**cosmological principle** Two assumptions that make up the basis of cosmology, namely that the universe is homogeneous and isotropic on sufficiently large scales. (p. 669)

**cosmological redshift** The component of the redshift of an object that is due only to the Hubble flow of the universe. (p. 620)

**cosmology** The study of the structure and evolution of the entire universe. (p. 668)

**cosmos** The universe.

**coudé focus** Focus produced far from the telescope using a series of mirrors. Allows the use of heavy and/or finely tuned equipment to analyze the image.

**crater** Bowl-shaped depression on the surface of a planet or moon, resulting from a collision with interplanetary debris. (p. 192)

**crescent** Appearance of the Moon (or a planet) when less than half of the body's hemisphere is visible from Earth.

**crest** Maximum departure of a wave above its undisturbed state.

**critical density** The cosmic density corresponding to the dividing line between a universe that recollapses and one that expands forever. (p. 674)

**critical universe** Universe in which the density of matter is exactly equal to the critical density. The universe is infinite in extent and has zero curvature. The expansion will continue forever, but will approach an expansion speed of zero. (p. 676)

**crust** Layer of Earth which contains the solid continents and the seafloor. (p. 162)

**C-type asteroid** See carbonaceous asteroid.

**cultural evolution** Change in the ideas and behavior of a society over time.

**current sheet** Flat sheet on Jupiter's magnetic equator where most of the charged particles in the magnetosphere lie due to the planet's rapid rotation.

## D

**D ring** Collection of very faint, thin rings, extending from the inner edge of the C ring down nearly to the cloud tops of Saturn. This region contains so few particles that it is completely invisible from Earth. (p. 301)

**dark dust cloud** A large cloud, often many parsecs across, which contains gas and dust in a ratio of about 10:1 gas atoms for every dust particle. Typical densities are a few tens or hundreds of millions of particles per cubic meter. (p. 459)

**dark energy** Generic name given to the unknown cosmic force field thought to be responsible for the observed acceleration of the Hubble expansion. (p. 679)

**dark halo** Region of a galaxy beyond the visible halo where dark matter is thought to reside. (p. 596)

**dark matter** Term used to describe the mass in galaxies and clusters whose existence we infer from rotation curves and other techniques, but that has not been confirmed by observations at any electromagnetic wavelength. (p. 597)

**dark matter particle** Particle undetectable at any electromagnetic wavelength, but can be inferred from its gravitational influence.

**daughter/fission theory** Theory suggesting that the Moon originated out of Earth.

**declination** Celestial coordinate used to measure latitude above or below the celestial equator on the celestial sphere.

**decoupling** Event in the early universe when atoms first formed, after which photons could propagate freely through space. (p. 699)

**deferent** A construct of the geocentric model of the solar system which was needed to explain observed planetary motions. A deferent is a large circle encircling Earth, on which an epicycle moves. (p. 38)

**degree** Unit of angular measure. There are 360 degrees in one complete circle.

**density** A measure of the compactness of the matter within an object, computed by dividing the mass of the object by its volume. Units are kilograms per cubic meter ( $\text{kg}/\text{m}^3$ ), or grams per cubic centimeter ( $\text{g}/\text{cm}^3$ ). (p. 139)

**detached binary** Binary system where each star lies within its respective Roche lobe.

**detector noise** Readings produced by an instrument even when it is not observing anything; produced by the electronic components within the detector itself.

**deuterium** A form of hydrogen with an extra neutron in its nucleus.

**deuterium bottleneck** Period in the early universe between the start of deuterium production and the time when the universe was cool enough for deuterium to survive.

**deuteron** An isotope of hydrogen in which a neutron is bound to the proton in the nucleus. Often called "heavy hydrogen" because of the extra mass of the neutron. (p. 411)

**differential rotation** The tendency for a gaseous sphere, such as a jovian planet or the Sun, to rotate at a different rate at the equator than at the poles. More generally, a condition where the angular speed varies with location within an object. (p. 267)

**differentiation** Variation in the density and composition of a body, such as Earth, with low-density material on the surface and higher density material in the core. (p. 170)

**diffraction** The ability of waves to bend around corners. The diffraction of light establishes its wave nature. (p. 67)

**diffraction grating** Sheet of transparent material with many closely spaced parallel lines ruled on it, designed to separate white light into a spectrum.

**diffraction-limited resolution** Theoretical resolution that a telescope can have due to diffraction of light at the telescope's aperture. Depends on the wavelength of radiation and the diameter of the telescope's mirror. (p. 107)

**direct motion** See prograde motion.

**distance modulus** Difference between the apparent and absolute magnitude of an object; equivalent to distance, by the inverse-square law.

**diurnal motion** Apparent daily motion of the stars, caused by Earth's rotation.

**DNA** Deoxyribonucleic acid, the molecule that carries genetic information and determines the characteristics of a living organism.

**Doppler effect** Any motion-induced change in the observed wavelength (or frequency) of a wave. (p. 73)

**double-line spectroscopic binary** Binary system in which spectral lines of both stars can be distinguished and seen to shift back and forth as the stars orbit one another.

**double-star system** System containing two stars in orbit around one another.

**Drake equation** Expression that gives an estimate of the probability that intelligence exists elsewhere in the Galaxy, based on a number of supposedly necessary conditions for intelligent life to develop. (p. 724)

**dust grain** An interstellar dust particle, roughly  $10^{-7}$  m in size, comparable to the wavelength of visible light. (p. 450)

**dust lane** A lane of dark, obscuring interstellar dust in an emission nebula or galaxy. (p. 455)

**dust tail** The component of a comet's tail that is composed of dust particles. (p. 346)

**dwarf** Any star with radius comparable to, or smaller than, that of the Sun (including the Sun itself). (p. 434)

**dwarf elliptical** Elliptical galaxy as small as 1 kiloparsec across, containing only a few million stars.

**dwarf galaxy** Small galaxy containing a few million stars.

**dwarf irregular** Small irregular galaxy containing only a few million stars.

**dwarf planet** A body that orbits the Sun and is massive enough that its own gravity has caused its shape to be approximately spherical, but which is insufficiently massive to have cleared other bodies from “the neighborhood” of its orbit. (p. 357)

**dynamo theory** Theory that explains planetary and stellar magnetic fields in terms of rotating, conducting material flowing in an object’s interior. (p. 182)

## E

**E ring** A faint ring, well outside the main ring system of Saturn, which was discovered by *Voyager* and is thought to be associated with volcanism on the moon Enceladus. (p. 301)

**Earth-crossing asteroid** An asteroid whose orbit crosses that of Earth. Earth-crossing asteroids are also called Apollo asteroids, after the first asteroid of this type discovered. (p. 343)

**earthquake** A sudden dislocation of rocky material near Earth’s surface.

**eccentricity** A measure of the flatness of an ellipse, equal to the distance between the two foci divided by the length of the major axis. (p. 45)

**eclipse** Event during which one body passes in front of another, so that the light from the occulted body is blocked. (p. 19)

**eclipse season** Time of the year when the Moon lies in the same plane as Earth and Sun, so that eclipses are possible. (p. 22)

**eclipse year** Time interval between successive orbital configurations in which the line of nodes of the Moon’s orbit points toward the Sun.

**eclipsing binary** Rare binary-star system that is aligned in such a way that from Earth we observe one star pass in front of the other, eclipsing the other star. (p. 441)

**ecliptic** The apparent path of the Sun, relative to the stars on the celestial sphere, over the course of a year. (p. 14)

**effective temperature** Temperature of a blackbody of the same radius and luminosity as a given star or planet.

**ejecta (planetary)** Material thrown outward by a meteoroid impact.

**ejecta (stellar)** Material thrown into space by a nova or supernova.

**electric field** A field extending outward in all directions from a charged particle, such as a proton or an electron. The electric field determines the electric force exerted by the particle on all other charged particles in the universe; the strength of the electric field decreases with increasing distance from the charge according to an inverse-square law. (p. 63)

**electromagnetic energy** Energy carried in the form of rapidly fluctuating electric and magnetic fields.

**electromagnetic force** Force (electric or magnetic) exerted between any two charged particles. (p. 413)

**electromagnetic radiation** Another term for light, electromagnetic radiation transfers energy and information from one place to another. (p. 60)

**electromagnetic spectrum** The complete range of electromagnetic radiation, from radio waves to gamma rays, including the visible spectrum. All types of electromagnetic radiation are basically the same phenomenon, differing only by wavelength, and all move at the speed of light. (p. 65)

**electromagnetism** The union of electricity and magnetism, which do not exist as independent quantities but are in reality two aspects of a single physical phenomenon. (p. 64)

**electron** An elementary particle with a negative electric charge; one of the components of the atom. (p. 63)

**electron degeneracy pressure** The pressure produced by the resistance of electrons to further compression once they are squeezed to the point of contact. (p. 500)

**electrostatic force** Force between electrically charged objects.

**electroweak force** Unification of the weak electromagnetic forces. (p. 413)

**element** Matter made up of one particular atom. The number of protons in the nucleus of the atom determines which element it represents. (p. 90)

**elementary particle** Technically, a particle that cannot be subdivided into component parts; however, the term is also often used to refer to particles such as protons and neutrons, which are themselves made up of quarks.

**ellipse** Geometric figure resembling an elongated circle. An ellipse is characterized by its degree of flatness, or eccentricity, and the length of its long axis. In general, bound orbits of objects moving under gravity are elliptical. (p. 45)

**elliptical galaxy** Category of galaxy in which the stars are distributed in an elliptical shape on the sky, ranging from highly elongated to nearly circular in appearance. (p. 611)

**elongation** Angular distance between a planet and the Sun.

**emission line** Bright line in a specific location of the spectrum of radiating material, corresponding to emission of light at a certain frequency. A heated gas in a glass container produces emission lines in its spectrum. (p. 80)

**emission nebula** A glowing cloud of hot interstellar gas. The gas glows as a result of one or more nearby young stars which ionize the gas. Since the gas is mostly hydrogen, the emitted radiation falls predominantly in the red region of the spectrum, because of the hydrogen-alpha emission line. (p. 453)

**emission spectrum** The pattern of spectral emission lines produced by an element. Each element has its own unique emission spectrum. (p. 81)

**empirical** Discovery based on observational evidence (rather than from theory).

**Encke gap** A small gap in Saturn’s A ring. (p. 298)

**energy flux** Energy per unit area per unit time radiated by a star (or recorded by a detector).

**epicycle** A construct of the geocentric model of the solar system which was necessary to explain observed planetary motions. Each

planet rides on a small epicycle whose center in turn rides on a larger circle (the deferent). (p. 38)

**epoch of inflation** Short period of unchecked cosmic expansion early in the history of the universe. During inflation, the universe swelled in size by a factor of about  $10^{50}$ . (p. 702)

**equinox** See vernal equinox, autumnal equinox.

**equivalence principle** The principle that there is no experimental way to distinguish between a gravitational field and an accelerated frame of reference. (p. 560)

**escape speed** The speed necessary for one object to escape the gravitational pull of another. Anything that moves away from a gravitating body with more than the escape speed will never return. (p. 54)

**euclidean geometry** Geometry of flat space.

**event horizon** Imaginary spherical surface surrounding a collapsing star, with radius equal to the Schwarzschild radius, within which no event can be seen, heard, or known about by an outside observer. (p. 559)

**evolutionary theory** A theory which explains observations in a series of gradual steps, explainable in terms of well-established physical principles.

**evolutionary track** A graphical representation of a star's life as a path on the Hertzsprung–Russell diagram. (p. 475)

**excited state** State of an atom when one of its electrons is in a higher energy orbital than the ground state. Atoms can become excited by absorbing a photon of a specific energy, or by colliding with a nearby atom. (p. 85)

**extinction** The dimming of starlight as it passes through the interstellar medium. (p. 450)

**extrasolar planet** Planet orbiting a star other than the Sun. (p. 368)

**extremophilic** Adjective describing organisms that can survive in very harsh environments. (p. 723)

**eyepiece** Secondary lens through which an observer views an image. This lens is often chosen to magnify the image.

## F

**F ring** Faint narrow outer ring of Saturn, discovered by *Pioneer 11* in 1979. The F ring lies just inside the Roche limit of Saturn and was found by *Voyager 1* to be made up of several ring strands apparently braided together. (p. 303)

**Fahrenheit** Temperature scale in which the freezing point of water is 32 degrees and the boiling point of water is 212 degrees.

**false vacuum** Region of the universe that remained in the “unified” state after the strong and electroweak forces separated; one possible cause of cosmic inflation at very early times.

**fault line** Dislocation on a planet's surface, often indicating the boundary between two plates.

**field line** Imaginary line indicating the direction of an electric or magnetic field.

**fireball** Large meteor that burns up brightly and sometimes explosively in Earth's atmosphere.

**firmament** Old-fashioned term for the heavens (i.e., the sky).

**fission** See nuclear fission. (p. 525)

**flare** Explosive event occurring in or near an active region on the Sun. (p. 406)

**flatness problem** One of two conceptual problems with the standard Big Bang model, which is that there is no natural way to explain why the density of the universe is so close to the critical density. (p. 701)

**fluidized ejecta** The ejecta blankets around some Martian craters, which apparently indicate that the ejected material was liquid at the time the crater formed.

**fluorescence** Phenomenon where an atom absorbs energy, then radiates photons of lower energy as it cascades back to the ground state; in astronomy, often produced as ultraviolet photons from a hot young star that are absorbed by a neutral gas, causing some of the gas atoms to become excited and give off an optical (red) glow.

**flyby** Unbound trajectory of a spacecraft around a planet or other body.

**focal length** Distance from a mirror or the center of a lens to the focus.

**focus** One of two special points within an ellipse, whose separation from each other indicates the eccentricity. In a bound orbit, planets orbit in ellipses with the Sun at one focus. (p. 45)

**forbidden line** A spectral line seen in emission nebulae, but not seen in laboratory experiments because, under laboratory conditions, collisions kick the electron in question into some other state before emission can occur.

**force** Action on an object that causes its momentum to change. The rate at which the momentum changes is numerically equal to the force. (p. 49)

**fragmentation** The breaking up of a large object into many smaller pieces (for example, as the result of high-speed collisions between planetesimals and protoplanets in the early solar system). (p. 150)

**Fraunhofer lines** The collection of over 600 absorption lines in the spectrum of the Sun, first categorized by Joseph Fraunhofer in 1812.

**frequency** The number of wave crests passing any given point in a unit time. (p. 61)

**full** When the full hemisphere of the Moon or a planet can be seen from Earth.

**full Moon** Phase of the Moon in which it appears as a complete circular disk in the sky.

**fusion** See nuclear fusion.

## G

**G ring** Faint, narrow ring of Saturn, discovered by *Pioneer 11* and lying just outside the F ring. (p. 301)

**galactic bulge** Thick distribution of warm gas and stars around the center of a galaxy. (p. 578)

**galactic cannibalism** A galaxy merger in which a larger galaxy consumes a smaller one.

**galactic center** The center of the Milky Way, or any other galaxy. The point about which the disk of a spiral galaxy rotates. (p. 585)



**galactic disk** Flattened region of gas and dust that bisects the galactic halo in a spiral galaxy. This is the region of active star formation. (p. 578)

**galactic epoch** Period from 100 million to 3 billion years after the Big Bang when large agglomerations of matter (galaxies and galaxy clusters) formed and grew.

**galactic habitable zone** Region of a galaxy in which conditions are conducive to the development of life.

**galactic halo** Region of a galaxy extending far above and below the galactic disk, where globular clusters and other old stars reside. (p. 578)

**galactic nucleus** Small, central, high-density region of a galaxy. Almost all the radiation from active galaxies is generated within the nucleus. (p. 601)

**galactic rotation curve** Plot of rotation speed versus distance from the center of a galaxy.

**Galactic year** Time taken for objects at the distance of the Sun (about 8 kpc) to orbit the center of the Galaxy, roughly 225 million years.

**galaxy** Gravitationally bound collection of a large number of stars. The Sun is a star in the Milky Way Galaxy. (p. 578)

**galaxy cluster** A collection of galaxies held together by their mutual gravitational attraction. (p. 617)

**Galilean moons** The four brightest and largest moons of Jupiter (Io, Europa, Ganymede, Callisto), named after Galileo Galilei, the seventeenth-century astronomer who first observed them. (p. 266)

**Galilean satellites** See Galilean moons.

**gamma ray** Region of the electromagnetic spectrum, far beyond the visible spectrum, corresponding to radiation of very high frequency and very short wavelength. (p. 60)

**gamma-ray burst** Object that radiates tremendous amounts of energy in the form of gamma rays, possibly due to the collision and merger of two neutron stars initially in orbit around one another. (p. 552)

**gamma-ray spectrograph** Spectrograph designed to work at gamma-ray wavelengths. Used to map the abundances of certain elements on the Moon and Mars.

**gaseous** Composed of gas.

**gas-exchange experiment** Experiment to look for life on Mars. A nutrient broth was offered to Martian soil specimens. If there were life in the soil, gases would be created as the broth was digested.

**gene** Sequence of nucleotide bases in the DNA molecule that determines the characteristics of a living organism.

**general theory of relativity** Theory proposed by Einstein to incorporate gravity into the framework of special relativity. (p. 560)

**geocentric model** A model of the solar system that holds that Earth is at the center of the universe and all other bodies are in orbit around it. The earliest theories of the solar system were geocentric. (p. 38)

**giant** A star with a radius between 10 and 100 times that of the Sun. (p. 434)

**giant elliptical** Elliptical galaxy up to a few megaparsecs across, containing trillions of stars.

**gibbous** Appearance of the Moon (or a planet) when more than half (but not all) of the body's hemisphere is visible from Earth.

**globular cluster** Tightly bound, roughly spherical collection of hundreds of thousands, and sometimes millions, of stars spanning about 50 parsecs. Globular clusters are distributed in the halos around the Milky Way and other galaxies. (p. 487)

**gluon** Particle that exerts or mediates the strong force in quantum physics.

**gradient** Rate of change of some quantity (e.g., temperature or composition) with respect to location in space.

**Grand Unified Theories (GUTs)** Class of theories describing the behavior of the single force that results from unification of the strong, weak, and electromagnetic forces in the early universe. (pp. 694, 695)

**granite** Igneous rock, containing silicon and aluminum, that makes up most of Earth's crust.

**granulation** Mottled appearance of the solar surface, caused by rising (hot) and falling (cool) material in convective cells just below the photosphere. (p. 396)

**gravitational force** Force exerted on one body by another due to the effect of gravity. The force is directly proportional to the masses of both bodies involved and inversely proportional to the square of the distance between them. (pp. 51, 413)

**gravitational instability theory** Theory that the jovian planets formed directly from the solar nebula via instabilities in the gas leading to gravitational contraction. See core-accretion theory. (p. 153)

**gravitational lensing** The effect induced on the image of a distant object by a massive foreground object. Light from the distant object is bent into two or more separate images. (p. 598)

**gravitational radiation** Radiation resulting from rapid changes in a body's gravitational field.

**gravitational redshift** A prediction of Einstein's general theory of relativity. Photons lose energy as they escape the gravitational field of a massive object. Because a photon's energy is proportional to its frequency, a photon that loses energy suffers a decrease in frequency, which corresponds to an increase, or redshift, in wavelength. (p. 562)

**graviton** Particle carrying the gravitational field in theories attempting to unify gravity and quantum mechanics.

**gravity** The attractive effect that any massive object has on all other massive objects. The greater the mass of the object, the stronger its gravitational pull. (p. 51)

**gravity assist** Using gravity to change the flight path of a satellite or spacecraft. (p. 142)

**gravity wave** Gravitational counterpart of an electromagnetic wave.

**great attractor** A huge accumulation of mass in the relatively nearby universe (within about 200 Mpc of the Milky Way).

**Great Dark Spot** Prominent storm system in the atmosphere of Neptune observed by *Voyager 2*, near the equator of the planet. The system was comparable in size to Earth. (p. 325)

**Great Red Spot** A large, high-pressure, long-lived storm system visible in the atmosphere of Jupiter. The Red Spot is roughly twice the size of Earth. (p. 268)

**great wall** Extended sheet of galaxies measuring at least 200 megaparsecs across; one of the largest known structures in the universe.

**greenhouse effect** The partial trapping of solar radiation by a planetary atmosphere, similar to the trapping of heat in a greenhouse. (p. 166)

**greenhouse gas** Gas (such as carbon dioxide or water vapor) that efficiently absorbs infrared radiation. (p. 166)

**ground state** The lowest energy state that an electron can have within an atom. (p. 84)

**GUT epoch** Period when gravity separated from the other three forces of nature.

**gyroscope** System of rotating wheels that allows a spacecraft to maintain a fixed orientation in space.

## H

**habitable zone** Three-dimensional zone of comfortable temperature (corresponding to liquid water) that surrounds every star. (p. 381)

**half-life** The amount of time it takes for half of the initial amount of a radioactive substance to decay into something else. (p. 172)

**Hayashi track** Evolutionary track followed by a protostar during the final pre-main-sequence phase before nuclear fusion begins.

**heat** Thermal energy, the energy of an object due to the random motion of its component atoms or molecules.

**heat death** End point of a bound universe, in which all matter and life are destined to be incinerated.

**heavy element** In astronomical terms, any element heavier than hydrogen and helium.

**heliocentric model** A model of the solar system that is centered on the Sun, with Earth in motion about the Sun. (p. 39)

**helioseismology** The study of conditions far below the Sun's surface through the analysis of internal "sound" waves that repeatedly cross the solar interior. (p. 393)

**helium-burning shell** Shell of burning helium gas surrounding a nonburning stellar core of carbon ash.

**helium capture** The formation of heavy elements by the capture of a helium nucleus. For example, carbon can form heavier elements by fusion with other carbon nuclei, but it is much more likely to occur by helium capture, which requires less energy. (p. 535)

**helium flash** An explosive event in the post-main-sequence evolution of a low-mass star. When helium fusion begins in a dense stellar core, the burning is explosive in nature. It continues until the energy released is enough to expand the core, at which point the star achieves stable equilibrium again. (p. 500)

**helium precipitation** Mechanism responsible for the low abundance of helium of Saturn's atmosphere. Helium condenses in the upper layers to form a mist, which rains down toward Saturn's interior, just as water vapor forms into rain in the atmosphere of Earth. (p. 297)

**helium shell flash** Condition in which the helium-burning shell in the core of a star cannot respond to rapidly changing conditions within it, leading to a sudden temperature rise and a dramatic increase in nuclear reaction rates.

**Hertzsprung–Russell (H–R) diagram** A plot of luminosity against temperature (or spectral class) for a group of stars. (p. 434)

**HI region** Region of space containing primarily neutral hydrogen. (p. 456)

**hierarchical merging** Widely accepted galaxy-formation scenario in which galaxies formed as relatively small objects in the early universe

and subsequently collided and merged to form the large galaxies observed today. The present Hubble type of a galaxy depends on the sequence of merger events in the galaxy's past. (p. 645)

**high-energy astronomy** Astronomy using X- or gamma-ray radiation rather than optical radiation.

**high-energy telescope** Telescope designed to detect X- and gamma-ray radiation. (p. 125)

**highlands** Relatively light-colored regions on the surface of the Moon that are elevated several kilometers above the maria. Also called terrae. (p. 192)

**high-mass star** Star with a mass more than 8 times that of the Sun; progenitor of a neutron star or black hole.

**HII region** Region of space containing primarily ionized hydrogen. (p. 456)

**homogeneity** Assumed property of the universe such that the number of galaxies in an imaginary large cube of the universe is the same no matter where in the universe the cube is placed. More generally, "the same everywhere." (p. 669)

**horizon problem** One of two conceptual problems with the standard Big Bang model, which is that some regions of the universe that have very similar properties are too far apart to have exchanged information within the age of the universe. (p. 700)

**horizontal branch** Region of the Hertzsprung–Russell diagram where post-main-sequence stars again reach hydrostatic equilibrium. At this point, the star is burning helium in its core and fusing hydrogen in a shell surrounding the core. (p. 500)

**hot dark matter** A class of candidates for the dark matter in the universe, composed of lightweight particles such as neutrinos, much less massive than the electron. (p. 706)

**hot Jupiter** A massive, gaseous planet orbiting very close to its parent star. (p. 375)

**hot longitudes (Mercury)** Two opposite points on Mercury's equator where the Sun is directly overhead at perihelion.

**Hubble classification scheme** Method of classifying galaxies according to their appearance, developed by Edwin Hubble. (p. 608)

**Hubble diagram** Plot of galactic recession velocity versus distance; evidence for an expanding universe.

**Hubble flow** Universal recession described by the Hubble diagram and quantified by Hubble's Law.

**Hubble's constant** The constant of proportionality that gives the relation between recessional velocity and distance in Hubble's law. (p. 620)

**Hubble's law** Law that relates the observed velocity of recession of a galaxy to its distance from us. The velocity of recession of a galaxy is directly proportional to its distance away. (p. 620)

**hydrocarbon** Molecule consisting solely of hydrogen and carbon.

**hydrogen envelope** An invisible sheath of gas engulfing the coma of a comet, usually distorted by the solar wind and extending across millions of kilometers of space. (p. 346)

**hydrogen shell burning** Fusion of hydrogen in a shell that is driven by contraction and heating of the helium core. Once hydrogen is depleted in the core of a star, hydrogen burning stops and the core

contracts due to gravity, causing the temperature to rise, heating the surrounding layers of hydrogen in the star, and increasing the burning rate there. (p. 497)

**hydrosphere** Layer of Earth that contains the liquid oceans and accounts for roughly 70 percent of Earth's total surface area. (p. 162)

**hydrostatic equilibrium** Condition in a star or other fluid body in which gravity's inward pull is exactly balanced by internal forces due to pressure. (p. 392)

**hyperbola** Curve formed when a plane intersects a cone at a small angle to the axis of the cone.

**hypernova** Explosion where a massive star undergoes core collapse and forms a black hole and a gamma-ray burst. *See* supernova.

## I

**igneous** Type of rock formed from molten material.

**image** The optical representation of an object produced when the object is reflected or refracted by a mirror or lens. (p. 100)

**impact theory (Moon)** Combination of the capture and daughter theories, suggesting that the Moon formed after an impact which dislodged some of Earth's mantle and placed it in orbit. (p. 210)

**inertia** The tendency of an object to continue moving at the same speed and in the same direction, unless acted upon by a force. (p. 50)

**inferior conjunction** Orbital configuration in which an inferior planet (Mercury or Venus) lies closest to Earth.

**inflation** *See* epoch of inflation.

**infrared** Region of the electromagnetic spectrum just outside the visible range, corresponding to light of a slightly longer wavelength than red light. (p. 60)

**infrared telescope** Telescope designed to detect infrared radiation. Many such telescopes are designed to be lightweight so that they can be carried above (most of) Earth's atmosphere by balloons, airplanes, or satellites. (p. 121)

**infrared waves** Electromagnetic radiation with wavelength in the infrared part of the spectrum.

**inhomogeneity** Deviation from perfectly uniform density; in cosmology, inhomogeneities in the universe are ultimately due to quantum fluctuations before inflation.

**inner core** The central part of Earth's core, thought to be solid, and composed mainly of nickel and iron. (p. 169)

**instability strip** Region of the Hertzsprung–Russell diagram where pulsating post-main-sequence stars are found.

**intensity** A basic property of electromagnetic radiation that specifies the amount or strength of the radiation.

**intercloud medium** Superheated bubbles of hot gas extending far into interstellar space.

**intercrater plains** Regions on the surface of Mercury that do not show extensive cratering but are relatively smooth. (p. 194)

**interference** The ability of two or more waves to interact in such a way that they either reinforce or cancel each other. (p. 67)

**interferometer** Collection of two or more telescopes working together as a team, observing the same object at the same time and at

the same wavelength. The effective diameter of an interferometer is equal to the distance between its outermost telescopes. (p. 118)

**interferometry** Technique in widespread use to dramatically improve the resolution of radio and infrared maps. Several telescopes observe the object simultaneously, and a computer analyzes how the signals interfere with each other. (p. 118)

**intermediate-mass black hole** Black hole having a mass 100–1000 times greater than the mass of the Sun.

**interplanetary matter** Matter in the solar system that is not part of a planet or moon—cosmic “debris.” (p. 143)

**interstellar dust** Microscopic dust grains that populate space between the stars, having their origins in the ejected matter of long-dead stars.

**interstellar gas cloud** A large cloud of gas found in the space among the stars.

**interstellar medium** The matter between stars, composed of two components, gas and dust, intermixed throughout all of space. (p. 450)

**intrinsic variable** Star that varies in appearance due to internal processes (rather than, say, interaction with another star).

**inverse-square law** The law that a field follows if its strength decreases with the square of the distance. Fields that follow the inverse-square law decrease rapidly in strength as the distance increases, but never quite reach zero. (p. 51)

**Io plasma torus** Doughnut-shaped region of energetic ionized particles, emitted by the volcanoes on Jupiter's moon Io and swept up by Jupiter's magnetic field.

**ion** An atom that has lost one or more of its electrons.

**ionization state** Term describing the number of electrons missing from an atom: I refers to a neutral atom, II refers to an atom missing one electron, and so on.

**ionized** State of an atom or molecule that has lost one or more of its electrons. (p. 84)

**ionosphere** Layer in Earth's atmosphere above about 80 km where the atmosphere is significantly ionized and conducts electricity. (p. 163)

**ion tail** Thin stream of ionized gas that is pushed away from the head of a comet by the solar wind. It extends directly away from the Sun. Often referred to as a plasma tail. (p. 346)

**irregular galaxy** A galaxy that does not fit into any of the other major categories in the Hubble classification scheme. (p. 612)

**isotopes** Nuclei containing the same number of protons but different numbers of neutrons. Most elements can exist in several isotopic forms. A common example of an isotope is deuterium, which differs from normal hydrogen by the presence of an extra neutron in the nucleus. (p. 412)

**isotropic** Looking the same in every direction. Often applied to the universe as part of the cosmological principle. (p. 669)

## J

**jet stream** Relatively strong winds in the upper atmosphere channeled into a narrow stream by a planet's rotation. Normally refers to a horizontal, high-altitude wind.

**joule** The SI unit of energy.

**jovian planet** One of the four giant outer planets of the solar system, resembling Jupiter in physical and chemical composition. (p. 140)

## K

**Kelvin scale** Temperature scale in which absolute zero is at 0 K; a change of 1 kelvin is the same as a change of 1 degree Celsius.

**Kelvin-Helmholtz contraction phase** Evolutionary track followed by a star during the protostar phase.

**Kepler's laws of planetary motion** Three laws, based on precise observations of the motions of the planets by Tycho Brahe, that summarize the motions of the planets about the Sun.

**kinetic energy** Energy of an object due to its motion.

**Kirchhoff's laws** Three rules governing the formation of different types of spectra. (p. 83)

**Kirkwood gaps** Gaps in the spacings of orbital semimajor axes of asteroids in the asteroid belt, produced by dynamical resonances with nearby planets, especially Jupiter. (p. 345)

**Kuiper belt** A region in the plane of the solar system outside the orbit of Neptune where most short-period comets are thought to originate. (p. 352)

**Kuiper-belt object** Small icy body orbiting in the Kuiper belt. (p. 353)

## L

**labeled-release experiment** Experiment to look for life on Mars. Radioactive carbon compounds were added to Martian soil specimens. Scientists looked for indications that the carbon had been eaten or inhaled.

**Lagrangian point** One of five special points in the plane of two massive bodies orbiting one another, where a third body of negligible mass can remain in equilibrium. (p. 314)

**lander** Spacecraft that lands on the object it is studying.

**laser ranging** Method of determining the distance to an object by firing a laser beam at it and measuring the time taken for the light to return.

**lava dome** Volcanic formation formed when lava oozes out of fissures in a planet's surface, creating the dome, and then withdraws, causing the crust to crack and subside. (p. 225)

**law of conservation of angular momentum** A fundamental law of physics that states that the total angular momentum of a system always remains constant in any physical process. This law ensures that a spinning gas cloud must spin faster as it contracts.

**law of conservation of mass and energy** A fundamental law of modern physics that states that the sum of mass and energy must always remain constant in any physical process. In fusion reactions, the lost mass is converted into energy, primarily in the form of electromagnetic radiation. (p. 411)

**laws of planetary motion** Three laws derived by Kepler describing the motion of the planets around the Sun. (p. 45)

**lens (eye)** The part of the eye that refracts light onto the retina.

**lens** Optical instrument made of glass or some other transparent material, shaped so that, as a parallel beam of light passes through it, the rays are refracted so as to pass through a single focal point.

**lepton** From the Greek word for *light*, referring in particle physics to low-mass particles such as electrons, muons, and neutrinos that interact via the weak force. (p. 694)

**lepton epoch** Period when the light elementary particles (leptons) were in thermal equilibrium with the cosmic radiation field.

**lidar** Light Detection and Ranging—a device that uses laser-ranging to measure distance.

**light** See electromagnetic radiation.

**light curve** The variation in brightness of a star with time. (p. 441)

**light element** In astronomical terms, hydrogen and helium.

**light-gathering power** Amount of light a telescope can view and focus, proportional to the area of the primary mirror.

**lighthouse model** The leading explanation for pulsars. A small region of the neutron star, near one of the magnetic poles, emits a steady stream of radiation that sweeps past Earth each time the star rotates. The period of the pulses is the star's rotation period. (p. 545)

**light pollution** Unwanted, upward-directed light from streets, homes and businesses, that scatters back to Earth from dust in the air, obscuring the faint objects that astronomers want to observe. (p. 111)

**light-year** The distance that light, moving at a constant speed of 300,000 km/s, travels in one year. One light-year is about 10 trillion kilometers. (p. 6)

**limb** Edge of a lunar, planetary, or solar disk.

**line of nodes** Intersection of the plane of the Moon's orbit with Earth's orbital plane.

**line of sight technique** Method of probing interstellar clouds by observing their effects on the spectra of background stars.

**linear momentum** Tendency of an object to keep moving in a straight line with constant velocity; the product of the object's mass and velocity.

**lithosphere** Earth's crust and a small portion of the upper mantle that make up Earth's plates. This layer of Earth undergoes tectonic activity. (p. 174)

**Local Bubble** The particular low-density intercloud region surrounding the Sun.

**Local Group** The small galaxy cluster that includes the Milky Way Galaxy. (p. 618)

**Local Supercluster** Collection of galaxies and clusters centered on the Virgo Cluster. See also supercluster.

**logarithm** The power to which 10 must be raised to produce a given number.

**logarithmic scale** Scale using the logarithm of a number rather than the number itself; commonly used to compress a large range of data into more manageable form.

**look-back time** Time in the past when an object emitted the radiation we see today.

**Lorentz contraction** Apparent contraction of an object in the direction in which it is moving. (p. 559)

**low-mass star** Star with a mass less than 8 times that of the Sun; progenitor of a white dwarf.

**luminosity** One of the basic properties used to characterize stars, luminosity is defined as the total energy radiated by star each second, at all wavelengths. (p. 391)

**luminosity class** A classification scheme that groups stars according to the width of their spectral lines. For a group of stars with the same temperature, luminosity class differentiates between supergiants, giants, main-sequence stars, and subdwarfs. (p. 438)

**lunar dust** *See* regolith.

**lunar eclipse** Celestial event during which the moon passes through the shadow of Earth, temporarily darkening its surface. (p. 19)

**lunar phase** The appearance of the Moon at different points along its orbit. (p. 19)

**Lyman-alpha forest** Collection of lines in an object's spectra starting at the redshifted wavelength of the object's own Lyman-alpha emission line and extending to shorter wavelengths; produced by gas in galaxies along the line of sight.

## M

**macroscopic** Large enough to be visible by the unaided eye.

**Magellanic Clouds** Two small irregular galaxies that are gravitationally bound to the Milky Way Galaxy. (p. 612)

**magnetic field** Field that accompanies any changing electric field and governs the influence of magnetized objects on one another. (p. 64)

**magnetic poles** Points on a planet where the planetary magnetic field lines intersect the planet's surface vertically.

**magnetism** The presence of a magnetic field.

**magnetometer** Instrument that measures magnetic field strength.

**magnetopause** Boundary between a planet's magnetosphere and the solar wind.

**magnetosphere** A zone of charged particles trapped by a planet's magnetic field, lying above the atmosphere. (p. 162)

**magnitude scale** A system of ranking stars by apparent brightness, developed by the Greek astronomer Hipparchus. Originally, the brightest stars in the sky were categorized as being of first magnitude, while the faintest stars visible to the naked eye were classified as sixth magnitude. The scheme has since been extended to cover stars and galaxies too faint to be seen by the unaided eye. Increasing magnitude means fainter stars, and a difference of five magnitudes corresponds to a factor of 100 in apparent brightness. (p. 426)

**main sequence** Well-defined band on the Hertzsprung–Russell diagram on which most stars are found, running from the top left of the diagram to the bottom right. (p. 435)

**main-sequence turnoff** Special point on the Hertzsprung–Russell diagram for a cluster, indicative of the cluster's age. If all the stars in the cluster are plotted, the lower mass stars will trace out the main sequence up to the point where stars begin to evolve off the main sequence toward the red giant branch. The point where stars are just beginning to evolve off is the main-sequence turnoff. (p. 513)

**major axis** The long axis of an ellipse.

**mantle** Layer of Earth just interior to the crust. (p. 162)

**mare** Relatively dark-colored and smooth region on the surface of the Moon (plural: *maria*). (p. 192)

**marginally bound universe** Universe that will expand forever but at an increasingly slow rate.

**mass** A measure of the total amount of matter contained within an object. (p. 50)

**mass-energy equivalence** Principle that mass and energy are not independent, but can be converted from one to the other according to Einstein's formula  $E = mc^2$ . (p. 410)

**mass function** Relation between the component masses of a single-line spectroscopic binary.

**massive compact halo object (MACHO)** Collective name for "stellar" candidates for dark matter, including brown dwarfs, white dwarfs, and low-mass red dwarfs.

**mass-luminosity relation** The dependence of the luminosity of a main-sequence star on its mass. The luminosity increases roughly as the mass raised to the third power.

**mass-radius relation** The dependence of the radius of a main-sequence star on its mass. The radius rises roughly in proportion to the mass.

**mass transfer** Process by which one star in a binary system transfers matter onto the other.

**matter** Anything having mass.

**matter era** (current) era following the radiation era when the universe is larger and cooler, and matter is the dominant constituent of the universe.

**matter-antimatter annihilation** Reaction in which matter and antimatter annihilate to produce high energy gamma rays. *See also* antiparticle.

**matter-dominated universe** A universe in which the density of matter exceeds the density of radiation. The present-day universe is matter dominated. (p. 691)

**Maunder minimum** Lengthy period of solar inactivity that extended from 1645 to 1715. (p. 405)

**mean solar day** Average length of time from one noon to the next, taken over the course of a year—24 hours.

**medium** Material through which a wave, such as sound, travels.

**mesosphere** Region of Earth's atmosphere lying between the stratosphere and the ionosphere, 50–80 km above Earth's surface. (p. 163)

**Messier object** Member of a list of "fuzzy" objects compiled by astronomer Charles Messier in the eighteenth century.

**metabolism** The daily utilization of food and energy by which organisms stay alive.

**metallic** Composed of metal or metal compounds.

**metamorphic** Rocks created from existing rocks exposed to extremes of temperature or pressure.

**meteor** Bright streak in the sky, often referred to as a "shooting star," resulting from a small piece of interplanetary debris entering Earth's atmosphere and heating air molecules, which emit light as they return to their ground states. (p. 358)

**meteor shower** Event during which many meteors can be seen each hour, caused by the yearly passage of Earth through the debris spread along the orbit of a comet. (p. 359)

**meteorite** Any part of a meteoroid that survives passage through the atmosphere and lands on the surface of Earth. (p. 359)

**meteoroid** Chunk of interplanetary debris prior to encountering Earth's atmosphere. (p. 358)

**meteoroid swarm** Pebble-sized cometary fragments dislodged from the main body, moving in nearly the same orbit as the parent comet. (p. 359)

**microlensing** Gravitational lensing by individual stars in a galaxy. (p. 659)

**micrometeoroid** Relatively small chunk of interplanetary debris ranging in size from a dust particle to a pebble. (p. 359)

**microquasar** Stellar sized source of energetic X and gamma radiation, powered by accretion onto a neutron star or black hole, somewhat like a quasar but on a much smaller scale. (p. 550)

**microsphere** Small droplet of protein-like material that resists dissolution in water.

**midocean ridge** Place where two plates are moving apart, allowing fresh magma to well up. (p. 176)

**Milky Way Galaxy** The spiral galaxy in which the Sun resides. The disk of our Galaxy is visible in the night sky as the faint band of light known as the Milky Way. (p. 578)

**millisecond pulsar** A pulsar whose period indicates that the neutron star is rotating nearly 1000 times each second. The most likely explanation for these rapid rotators is that the neutron star has been spun up by drawing in matter from a companion star. (p. 550)

**molecular cloud** A cold, dense interstellar cloud which contains a high fraction of molecules. It is thought that the relatively high density of dust particles in these clouds plays an important role in the formation and preservation of the molecules. (p. 463)

**molecular cloud complex** Collection of molecular clouds that spans as much as 50 parsecs and may contain enough material to make millions of Sun-size stars. (p. 465)

**molecule** A tightly bound collection of atoms held together by the atoms' electromagnetic fields. Molecules, like atoms, emit and absorb photons at specific wavelengths. (p. 91)

**molten** In liquid form due to high temperatures.

**moon** A small body in orbit around a planet.

**mosaic (photograph)** Composite photograph made up of many smaller images.

**M-type asteroid** Asteroid containing large fractions of nickel and iron.

**multiple star system** Group of two or more stars in orbit around one another.

**muon** A type of lepton (along with the electron and tau).

## N

**naked singularity** A singularity that is not hidden behind an event horizon.

**nanobacterium** Very small bacterium with diameter in the nanometer range.

**nanometer** One billionth of a meter.

**neap tide** The smallest tide, occurring when the Earth–Moon line is perpendicular to the Earth–Sun line at the first and third quarters.

**nebula** General term used for any “fuzzy” patch on the sky, either light or dark. (p. 453)

**nebular theory** One of the earliest models of solar system formation, dating back to Descartes, in which a large cloud of gas began to collapse under its own gravity to form the Sun and planets. (p. 148)

**nebulousity** “Fuzziness,” usually in the context of an extended or gaseous astronomical object.

**neon-oxygen white dwarf** White dwarf formed from a low-mass star with a mass close to the “high-mass” limit, in which neon and oxygen form in the core.

**neutrino** Virtually massless and chargeless particle that is one of the products of fusion reactions in the Sun. Neutrinos move at close to the speed of light, and interact with matter hardly at all. (p. 411)

**neutrino oscillations** Likely solution to the solar neutrino problem, in which the neutrino has a very tiny mass. In this case, the correct number of neutrinos can be produced in the solar core, but on their way to Earth some can “oscillate,” or become transformed into other particles, and thus go undetected. (p. 415)

**neutron** An elementary particle with roughly the same mass as a proton, but which is electrically neutral. Along with protons, neutrons form the nuclei of atoms. (p. 90)

**neutron capture** The primary mechanism by which very massive nuclei are formed in the violent aftermath of a supernova. Instead of fusion of like nuclei, heavy elements are created by the addition of more and more neutrons to existing nuclei. (p. 536)

**neutron degeneracy pressure** Pressure due to the Pauli exclusion principle, arising when neutrons are forced to come into close contact. (p. 544)

**neutron spectrometer** Instrument designed to search for water ice by looking for hydrogen.

**neutron star** A dense ball of neutrons that remains at the core of a star after a supernova explosion has destroyed the rest of the star. Typical neutron stars are about 20 km across, and contain more mass than the Sun. (p. 544)

**neutronization** Process occurring at high densities, in which protons and electrons are crushed together to form neutrons and neutrinos.

**new Moon** Phase of the Moon during which none of the lunar disk is visible.

**Newtonian mechanics** The basic laws of motion, postulated by Newton, which are sufficient to explain and quantify virtually all of the complex dynamical behavior found on Earth and elsewhere in the universe. (p. 49)

**Newtonian telescope** A reflecting telescope in which incoming light is intercepted before it reaches the prime focus and is deflected into an eyepiece at the side of the instrument. (p. 103)

**nodes** Two points on the Moon's orbit when it crosses the ecliptic.

**nonrelativistic** Speed that is much less than the speed of light.

**nonthermal spectrum** Continuous spectrum not well described by a blackbody.

**normal galaxy** Galaxy whose overall energy emission is consistent with the summed light of many stars. (p. 622)

**north celestial pole** Point on the celestial sphere directly above Earth's North Pole.

**Northern and Southern Lights** Colorful displays produced when atmospheric molecules, excited by charged particles from the Van Allen belts, fall back to their ground state.

**nova** A star that suddenly increases in brightness, often by a factor of as much as 10,000, then slowly fades back to its original luminosity. A nova is the result of an explosion on the surface of a white-dwarf star, caused by matter falling onto its surface from the atmosphere of a binary companion. (p. 522)

**nuclear binding energy** Energy that must be supplied to split an atomic nucleus into neutrons and protons.

**nuclear epoch** Period when protons and neutrons fused to form heavier nuclei.

**nuclear fission** Mechanism of energy generation used in nuclear reactors on Earth, in which heavy nuclei are split into lighter ones, releasing energy in the process. (p. 525)

**nuclear fusion** Mechanism of energy generation in the core of the Sun, in which light nuclei are combined, or fused, into heavier ones, releasing energy in the process. (p. 410)

**nuclear reaction** Reaction in which two nuclei combine to form other nuclei, often releasing energy in the process. *See also* fusion.

**nucleotide base** An organic molecule, the building block of genes that pass on hereditary characteristics from one generation of living creatures to the next. (p. 718)

**nucleus** Dense, central region of an atom, containing both protons and neutrons, and orbited by one or more electrons. (p. 84); The solid region of ice and dust that composes the central region of the head of a comet. (p. 346); The dense central core of a galaxy. (p. 346)

## O

**obscuration** Blockage of light by pockets of interstellar dust and gas.

**Olbers's paradox** A thought experiment suggesting that if the universe were homogeneous, infinite, and unchanging, the entire night sky would be as bright as the surface of the Sun. (p. 670)

**Oort cloud** Spherical halo of material surrounding the solar system out to a distance of about 50,000 AU; where most comets reside. (p. 353)

**opacity** A quantity that measures a material's ability to block electromagnetic radiation. Opacity is the opposite of transparency. (p. 66)

**open cluster** Loosely bound collection of tens to hundreds of stars, a few parsecs across, generally found in the plane of the Milky Way. (p. 486)

**open universe** Geometry that the universe would have if the density of matter were less than the critical value. In an open universe there is not enough matter to halt the expansion of the universe. An open universe is infinite in extent. (p. 676)

**opposition** Orbital configuration in which a planet lies in the opposite direction from the Sun, as seen from Earth.

**optical double** Chance superposition in which two stars appear to lie close together but are actually widely separated.

**optical telescope** Telescope designed to observe electromagnetic radiation at optical wavelengths.

**orbital** One of several energy states in which an electron can exist in an atom. (p. 84)

**orbital period** Time taken for a body to complete one full orbit around another.

**orbiter** Spacecraft that orbits an object to make observations.

**organic compound** Chemical compound (molecule) containing a significant fraction of carbon atoms; the basis of living organisms.

**outer core** The outermost part of Earth's core, thought to be liquid and composed mainly of nickel and iron. (p. 169)

**outflow channel** Surface feature on Mars, evidence that liquid water once existed there in great quantity; thought to be the relics of catastrophic flooding about 3 billion years ago. Found only in the equatorial regions of the planet. (p. 245)

**outgassing** Production of atmospheric gases (carbon dioxide, water vapor, methane, and sulphur dioxide) by volcanic activity.

**ozone layer** Layer of Earth's atmosphere at an altitude of 20–50 km where incoming ultraviolet solar radiation is absorbed by oxygen, ozone, and nitrogen in the atmosphere. (p. 164)

## P

**pair production** Process in which two photons of electromagnetic radiation give rise to a particle–antiparticle pair. (p. 691)

**parallax** The apparent motion of a relatively close object with respect to a more distant background as the location of the observer changes. (p. 25)

**parsec** The distance at which a star must lie in order for its measured parallax to be exactly 1 arc second; 1 parsec equals 206,000 AU. (p. 423)

**partial eclipse** Celestial event during which only a part of the occulted body is blocked from view. (p. 20)

**particle** A body with mass but of negligible dimension.

**particle accelerator** Device used to accelerate subatomic particles to relativistic speeds.

**particle–antiparticle pair** Pair of particles (e.g., an electron and a positron) produced by two photons of sufficiently high energy.

**particle detector** Experimental equipment that allows particles and antiparticles to be detected and identified.

**Pauli exclusion principle** A rule of quantum mechanics that prohibits electrons in dense gas from being squeezed too close together.

**penumbra** Portion of the shadow cast by an eclipsing object in which the eclipse is seen as partial. (p. 21); The outer region of a sunspot, surrounding the umbra, which is not as dark and not as cool as the central region. (p. 401)

**perihelion** The closest approach to the Sun of any object in orbit about it.

**period** The time needed for an orbiting body to complete one revolution about another body. (pp. 47, 61)

**period–luminosity relation** A relation between the pulsation period of a Cepheid variable and its absolute brightness. Measurement of the pulsation period allows the distance of the star to be determined. (p. 583)

**permafrost** Layer of permanently frozen water ice thought to lie just under the surface of Mars. (p. 247)

**phase** Appearance of the sunlit face of the Moon at different points along its orbit, as seen from Earth. (p. 18)

**photodisintegration** Process occurring at high temperatures, in which individual photons have enough energy to split a heavy nucleus (e.g., iron) into lighter nuclei.

**photoelectric effect** Emission of an electron from a surface when a photon of electromagnetic radiation is absorbed. (p. 88)

**photoevaporation** Process in which a cloud in the vicinity of a new-born hot star is dispersed by the star's radiation.

**photometer** A device that measures the total amount of light received in all or part of the image. (p. 111)

**photometry** Branch of observational astronomy in which the brightness of a source is measured through each of a set of standard filters. (pp. 111, 429)

**photomicrograph** Photograph taken through a microscope.

**photon** Individual packet of electromagnetic energy that makes up electromagnetic radiation. (p. 85)

**photosphere** The visible surface of the Sun, lying just above the uppermost layer of the Sun's interior, and just below the chromosphere. (p. 390)

**photosynthesis** Process by which plants manufacture carbohydrates and oxygen from carbon dioxide and water using chlorophyll and sunlight as the energy source.

**pixel** One of many tiny picture elements, organized into a two-dimensional array, making up a digital image. (p. 109)

**Planck curve** See blackbody curve.

**Planck epoch** Period from the beginning of the universe to roughly  $10^{-43}$  second, when the laws of physics are not understood.

**Planck's constant** Fundamental physical constant relating the energy of a photon to its radiation frequency (color).

**planet** One of eight major bodies that orbit the Sun, visible to us by reflected sunlight.

**planetary nebula** The ejected envelope of a red-giant star, spread over a volume roughly the size of our solar system. (p. 503)

**planetary ring system** Material organized into thin, flat rings encircling a giant planet, such as Saturn.

**planetary transit** Orbital configuration in which an inferior planet (e.g., Mercury or Venus, as seen from Earth) is observed to pass directly in front of the Sun. (p. 372)

**planetesimal** Term given to objects in the early solar system that had reached the size of small moons, at which point their gravitational fields were strong enough to begin influencing their neighbors. (p. 150)

**plasma** A gas in which the constituent atoms are completely ionized.

**plate tectonics** The motions of regions of Earth's lithosphere, which drift with respect to one another. Also known as continental drift. (p. 174)

**plutino** Kuiper-belt object whose orbital period (like that of Pluto) is in a 3:2 resonance with the orbit of Neptune.

**plutoid** A dwarf planet orbiting beyond Neptune. (p. 357)

**polarity** A measure of the direction of the solar magnetic field in a sunspot. Conventionally, lines coming out of the surface are labeled "S" while those going into the surface are labeled "N." (p. 402)

**polarization** The alignment of the electric fields of emitted photons, which are generally emitted with random orientations. (p. 453)

**polar vortex** Long-lived pattern of circulating winds around the pole of a planet. (p. 230)

**Population I and II stars** Classification scheme for stars based on the abundance of heavy elements. Within the Milky Way, Population I refers to young disk stars and Population II refers to old halo stars.

**positron** Atomic particle with properties identical to those of a negatively charged electron, except for its positive charge. This positron is the antiparticle of the electron. Positrons and electrons annihilate one another when they meet, producing pure energy in the form of gamma rays. (p. 411)

**prebiotic compound** Molecule that can combine with others to form the building blocks of life.

**precession** The slow change in the direction of the rotation axis of a spinning object, caused by some external gravitational influence. (p. 17)

**primary atmosphere** The chemical components that would have formed Earth's atmosphere.

**primary mirror** Mirror placed at the prime focus of a telescope (see prime focus).

**prime focus** The point in a reflecting telescope where the mirror focuses incoming light to a point. (p. 100)

**prime-focus image** Image formed at the prime focus of a telescope.

**primeval fireball** Hot, dense state of the universe at very early times, just after the Big Bang. (p. 670)

**primordial matter** Matter created during the early, hot epochs of the universe.

**primordial nucleosynthesis** The production of elements heavier than hydrogen by nuclear fusion in the high temperatures and densities that existed in the early universe. (p. 697)

**principle of cosmic censorship** A proposition to separate the unexplained physics near a singularity from the rest of the well-behaved universe. The principle states that nature always hides any singularity, such as a black hole, inside an event horizon, which insulates the rest of the universe from seeing it.

**progenitor** "Ancestor" star of a given object; for example, the star that existed before a supernova explosion is the supernova's progenitor.

**prograde motion** Motion across the sky in the eastward direction.

**prominence** Loop or sheet of glowing gas ejected from an active region on the solar surface, which then moves through the inner parts of the corona under the influence of the Sun's magnetic field. (p. 405)

**proper motion** The angular movement of a star across the sky, as seen from Earth, measured in seconds of arc per year. This movement is a result of the star's actual motion through space. (p. 424)

**protein** Molecule made up of amino acids that controls metabolism.



**proton** An elementary particle carrying a positive electric charge, a component of all atomic nuclei. The number of protons in the nucleus of an atom dictates what type of atom it is. (p. 63)

**proton-proton chain** The chain of fusion reactions, leading from hydrogen to helium, that powers main-sequence stars. (p. 412)

**protoplanet** Clump of material, formed in the early stages of solar system formation, that was the forerunner of the planets we see today. (p. 150)

**protostar** Stage in star formation when the interior of a collapsing fragment of gas is sufficiently hot and dense that it becomes opaque to its own radiation. The protostar is the dense region at the center of the fragment. (p. 474)

**protostellar disk** Swirling disk of gas and dust within which a star (and possibly a planetary system) forms. The “solar nebula,” in the case of our Sun. (p. 475)

**protosun** The central accumulation of material in the early stages of solar system formations, the forerunner of the present-day Sun. (p. 150)

**Ptolemaic model** Geocentric solar system model, developed by the second-century astronomer Claudius Ptolemy. It predicted with great accuracy the positions of the then known planets. (p. 38)

**pulsar** Object that emits radiation in the form of rapid pulses with a characteristic pulse period and duration. Charged particles, accelerated by the magnetic field of a rapidly rotating neutron star, flow along the magnetic field lines, producing radiation that beams outward as the star spins on its axis. (p. 545)

**pulsating variable star** A star whose luminosity varies in a predictable, periodic way. (p. 581)

**P-waves** Pressure waves from an earthquake that travel rapidly through liquids and solids.

**pyrolic-release experiment** Experiment to look for life on Mars. Radioactively tagged carbon dioxide was added to Martian soil specimens. Scientists looked for indications that the radioactive material had been absorbed.

## Q

**quantization** The fact that light and matter on small scales behave in a discontinuous manner, and manifest themselves in the form of tiny “packets” of energy, called quanta. (p. 85)

**quantum fluctuation** Temporary random change in the amount of energy at a point in space.

**quantum gravity** Theory combining general relativity with quantum mechanics.

**quantum mechanics** The laws of physics as they apply on atomic scales. (p. 85)

**quark** A fundamental matter particle that interacts via the strong force; basic constituent of protons and neutrons. (p. 694)

**quark epoch** Period when all heavy elementary particles (composed of quarks) were in thermal equilibrium with the cosmic radiation field.

**quarter Moon** Lunar phase in which the Moon appears as a half disk.

**quasar** Starlike radio source with an observed redshift that indicates an extremely large distance from Earth. The brightest nucleus of a distant active galaxy. (p. 629)

**quasar feedback** The idea that some of the energy released by a quasar in an active galactic nucleus heats and ejects the gas in the surrounding galaxy, shutting off the quasar fuel supply and suppressing star formation in the galaxy, connecting the growth of the central black hole to the properties of the host galaxy. (p. 654)

**quasi-stellar object (QSO)** *See* quasar.

**quiescent prominence** Prominence that persists for days or weeks, hovering high above the solar photosphere.

**quiet Sun** The underlying predictable elements of the Sun’s behavior, such as its average photospheric temperature, which do not significantly change in time.

**quintessence** Dark energy candidate that evolves in a way that depends on the density of matter and radiation in the universe, and which may provide a natural explanation for why dark matter seems to be just now emerging as the dominant force in the universe.

## R

**radar** Acronym for radio detection and ranging. Radio waves are bounced off an object, and the time taken for the echo to return indicates its distance. (p. 48)

**radial motion** Motion along a particular line of sight, which induces apparent changes in the wavelength (or frequency) of radiation received.

**radial velocity** Component of a star’s velocity along the line of sight.

**radian** Angular measure equivalent to  $180/\pi = 57.3$  degrees.

**radiant** Constellation from which a meteor shower appears to come.

**radiation** A way in which energy is transferred from place to place in the form of a wave. Light is a form of electromagnetic radiation. (p. 60)

**radiation darkening** The effect of chemical reactions that result when high-energy particles strike the icy surfaces of objects in the outer solar system. The reactions lead to a buildup of a dark layer of material. (p. 329)

**radiation era** The first few thousand years after the Big Bang when the universe was small, dense, and dominated by radiation.

**radiation zone** Region of the Sun’s interior where extremely high temperatures guarantee that the gas is completely ionized. Photons only occasionally interact with electrons, and travel through this region with relative ease. (p. 390)

**radiation-dominated universe** Early epoch in the universe, when the equivalent density of radiation in the cosmos exceeded the density of matter. (p. 691)

**radio** Region of the electromagnetic spectrum corresponding to the longest wavelength radiation. (p. 60)

**radioactivity** The release of energy by rare, heavy elements when their nuclei decay into lighter nuclei. (p. 171)

**radio galaxy** Type of active galaxy that emits most of its energy in the form of long-wavelength radiation. (p. 625)

**radiograph** Image made from observations at radio wavelengths.

**radio lobe** Roundish extended region of radio-emitting gas, lying well beyond the center of a radio galaxy. (p. 625)

**radio telescope** Large instrument designed to detect radiation from space at radio wavelengths. (p. 114)

**radio wave** Electromagnetic radiation with wavelength in the radio part of the spectrum (p. 76)

**radius–luminosity–temperature relationship** A mathematical proportionality, arising from Stefan's law, which allows astronomers to indirectly determine the radius of a star once its luminosity and temperature are known. (p. 433)

**rapid mass transfer** Mass transfer in a binary system that proceeds at a rapid and unstable rate, transferring most of the mass of one star onto the other.

**ray** The path taken by a beam of radiation.

**Rayleigh scattering** Scattering of light by particles in the atmosphere.

**recession velocity** Rate at which two objects are separating from one another.

**recombination** See decoupling. (p. 699)

**recurrent nova** Star that “goes nova” several times over the course of a few decades.

**reddening** Dimming of starlight by interstellar matter, which tends to scatter higher-frequency (blue) components of the radiation more efficiently than the lower-frequency (red) components. (p. 451)

**red dwarf** Small, cool faint star at the lower-right end of the main sequence on the Hertzsprung–Russell diagram. (p. 436)

**red giant** A giant star whose surface temperature is relatively low so that it glows red. (pp. 434)

**red-giant branch** The section of the evolutionary track of a star corresponding to intense hydrogen shell burning, which drives a steady expansion and cooling of the outer envelope of the star. As the star gets larger in radius and its surface temperature cools, it becomes a red giant. (p. 499)

**red-giant region** The upper-right corner of the Hertzsprung–Russell diagram, where red-giant stars are found. (p. 436)

**redshift** Motion-induced change in the wavelength of light emitted from a source moving away from us. The relative recessional motion causes the wave to have an observed wavelength longer (and hence redder) than it would if it were not moving.

**redshift survey** Three-dimensional survey of galaxies, using redshift to determine distance.

**red supergiant** An extremely luminous red star. Often found on the asymptotic-giant branch of the Hertzsprung–Russell diagram. (p. 434)

**reflecting telescope** A telescope which uses a curved mirror to focus light from a distant object. (p. 100)

**reflection nebula** Bluish nebula caused by starlight scattering from dust particles in an interstellar cloud located just off the line of sight between Earth and a bright star. (p. 455)

**refracting telescope** A telescope that uses a lens to gather and focus light from a distant object. (p. 100)

**refraction** The tendency of a wave to bend as it passes from one transparent medium to another. (p. 100)

**regolith** Surface dust on the moon, several tens of meters thick in places, caused by billions of years of meteoritic bombardment.

**relativistic** Speeds comparable to the speed of light.

**relativistic fireball** Leading explanation of a gamma-ray burst in which an expanding region of superhot gas radiates in the gamma-ray part of the spectrum.

**remnant** The object left behind after a supernova explosion. Can refer to (1) the expanding and cooling shell of glowing gas resulting from the event, or (2) the neutron star or black hole that remains at the center of the explosion. (p. 544)

**residual cap** Portion of Martian polar ice caps that remains permanently frozen, undergoing no seasonal variations. (p. 249)

**resonance** Circumstance in which two characteristic times are related in some simple way, for example, an asteroid with an orbital period exactly half that of Jupiter. (p. 198)

**retina (eye)** The back part of the eye onto which light is focused by the lens.

**retrograde motion** Backward, westward loop traced out by a planet with respect to the fixed stars. (p. 37)

**revolution** Orbital motion of one body about another, such as Earth about the Sun. (p. 14)

**revolving** See revolution. (p. 13)

**Riemannian geometry** Geometry of positively curved space (e.g., the surface of a sphere).

**right ascension** Celestial coordinate used to measure longitude on the celestial sphere. The zero point is the position of the Sun at the vernal equinox.

**rille** A ditch on the surface of the Moon where molten lava flowed in the past. (p. 206)

**ring** See planetary ring system.

**ringlet** Narrow region in Saturn's planetary ring system where the density of ring particles is high. *Voyager* discovered that the rings visible from Earth are actually composed of tens of thousands of ringlets. (p. 300)

**Roche limit** Often called the tidal stability limit, the Roche limit gives the distance from a planet at which the tidal force (due to the planet) between adjacent objects exceeds their mutual attraction. Objects within this limit are unlikely to accumulate into larger objects. The rings of Saturn occupy the region within Saturn's Roche limit. (p. 299)

**Roche lobe** An imaginary surface around a star. Each star in a binary system can be pictured as being surrounded by a teardrop-shaped zone of gravitational influence, the Roche lobe. Any material within the Roche lobe of a star can be considered to be part of that star. During evolution, one member of the binary system can expand so that it overflows its own Roche lobe and begins to transfer matter onto the other star. (p. 516)

**rock** Material made predominantly from compounds of silicon and oxygen.

**rock cycle** Process by which surface rock on Earth is continuously redistributed and transformed from one type into another. (p. 174)

**rotation** Spinning motion of a body about an axis. (p. 12)

**rotation curve** Plot of the orbital speed of disk material in a galaxy against its distance from the galactic center. Analysis of rotation curves of spiral galaxies indicates the existence of dark matter. (p. 596)

**R-process** “Rapid” process in which many neutrons are captured by a nucleus during a supernova explosion.

**RR Lyrae variable** Variable star whose luminosity changes in a characteristic way. All RR Lyrae stars have more or less the same average luminosity. (p. 581)

**runaway greenhouse effect** A process in which the heating of a planet leads to an increase in its atmosphere’s ability to retain heat and thus to further heating, causing extreme changes in the temperature of the surface and the composition of the atmosphere. (p. 232)

**runoff channel** River-like surface feature on Mars, evidence that liquid water once existed there in great quantities. They are found in the southern highlands and are thought to have been formed by water that flowed nearly 4 billion years ago. (p. 245)

## S

**S0 galaxy** Galaxy that shows evidence of a thin disk and a bulge, but which has no spiral arms and contains little or no gas. (p. 612)

**Sagittarius A/Sgr A\*** Strong radio source corresponding to the supermassive black hole at the center of the Milky Way. (p. 601)

**Saros cycle** Time interval between successive occurrences of the “same” solar eclipse, equal to 18 years, 11.3 days.

**satellite** A small body orbiting another larger body.

**SB0 galaxy** An S0-type galaxy whose disk shows evidence of a bar. (p. 612)

**scalar field** Class of quantum-mechanical fields in Grand Unified Theories whose interactions with particles determine those particles’ properties. Inflation in the early universe is thought to have occurred when one or more scalar fields temporarily increased in energy above their normal states. (p. 701)

**scarp** Surface feature on Mercury thought to be the result of cooling and shrinking of the crust forming a wrinkle on the face of the planet. (p. 206)

**Schmidt telescope** Type of telescope having a very wide field of view, allowing large areas of the sky to be observed at once.

**Schwarzschild radius** The distance from the center of an object such that, if all the mass were compressed within that region, the escape speed would equal the speed of light. Once a stellar remnant collapses within this radius, light cannot escape and the object is no longer visible. (p. 557)

**scientific method** The set of rules used to guide science, based on the idea that scientific “laws” be continually tested, and modified or replaced if found inadequate. (p. 8)

**scientific notation** Expressing large and small numbers using power-of-10 notation.

**seasonal cap** Portion of Martian polar ice caps that is subject to seasonal variations, growing and shrinking once each Martian year. (p. 249)

**seasons** Changes in average temperature and length of day that result from the tilt of Earth’s (or any planet’s) axis with respect to the plane of its orbit. (p. 16)

**secondary atmosphere** The chemicals that composed Earth’s atmosphere after the planet’s formation, once volcanic activity outgassed chemicals from the interior.

**sedimentary** Type of rock formed from the buildup of sediment.

**seeing** Term used to describe the ease with which good telescopic observations can be made from Earth’s surface, given the blurring effects of atmospheric turbulence. (p. 111)

**seeing disk** Roughly circular region on a detector over which a star’s point-like images is spread, due to atmospheric turbulence. (p. 111)

**seismic wave** A wave that travels outward from the site of an earthquake through Earth. (p. 168)

**seismology** The study of earthquakes and the waves they produce in Earth’s interior.

**seismometer** Equipment designed to detect and measure the strength of earthquakes (or quakes on any other planet).

**selection effect** Observational bias in which a measured property of a collection of objects is due to the way in which the measurement was made, rather than being intrinsic to the objects themselves. (p. 279)

**self-propagating star formation** Mode of star formation in which shock waves produced by the formation and evolution of one generation of stars triggers the formation of the next. (p. 594)

**semi-detached** Binary system where one star lies within its Roche lobe but the other fills its Roche lobe and is transferring matter onto the first star.

**semimajor axis** One-half of the major axis of an ellipse. The semimajor axis is the way in which the size of an ellipse is usually quantified. (p. 45)

**SETI** Acronym for search for extraterrestrial intelligence. (p. 729)

**Seyfert galaxy** Type of active galaxy whose emission comes from a very small region within the nucleus of an otherwise normal-looking spiral system. (p. 624)

**shepherd satellite** Satellite whose gravitational effect on a ring helps preserve the ring’s shape. Examples are two satellites of Saturn, Prometheus and Pandora, whose orbits lie on either side of the F ring. (p. 304)

**shield volcano** A volcano produced by repeated nonexplosive eruptions of lava, creating a gradually sloping, shield-shaped low dome. Often contains a caldera at its summit. (p. 225)

**shock wave** Wave of matter, which may be generated by a newborn star or supernova, that pushes material outward into the surrounding molecular cloud. The material tends to pile up, forming a rapidly moving shell of dense gas. (p. 484)

**short-period comet** Comet with an orbital period of less than 200 years.

**SI** Système International, the international system of metric units used to define mass, length, time, etc.

**sidereal day** The time needed between successive risings of a given star. (p. 13)

**sidereal month** Time required for the Moon to complete one trip around the celestial sphere. (p. 18)

**sidereal year** The time required for the constellations to complete one cycle around the sky and return to their starting points, as seen

from a given point on Earth. Earth's orbital period around the Sun is one sidereal year. (p. 17)

**single-line spectroscopic binary** Binary system in which one star is too faint for its spectrum to be distinguished, so only the spectrum of the brighter star can be seen to shift back and forth as the stars orbit one another.

**singularity** A point in the universe where the density of matter and the gravitational field are infinite, such as at the center of a black hole. (p. 563)

**sister/coformation theory (Moon)** Theory suggesting that the Moon formed as a separate object close to Earth.

**soft landing** Use of rockets, parachutes, or packaging to break the fall of a space probe as it lands on a planet.

**solar constant** The amount of solar energy reaching Earth's atmosphere per unit area per unit time, approximately  $1400 \text{ W/m}^2$ . (p. 390)

**solar core** The region at the center of the Sun, with a radius of nearly 200,000 km, where powerful nuclear reactions generate the Sun's energy output.

**solar cycle** The 22-year period that is needed for both the average number of spots and the Sun's magnetic polarity to repeat themselves. The Sun's polarity reverses on each new 11-year sunspot cycle. (p. 404)

**solar day** The period of time between the instant when the Sun is directly overhead (i.e., noon) to the next time it is directly overhead. (p. 13)

**solar eclipse** Celestial event during which the new Moon passes directly between Earth and the Sun, temporarily blocking the Sun's light. (p. 20)

**solar interior** The region of the Sun between the solar core and the photosphere.

**solar maximum** Point of the sunspot cycle during which many spots are seen. They are generally confined to regions in each hemisphere, between about 15 and 20 degrees latitude.

**solar minimum** Point of the sunspot cycle during which only a few spots are seen. They are generally confined to narrow regions in each hemisphere at about 25–30 degrees latitude.

**solar nebula** The swirling gas surrounding the early Sun during the epoch of solar system formation, also referred to as the primitive solar system. (p. 148)

**solar neutrino problem** The discrepancy between the theoretically predicted flux of neutrinos streaming from the Sun as a result of fusion reactions in the core and the flux that is actually observed. The observed number of neutrinos is only about half the predicted number. (p. 414)

**solar system** The Sun and all the bodies that orbit it—Mercury, Venus, Earth, Mars, Jupiter, Saturn, Uranus, Neptune, their moons, the asteroids, the comets, and trans-Neptunian objects. (p. 136)

**solar wind** An outward flow of fast-moving charged particles from the Sun. (p. 390)

**solstice** See summer solstice, winter solstice.

**sotropic** Assumed property of the universe such that the universe looks the same in every direction.

**south celestial pole** Point on the celestial sphere directly above Earth's South Pole. (p. 12)

**spacetime** Single entity combining space and time in special and general relativity. (p. 558)

**spatial resolution** The dimension of the smallest detail that can be seen in an image.

**special theory of relativity** Theory proposed by Einstein to deal with the preferred status of the speed of light. (p. 557)

**speckle interferometry** Technique whereby many short-exposure images of a star are combined to make a high-resolution map of the star's surface.

**spectral class** Classification scheme, based on the strength of stellar spectral lines, which is an indication of the temperature of a star. (p. 431)

**spectral window** Wavelength range in which Earth's atmosphere is transparent.

**spectrograph** Instrument used to produce detailed spectra of stars. Usually, a spectrograph records a spectrum on a CCD detector, for computer analysis.

**spectrometer** Instrument used to produce detailed spectra of stars. Usually, a spectrograph records a spectrum on a photographic plate, or more recently, in electronic form on a computer. (p. 111)

**spectroscope** Instrument used to view a light source so that it is split into its component colors. (p. 80)

**spectroscopic binary** A binary-star system which appears as a single star from Earth, but whose spectral lines show back-and-forth Doppler shifts as two stars orbit one another. (p. 440)

**spectroscopic parallax** Method of determining the distance to a star by measuring its temperature and then determining its absolute brightness by comparing with a standard Hertzsprung–Russell diagram. The absolute and apparent brightness of the star give the star's distance from Earth. (p. 437)

**spectroscopy** The study of the way in which atoms absorb and emit electromagnetic radiation. Spectroscopy allows astronomers to determine the chemical composition of stars. (p. 83)

**spectrum** The separation of light into its component colors.

**speed** Distance moved per unit time, independent of direction. See also velocity.

**speed of light** The fastest possible speed, according to the currently known laws of physics. Electromagnetic radiation exists in the form of waves or photons moving at the speed of light. (p. 64)

**spicule** Small solar storm that expels jets of hot matter into the Sun's lower atmosphere.

**spin-orbit resonance** State that a body is said to be in if its rotation period and its orbital period are related in some simple way.

**spiral arm** Distribution of material in a galaxy forming a pinwheel-shaped design, beginning near the galactic center. (p. 592)

**spiral density wave** (1) A wave of matter formed in the plane of planetary rings, similar to ripples on the surface of a pond, which wrap around the rings forming spiral patterns similar to grooves in a record disk. Spiral density waves can lead to the appearance of ringlets. (p. 592) (2) Proposed explanation for the existence of galactic spiral arms, in which coiled waves of gas compression move through the galactic disk, triggering star formation.

**spiral galaxy** Galaxy composed of a flattened, star-forming disk component which may have spiral arms and a large central galactic bulge. (pp. 580, 608)

**spiral nebula** Historical name for spiral galaxies, describing their appearance.

**spring tide** The largest tides, occurring when the Sun, Moon, and Earth are aligned at new and full Moon.

**S-process** “Slow” process in which neutrons are captured by a nucleus; the rate is typically one neutron capture per year.

**standard candle** Any object with an easily recognizable appearance and known luminosity, which can be used in estimating distances. Supernovae, which all have the same peak luminosity (depending on type), are good examples of standard candles and are used to determine distances to other galaxies. (p. 615)

**standard solar model** A self-consistent picture of the Sun, developed by incorporating the important physical processes that are thought to be important in determining the Sun’s internal structure into a computer program. The results of the program are then compared with observations of the Sun and modifications are made to the model. The standard solar model, which enjoys widespread acceptance, is the result of this process. (p. 392)

**star** A glowing ball of gas held together by its own gravity and powered by nuclear fusion in its core. (p. 390)

**starburst galaxy** Galaxy in which a violent event, such as near-collision, has caused an intense episode of star formation in the recent past. (p. 644)

**star cluster** A grouping of anywhere from a dozen to a million stars that formed at the same time from the same cloud of interstellar gas. Stars in clusters are useful to aid our understanding of stellar evolution because, within a given cluster, stars are all roughly the same age and chemical composition and lie at roughly the same distance from Earth. (p. 486)

**Stefan’s law** Relation that gives the total energy emitted per square centimeter of its surface per second by an object of a given temperature. Stefan’s law shows that the energy emitted increases rapidly with an increase in temperature, proportional to the temperature raised to the fourth power. (p. 70)

**stellar epoch** Most recent period when stars, planets, and life have appeared in the universe.

**stellar nucleosynthesis** The formation of heavy elements by the fusion of lighter nuclei in the hearts of stars. Except for hydrogen and helium, all other elements in our universe resulted from stellar nucleosynthesis. (p. 531)

**stellar occultation** The dimming of starlight produced when a solar system object such as a planet, moon, or ring passes directly in front of a star. (p. 332)

**stratosphere** The portion of Earth’s atmosphere lying above the troposphere, extending up to an altitude of 40–50 km. (p. 163)

**string theory** Theory that interprets all particles and forces in terms of particular modes of vibration of submicroscopic strings. (p. 694)

**strong nuclear force** Short-range force responsible for binding atomic nuclei together. The strongest of the four fundamental forces of nature. (p. 411)

**S-type asteroid** Asteroid made up mainly of silicate or rocky material.

**subatomic particle** Particle smaller than the size of an atomic nucleus.

**subduction zone** Place where two plates meet and one slides under the other. (p. 176)

**subgiant** Star on the subgiant branch of the Hertzsprung–Russell diagram. (p. 498)

**subgiant branch** The section of the evolutionary track of a star corresponding to changes that occur just after hydrogen is depleted in the core, and core hydrogen burning ceases. Shell hydrogen burning heats the outer layers of the star, which causes a general expansion of the stellar envelope. (p. 498)

**sublimation** Process by which element changes from the solid to the gaseous state, without becoming liquid.

**summer solstice** Point on the ecliptic where the Sun is at its northernmost point above the celestial equator, occurring on or near June 21. (p. 15)

**sunspot** An Earth-sized dark blemish found on the surface of the Sun. The dark color of the sunspot indicates that it is a region of lower temperature than its surroundings. (p. 401)

**sunspot cycle** The fairly regular pattern that the number and distribution of sunspots follows, in which the average number of spots reaches a maximum every 11 or so years, then fall off to almost zero. (p. 404)

**supercluster** Grouping of several clusters of galaxies into a larger, but not necessarily gravitationally bound, unit. (p. 656)

**super-Earth** Extrasolar planet with mass between 2 and 10 Earth masses. Both rocky and ocean super-Earths have been detected. (p. 374)

**superforce** An attempt to combine the strong and electroweak forces into one single force.

**supergiant** A star with a radius between 100 and 1000 times that of the Sun. (p. 434)

**supergranulation** Large-scale flow pattern on the surface of the Sun, consisting of cells measuring up to 30,000 km across, thought to be the imprint of large convective cells deep in the solar interior. (p. 397)

**superior conjunction** Orbital configuration in which an inferior planet (Mercury or Venus) lies farthest from Earth (on the opposite side of the Sun).

**supermassive black hole** Black hole having a mass a million to a billion times greater than the mass of the Sun; usually found in the central nucleus of a galaxy. (p. 569)

**supernova** Explosive death of a star, caused by the sudden onset of nuclear burning (Type I), or an enormously energetic shock wave (Type II). One of the most energetic events of the universe, a supernova may temporarily outshine the rest of the galaxy in which it resides. (p. 526)

**supernova remnant** The scattered glowing remains from a supernova that occurred in the past. The Crab Nebula is one of the best-studied supernova remnants. (p. 528)

**supersymmetric relic** Massive particle that should have been created in the Big Bang if supersymmetry is correct.

**supersymmetry** Fundamental symmetry of the universe in which all particles that are acted on by forces (e.g., quarks and electrons) are paired with particles that transmit forces (e.g., gluons and photons). It predicts the existence of supersymmetric partners for all known elementary particles. (p. 694)

**surface gravity** The acceleration due to gravity at the surface of a star or planet.

**S-waves** Shear waves from an earthquake, which can travel only through solid material and move more slowly than P-waves.

**synchronous orbit** State of an object when its period of rotations is exactly equal to its average orbital period. The Moon is in a synchronous orbit, and so presents the same face toward Earth at all times. (p. 195)

**synchrotron radiation** Type of nonthermal radiation produced by high-speed charged particles, such as electrons, as they are accelerated in a strong magnetic field. (p. 633)

**synodic month** Time required for the Moon to complete a full cycle of phases. (p. 19)

## T

**T Tauri star** Protostar in the late stages of formation, often exhibiting violent surface activity. T Tauri stars have been observed to brighten noticeably in a short period of time, consistent with the idea of rapid evolution during this final phase of stellar formation. (p. 476)

**tail** Component of a comet that consists of material streaming away from the main body, sometimes spanning hundreds of millions of kilometers. May be composed of dust or ionized gases. (p. 345)

**tau** A type of lepton (along with the electron and muon).

**tectonic fracture** Crack on a planet's surface, in particular on the surface of Mars, caused by internal geological activity.

**telescope** Instrument used to capture as many photons as possible from a given region of the sky and concentrate them into a focused beam for analysis. (p. 100)

**temperature** A measure of the amount of heat in an object, and an indication of the speed of the particles that comprise it. (p. 68)

**tenuous** Thin, having low density.

**terminator** The line separating night from day on the surface of the Moon or a planet.

**terrae** See highlands.

**terrestrial planet** One of the four innermost planets of the solar system, resembling Earth in general physical and chemical properties. (p. 140)

**theoretical model** An attempt to construct a mathematical explanation of a physical process or phenomenon, within the assumptions and confines of a given theory. In addition to providing an explanation of the observed facts, the model generally makes new predictions that can be tested by further observation or experimentation. (p. 8)

**theories of relativity** Einstein's theories, on which much of modern physics rests. Two essential facts of the theory are that nothing can travel faster than the speed of light and that everything, including light, is affected by gravity.

**theory** A framework of ideas and assumptions used to explain some set of observations and make predictions about the real world. (p. 8)

**thermal equilibrium** Condition in which new particle-antiparticle pairs are created from photons at the same rate as pairs annihilate one another to produce new photons.

**thick disk** Region of a spiral galaxy where an intermediate population of stars resides, younger than the halo stars but older than stars in the disk.

**threshold temperature** Critical temperature above which pair production is possible and below which pair production cannot occur.

**tidal bulge** Elongation of Earth caused by the difference between the gravitational force on the side nearest the Moon and the force on the side farthest from the Moon. The long axis of the tidal bulge points toward the Moon. More generally, the deformation of any body produced by the tidal effect of a nearby gravitating object. (p. 183)

**tidal force** The variation in one body's gravitational force from place to place across another body—for example, the variation of the Moon's gravity across Earth. (p. 183)

**tidal locking** Circumstance in which tidal forces have caused a moon to rotate at exactly the same rate at which it revolves around its parent planet, so that the moon always keeps the same face turned toward the planet.

**tidal stability limit** The minimum distance within which a moon can approach a planet before being torn apart by the planet's tidal force.

**tides** Rising and falling motion of terrestrial bodies of water, exhibiting daily, monthly, and yearly cycles. Ocean tides on Earth are caused by the competing gravitational pull of the Moon and Sun on different parts of Earth. (p. 182)

**time dilation** A prediction of the theory of relativity, closely related to the gravitational redshift. To an outside observer, a clock lowered into a strong gravitational field will appear to run slow. (p. 563)

**total eclipse** Celestial event during which one body is completely blocked from view by another. (p. 21)

**transit** See planetary transit.

**transition zone** The region of rapid temperature increases that separates the Sun's chromosphere from the corona. (p. 390)

**trans-Neptunian object** A small, icy body orbiting beyond the orbit of Neptune. Pluto and Eris are the largest currently known examples. (p. 353)

**transverse motion** Motion perpendicular to a particular line of sight, which does not result in Doppler shift in radiation received.

**transverse velocity** Component of star's velocity perpendicular to the line of sight.

**triangulation** Method of determining distance based on the principles of geometry. A distant object is sighted from two well-separated locations. The distance between the two locations and the angle between the line joining them and the line to the distant object are all that are necessary to ascertain the object's distance. (p. 24)

**triple-alpha process** The creation of carbon-12 by the fusion of three helium-4 nuclei (alpha particles). Helium-burning stars occupy a region of the Hertzsprung-Russell diagram known as the horizontal branch. (p. 499)

**triple star system** Three stars that orbit one another, bound together by gravity.

**Trojan asteroid** One of two groups of asteroids which orbit at the same distance from the Sun as Jupiter, 60 degrees ahead of and behind the planet. (p. 344)

**tropical year** The time interval between one vernal equinox and the next. (p. 17)

**troposphere** The portion of Earth's atmosphere from the surface to about 12 km. (p. 163)

**trough** Maximum departure of a wave below its undisturbed state.

**true space motion** True motion of a star, taking into account both its transverse and radial motion according to the Pythagorean theorem.

**Tully-Fisher relation** A relation used to determine the absolute luminosity of a spiral galaxy. The rotational velocity, measured from the broadening of spectral lines, is related to the total mass, and hence the total luminosity. (p. 616)

**turnoff mass** The mass of a star that is just now evolving off the main sequence in a star cluster.

**21-centimeter line** Spectral line in the radio region of the electromagnetic spectrum, associated with a change of spin of the electron in a hydrogen atom.

**21-centimeter radiation** Radio radiation emitted when an electron in the ground state of a hydrogen atom flips its spin to become antiparallel to the spin of the proton in the nucleus. (p. 462)

**twin quasar** Quasar that is seen twice at different locations in the sky due to gravitational lensing.

**Type I supernova** One possible explosive death of a star. A white dwarf in a binary-star system can accrete enough mass that it cannot support its own weight. The star collapses and temperatures become high enough for carbon fusion to occur. Fusion begins throughout the white dwarf almost simultaneously and an explosion results. (p. 527)

**Type II supernova** One possible explosive death of a star, in which the highly evolved stellar core rapidly implodes and then explodes, destroying the surrounding star. (p. 527)

## U

**ultraviolet** Region of the electromagnetic spectrum, just beyond the visible range, corresponding to wavelengths slightly shorter than blue light. (p. 60)

**ultraviolet telescope** A telescope that is designed to collect radiation in the ultraviolet part of the spectrum. Earth's atmosphere is partially opaque to these wavelengths, so ultraviolet telescopes are put on rockets, balloons, and satellites to get high above most or all of the atmosphere. (p. 123)

**umbra** Central region of the shadow cast by an eclipsing body (p. 21); The central region of a sunspot, which is its darkest and coolest part. (p. 401)

**unbound** An orbit which does not stay in a specific region of space, but where an object escapes the gravitational field of another. Typical unbound orbits are hyperbolic in shape. (p. 54)

**unbound trajectory** Path of an object with launch speed high enough that it can escape the gravitational pull of a planet.

**uncompressed density** The density a body would have in the absence of any compression due to its own gravity.

**universe** The totality of all space, time, matter, and energy. (p. 6)

**unstable nucleus** Nucleus that cannot exist indefinitely, but rather must eventually decay into other particles or nuclei.

**upwelling** Upward motion of material having temperature higher than the surrounding medium.

**Urey-Miller experiment** Groundbreaking experiment that simulated conditions on the early Earth and tested for the occurrence of chemical evolution—the production of complex molecules similar to those found in all living things on our planet. (p. 718)

## V

**vacuum energy** Energy associated with empty space when a scalar field temporarily has higher than normal energy. (p. 702)

**Van Allen belt** One of at least two doughnut-shaped regions of magnetically trapped, charged particles high above Earth's atmosphere. (p. 180)

**variable star** A star whose luminosity changes with time. (p. 581)

**velocity** Displacement (distance plus direction) per unit time. *See also* speed. (p. 50)

**vernal equinox** Date on which the Sun crosses the celestial equator moving northward, occurring on or near March 21. (p. 17)

**visible light** The small range of the electromagnetic spectrum that human eyes perceive as light. The visible spectrum ranges from about 400–700 nm, corresponding to blue through red light. (p. 60)

**visible spectrum** The small range of the electromagnetic spectrum that human eyes perceive as light. The visible spectrum ranges from about 4000–7000 angstroms, corresponding to blue through red light.

**visual binary** A binary-star system in which both members are resolvable from Earth. (p. 440)

**void** Large, relatively empty region of the universe around which superclusters and “walls” of galaxies are organized. (p. 657)

**volcano** Upwelling of hot lava from below Earth's crust to the planet's surface.

## W

**wane** (referring to the Moon or a planet) To shrink. The Moon appears to wane, or shrink in size, for two weeks after full Moon.

**warm longitudes (Mercury)** Two opposite points on Mercury's equator where the Sun is directly overhead at aphelion. Cooler than the hot longitudes by 150 degrees.

**water hole** The radio interval between 18 cm and 21 cm, the respective wavelengths at which hydroxyl (OH) and hydrogen (H) radiate, in which intelligent civilizations might conceivably send their communication signals. (p. 731)

**water volcano** Volcano that ejects water (molten ice) rather than lava (molten rock) under cold conditions.

**watt/kilowatt** Unit of power: 1 watt (W) is the emission of 1 joule (J) per second; 1 kilowatt (kW) is 1000 watts.

**wave** A pattern that repeats itself cyclically in both time and space. Waves are characterized by the speed at which they move, their frequency, and their wavelength. (p. 60)

**wave period** The amount of time required for a wave to repeat itself at a specific point in space.

**wave theory of radiation** Description of light as a continuous wave phenomenon, rather than as a stream of individual particles. (p. 65)

**wavelength** The distance from one wave crest (or trough) to the next, at a given instant in time. (p. 61)

**wax** (referring to the Moon or a planet) To grow. The Moon appears to wax, or grow in size, for two weeks after new Moon.

**weak nuclear force** Short-range force, weaker than both electromagnetism and the strong force, but much stronger than gravity; responsible for certain nuclear reactions and radioactive decays. (p. 412)

**weakly interacting massive particle (WIMP)** Class of subatomic particles that might have been produced early in the history of the universe; dark-matter candidates.

**weight** The gravitational force exerted on you by Earth (or the planet on which you happen to be standing). (p. 49)

**weird terrain** A region on the surface of Mercury with oddly rippled features. This feature is thought to be the result of a strong impact that occurred on the other side of the planet, and sent seismic waves traveling around the planet, converging in the weird region.

**white dwarf** A dwarf star with sufficiently high surface temperature that it glows white. (pp. 434, 505)

**white-dwarf region** The bottom-left corner of the Hertzsprung–Russell diagram, where white-dwarf stars are found. (p. 436)

**white oval** Light-colored region near the Great Red Spot in Jupiter's atmosphere. Like the red spot, such regions are apparently rotating storm systems. (p. 272)

**Wien's law** Relation between the wavelength at which a black-body curve peaks and the temperature of the emitter. The peak

wavelength is inversely proportional to the temperature, so the hotter the object, the bluer its radiation. (p. 70)

**winter solstice** Point on the ecliptic where the Sun is at its southernmost point below the celestial equator, occurring on or near December 21. (p. 16)

**wispy terrain** Prominent light-colored streaks on Rhea, one of Saturn's moons.

## X

**X-ray** Region of the electromagnetic spectrum corresponding to radiation of high frequency and short wavelength, far beyond the visible spectrum. (p. 60)

**X-ray burster** X-ray source that radiates thousands of times more energy than our Sun in short bursts lasting only a few seconds. A neutron star in a binary system accretes matter onto its surface until temperatures reach the level needed for hydrogen fusion to occur. The result is a sudden period of rapid nuclear burning and release of energy. (p. 550)

**X-ray nova** Nova explosion detected at X-ray wavelengths.

## Z

**Zeeman effect** Broadening or splitting of spectral lines due to the presence of a magnetic field.

**zero-age main sequence** The region on the Hertzsprung–Russell diagram, as predicted by theoretical models, where stars are located at the onset of nuclear burning in their cores. (p. 477)

**zodiac** The 12 constellations on the celestial sphere through which the Sun appears to pass during the course of a year. (p. 14)

**zonal flow** Alternating regions of westward and eastward flow, roughly symmetrical about the equator of Jupiter, associated with the belts and zones in the planet's atmosphere. (p. 269)

**zone** Bright, high-pressure region in the atmosphere of a jovian planet, where gas flows upward. (p. 268)



*This page intentionally left blank*

# Answers to Check Questions

## Concept Checks and Process of Science Checks

### Chapter 1

**1.1 (p. 9)** A theory can never become proven “fact,” because it can always be invalidated, or forced to change, by a single contradictory observation. However, once a theory’s predictions have been repeatedly verified by experiments over many years, it is often widely regarded as “true.”

**1.2 (p. 13)** (1) Because the celestial sphere provides a natural means of specifying the locations of stars on the sky. Celestial coordinates are directly related to Earth’s orientation in space, but are independent of Earth’s rotation. (2) Distance information is lost. **1.3 (p. 17)** (1) In the Northern Hemisphere, summer occurs when the Sun is near its highest (northernmost) point on the celestial sphere, or, equivalently, when Earth’s North Pole is “tipped” toward the Sun, the days are longest and the Sun is highest in the sky. Winter occurs when the Sun is lowest in the sky (near its southernmost point on the celestial sphere) and the days are shortest. (2) We see different constellations because Earth has moved halfway around its orbit between these seasons and the darkened hemisphere faces an entirely different group of stars. **1.4 (p. 23)** (1) The angular size of the Moon would remain the same, but that of the Sun would be halved, making it easier for the Moon to eclipse the Sun. We would expect to see total or partial eclipses, but no annular ones. (2) If the distance is halved, the angular size of the Sun would double and total eclipses would never be seen, only partial or annular ones. **1.5 (p. 27)** Because astronomical objects are too distant for direct (“measuring tape”) measurements, so we must rely on indirect means and mathematical reasoning.

### Chapter 2

**2.1 (p. 41)** In the geocentric view, retrograde motion is the real backward motion of a planet as it moves on its epicycle. In the heliocentric view, the backward motion is only apparent, caused by Earth “overtaking” the planet in its orbit.

**2.2 (p. 43)** Mainly simplicity and elegance. Both theories made testable predictions, but until Newton developed his laws, neither could explain *why* the planets move as they do. However, Copernicus’s model was much simpler than the Ptolemaic version, which became more and more convoluted as observations improved. **2.3 (p. 47)** Because the laws were derived using only the orbits of the planets Mercury through Saturn, before the outermost planets were known.

**2.4 (p. 49)** Because Kepler determined the overall geometry of the solar system by triangulation using Earth’s orbit as a baseline, so all distances were known only relative to the scale of Earth’s orbit—the astronomical unit. **2.5 (p. 54)** Kepler’s

laws were accurate descriptions of planetary motion based on observations, but they contained no insight into *why* the planets orbited the Sun or why the orbits are as they are. Newtonian mechanics explained the orbits in terms of universal laws, and in addition made detailed predictions about the motion of other bodies in the cosmos—moons, comets, other stars—that were hitherto impossible to make. **2.6 (p. 55)** In the absence of any force, a planet would move in a straight line with constant velocity (Newton’s first law) and therefore tends to move along the tangent to its orbital path. The Sun’s gravity causes the planet to accelerate toward the Sun (Newton’s second law), bending its trajectory into the orbit we observe.

### Chapter 3

**3.1 (p. 65)** Light displays properties that are characteristic of wave motion in other contexts. Light exhibits diffraction as it passes a corner or through a narrow opening, and two light rays can interfere (reinforce or cancel) one another. These effects are small in everyday experience, but are easily measured in the laboratory. They are consistent with the wave theory, and inconsistent with a particle description of radiation. **3.2 (p. 68)** All are electromagnetic radiation and travel at the speed of light. In physical terms, they differ only in frequency (or wavelength), although their effects on our bodies (or our detectors) differ greatly. **3.3 (p. 73)** As the switch is turned and the temperature of the filament rises, the bulb’s brightness increases rapidly, by Stefan’s law, and its color shifts, by Wien’s law, from invisible infrared to red to yellow to white.

**3.4 (p. 75)** According to *More Precisely* 2-2, measuring masses in astronomy usually entails measuring the orbital speed of one object—a companion star, or a planet, perhaps—around another. In most cases, the Doppler effect is an astronomer’s only way of making such measurements.

### Chapter 4

**4.1 (p. 83)** They are characteristic frequencies (wavelengths) at which matter absorbs or emits photons of electromagnetic radiation. They are unique to each atom or molecule, and thus provide a means of identifying the gas producing them. **4.2 (p. 87)** On macroscopic (everyday) scales, light exhibits wave properties, as discussed in Chapter 3. However, on microscopic scales, light displays particle characteristics, with the particles (photons) carrying specific amounts of energy—the photoelectric effect and atomic spectra rely on this fact. Thus scientists have to conclude that light behaves as *both* a wave and a

particle, depending on circumstances.

**4.3 (p. 90)** Spectral lines correspond to transitions between specific orbitals within an atom. The structure of an atom determines the energies of these orbitals, hence the possible transitions, and hence the energies (colors) of the photons involved. **4.4 (p. 92)** In addition to changes involving electron energies, changes involving a molecule’s vibration or rotation can also result in emission or absorption of radiation. **4.5 (p. 95)** Because with few exceptions, spectral analysis is the only way we have of determining the physical conditions—composition, temperature, density, velocity, etc.—in a distant object. Without spectral analysis, astronomers would know next to nothing about the properties of stars and galaxies.

### Chapter 5

**5.1 (p. 105)** Because reflecting telescopes are easier to design, build, and maintain than refracting instruments. **5.2 (p. 108)** The need to gather as much light as possible, and the need to achieve the highest possible angular resolution.

**5.3 (p. 111)** First, photographic plates are inefficient; CCDs are almost always used. Second, because they want to make detailed measurements of brightness, variability, and spectra, which require the use of other, non-imaging detectors.

**5.4 (p. 114)** To reduce or overcome the effects of atmospheric absorption, instruments are placed on high mountains or in space. To compensate for atmospheric turbulence, adaptive optics techniques probe the air above the observing site and adjust the mirror surface accordingly to try to recover the undistorted image. **5.5 (p. 118)** Radio observations allow us to see objects whose visible light is obscured by intervening matter, or which simply do not emit most of their energy in the visible portion of the spectrum. **5.6 (p. 121)** The long wavelength of radio radiation. Astronomers use the largest radio telescopes possible and interferometry, which combines the signals from two or more separate telescopes to create the effect of a single instrument of much larger diameter. **5.7 (p. 125)** Many astronomical objects have very different appearances when viewed at different wavelengths. Some emit only in certain bands, others are partly or wholly obscured by intervening material. By observing at many wavelengths, astronomers gain a much more complete picture of the object under study. **5.8 (p. 128)** Benefits: they are above the atmosphere, so they are unaffected by seeing or absorption; they can also make round-the-clock observations. Drawbacks: cost, smaller size, inaccessibility, vulnerability to damage by radiation and cosmic rays.

## Chapter 6

**6.1 (p. 137)** They provide new tests of our theories of solar system formation, and new examples of the sorts of planetary systems that are possible.

**6.2 (p. 139)** By applying the laws of geometry and Newtonian mechanics to observations made from Earth and (more recently) by visiting spacecraft.

**6.3 (p. 140)** The planets orbit in very nearly the same plane—the ecliptic. Viewed from outside the system, only the orbit of Mercury would deviate noticeably from this plane. **6.4 (p. 143)** Because the two classes of planet differ in almost every physical property—orbit, mass, radius, composition, existence of rings, and number of moons. **6.5 (p. 144)** Because it is thought to be much less evolved than material now found in planets, and hence a better indicator of conditions in the early solar system. **6.6 (p. 152)** The shapes and orientations of the orbits were set by the fact that the planets formed in a spinning disk; the composition differences result from the differing materials available to form planets at various distances from the heat of the Sun. **6.7 (p. 155)** Because the jovian planets could start to form only after the solar nebula had formed, but their formation was terminated when the Sun reached the T Tauri phase and dispersed the disk. **(p. 156)** Yes—if a star formed with a disk of matter around it then the basic processes of condensation and accretion would probably have occurred there too, even if it doesn't have planets like Earth.

## Chapter 7

**7.1 (p. 168)** It raises Earth's average surface temperature above the freezing point of water, which was critical for the development of life on our planet. Should the greenhouse effect continue to strengthen, however, it may conceivably cause catastrophic climate changes on Earth. **7.2 (p. 172)** We could still construct models based on Earth's gravity and other surface properties, but we would have far less detailed direct (from volcanoes) or indirect (from seismic studies following earthquakes) information on our planet's interior. **7.3 (p. 180)** Convection currents in the upper mantle cause portions of Earth's crust—plates—to slide around on the surface. As the plates move and interact, they are responsible for volcanism, earthquakes, the formation of mountain ranges and ocean trenches, and the creation and destruction of oceans and continents. **7.4 (p. 182)** It tells us that the planet has a conducting, liquid core in which the magnetic field is continuously generated. **7.5 (p. 184)** A tidal force is the *variation* in one body's gravitational force from place to place across another. Tidal forces tend to deform a body, rather than causing an overall acceleration, and they decrease proportional to the inverse cube, rather than the inverse square, of the distance.

## Chapter 8

**8.1 (p. 192)** Both bodies have substantially lower escape speeds than Earth, and any atmosphere they may have once had has escaped into space long

ago due to their high temperatures. **8.2 (p. 194)** The maria are younger, denser, and much less heavily cratered than the highlands. **8.3 (p. 201)** In the case of the Moon, Earth's tidal force has slowed the spin to the point where the rotation rate is now exactly synchronized with the Moon's orbital period around Earth. For Mercury, the Sun's tidal force has caused the rotation period to become exactly  $\frac{2}{3}$  of the orbital period, and the rotation axis to be exactly perpendicular to the planet's orbit plane. A synchronous orbit is not possible because of Mercury's eccentric orbit around the Sun. **8.4 (p. 206)** Heavy bombardment long ago created the basins which later filled with lava to form the maria. Subsequent impacts created virtually all the lunar craters, large and small, we see today. Meteoritic bombardment is the main agent of lunar erosion, although the rate is much smaller than the erosion rate on Earth. **8.5 (p. 208)** They are thought to have formed when the planet's core cooled and contracted, causing the crust to crumple. Faults on Earth are the result of tectonic activity. **8.6 (p. 210)** Generation of a magnetic field is thought to require a rapidly rotating body with a conducting liquid core. Neither the Moon nor Mercury rotates rapidly and, while Mercury's core may still be partly liquid, the Moon's probably is not. **8.7 (p. 211)** The impact theory holds that a collision created the Moon essentially from Earth's mantle, accounting for the composition similarities. If the collision occurred after Earth had already differentiated and the dense material had formed a core, relatively little of this material would have found its way into the newborn Moon.

## Chapter 9

**9.1 (p. 220)** It is very slow and retrograde. The reason is unknown, but may simply be a matter of chance. **9.2 (p. 221)** Because they were observing in the optical and could not see the surface. Their measurements pertained to the upper atmosphere, above the planet's reflective cloud layers. **9.3 (p. 228)** No—they are mostly shield volcanoes, where lava upwells through a "hot spot" in the crust. There appears to be no plate tectonic activity on Venus. **9.4 (p. 232)** Given Venus's other bulk similarities to Earth, the planet might well have had an Earth-like climate. **9.5 (p. 233)** The dynamo model of planetary magnetism implies that both a conducting liquid core and rapid rotation are needed to generate a magnetic field. Venus's rotation is the slowest of any planet in the solar system.

## Chapter 10

**10.1 (p. 239)** Because they occur when Martian opposition happens to coincide approximately with Martian perihelion. Such an alignment happens every 7 Martian synodic years. **10.2 (p. 240)** Mars does have seasons, since its rotation axis is inclined to its orbit plane in much the same way as Earth's is. However, the Martian seasons are also affected by the planet's eccentric orbit. The appearance of the planet

changes seasonally, although the changes have nothing to do with growing cycles, as was once thought. **10.3 (p. 244)** The lowlands are much less heavily cratered, implying that they have been resurfaced by volcanism (or smoothed by erosion) since the highlands formed. **10.4 (p. 249)** Before spacecraft visited Mars, the prevailing scientific view was that Mars was a dry, geologically dead world, unlikely ever to have been suitable for life. However, as more and more data have accumulated from spacecraft in orbit and on the planet's surface, scientists have realized that Mars has had a varied, and much wetter, past than was previously thought. Evidence exists for large quantities of flowing and standing water in the past, and water ice has been found at the poles and under the Martian surface. **10.5 (p. 256)** Some of it may have escaped into space. Of the rest, some exists in the form of permafrost (or perhaps liquid water) below the surface. The rest is contained in the Martian polar caps. **10.6 (p. 259)** Some of it was lost due to the planet's weak gravity, aided by violent meteoritic impacts. The rest became part of the planet's surface rocks, the polar caps, or the subsurface permafrost. **10.7 (p. 260)** The planet's small size, which allowed the planet's internal heat to escape, effectively shutting down the processes that drive mantle convection, volcanism, and plate tectonics. **10.8 (p. 261)** They are much smaller, and orbit much closer to the parent planet. In addition, they seem to have a different formation history—their composition differences from Mars suggest that they were captured by the planet long after they (and Mars) formed; for Earth's Moon, this scenario does not seem to work.

## Chapter 11

**11.1 (p. 268)** The magnetic field is generated by the motion of electrically conducting liquid in the deep interior, and therefore presumably shares the rotation of that region of the planet. **11.2 (p. 274)** Like weather systems on Earth, the belts and zones are regions of high and low pressure and are associated with convective motion. However, unlike storms on Earth, they wrap all the way around the planet because of Jupiter's rapid rotation. In addition, the clouds are arranged in three distinct layers and the bright colors are the result of cloud chemistry unlike anything operating in Earth's atmosphere. Jupiter's spots are somewhat similar to hurricanes on Earth, but they are far larger and longer-lived. **11.3 (p. 277)** By constructing theoretical models constrained by detailed spacecraft observations of the planet's bulk properties. **11.4 (p. 279)** Because Jupiter rotates more rapidly than Earth and because the volume of conducting fluid responsible for the field is much greater. Both these factors result in a much stronger planetary magnetic field. **11.5 (p. 285)** Because liquid water is thought to have played a critical role in the appearance of life on Earth, and is presumed to be similarly important elsewhere in the solar system. **11.6 (p. 287)** Jupiter's gravitational field, via its tidal effect on the moons.

## Chapter 12

**12.1 (p. 293)** Because Earth crossed Saturn's ring plane in 1995. The images from before then see the rings from above, those from later see the rings from below. **12.2 (p. 296)** Saturn's cloud and haze layers are thicker than those on Jupiter because of Saturn's lower gravity, so we usually see only the upper level of the atmosphere even though the same basic features and cloud layers are there. **12.3 (p. 297)** Roughly half of it has precipitated into the planet's interior, reducing the helium fraction in the outer layers (and releasing gravitational energy as it fell). **12.4 (p. 304)** The Roche limit is the radius inside of which a moon will be torn apart by tidal forces. Planetary rings are found inside the Roche limit, (most) moons outside. Orbital resonances between ring particles and moons are responsible for much of the fine structure in the rings. They may also maintain the orbits of shepherd satellites that keep some ring features sharp. **12.5 (p. 310)** It has a thick atmosphere (denser than Earth's), unlike any other moon in the solar system, and liquid methane on its surface. **12.6 (p. 314)** Because they are tidally locked by Saturn's gravity into synchronous orbits, and therefore all have permanently leading and trailing faces that interact differently with the environment around the planet.

## Chapter 13

**13.1 (p. 321)** Uranus's orbit was observed to deviate from a perfect ellipse, leading astronomers to try to compute the mass and location of the body responsible for those discrepancies. That body was Neptune. **13.2 (p. 323)** For unknown reasons, Uranus rotates "on its side"—with its rotation axis almost in the plane of the ecliptic. **13.3 (p. 326)** Neptune lies far from the Sun and is very cold, so the source of the energy for these atmospheric phenomena is not known. Uranus, closer to the Sun but with a similar atmospheric temperature, shows much less activity. **13.4 (p. 328)** In each case, the field is significantly offset from the planet's center and inclined to the rotation axis. **13.5 (p. 332)** Triton shows evidence for surface activity—nitrogen geysers and water volcanoes—that seems to have erased most of its impact craters. **13.6 (p. 334)** Both are thin rings that require shepherd satellites to prevent them from spreading out and dispersing.

## Chapter 14

**14.1 (p. 343)** Similarities: all orbit in the inner solar system, and are solid bodies of generally "terrestrial" composition. Differences: asteroids are much smaller than the terrestrial planets, and their orbits are much less regular. **14.2 (p. 345)** Because most Earth-crossing asteroids will eventually come very close to or even collide with our planet, with potentially catastrophic results. **14.3 (p. 351)** Asteroids are generally rocky. Comets are predominantly made of ice, with some dust and other debris mixed in. **14.4 (p. 353)** Most comets never come close enough to the Sun for us to see

them. **14.5 (p. 358)** Because of its similarities to the known Kuiper belt objects, as well as to the icy moons of the outer planets, it is now regarded as the largest member of the Kuiper belt, rather than as the smallest planet. **14.6 (p. 362)** They are fragments of comets or asteroids orbiting the Sun in interplanetary space. They are important to planetary scientists because they generally consist of ancient material, and contain vital information on conditions in the early solar system.

## Chapter 15

**15.1 (p. 370)** Because it must explain the general "orderly" features of solar system architecture, while still accommodating the fact that there are exceptions to many of them. **15.2 (p. 373)** The two major techniques that have led to the discovery of almost all extrasolar planets are radial velocity measurements, where astronomers detect the back and forth movement of the parent star, and transits, where the planet temporarily blocks the light of the star. We don't have many images because, in almost all cases, the planet is lost in the star's glare. **15.3 (p. 379)** (1) Only planets significantly more massive than Earth have so far been detected; (2) The planets' orbits are often quite eccentric; (3) The "hot Jupiters" orbit very close to their parent star. **15.4 (p. 382)** Hot Jupiters are explained as a consequence of giant planet migration in the nebular disk; eccentric Jupiters could be the result of gravitational interactions between planets or due to the tidal effects of other stars; the lack of Earth-sized planets is thought to be due to observational limitations.

## Chapter 16

**16.1 (p. 392)** When we simply multiply the solar constant by the total area to obtain the solar luminosity, we are implicitly assuming that the same amount of energy reaches every square meter of the large sphere in Figure 16.3. **16.2 (p. 397)** The energy may be carried in the form of (1) radiation, where energy travels in the form of light, and (2) convection, where energy is carried by physical motion of upwelling solar gas. **16.3 (p. 400)** The spectrum shows (1) emission lines of (2) highly ionized elements, implying high temperature. **16.4 (p. 405)** There is a strong field, with a well-defined east-west organization, just below the surface. The field direction in the southern hemisphere is opposite that in the north. However, the details of the fields are very complex. **16.5 (p. 409)** Because high-energy particles from flares and coronal mass ejections can affect Earth's magnetosphere and atmosphere, disrupting communications and power transmission on our planet. On longer time scales, it appears that long-term changes in the solar cycle may also be correlated with climate variations on Earth. **16.6 (p. 414)** Because the Sun shines by nuclear fusion, which converts mass into energy as hydrogen turns into helium. **16.7 (p. 415)** When theory and observation come into conflict, observers must repeat and extend their experiments to check and improve

the results, while theorists must look for errors or omissions in their calculations. In the case of the solar neutrino problem, new observations reinforced the disagreement with the theoretical calculations. It turned out that the calculations were also correct as far as they went, but the theory was found to be incomplete, as it omitted neutrino oscillations, which accounted for the discrepancy and resolved the problem.

## Chapter 17

**17.1 (p. 425)** (1) Because stars are so far away that their parallaxes relative to any baseline on Earth are too small to measure accurately. (2) Because the transverse component must be determined by measuring the star's proper motion, which decreases as the star's distance increases, and is too small to measure for most distant stars. **17.2 (p. 427)** Nothing—we need to know their distances before the luminosities (or absolute magnitudes) can be determined. **17.3 (p. 431)** Because temperature controls which excited states the star's atoms and ions are in, and hence which atomic transitions are possible. **17.4 (p. 434)** Yes, using the radius-luminosity-temperature relationship, but only if we can find a method of determining the luminosity that doesn't depend on the inverse-square law (Sec. 17.3). **17.5 (p. 437)** Because giants are intrinsically very luminous, and can be seen to much greater distances than the more common main-sequence stars or white dwarfs. **17.6 (p. 439)** All stars would be further away, but their measured spectral types and apparent brightnesses would be unchanged, so their luminosities would be greater than previously thought. The main sequence would therefore move vertically upward in the H-R diagram. (We would then use larger luminosities in the method of spectroscopic parallax, so distances inferred by that method would also increase.) **17.7 (p. 444)** We don't—we assume that their masses are the same as similar stars found in binaries.

## Chapter 18

**18.1 (p. 453)** Because the scale of interstellar space is so large that even very low densities can add up to a large amount of obscuring matter along the line of sight to a distant star. **18.2 (p. 459)** Because the UV light is absorbed by hydrogen gas in the surrounding cloud, ionizing it to form the emission nebulae. The red light is H radiation; part of the visible hydrogen spectrum emitted as electrons and protons recombine to form hydrogen atoms. **18.3 (p. 461)** By studying absorption lines caused by atoms and molecules in the clouds, and the general extinction due to dust, it is possible to map out a cloud's properties—so long as enough stars are conveniently located behind it. **18.4 (p. 463)** Because most of the interstellar matter in the galactic disk is made up of atomic hydrogen, and 21-centimeter radiation provides a probe of that gas largely unaffected by interstellar absorption. **18.5 (p. 465)** Because the main constituent, hydrogen, is very hard to observe, so

astronomers must use other molecules as tracers of the cloud's properties.

## Chapter 19

**19.1 (p. 472)** The competing effects of gravity, which tends to make an interstellar cloud collapse, and heat (pressure), which opposes that collapse.

**19.2 (p. 477)** (1) The existence of a photosphere, meaning that the inner part of the cloud becomes opaque to its own radiation, signaling the slowing of the collapse phase. (2) Nuclear fusion in the core and equilibrium between pressure and gravity. **19.3 (p. 478)** No—different parts of the main sequence correspond to stars of different masses. A typical star stays at roughly the same location on the main sequence of most of its lifetime. **19.4 (p. 483)** We assume that we observe objects at many different evolutionary stages, and that the snapshot therefore provides a representative sample of the evolutionary stages that stars go through. **19.5 (p. 486)** Star formation may be triggered by some external event, which might cause several interstellar clouds to start contracting at once. Alternatively, the shock wave produced when an emission nebula forms may be sufficient to send another nearby part of the same cloud into collapse. **19.6 (p. 491)** Because stars form at different rates—high-mass stars reach the main sequence and start disrupting the parent cloud long before lower-mass stars have finished forming.

## Chapter 20

**20.1 (p. 496)** We can't observe a single star evolve, but we can observe large numbers of stars at different stages of their lives, and hence build up an accurate statistical picture of stellar evolutionary tracks. **20.2 (p. 502)** Because the nonburning inner core, unsupported by fusion, begins to shrink, releasing gravitational energy, heating the overlying layers and causing them to burn more vigorously, thus increasing the luminosity. **20.3 (p. 509)** Because the core's contraction is halted by the pressure of degenerate (tightly packed) electrons before it reaches a temperature high enough for the next stage of fusion to begin. In fact, this statement is true whether the "next round" is hydrogen fusion (brown dwarf), helium fusion (helium white dwarf), or carbon fusion (carbon-oxygen white dwarf). **20.4 (p. 512)** Fusion in high-mass stars is not halted by electron degeneracy pressure. Temperatures are always high enough that each new burning stage can start before degeneracy becomes important. Such stars continue to fuse more and more massive nuclei, faster and faster, eventually exploding in a supernova. **20.5 (p. 515)** Because a star cluster gives us a "snapshot" of stars of many different masses, but of the same age and initial composition, allowing us to directly test the predictions of the theory. **20.6 (p. 517)** Because many, if not most, stars are found in binaries, and stars in binaries can follow evolutionary paths quite different from those they would follow if single.

## Chapter 21

**21.1 (p. 523)** No, because it is of low mass and not a member of a binary-star system. **21.2 (p. 526)** Because iron cannot fuse to produce energy. As a result, no further nuclear reactions are possible, and the core's equilibrium cannot be restored.

**21.3 (p. 530)** Because the two types of supernova differ in their spectra and their light curves, making it impossible to explain them in terms of a single phenomenon. **21.4 (p. 538)** Because they are readily formed by helium capture, a process common in evolved stars. Other elements (with masses not multiples of four) had to form via less common reactions involving proton and neutron capture. **21.5 (p. 539)** Because it is responsible for creating and dispersing all the heavy elements out of which we are made. In addition, it may also have played an important role in triggering the collapse of the interstellar cloud from which our solar system formed.

## Chapter 22

**22.1 (p. 545)** No—only Type II supernovae. According to theory, the rebounding central core of the original star in a Type II supernova becomes a neutron star. **22.2 (p. 548)** Because (1) not all supernovae form neutron stars, (2) the pulses are beamed, so not all pulsing neutron stars are visible from Earth, and (3) pulsars spin down and become too faint to observe after a few tens of millions of years. **22.3 (p. 552)** Some X-ray sources are binaries containing accreting neutron stars, which may be in the process of being spun up to form millisecond pulsars. **22.4 (p. 555)** They are energetic bursts of gamma rays, roughly isotropically distributed on the sky, occurring about once per day. They pose a challenge because they are very distant, and hence extremely luminous, but their energy originates in a region less than a few hundred kilometers across. There also appear to be two distinct types, with different energy generation mechanisms. **22.5 (p. 561)** Newton's theory describes gravity as a force produced by a massive object that influences all other massive objects. Einstein's relativity describes gravity as a curvature of space-time produced by a massive object; that curvature then determines the trajectories of all particles—matter or radiation—in the universe. **22.6 (p. 564)** Because the object would appear to take infinitely long to reach the event horizon, and its light would be infinitely redshifted by the time it got there. **22.7 (p. 571)** By observing their gravitational effects on other objects, and from the X-rays emitted as matter plunges toward the event horizon.

## Chapter 23

**23.1 (p. 578)** The Milky Way is the thin plane of the Galactic disk, seen from within. When our line of sight lies in the plane of the Galaxy, we see many stars blurring into a continuous band. In other directions, we see darkness. **23.2 (p. 586)** No, because even the brightest Cepheids are unobservable at distances of more than a kiloparsec or so through the obscuration

of interstellar dust. **23.3 (p. 589)** Because the compositions, ages, and orbits of the two classes of stars are quite different from one another. **23.4 (p. 590)** The halo formed early in our Galaxy's history, before gas and dust had formed a spinning, flattened disk. Disk stars are still forming today. Consequently, halo stars are old and move in more or less random three-dimensional orbits, while the disk contains stars of all ages, all moving in roughly circular orbits around the Galactic center. **23.5 (p. 595)** Because differential rotation would destroy the spiral structure within a few hundred million years. **23.6 (p. 599)** Scientists are unwilling to suggest new physics to explain observations, and competing theories of galactic rotation curves and "missing mass" on large scales do exist. However, despite their reservations, most astronomers have concluded that dark matter, introduced as material with gravity but no interaction with electromagnetic radiation, best fits the observed facts. Several theories exist to explain how dark matter might have been formed in the early universe, and they lead to experimental tests that should some day detect dark matter, if it exists. **23.7 (p. 602)** Observations of rapidly moving stars and gas, and the variability of the radiation emitted suggest the presence of a roughly 4 million-solar-mass black hole.

## Chapter 24

**24.1 (p. 614)** Most galaxies are not large spirals—the most common galaxy types are dwarf ellipticals and dwarf irregulars. **24.2 (p. 619)** Because distance-measurement techniques ultimately rely upon the existence of very bright objects whose luminosities can be inferred by other means. Such objects become increasingly hard to find and calibrate the farther we look out into intergalactic space. **24.3 (p. 621)** It doesn't use the inverse-square law. The other methods all provide a way of determining the luminosity of a distant object, which then is converted to a distance using the inverse-square law. Hubble's law gives a direct connection between redshift and distance. **24.4 (p. 628)** It means that the energy source cannot simply be the summed energy of a huge number of stars—some other mechanism must be at work. **24.5 (p. 629)** When quasars were discovered, they were thought to be faint, relative nearby stars, or starlike objects, although their unusual spectra posed problems for astronomers. Once astronomers realized that their odd spectra actually meant that they had very large redshifts, it became clear that quasars were actually among the most distant—and hence the most luminous—objects in the entire universe. **24.6 (p. 634)** The energy is generated in an accretion disk in the central nucleus of the visible galaxy, then transported by jets out of the galaxy and into the lobes, where it is eventually emitted by the synchrotron process in the form of radio waves.

## Chapter 25

**25.1 (p. 642)** First, that the galaxies are gravitationally bound to the cluster. Second, and

more fundamentally, that the laws of physics as we know them in the solar system—gravity, atomic structure, the Doppler effect—all apply on very large scales, and to systems possibly containing a lot of dark matter. **25.2 (p. 645)** Isolated galaxies do change in time as stars form and evolve, but collisions and mergers between galaxies are the main way in which galaxies grow. The large galaxies we see today are the result of repeated collisions between smaller galaxies in the past. **25.3 (p. 650)** Stars form by collapse and fragmentation of a large interstellar cloud and subsequently evolve largely in isolation. Galaxies form by mergers of smaller objects, and interactions with other galaxies play a major role in their evolution. **25.4 (p. 656)** Probably not. There may be galaxies (especially the smaller ones) that do not harbor supermassive central black holes, and in any case, only galaxies in clusters are likely to experience the encounters that trigger activity. **25.5 (p. 662)** Light traveling from these objects to Earth is influenced by cosmic structure all along the line of sight. Light rays are deflected by the gravitational field of intervening concentrations of mass, and gas along the line of sight produces absorption lines whose redshifts tell us the distance at which each feature formed. The light received on Earth thus gives astronomers a “core sample” through the universe, from which detailed information can be extracted.

## Chapter 26

**26.1 (p. 669)** On very large scales—more than 300 Mpc—the distribution of galaxies seems to be roughly the same everywhere and in all directions. **26.2 (p. 673)** Because, tracing the motion backwards in time, it implies that all galaxies, and in fact everything in the entire universe, were located at a single point at the same instant in the past. **26.3 (p. 675)** The universe can expand forever, in which case we die a cold death in which all activity gradually fades away, or the expansion can stop and the universe will recollapse to a fiery Big Crunch. **26.4 (p. 676)** A low-density

universe has negative curvature, a critical density universe is spatially flat (Euclidean), and a high-density universe has positive curvature (and is finite in extent). **26.5 (p. 680)** There doesn't seem to be enough matter to halt the collapse and, in addition, the observed cosmic acceleration suggests the existence of a large-scale repulsive force in the cosmos that also opposes recollapse. **26.6 (p. 683)** The observed acceleration of the expansion of the universe implies that some nongravitational force must be at work. Dark energy is our current best theory of the cause of that force. In addition, galactic and cosmological observations indicate that the universe is spatially flat, and hence has critical density, but that the density of matter (mostly dark) cannot account for the total. The amount of dark energy needed to account for the cosmic acceleration is in good agreement with the required extra density—about 70 percent of the critical value. **26.7 (p. 685)** At the time of the Big Bang. It is the electromagnetic remnant of the primeval fireball.

## Chapter 27

**27.1 (p. 693)** It means that the total mass-energy density of the universe, which today is made up almost entirely of matter and dark energy, was once comprised almost entirely of radiation. We know this because, going back in time toward the Big Bang, the dark energy density stays constant as the universe contracts, the matter density increases because the volume shrinks, but the radiation density increases even faster because of the cosmological redshift. Thus, at sufficiently early times, radiation was the dominant component of the cosmos. **27.2 (p. 697)** Once the temperature of the expanding, cooling universe dropped below the point where particle-antiparticle pairs could no longer be created from the radiation background, the particles separated out of the radiation field. Particles and antiparticles annihilated one another, and any leftover “frozen out” matter has survived to the present day. **27.3 (p. 700)** Because the amount of

deuterium observed in the universe today implies that the present density of normal matter is at most a few percent of the critical value—much less than the density of dark matter inferred from dynamical studies. **27.4 (p. 704)** Inflation implies that the universe is flat, and hence that the total cosmic density equals the critical value. However, the matter density seems to be only about one third of the critical value, and the density of electromagnetic radiation (the microwave background) is a tiny fraction of the critical density. The remaining density may be in the form of the “dark energy” thought to be powering the accelerating cosmic expansion (Section 27.3). **27.5 (p. 706)** Because it could clump to form the large density fluctuations that would ultimately become galaxies and clusters, without leaving a correspondingly large imprint in the microwave background. After it decoupled from the radiation background, normal matter flowed into the dark matter structures, forming the galaxies we see. **27.6 (p. 710)** They allow us to measure the value of  $\Omega_0$ —and in fact imply that it is very close to 1.

## Chapter 28

**28.1 (p. 722)** The formation of complex molecules from simple ingredients by nonbiological processes has been repeatedly demonstrated, but no living cell or self-replicating molecule has ever been created. **28.2 (p. 724)** Mars remains the most likely site, although Europa and Titan also have properties that might have been conducive to the emergence of living organisms. **28.3 (p. 729)** It breaks a complex problem up into simpler “astronomical,” “biochemical,” “anthropological,” and “cultural” pieces, which can be analyzed separately. It also identifies the types of stars where a search might be most fruitful. **28.4 (p. 732)** It is in the radio part of the spectrum, where Galactic absorption is least, at a region where natural Galactic background “static” is minimized, and in a portion of the spectrum characterized by lines of hydrogen and hydroxyl, both of which would likely have significance to a technological civilization.

# Answers to Self-Test Questions

## Chapter 1

**Multiple Choice:** 1.1 b, 1.2 b, 1.3 d, 1.4 a, 1.5 c, 1.6 a, 1.7 c, 1.8 c, 1.9 a, 1.10 d

**Odd-Numbered Problems:** 1.1 (c) the Moon (384,000 km) 1.3 Scorpio 1.5 (a) 57,300 km; (b)  $3.44 \times 10^6$  km; (c)  $2.06 \times 10^8$  km 1.7 391.

## Chapter 2

**Multiple Choice:** 2.1 a, 2.2 d, 2.3 b, 2.4 c, 2.5 c, 2.6 a, 2.7 c, 2.8 b, 2.9 c, 2.20 a

**Odd-Numbered Problems:** 2.1 (a) 110 km; (b) 44,000 km; (c) 370,000 km 2.3 146 days 2.5  $8.1''$ , if Mercury is at aphelion and Earth is at perihelion at the point of closest approach 2.7 1.7 km/s; 2.4 km/s.

## Chapter 3

**Multiple Choice:** 3.1 a, 3.2 c, 3.3 b, 3.4 b, 3.5 d, 3.6 a, 3.7 b, 3.8 d, 3.9 a, 3.10 b

**Odd-Numbered Problems:** 3.1 1480 m/s 3.3 920 W, assuming my skin temperature is 300 K ( $27^\circ\text{C}$ ) and my surface area is roughly  $2\text{ m}^2$ ; of course, I also absorb energy from my surroundings, so the net energy loss is less—about  $2\sigma(300^4 - 290^4) = 120\text{ W}$  if my surroundings are 10 K cooler than me 3.5 2.9 microns 3.7 300 km/s moving away.

## Chapter 4

**Multiple Choice:** 4.1 c, 4.2 c, 4.3 d, 4.4 b, 4.5 b, 4.6 b, 4.7 b, 4.8 b, 4.9 b, 4.10 d

**Odd-Numbered Problems:** 4.1 2.8 eV, 6.2 eV 4.3 (a) 620 nm; (b) 12,400 nm; (c) 0.25 nm 4.5 the first six Balmer lines, ranging in wavelength from 656 nm (H $\alpha$ ) to 410 nm (H $\zeta$ ) 4.7 122.8 nm.

## Chapter 5

**Multiple Choice:** 5.1 c, 5.2 d, 5.3 d, 5.4 b, 5.5 c, 5.6 c, 5.7 a, 5.8 d, 5.9 b, 5.10 c

**Odd-Numbered Problems:** 5.1 0.29 arc seconds; 6.8 pixels 5.3 6.7 minutes; 1.7 minutes 5.5 15.3 m 5.7 5.5 km; 92 m; 1.8 m.

## Chapter 6

**Multiple Choice:** 6.1 a, 6.2 d, 6.3 a, 6.4 c, 6.5 a, 6.6 b, 6.7 c, 6.8 d, 6.9 c, 6.10 c

**Odd-Numbered Problems:** 6.1 Mercury: 0.31 AU, 0.47 AU; Mars: 1.38 AU, 1.67 AU 6.3  $7 \times 10^{20}$ , 0.012 percent of Earth's mass 6.5 (a) 62.5 years; (b) 9 days 6.7 481,000.

## Chapter 7

**Multiple Choice:** 7.1 b, 7.2 d, 7.3 a, 7.4 c, 7.5 b, 7.6 b, 7.7 c, 7.8 b, 7.9 d, 7.10 b

**Odd-Numbered Problems:** 7.1  $5.0 \times 10^{18}$  kg;  $8.4 \times 10^{-7}$  times Earth's mass 7.3 (a)  $8.4 \times 10^{-3}$ ; (b) 0.16; (c) 0.82; (d) 0.014 7.5 100 Myr 7.7 1.9 billion years.

## Chapter 8

**Multiple Choice:** 8.1 a, 8.2 b, 8.3 b, 8.4 c, 8.5 b, 8.6 a, 8.7 b, 8.8 d, 8.9 a, 8.10 a

**Odd-Numbered Problems:** 8.1 8.5 minutes 8.3  $1.71^\circ$ ,  $1.13^\circ$  8.5 1.7 hours 8.7 need at least  $4.8 \times 10^5$  such craters, requiring at least 4.8 trillion years; rate would have to increase by a factor of about 1000.

## Chapter 9

**Multiple Choice:** 9.1 a, 9.2 c, 9.3 c, 9.4 b, 9.5 b, 9.6 c, 9.7 b, 9.8 c, 9.9 a, 9.10 a

**Odd-Numbered Problems:** 9.1 (a) 35.1 arc seconds; (b) 22.9 arc second; (c) 9.7 arc seconds 9.3 470 seconds, 280 seconds 9.5 yes—the smallest detectable feature would be about 20 km across, much smaller than the largest impact features 9.7 400 km/h, 250 mph.

## Chapter 10

**Multiple Choice:** 10.1 b, 10.2 c, 10.3 b, 10.4 c, 10.5 b, 10.6 d, 10.7 a, 10.8 c, 10.9 b, 10.10 a

**Odd-Numbered Problems:** 10.1  $41^\circ$  10.3 59 lbs, assuming you weigh 150 lbs on Earth 10.5  $3.2 \times 10^{16}$  kg, taking Earth's atmospheric mass to be  $5.0 \times 10^{18}$  kg; 2.8 times the seasonal polar cap mass of  $1.2 \times 10^{16}$  kg 10.7  $1.3 \times 10^{20}$  kg; 3900 times the current atmospheric mass.

## Chapter 11

**Multiple Choice:** 11.1 c, 11.2 b, 11.3 a, 11.4 c, 11.5 a, 11.6 d, 11.7 b, 11.8 a, 11.9 a, 11.10 c

**Odd-Numbered Problems:** 11.1 2.5 times greater 11.3 62 days 11.5 1.2% of the actual mass 11.7 4800 times greater; ratio for Earth–Moon is 81.

## Chapter 12

**Multiple Choice:** 12.1 c, 12.2 c, 12.3 d, 12.4 c, 12.5 d, 12.6 d, 12.7 b, 12.8 c, 12.9 a, 12.10 c

**Odd-Numbered Problems:** 12.1  $47''$  12.3  $7.3 \times 10^{22}$  kg;  $1.3 \times 10^{-4}$  times the actual mass, 1.2 percent of Earth's mass 12.5  $6300\text{ kg/m}^3$ , assuming a radius of 15,000 km and a mass of 15 Earth masses 12.7  $1.1 \times 10^{18}$  particles.

## Chapter 13

**Multiple Choice:** 13.1 b, 13.2 b, 13.3 c, 13.4 d, 13.5 c, 13.6 d, 13.7 d, 13.8 a, 13.9 a, 13.10 d

**Odd-Numbered Problems:** 13.1  $1.7'$ ,  $13.2'$ ; yes

13.3 10,000 km/h—it moves halfway around the planet in just over 8 hours; this is half the rotation period, it's mostly rotation 13.5 0.42 Earth–Moon masses 13.7 yes, at 37 K, an escape speed of 1.08 km/s would be needed; Triton's escape speed is 1.46 km/s.

## Chapter 14

**Multiple Choice:** 14.1 c, 14.2 d, 14.3 a, 14.4 c, 14.5 a, 14.6 d, 14.7 b, 14.8 d, 14.9 d, 14.10 d

**Odd-Numbered Problems:** 14.1 (a) a 150-lb person would weigh 4.8 lbs; (b) 0.41 km/s 14.3 21 hours, taking the middle of the stated mass range and assuming a circular orbit 14.5  $3 \times 10^{12}$  kg; 0.06 percent 14.7 5 trillion.

## Chapter 15

**Multiple Choice:** 15.1 a, 15.2 c, 15.3 c, 15.4 a, 15.5 d, 15.6 d, 15.7 c, 15.8 a, 15.9 b, 15.10 d

**Odd-Numbered Problems:** 15.1 0.039 AU 15.3 period is 31.6 years; we could need up to twice this time if we happen to just miss a transit at the start of our observing period 15.5 0.3 AU 15.7  $1.7\text{ g} = 16.3\text{ m/s}^2$ ;  $0.9\text{ g} = 8.9\text{ m/s}^2$ .

## Chapter 16

**Multiple Choice:** 16.1 c, 16.2 b, 16.3 c, 16.4 a, 16.5 a, 16.6 b, 16.7 b, 16.8 c, 16.9 c, 16.10 b

**Odd-Numbered Problems:** 16.1 14,600  $\text{W/m}^2$ ; 52  $\text{W/m}^2$  16.3 (a) 3000 km; (b) 1460; (c) 167 minutes 16.5 36% as much light as the photosphere 16.7 10,800,000 km, for a coronal temperature of 1,000,000 K.

## Chapter 17

**Multiple Choice:** 17.1 a, 17.2 d, 17.3 d, 17.4 b, 17.5 c, 17.6 b, 17.7 c, 17.8 b, 17.9 c, 17.10 d

**Odd-Numbered Problems:** 17.1 83 pc; 0.36 arc seconds 17.3 80 solar luminosities 17.5 B is three times farther away 17.7 (a) 100 pc; (b) 4200 pc; (c) 170,000 pc; (d) 1,000,000 pc.

## Chapter 18

**Multiple Choice:** 18.1 a, 18.2 d, 18.3 c, 18.4 a, 18.5 d, 18.6 c, 18.7 b, 18.8 d, 18.9 a, 18.10 a

**Odd-Numbered Problems:** 18.1 1.9 grams 18.3  $7.1 \times 10^{20}\text{ m}^3$ , or a cube of side 8900 km 18.5 9.03 magnitudes 18.7 escape speeds (km/s): 1.8, 1.1, 1.1, 0.80; average molecular speeds: 13.6, 14.0, 14.6, 14.2; no.

## Chapter 19

**Multiple Choice:** 19.1 c, 19.2 a, 19.3 d, 19.4 b,

19.5 b, 19.6 a, 19.7 a, 19.8 b, 19.9 a, 19.10 b  
**Odd-Numbered Problems:** 19.1 yes, barely: the escape speed is 0.93 km/s, molecular speed is

0.35 km/s **19.3** luminosity decreases by a factor of 7900; absolute magnitude increases by 9.7  
**19.5** 8.3 magnitudes **19.7**  $10^{-6}$  solar luminosities.

## Chapter 20

**Multiple Choice:** **20.1** c, **20.2** b, **20.3** a, **20.4** b, **20.5** a, **20.6** c, **20.7** a, **20.8** b, **20.9** b, **20.10** d  
**Odd-Numbered Problems:** **20.1**  $9.9 \times 10^{26}$  kg destroyed,  $8.9 \times 10^{43}$  J; emitted **20.3** (a) 3370; (b) 84,300 solar luminosities **20.5** (a) 2.9 solar masses; (b) 1.7 solar masses **20.7** core density is  $3.5 \times 10^7$  kg/m<sup>3</sup>; solar core density is  $1.5 \times 10^5$  kg/m<sup>3</sup>, 230 times less; envelope density is  $5.7 \times 10^{-4}$  kg/m<sup>3</sup>,  $6.1 \times 10^{13}$  times less.

## Chapter 21

**Multiple Choice:** **21.1** a, **21.2** d, **21.3** d, **21.4** b, **21.5** d, **21.6** c, **21.7** c, **21.8** b, **21.9** b, **21.10** c  
**Odd-Numbered Problems:** **21.1**  $m = 20$ , 3.2 Mpc **21.3** 0.44 pc; no—there are no O or B stars (in fact no stars at all) within that distance of us. For the Moon: 320 pc; yes, but rarely **21.5**  $1.2 \times 10^{44}$  J; about 10 times greater than a supernova's electromagnetic output, and about 1/10 the output in the form of neutrinos **21.7** roughly 1000 km/s; not a very good assumption—the nebula is moving too fast to be affected much by gravity, although it is probably slowing down as it runs into the interstellar medium.

## Chapter 22

**Multiple Choice:** **22.1** b, **22.2** c, **22.3** a, **22.4** b, **22.5** c, **22.6** d, **22.7** b, **22.8** b, **22.9** c, **22.10** b

**Odd-Numbered Problems:** **22.1** 1 million revolutions per day, or 11.6 revolutions per second **22.3**  $1.9 \times 10^{12}$  m/s<sup>2</sup>, or 200 billion Earth gravities; 190,000 km/s, or 64 percent of the speed of light; 306,000 km/s **22.5**  $6.3 \times 10^4$  **22.7** (a) 26 microarcsec; (b) 0.016 arcsec; (c) 4.1 arcmin; (d) 8100 AU = 0.039 pc.

## Chapter 23

**Multiple Choice:** **23.1** d, **23.2** b, **23.3** d, **23.4** b, **23.5** b, **23.6** c, **23.7** a, **23.8** c, **23.9** d, **23.10** c  
**Odd-Numbered Problems:** **23.1** 2.0", much less than the angular diameter of Andromeda **23.3** 100 kpc **23.5** 0.014"/yr; proper motion has in fact been measured for several globular clusters **23.7** (a) 3.9 kpc; (b) 19.7 kpc.

## Chapter 24

**Multiple Choice:** **24.1** c, **24.2** c, **24.3** a, **24.4** c, **24.5** c, **24.6** c, **24.7** c, **24.8** b, **24.9** a, **24.10** b  
**Odd-Numbered Problems:** **24.1** 320 Mpc **24.3** 14,000 km/s, 57 Mpc; 12,000 km/s, 67 Mpc; 16,000 km/s, 50 Mpc **24.5**  $M = -22.5$ ;  $1.5 \times 10^8$  solar luminosities **24.7**  $1.5 \times 10^8$  solar masses.

## Chapter 25

**Multiple Choice:** **25.1** d, **25.2** b, **25.3** c, **25.4** a, **25.5** a, **25.6** b, **25.7** a, **25.8** d, **25.9** b, **25.10** a  
**Odd-Numbered Problems:** **25.1** 6.5 billion years **25.3**  $2.6 \times 10^{14}$  solar masses; reasonable to within a factor of perhaps 2–3 **25.5**  $9.4 \times 10^{10}$  solar masses **25.7**  $1.8 \times 10^{12}$  solar masses.

## Chapter 26

**Multiple Choice:** **26.1** a, **26.2** d, **26.3** b, **26.4** b, **26.5** c, **26.6** d, **26.7** c, **26.8** c, **26.9** d, **26.10** d  
**Odd-Numbered Problems:** **26.1** 1000 Mpc **26.3**  $4 \times 10^8$  **26.5** 1200 km/s **26.7** (a) 30,000 tons; (b) 2.8 pc.

## Chapter 27

**Multiple Choice:** **27.1** a, **27.2** d, **27.3** a, **27.4** b, **27.5** a, **27.6** c, **27.7** b, **27.8** a, **27.9** b, **27.10** c  
**Odd-Numbered Problems:** **27.1** 16 kpc **27.3** (a) matter, factor of 5.5; (b) radiation, factor of 50,000 **27.5**  $3 \times 10^{-12}$  m; hard X-ray/gamma ray **27.7** 12.7 Mpc.

## Chapter 28

**Multiple Choice:** **28.1** a, **28.2** b, **28.3** c, **28.4** c, **28.5** c, **28.6** b, **28.7** d, **28.8** d, **28.9** b, **28.10** a  
**Odd-Numbered Problems:** **28.1** 6.3 seconds (for a 20-year-old reader); 20.9 seconds (in 2011); 73 seconds; 2.7 minutes; 240 days **28.3** 27 AU **28.5** (a) 0.2; (b) 20; (c) 2000 **28.7**  $1.43 \times 10^9$  to  $1.67 \times 10^9$  Hz;  $2.4 \times 10^6$  channels.



*This page intentionally left blank*

# Photo Credits/Text Permissions

## Photos

### Chapter 1

**Part Opener 1a** MASS/UMass/IPAC-Caltech/NASA/NS **Part Opener 1c** California Institute of Technology/Palomar/Hale Observatory **Chapter Opener** ESO/Y. Beletsky **1.1** Jerry Lodriguss/Science Source **1.2** NASA **1.3** AURA **1.4** Robert Gendler/Science Source **1.5** NASA **1.7a–e** Glen Schneider **1.8** Pere Sanz/Alamy **1.12** AURA **1.20a–g** UC/Lick Observatory **1.22** Glen Schneider **1.23** M. Tsavkova/B. Angelov **1.24a** NOAA **1.24a–c** Glen Schneider **1.25** Glen Schneider **p. 29** Jerry Lodriguss/Science Source, NASA, AURA

### Chapter 2

**Chapter Opener** Sheila Terry/Science Source **2.1a** Sergio Pitamitz/SuperStock **2.1b** David Nunuk/All Canada Photos/SuperStock **2.2a** G. Gerster/Science Source **2.2b** Flonline digitale Bildagentur GmbH/Alamy **2.2c** Harrison Lapahie, Jr. **2.3** Istanbul University Library, Istanbul, Turkey/The Bridgeman Art Library **2.4** Boston Museum of Science **2.8** E. Lessing/Art Resource **2.10** Art Resource **2.11** Galilei, Galileo. “Galilean Moons.” Sidereus Nuncius. 1610. **2.12a–b** R. Beebe/Courtesy of New Mexico State University **2.13** E. Lessing/Art Resource **2.14** Newberry Library/SuperStock **2.17** P. Jones **2.19** Sheila Terry/Science Source

### Chapter 3

**Chapter Opener** NASA/JPL-Caltech **3.1** Robert Gendler/Science Source **3.11a** ESO **3.11b** A. Noriega-Crespo/AURA **3.11c** SST **3.11d** Galax/NASA

### Chapter 4

**Chapter Opener** NOAO/AURA **4.3** Wabash Instrument Corporation **4.4** AURA **4.6** Mattson, Barbara; Lochner, Jim; Gibb, Meredith; and Newman, Phil. “Absorption Spectra” Imagine Science. September 6, 2006. Accessed <http://imagine.gsfc.nasa.gov/YBA/M31-velocity/spectra-more.html> **4.10** NASA **4.12** ESO **4.14** Bausch & Lomb, Inc. **4.15** Chaisson, Eric. “Doppler Shift of Pure Hydrogen” **p. 96** NASA

### Chapter 5

**Chapter Opener** L. Calçada/ESO **5.1** Richard Megna/Fundamental Photographs **5.7a–b** W. M. Keck Observatory **5.08a** Dana Berry **5.8b–1** George Jacoby, Bruce Bohannon, Mark Hanna/NOAO/AURA/NSF **5.8b–2** NASA, ESA,

K. Kuntz (JHU), F. Bresolin (University of Hawaii), J. Trauger (Jet Propulsion Lab), J. Mould (NOAO), Y.-H. Chu (University of Illinois, Urbana), and STScI **5.9a–b** AURA **5.10a** R. Wainscoat; R. Underwood/Keck Observatory **5.10b** Subaru Telescope/National Astronomical Observatory of Japan NAOJ **5.11** ESO **5.13a–d** AURA **5.14a** MIT Lincoln Lab **5.14b** NOAO **5.14d** AURA **5.15a–c** AURA/NASA **5.17** ESO **5.18a–b** ESO **5.19** Lick Observatory **5.20a–1** NOAO **5.20a–2** NOAO **5.20b** MIT Lincoln Laboratory **5.21** NRAO **5.22a–c** D. Parker/T. Acevedo/NAIC; Cornell **5.23** MIT **5.24** AURA **5.25a–b** NRAO **5.27a** ALMA (ESO/NAOJ/NRAO) **5.27b** NASA, ESA, and the Hubble Heritage Team (STScI/AURA)-ESA/Hubble Collaboration **5.28** E. Simison/Sea West Enterprises **5.29a–b** Lick Observatory **5.29c** NASA **5.29d** Smithsonian Astrophysical Observatory **5.30a** ESO/NASA/JPL-Caltech/S. Kraus **5.30b** NASA/JPL-Caltech **5.31a** ESA **5.31b** ESO **p. 124** NRAO **p. 124** NRAO, A. C. Boley (University of Florida, Sagan Fellow), M. J. Payne, E. B. Ford, M. Shabran (University of Florida), S. Corder (North American ALMA Science Center, National Radio Astronomy Observatory), and W. Dent (ALMA, Chile), NRAO/AUI/NSF; NASA, ESA, P. Kalas, J. Graham, E. Chiang, E. Kite (University of California, Berkeley), M. Clampin (NASA Goddard Space Flight Center), M. Fitzgerald (Lawrence Livermore National Laboratory), and K. Stapelfeldt and J. Krist (NASA Jet Propulsion Laboratory) **5.32a** NASA **5.32b** GALEX **5.34** NASA **5.35** NASA **5.35** CXC/SAO **5.36a–b** NASA **5.37a** ESA **5.37b** UMass/Caltech **5.37c** A. Mellinger **5.37d** MPI **5.37e** NASA **p. 129** AURA, MIT Lincoln Lab, NRAO **p. 130** ESO/NASA/JPL-Caltech/S. Kraus, NRAO, NASA

### Chapter 6

**Part Opener 2a** Lowell Observatory Archives **Part Opener 2b–c** NASA **Part Opener 2d** NASA, ESA, The Hubble Heritage Team (STScI/AURA), J. Bell (Cornell Univ.), and M. Wolff (Space Sci Inst.) **Part Opener 2e** NASA Jet Propulsion Library **Part Opener 2f** NASA/JPL-Caltech/MSSS **Chapter Opener** JAXA **6.1** Museo della Scienza; Scala/Art Resource, NY **6.2** Birr Scientific & Heritage Foundation **6.3** NASA **6.4** NASA/Jet Propulsion Laboratory **6.6** Jerry Lodriguss/Science Source **6.8a** NASA/JPL-Caltech/UCAL/MPS/DLR/IDA **6.8b** S.Deiries/ESO **6.8c** NASA **p. 146** NASA/Johns Hopkins University Applied Physics Laboratory/Carnegie Institution of Washington **p. 146** Sovfoto/Eastfoto, NASA, NASA **p. 147** NASA/Jet Propulsion Laboratory, NASA, NASA/Jet Propulsion Laboratory, NASA **6.10a** NASA Jet Propulsion Laboratory **6.10b** Dana Berry **p. 149** Gamma-Rapho via Getty

Images **6.11** David Malin/Anglo-Australian Telescope **6.12** NASA **p. 156** NASA

### Chapter 7

**Chapter Opener** NASA/NOAA/GSFC/Suomi NPP/VIIRS/Norman Kuring **7.10a** Douglas Peebles Photography/Alamy **7.10b** AP Photo/Jack Smith **7.10c** Haruyoshi Yamaguchi/Corbis **7.13a** Earth Satellite Corporation/Science Photo Library/Science Source **7.13b** Michael Klesius/National Geographic Image Collection **7.14** David Parker/Science Source **7.20a** University Corporation for Atmospheric Research NCAR **7.20b** NASA **p. 165** Science Source

### Chapter 8

**Chapter Opener** NASA **8.1b** John Sanford/Science Source **8.2a–b** R. Beebe/Courtesy of New Mexico State University **8.3** UC/Lick Observatory **8.4a–c** California Inst. of Technology/Palomar/Hale Observatory **8.5** NASA **8.6** NASA **8.7** California Inst. of Technology/Palomar/Hale Observatory **8.8** NASA/Johns Hopkins University Applied Physics Laboratory/Carnegie Institution of Washington **8.9** NASA **8.14a–b** NASA **8.15a–b** NASA **8.16** NASA/John F. Kennedy Space Center **8.17** NASA **8.18a–b** U.S. Geological Survey, Denver **8.19** NASA **8.20** NASA **8.21a** NASA **8.21b** NASA/Johns Hopkins University Applied Physics Laboratory/Carnegie Institution of Washington **8.22** NASA **8.26a–f** Professor Willie Benz **8.27a–c** U.S. Geological Survey, Denver **p. 198** NASA **p. 199** NASA **p. 213** Lick Observatory Publications Office **p. 214** Professor Willie Benz, U.S. Geological Survey, Denver

### Chapter 9

**Chapter Opener** NASA/Jet Propulsion Laboratory **9.1** Jerry Schad/Science Source **9.2a–c** UC/Lick Observatory **9.5** AURA **9.6a** NASA **9.6b** ESA **9.7a–b** NASA **9.8c** NASA **9.9a–b** NASA **9.10** NASA **9.11** NASA **9.12a–b** NASA **9.13a–b** NASA **9.14** NASA **9.15a–b** NASA **9.16a–b** Russian Space Agency **9.18** NASA **9.19a** ESA **9.19b** NASA **p. 233** Jerry Schad/Science Source, AURA, NASA **p. 234** NASA

### Chapter 10

**Chapter Opener** NASA **10.2a** CNRS **10.2b–c** NASA **10.3** NASA **10.4** NASA **10.5** ESA **10.6a–b** NASA **10.7** NASA **10.8a** ESA **10.8b** NASA/CXC **10.9** NASA Jet Propulsion Laboratory **10.10a–b** NASA **10.11a–c** NASA **10.12a–b** NASA **10.13a** NASA Jet Propulsion Laboratory **10.13b** NASA **10.13c** Gordon Osinski/Canadian Space Agency **10.14** NASA **p. 250** NASA, NASA, NASA, NASA, NASA, NASA **10.15a–b** ESA **10.15(insert)** NASA

10.16 NASA 10.17 NASA 10.18a NASA/JPL/  
Cornell 10.18b NASA 10.18c NASA Jet Propulsion  
Laboratory 10.19 NASA 10.20a–b NASA/  
JPL-Caltech/MSSS 10.22a–b NASA 10.24a–b  
Kees Veenbos 10.25a ESA 10.25b NASA p. 261  
CNRS, NASA, NASA, NASA, NASA Jet Propul-  
sion Laboratory p. 262 NASA, NASA

## Chapter 11

**Chapter Opener** NASA Jet Propulsion Labora-  
tory 11.2a–b NASA 11.2c AURA 11.4a–c NASA 11.5  
NASA/Kennedy Space Center 11.6 NASA/Kennedy  
Space Center 11.8 NASA 11.9a–b NASA 11.10 NASA  
11.11a–b NASA p. 275 NASA, NASA, Space Tele-  
scope Science Institute 11.14 NASA 11.16 NASA 11.17  
NASA 11.19a–b NASA 11.20a–c NASA 11.22a–d  
NASA 11.23a–c NASA 11.24a–b NASA 11.25 NASA  
p. 287 NASA p. 288 NASA/Kennedy Space Center

## Chapter 12

**Chapter Opener** NASA Jet Propulsion Labora-  
tory 12.1 NASA/Johnson Space Center 12.2 NASA/  
Jet Propulsion Laboratory 12.4a–b NASA 12.6a–d  
NASA 12.7 NASA 12.9 NASA 12.13a–d NASA  
12.14a–b NASA/Jet Propulsion Laboratory 12.15a  
NASA Jet Propulsion Laboratory 12.15b NASA 12.16  
NASA 12.17a–b NASA Jet Propulsion Laboratory  
12.18a–b NASA 12.18c CFHT 12.19b NASA 12.20a  
NASA 12.20b ESA 12.21a–b NASA 12.22 NASA/  
ESA 12.24a UC/Lick Observatory 12.24b–g NASA  
12.24h NASA Jet Propulsion Laboratory 12.25a  
NASA/JPL/Space Science Institute 12.26 SST/JPL  
p. 315 NASA Jet Propulsion Laboratory, NASA  
Jet Propulsion Laboratory, NASA Jet Propulsion  
Laboratory, NASA, NASA/ESA, Lick Observatory  
Publications Office, NASA, NASA, NASA, NASA,  
NASA, NASA Jet Propulsion Laboratory

## Chapter 13

**Chapter Opener** L. Sromovsky and P. Fry,  
W. M. Keck Observatory 13.1 UC/Lick Observatory  
13.2 NASA 13.3 UC/Lick Observatory 13.4a–b  
NASA 13.5a–b NASA 13.5c NASA/John F. Ken-  
nedy Space Center 13.5d–e NASA 13.7a–d NASA  
13.8a–b NASA 13.8c–e NASA/John F. Kennedy  
Space Center 13.11a–g NASA 13.11h NASA/Jet  
Propulsion Laboratory 13.12a–b NASA 13.13 NASA  
13.14a–b NASA 13.16a NASA 13.17 NASA 13.18  
NASA p. 335 NASA, NASA, NASA, NASA

## Chapter 14

**Chapter Opener** S. Deiries/ESO 14.1 NASA/  
SWRI 14.2a–c NASA 14.3a–b NASA 14.4a–b  
NASA/JPL-Caltech/UCLA/MPS/DLR/IDA 14.5  
Xinhuashidian 14.8 National Optical Astronomy  
Observatories (NOAO) 14.9a U.S. Naval Obser-  
vatory 14.9b Walter Pacholka/Science Source  
14.11a–b California Institute of Technology/  
Palomar/Hale Observatory 14.11c Mt. Stromlo  
and Siding Springs Observatories 14.13 European  
Space Agency p. 350 David A. Hardy/Science  
Source 14.14a–c NASA Jet Propulsion Laboratory  
14.15 NASA Jet Propulsion Laboratory 14.18a U.S.  
Naval Observatory 14.18b NASA 14.20 NASA

14.21a–c LPL/Keck 14.22a–c NASA 14.24a Pekka  
Parviainen/Science Photo Library/Science Source  
14.24b Jerry Lodriguss/Science Source 14.27  
NASA 14.28 Sovfoto/Eastfoto 14.29a Jonathan  
Blair/Corbis 14.29b Jon Mandaville/Saudi  
Aramco World/PADIA 14.30a–b Science  
Graphics p. 363 NASA, NASA, National  
Optical Astronomy Observatories (NOAO),  
NASA, Sovfoto/Eastfoto

## Chapter 15

**Chapter Opener** NASA 15.1 ESO p. 376  
D. DeMartin/ESO, L. Cal, ada/Nick Risinger/  
ESO 15.9a L. Cal, ada/ESO 15.9b M. Kornmesser/  
Nick Risinger/ESO 15.10b T. Pyle/NASA 15.13  
Christine Pulliam and David Aguilar p. 383 ESO,  
M. Kornmesser/Nick Risinger/ESO

## Chapter 16

**Chapter Opener** NASA 16.1 NOAO 16.3 NASA  
16.5 “Solar Oscillations” GONG. National Solar  
Observatory. p. 395 NASA 16.8 SST/Royal Swed-  
ish Academy of Sciences 16.9 California Inst. of  
Technology/Palomar/Hale Observatory 16.10  
Glen Schneider 16.11 NASA 16.12 National Solar  
Observatory 16.14 California Inst. of Technology/  
Palomar/Hale Observatory 16.15a California Inst.  
of Technology/Palomar/Hale Observatory 16.15b  
SST/Royal Swedish Academy of Sciences 16.16b  
NASA 16.18 NASA 16.21a NASA Headquarters/  
European Space Agency 16.21b NASA 16.22  
USAF 16.23 European Space Agency 16.24a–d  
Lockheed Martin p. 409 Hendrik Avercamp  
(1585–1634), “Winter Games.” Oil on Canvas.  
Location: Rijksmuseum, Amsterdam, Holland/  
Bridgeman Art Library 16.25 NCAR High Alti-  
tude Observatory 16.28a SNO 16.28b Lawrence  
Berkeley National Laboratory p. 417 NASA, Glen  
Schneider, NASA, NASA, Lawrence Berkeley  
National Laboratory

## Chapter 17

**Chapter Opener** NASA/ESA 17.3a–b Harvard  
College Observatory 17.7b Getty Images 17.7c  
California Inst. of Technology/Palomar/Hale  
Observatory 17.7d UC/Lick Observatory 17.7e  
Association of Universities for Research in  
Astronomy 17.8a Pere Sanz/Alamy 17.8b J.  
Trauger/NASA/Jet Propulsion Laboratory 17.11a  
NASA 17.11b NASA/SAO 17.19a–c Harvard  
College Observatory p. 445a–c Harvard College  
Observatory

## Chapter 18

**Chapter Opener** Space Telescope Science Institute  
18.1a European Southern Observatory 18.2a–c Eu-  
ropean Southern Observatory 18.4 ESO/S. Guisard  
18.5, 18.5(insert) Harvard College Observatory  
18.6 Robert Gendler/Photo Researchers, Inc. 18.7a  
AURA 18.7b NASA/JPL-Caltech/J. Rho (SSC/  
Caltech) 18.9a ESA 18.9b AURA 18.9c NASA Jet  
Propulsion Laboratory 18.9d James M. Moran/  
NASA 18.10a NASA 18.10b NASA/John F. Kennedy  
Space Center 18.10c ESO 18.12a–b Charles J. Lada/

Harvard Smithsonian Center for Astrophysics  
18.13 Robert Gendler/Jim Misti/Steve Mazlin/Photo  
Researchers, Inc. 18.14a Royal Observatory of  
Belgium 18.14b European Southern Observatory  
18.19 AURA 18.20 AURA p. 465 European  
Southern Observatory p. 466 NASA Jet Propulsion  
Laboratory, Charles J. Lada/Harvard Smithsonian  
Center for Astrophysics, NASA

## Chapter 19

**Chapter Opener** STScI; ESA 19.1 NASA/ESA  
p. 479 NASA, NASA 19.8a AURA 19.8b–c  
NASA/Jet Propulsion Laboratory 19.9a Pere Sanz/  
Alamy 19.9b–c NASA Jet Propulsion Laboratory  
19.9d–e NASA 19.10a–b NASA 19.11d SST 19.12a  
NASA Goddard Laboratory for Atmospheres  
19.12b Dana Berry 19.13 NASA 19.13(insert)  
NASA 19.15 NASA 19.16 European Southern  
Observatory 19.16(insert) NASA Goddard Labo-  
ratory for Atmospheres 19.17a AURA 19.18a P.  
Seitzer 19.19a I. Bonnell and M. Bate 19.20a Lick  
Observatory Publications Office 19.20b NASA/Jet  
Propulsion Laboratory p. 490 European Southern  
Observatory, European Space Agency, NASA  
p. 491 NASA, NASA, NASA, NASA/Jet Propul-  
sion Laboratory p. 492 NASA, P. Seitzer

## Chapter 20

**Chapter Opener** STScI 20.9a AURA 20.9c  
NASA 20.10a AURA 20.10b Space Telescope  
Science Institute 20.10c NASA 20.12 California  
Inst. of Technology/Palomar/Hale Observatory  
20.13 AURA, NASA 20.14 NASA 20.15 NASA  
p. 511 NASA, John F. Kennedy Space Center/NASA,  
John F. Kennedy Space Center/NASA, John F.  
Kennedy Space Center/NASA, John F. Kennedy  
Space Center/NASA 20.18c NASA 20.19 National  
Optical Astronomy Observatories 20.20 ESO  
20.20(insert) NASA p. 518 Space Telescope Science  
Institute, ESO/NASA, ESO/NASA

## Chapter 21

**Chapter Opener** STScI 21.1a–b UC/Lick Obser-  
vatory 21.3 Dana Berry 21.4a California Inst. of  
Technology/Palomar/Hale Observatory 21.4b–d  
ESA 21.7a–b AURA 21.10b NASA 21.11 Harvard  
College Observatory 21.12a–b David Malin  
p. 533a–f NASA p. 533 Space Telescope Science  
Institute 21.18 NASA/CXC 21.19a European  
Southern Observatory 21.19b,d John F. Kennedy  
Space Center/NASA 21.19c National Optical  
Astronomy Observatory (NOAO) p. 539 NASA  
p. 540 Jet Propulsion Laboratory/NASA,  
Harvard-Smithsonian Center for Astrophysics,  
European Southern Observatory, John F. Kennedy  
Space Center/NASA, National Optical Astronomy  
Observatory (NOAO), John F. Kennedy Space  
Center/NASA

## Chapter 22

**Chapter Opener** John F. Kennedy Space  
Center/NASA 22.1 NASA 22.4a European  
Southern Observatory 22.4b John F. Kennedy  
Space Center/NASA 22.4c Lick Observatory

22.5a–b NASA 22.6a Smithsonian Astrophysical Observatory 22.6b–c NASA 22.7a–d National Radio Astronomy Observatory (NRAO) 22.9 ESO; NASA 22.9b ESO; NASA 22.12a–b NASA; ESO 22.14a–b European Southern Observatory 22.22a Harvard-Smithsonian Center for Astrophysics 22.22b NASA 22.23 L. J. Chaisson p. 568 NASA 22.24 National Radio Astronomy Observatory 22.25 Subaru; NASA 22.25(insert) Subaru; NASA 22.26 NASA p. 571 NASA, Smithsonian Astrophysical Observatory, NASA

## Chapter 23

**Part Opener** 4a Corbis/Bettmann 4b Edwin Hubble, 1923/Carnegie Institution of Washington 4c–d Harvard College Observatory **Chapter Opener** European Southern Observatory 23.1 Axel Mellinger, Central Michigan University 23.2a Robert Gendler/Photo Researchers, Inc. 23.2b California Inst. of Technology/Palomar/Hale Observatory 23.2c NASA 23.3a NASA 23.5c Harvard College Observatory p. 584 Harvard College Observatory 23.11 U MASS/CALTECH 23.16 Jet Propulsion Laboratory/NASA 23.18 AURA 23.19 R. Gendler 23.22 AAT, NASA 23.23 Association of Universities for Research in Astronomy, Inc. 23.24a AURA 23.24b European Southern Observatory 23.25a Jet Propulsion Laboratory/NASA 23.25b National Radio Astronomy Observatory (NRAO) 23.25c John F. Kennedy Space Center/NASA 23.25d National Radio Astronomy Observatory 23.26 European Southern Observatory 23.27a–f L. J. Chaisson p. 604 John F. Kennedy Space Center/NASA

## Chapter 24

**Chapter Opener** NRAO/STSci 24.1a AURA 24.1b NRAO/STSci 24.2a–b NASA 24.2c David Malin Images 24.3a NASA 24.4b David Malin Images 24.4a NASA 24.4c European Southern Observatory 24.5a–b Association of Universities for Research in Astronomy 24.5c Robert Gendler, MD 24.6a–b California Inst. of Technology/Palomar/Hale Observatory 24.7a Mount Stromlo & Siding Spring Observatory/Photo Researchers, Inc. 24.7b–c Royal Observatory, Edinburgh/Science Photo/Photo Researchers, Inc. 24.8a–b NASA 24.10 NASA 24.10(inserts a–f) NASA 24.11b Kennedy Space Center/NASA Headquarters 24.13a Matt BenDaniel 24.13b NASA 24.14 Matt BenDaniel 24.14(insert) Association of Universities for Research in Astronomy 24.15 John F. Kennedy Space Center/NASA 24.16a–j California Inst. of Technology/Palomar/Hale Observatory 24.20 NASA 24.21 NASA 24.22a ESO 24.22b NRAO 24.22(insert) SAO 24.23a National Optical Astronomy Observatories 24.23b National Radio Astronomy Observatory 24.24 Harvard-Smithsonian Center for Astrophysics 24.26a National Optical Astronomy Observatories 24.26b Association of Universities for Research in Astronomy 24.26c NASA 24.27a–b Association of Universities for Research in Astronomy 24.28 Adapted from Palomar/Caltech 24.29 Sloan Digital Sky Survey SDSS 24.30 NRAO 24.32a–b NASA 24.33a–b NASA p. 635 NASA, Matt BenDaniel, NASA, NASA

## Chapter 25

**Chapter Opener** Space Telescope Science Institute 25.1 NASA/Kennedy Space Center 25.3a–b NASA 25.4a National Radio Astronomy Observatory 25.4b California Inst. of Technology/Palomar/Hale Observatory 25.5 NASA 25.6 NASA 25.7 W. Keel 25.8a AURA; NASA; J. Barnes 25.8b Association of Universities for Research in Astronomy, Inc. 25.9a AURA; NASA; J. Barnes 25.9b NASA 25.10a NASA 25.12a NASA/ESA 25.12b Space Telescope Science Institute 25.13b Vasily Belokurov p. 651 Fermilab Visual Media Services, Sloan Digital Sky Survey SDSS 25.16 NASA 25.18a–b NASA 25.19 NASA 25.25 NASA 25.27a NASA 25.27b Dana Berry 25.28a NASA 25.28b Dana Berry 25.29a–b J. A. Tyson, Alcatel-Lucent; NOAO 25.30 NASA p. 663 NASA, NASA, NASA

## Chapter 26

**Chapter Opener** NASA 26.11a–f NASA 26.16 ALCATEL-LUCENT Technologies Bell Labs 26.19 NASA

## Chapter 27

**Chapter Opener** CERN 27.2 Fermi National Laboratory 27.15 V. Springel 27.16 NASA 27.17a ESA and the Planck Collaboration 27.17b NASA 27.19 Z. Rostomian/SDSS p. 711 NASA

## Chapter 28

**Chapter Opener** Dana Berry 28.1 Dana Berry p. 717 Robley C. Williams 28.4 NASA 28.5a Elso Barghoorn 28.5b Eric Chaisson 28.6 NASA 28.7 Harvard-Smithsonian Center for Astrophysics 28.9 NASA/JPL-Caltech/Malin Space Science Systems 28.10a Woods Hole Oceanographic Institute 28.10b Eric Chaisson 28.15 NASA 28.18a Seth Shostak 28.18b SETI Institute p. 733 Dana Berry, Harvard-Smithsonian Center for Astrophysics, Eric Chaisson

## Covers

Student Edition: ESO/S. Guisard ([www.eso.org/~sguisard](http://www.eso.org/~sguisard))  
Volume 1: Solar System: NASA/JPL-Caltech/MSSS/Mastcam  
Volume 2: Stars and Galaxies: ESO/F. Comeron

## Text

p. 22 “Periodicity of Solar Eclipses.” NASA. January 12, 2012. Accessed at <http://eclipse.gsfc.nasa.gov/SEsaros/SEperiodicity.html>

p. 24 **Figure 1.27** “Eclipse Tracks.” Annular and Hybrid Solar Eclipse Paths: 2001–2025. Accessed at <http://eclipse.gsfc.nasa.gov/SEatlas/SEatlas.html>

p. 47 **Table 2.1** “Solar System Exploration—Planets.” NASA. November 7, 2012. Accessed at <http://solarsystem.nasa.gov/planets/>

p. 135 **Table 6.1** “Planets.” NASA. November 7, 2012. Accessed at <http://solarsystem.nasa.gov/planets/>

pp. 146–147 **Discovery 6-2** “Solar System Exploration: Missions.” NASA. November 7, 2012. Accessed at <http://solarsystem.nasa.gov/missions/>

pp. 180–181 Stern, David, and Ptak, Andy. “Ask an Astrophysicist: What Are the Van Allen Belts?” Imagine the Universe! NASA. February 28, 1997. Accessed at [http://imagine.gsfc.nasa.gov/docs/ask\\_astro/answers/970228a.html](http://imagine.gsfc.nasa.gov/docs/ask_astro/answers/970228a.html)

p. 197 **More Precisely 8-1** “Mercury: Planetary Science.” National Astronomy and Ionosphere Center/Arecibo Observatory. NASA.

p. 198 **Discovery 8-1** Garber, Steve. “The Decision to Go to the Moon: President John F. Kennedy’s May 25, 1961 Speech before a Joint Session of Congress.” NASA History Office. August 7, 2012.

p. 228 “Venera 13 & 14 Bibliography.” NASA’s HEASARC: Observatories. NASA. June 26, 2003. Accessed at [http://heasarc.gsfc.nasa.gov/docs/heasarc/missions/biblio/venera1314\\_biblio.html](http://heasarc.gsfc.nasa.gov/docs/heasarc/missions/biblio/venera1314_biblio.html)

p. 229 “A curious cold layer around the atmosphere of Venus.” Science Programme, European Space Agency. October 1, 2012. Accessed at <http://sci.esa.int/science-e/www/object/index.cfm?fobjectid=50884>

p. 246 “Mars Global Surveyor.” Lunar Planetary Science. NASA. November 7, 1996. Accessed at <http://nssdc.gsfc.nasa.gov/planetary/marsurvr.html>

p. 248 “Mars Reconnaissance Orbiter.” NASA Jet Propulsion Laboratory. 2005. Accessed at <http://mars.jpl.nasa.gov/mro/>

p. 255 “Phoenix Mars Lander.” Missions. NASA. September 9, 2010. Accessed at [http://www.nasa.gov/mission\\_pages/phoenix/main/index.html](http://www.nasa.gov/mission_pages/phoenix/main/index.html)

p. 271 “Voyager, The Interstellar Mission.” Jet Propulsion Laboratory. California Technical Institute. NASA. 1977–1989.

p. 273 “Hubble Space Telescope.” Hubble Missions. NASA. 2008.

p. 274 **Discovery 11-1** “Comet Shoemaker-Levy Collision with Jupiter.” Jet Propulsion Laboratory. California Institute of Technology. NASA. March 14, 2011.

p. 286 “Galileo Legacy.” Solar System Exploration. NASA. September 3, 2003. Accessed at <http://solarsystem.nasa.gov/galileo/>

p. 298 **Table 12.1** “Solar System Exploration: Saturn: Rings.” NASA. April 18, 2012.

p. 303 **Table 12.2** “Solar System Exploration: Saturn: Moons.” NASA. April 18, 2012.

p. 306 **Discovery 12.1** Piazza, Enrico. “Cassini Solstice Mission.” Jet Propulsion Laboratory. California Institute of Technology. Accessed at <http://saturn.jpl.nasa.gov/mission/introduction/>

p. 309 “Cassini: Unlocking Saturn’s Secrets.” NASA. Accessed at [http://www.nasa.gov/mission\\_pages/cassini/media/cassini-20080730.html](http://www.nasa.gov/mission_pages/cassini/media/cassini-20080730.html)

- p. 312** “Cassini: Unlocking Saturn’s Secrets.” NASA. Accessed at [http://www.nasa.gov/mission\\_pages/cassini/timeline/index.html](http://www.nasa.gov/mission_pages/cassini/timeline/index.html)
- p. 313** Clavin, Whitney. “NASA Space Telescope Discovers Largest Ring Around Saturn.” NASA. October 6, 2009. Accessed at <http://www.spitzer.caltech.edu/news/966-ssc2009-19.NASA-Space-Telescope-Discovers-Largest-Ring-Around-Saturn>
- p. 313** “About Saturn and Its Moons-Iapetus.” NASA. Accessed at <http://saturn.jpl.nasa.gov/science/moons/iapetus/>
- p. 326** “Voyager, The Interstellar Mission: Uranus.” NASA. October 8, 2010. Accessed at <http://voyager.jpl.nasa.gov/science/uranus.html>
- p. 326** “Voyager, The Interstellar Mission: Uranus.” NASA. October 8, 2010. Accessed at <http://voyager.jpl.nasa.gov/science/uranus.html>
- p. 328** “Solar System Exploration: Neptune and Uranus.” NASA. 2012.
- p. 331** “Voyager, The Interstellar Mission: Triton.” NASA. October 18, 2010. Accessed at [http://voyager.jpl.nasa.gov/science/neptune\\_triton.html](http://voyager.jpl.nasa.gov/science/neptune_triton.html)
- p. 332** “Voyager, The Interstellar Mission: Triton.” NASA. October 18, 2010. Accessed at [http://voyager.jpl.nasa.gov/science/neptune\\_triton.html](http://voyager.jpl.nasa.gov/science/neptune_triton.html)
- p. 332** “Voyager, The Interstellar Mission.” NASA. October 18, 2010. Accessed at <http://voyager.jpl.nasa.gov/>
- p. 333** “Voyager, The Interstellar Mission: Uranus.” NASA. October 18, 2010. Accessed at <http://voyager.jpl.nasa.gov/science/uranus.html>
- p. 333 Table 13.2** “Uranus: Rings.” Solar System Exploration. NASA. December 14, 2012. Accessed at <http://solarsystem.nasa.gov/planets/profile.cfm?Object=Uranus&Display=Rings>
- p. 334 Table 13.3** “Neptune: Rings.” Solar System Exploration. NASA. December 14, 2012. Accessed at <http://solarsystem.nasa.gov/planets/profile.cfm?Object=Neptune&Display=Rings>
- p. 342** “NEAR.” Near Earth Object Program. NASA. December 17, 2012. Accessed at <http://neo.jpl.nasa.gov/missions/near.html>
- p. 342** “NEAR.” Near Earth Object Program. NASA. December 17, 2012. Accessed at <http://neo.jpl.nasa.gov/missions/near.html>
- p. 342** “Dawn: Journey to the Asteroid Belt.” NASA. December 6, 2012. Accessed at [http://www.nasa.gov/mission\\_pages/dawn/main/index.html](http://www.nasa.gov/mission_pages/dawn/main/index.html)
- p. 344** “NEAR.” Earth Objects Program. NASA. December 17, 2012. Accessed at <http://neo.jpl.nasa.gov/faq/#pha>
- p. 350** “Stardust: NASA’s Comet Sample Return Mission.” November 3, 2009. Accessed at <http://stardust.jpl.nasa.gov/overview/index.html>
- p. 351** “Deep Impact: Mission to a Comet.” NASA. [http://www.nasa.gov/mission\\_pages/deepimpact/main/index.html](http://www.nasa.gov/mission_pages/deepimpact/main/index.html)
- p. 357** “IAU 2006 General Assembly: Result of the IAU Resolution Votes.” International Astronomical Union. August 24, 2006. Accessed at [http://www.iau.org/public\\_press/news/detail/iau0603/](http://www.iau.org/public_press/news/detail/iau0603/)
- p. 359 Table 14.1** “Watch the Skies.” NASA. December 6, 2012. Accessed at <http://www.nasa.gov/topics/solarsystem/features/watchtheskies/index.html>
- p. 372** “Space Science: COROT Overview.” European Space Agency. May 8, 2007. Accessed at [http://www.esa.int/Our\\_Activities/Space\\_Science/COROT\\_overview](http://www.esa.int/Our_Activities/Space_Science/COROT_overview)
- p. 374** “Kepler: A Search for Habitable Planets.” NASA. December 9, 2012. Accessed at <http://kepler.nasa.gov/>
- p. 377** “ESA, Missions: COROT.” European Space Agency. October 27, 2009. Accessed at <http://sci.esa.int/science-e/www/object/index.cfm?fobjectid=31706>
- p. 387** Payne-Gaposchkin, Cecilia H. “Henry Norris Russell Prize Lecture of the American Astronomical Society, ‘Fifty Years of Novae,’ ” *The Astronomical Journal*, Volume 82, Number 9, September 1977.
- p. 395** “SOHO: Solar and Heliospheric Observatory.” NASA. Accessed at <http://sohowww.nascom.nasa.gov/home.html>
- p. 398 Table 16.2** “SOHO: Solar and Heliospheric Observatory.” NASA. Accessed at <http://sohowww.nascom.nasa.gov/home.html>
- p. 399** “SOHO: Solar and Heliospheric Observatory.” NASA. Accessed at <http://sohowww.nascom.nasa.gov/home.html>
- p. 407** “The Skylab Missions.” Solar Physics: Marshall Space Flight Center. NASA. September 2011. Accessed at <http://solarscience.msfc.nasa.gov/Skylab.shtml>
- p. 408** “Ulysses.” Jet Propulsion Laboratory. NASA. December 18, 2012.
- p. 498 Table 20.1** “Star Formation.” Astrophysics. NASA Science. October 16, 2012.
- p. 505 Table 20.2** “White Dwarfs.” Goddard Space Flight Center. NASA.
- p. 512 Table 20.3** “Stellar Evolution-The Birth, Life and Death of a Star.” NASA. April 10, 2009.
- p. 532 Discovery 21.1** “Supernova 1987A: Fast Forward to the Past.” NASA. August 8, 2005.
- p. 551** “Millisecond Pulsar.” The Arecibo Observatory. National Science Foundation.
- p. 552** “The CGRO Mission (1991–2000).” NASA. November 30, 2005.
- p. 553** “SWIFT Mission.” NASA. December 28, 2012.
- p. 557** “Black Holes.” NASA.
- p. 557** “The Michelson-Morley Experiment.” Georgia State University, Department of Physics. Accessed at <http://hyperphysics.phy-astr.gsu.edu/hbase/relativ/mmhist.html>
- p. 574** “‘Great Debate.’ Obituary of Harlow Shapley.” NASA. 1972. Accessed at [http://apod.nasa.gov/htmltest/gifcity/shapley\\_obit.html](http://apod.nasa.gov/htmltest/gifcity/shapley_obit.html)
- p. 575** Mayall, N.U. “EDWIN POWELL HUBBLE.” NASA SpaceLink. 1996. Accessed at [http://apod.nasa.gov/diamond\\_jubilee/hubble\\_nas.html](http://apod.nasa.gov/diamond_jubilee/hubble_nas.html)
- p. 580** “Biography: William Herschel (1738–1822).” Space Telescope Science Institute. Accessed at <http://amazing-space.stsci.edu/resources/explorations/groundup/lesson/bios/herschel-wm/>
- p. 583** “Henrietta Swan Leavitt.” Deaf Scientist Corner. Texas Woman’s University. Accessed at [http://www.twu.edu/dsc/swan\\_leavittI.htm](http://www.twu.edu/dsc/swan_leavittI.htm)
- p. 589 Table 23.1** “Components of the Galaxy.” Physics Department. The University of Tennessee, Knoxville. Accessed at <http://csep10.phys.utk.edu/astr162/lect/milkyway/components.html>
- p. 594 Discovery 23.2** Danforth, Charles. “The Origins of Spiral Arms.” The University of Colorado at Boulder. May 28, 1998. Accessed at <http://casa.colorado.edu/~danforth/science/spiral/>
- p. 597** “Hubble Sees an Unexpected Population of Young-Looking Stars.” NASA. Accessed at [http://www.nasa.gov/mission\\_pages/hubble/science/ngc6362.html](http://www.nasa.gov/mission_pages/hubble/science/ngc6362.html)
- p. 614 Table 24.1** “What Types of Galaxies Do We Find in the Universe?” University of Oregon.
- p. 623 Table 24.2** “Project II: Distance Measures in Cosmology.” Astronomy Department. University of Florida.
- p. 624** “Astronomy Bio ... Carl Seyfert.” The Lake County Astronomical Society.
- p. 651 Discovery 25.1** “The Sloan Digital Sky Survey.” The Sloan Digital Sky Survey: Mapping the Universe. Alfred P. Sloan Foundation. July 7, 2012. Accessed at <http://www.sdss.org/>
- p. 653** “Chandra X-Ray Observatory-Black Holes.” NASA via Harvard University-Smithsonian Center for Astrophysics. Accessed at [http://chandra.harvard.edu/learn\\_bh.html](http://chandra.harvard.edu/learn_bh.html)
- p. 709** “COBE: Cosmic Background Explorer.” NASA. June 26, 2008. Accessed at <http://lambda.gsfc.nasa.gov/product/cobe/>
- p. 722** “Mars Exploration Rovers.” NASA. Accessed at <http://marsrovers.jpl.nasa.gov/mission/>
- p. 722** “Exploring the Planets-Mars.” NASA. Accessed at <http://airandspace.si.edu/etp/mars/explore.html>
- p. 29** “Mars Science Laboratory.” NASA. Accessed at [http://www.nasa.gov/mission\\_pages/msl/index.html](http://www.nasa.gov/mission_pages/msl/index.html)

# Index

**Note: Page numbers followed by n indicate footnote material.**

## A

A 2218 galaxy cluster, 661  
A ring, 298, 300, 301, 303  
A0620-00 (X-ray binary system), 569  
Abell 39 planetary nebula, 504  
Abell 1689, 618  
Abell 1835, 642  
Aberration  
  chromatic, 102  
  of starlight, 42  
Absolute brightness, 425. *See also*  
  Luminosity  
Absolute magnitude, 427, 430  
Absolute temperature, 70  
Absolute zero, 69  
Absorption lines  
  atomic hydrogen, 658–659  
  quasar, 658–659  
Absorption (Fraunhofer) lines, 82–83,  
  397–398  
Absorption spectra, 460–461  
AC114 quasar, 659  
Acceleration, 50  
  cosmic, 678–679  
Accretion, 150, 368, 631  
  jovian planet formation and, 152–153  
Accretion disk, 522–523, 631, 633  
Accretion flows, 631  
Acoustic oscillations, 709–710  
Active galaxies/active galactic nuclei,  
  622–634, 654, 655–656. *See also*  
  Galactic nucleus  
  central engine of, 630–634  
  energy emission, 633–634  
  energy production, 630–632  
  defined, 622, 624  
  evolution of, 655  
  properties of, 623–624, 630  
  quasars vs., 629  
  radiation emitted by, 622–624  
  radio galaxies, 569, 624, 625–628,  
  641–642  
  scientific method and, 656  
  Seyfert galaxies, 624–625  
  starburst galaxies, 623–624  
  supermassive black holes at center of,  
  569, 601–603, 624  
Active optics, 112  
Active prominences, solar, 405  
Active regions, solar, 405–406  
Active Sun, 405–409  
  active regions, 405–406  
  changing solar corona, 407–410  
  X-ray images of, 407, 408  
Adams, John, 321  
Adams ring (Neptune), 334  
Adaptive optics, 112–114, 432  
Adrastea (moon), 287  
African plate, 175  
AG Carinae, 511  
Ahnighito (meteorite), 361  
Aine (corona), 226  
Air, gravity and, 196–197  
Aldebaran, 432, 433, 436  
Algal, 515–516  
  evolution of, 516  
  future of, 517  
ALH84001 (meteorite), 250–251  
Allen Telescope Array, 731–732  
Almagest (Ptolemy), 38–39  
Alpha Centauri, 376, 389, 423, 424–425, 430,  
  432, 444, 729  
Alpha Centauri A, 376  
Alpha Centauri B, 376  
Alpha Centauri C (Proxima Centauri), 376  
Alpha particles, 499  
Alpha process, 535–536  
Alpha ring (Uranus), 333  
ALSEP (Apollo Lunar Surface Experiments  
  Package), 198

Alvin submarine, 723  
AM radio signals, 164, 730  
Amalthea (satellite), 279  
Amino acids, 718, 719, 720  
Ammonia, 724  
Amor (Mars-crossing) asteroid, 340, 343  
Amplitude, 61  
Andromeda galaxy, 60, 65, 106, 108, 569,  
  574, 575, 578, 579, 581, 586, 588, 617, 645  
Andromeda Nebula, 571  
Ångström, 63  
Ångström, Anders, 63  
Angular diameter, 28  
Angular measurement, 13  
Angular momentum, 149  
  conservation of, 145, 149, 368, 544  
Angular resolution, 106–108, 111  
  interferometry and, 118, 119  
  radio telescopes and, 116–117  
Annular eclipse, 21–22  
Antarctic ozone hole, 164  
Antarctic plate, 175  
Antenna, “sugar-scoop,” 683  
Antennae (NGC 4038/4039) galaxies, 644  
Antennae galaxies, 120  
Antiparticle, 411  
Apennine Mountains, 206  
Aphelion, 46, 190, 199–200  
Aphrodite Terra, 222, 223, 224, 225, 226  
Apollo (Earth-crossing) asteroid, 340,  
  343–344  
Apollo Lunar Surface Experiments Package  
  (ALSEP), 198  
Apollo program, 180, 189, 190, 192, 193, 194,  
  202, 203, 204, 206, 209, 212  
  description of, 198–199  
Apophis (MN4) (asteroid), 344  
Apparent brightness, 425–426, 581, 582,  
  586, 598  
Apparent magnitude, 426, 427  
Apparent motion, 39  
Arabian plate, 175  
Arago ring (Neptune), 334  
Arc (of ring), 304  
Arc minute, 13  
Arc second, 13  
Arcturus, 432, 499, 588  
Arecibo radio telescope, 116–117, 197, 221  
Ariel (moon), 328, 329  
Aristarchus of Samos, 39  
Aristotle, 9, 38, 39, 41, 50  
Armstrong, Neil, 198, 199  
Arp 273 galaxy, 638, 639  
Assumptions of mediocrity, 717  
Asteroid belt, 136, 145, 155, 340, 344, 345  
Asteroids, 134–135, 143, 144, 155–156, 338,  
  339, 340–345  
  Earth-crossing, 343–344  
  naming of, 344n  
  near-miss events with, 343–344  
  observations from space, 341–343  
  orbital resonances of, 344–345  
  properties of, 138, 340, 341  
  stray, 360–361  
  types of, 341  
Asthenosphere, 174, 176  
Astrologer, 11  
Astronomical unit (AU), 47, 48  
Astronomy  
  ancient, 34–36  
  defined, 4  
  high-resolution, 111–114  
  learning from history, 508  
  modern, birth of, 41–42  
  ultraviolet, 123, 125  
Asymptote, 502n  
Asymptotic-giant branch, 501, 502, 502n, 512  
Atacama Large Millimeter Array (ALMA),  
  120, 124  
Athen asteroids, 343  
Atlantic Ocean, 174–175, 176  
Atlas (moon), 303, 304  
Atmosphere(s), 162–168. *See also specific*  
  *planets, stars, and other bodies*  
  bands in, 268–270

blue sky, reasons for, 165  
composition of, 197  
gravity and, 196–197  
heat and, 166–168  
opacity of, 66–67, 68  
origin of, 168  
ozone in, 164, 166  
primary, 168  
secondary, 168  
solar, 397–400  
structure of, 163–164  
surface heating and, 166–168  
of terrestrial planets, 141  
turbulence in, 111, 112  
of Venus, 228–232  
Atmospheric blurring, 111–112  
Atmospheric circulation on Earth, 409  
Atom(s), 84–87  
  classical, 85  
  excitation of, 85, 89  
  first, 699–700  
  formation of, 697–700  
  ionized, 84, 455–456  
  modern, 85  
  motions of, 472  
  nucleus of, 84, 172  
  spectra of, 87  
  structure of, 84–85  
  thermal motions of, 94  
Atomic epoch, 696, 705  
Aurora/aurorae  
  on Earth, 180–181, 182  
  on Jupiter, 277, 278  
  on Saturn, 297  
Autolycus crater, 205  
Autumnal equinox, 16, 17

## B

B ring, 298, 300, 301  
Background noise, 110  
Background radiation, ripples in, 707–709  
Balmer, Johann, 86  
Balmer (hydrogen) series, 86  
Bands, atmospheric, 268–270  
Barnard, E. E., 279  
Barnard 68 cloud, 71, 451, 458, 538  
Barnard 86 cloud, 150  
Barnard's star, 423, 424, 427, 431, 432, 439  
Barred-spiral galaxies, 609–611, 614  
Barringer Meteor Crater, 204, 205, 360  
Baryon acoustic oscillations, 709  
Basalt, 170  
Baseline of interferometer, 120  
Baseline triangle, 24  
Becklin–Neugebauer object, 481–482  
Bell, Jocelyn, 545  
Belts, in atmospheric bands, 268–270  
Berry, Dana, 715  
Beryllium-8, 499  
Bessel, Friedrich, 42  
Beta Persei, 515–516  
Beta Pictoris, 148  
Beta ring (Uranus), 333  
Betelgeuse, 427, 428, 431, 432, 434, 436, 439,  
  510, 512  
Big Bang theory, 671–673, 677, 680, 682,  
  683, 685, 693  
  early universe and, 690–692, 700, 701  
  galaxy formation rate after, 647, 648  
  quantum fluctuation and, 702  
“Big Crunch,” 674  
Big Horn Medicine Wheel, 34  
Binary black hole, 567, 653, 654  
Binary galaxy systems, 640, 641  
Binary pulsar, 568  
Binary-star systems, 440–441  
  black holes in, 564–567, 569  
  brown dwarfs in, 479  
  double-pulsar, 568  
  gravitational radiation from,  
  554, 568  
  mass in, 441–442  
  measuring, 443  
  neutron stars and, 548–552

planets in, 727  
  stellar evolution in, 515–517  
Biochemistries, alternative, 724  
Biological evolution, 721  
Bipolar flow/jets, 483–484  
Black dwarfs, 507  
Black hole(s), 555–557  
  binary, 567, 653, 654  
  in binary-star systems, 564–567, 569  
  in centers of galaxies, 569  
  curved space and, 560–561  
  defined, 556  
  distinguishing neutron star from,  
  569–571  
  escape speed of, 556  
  event horizon of, 557–558, 562–563  
  as final stage of stellar evolution,  
  555–556  
  in galaxies, 569  
  heating of, 561–562  
  intermediate-mass, 569, 570  
  masses of, 652, 653  
  observational evidence for, 464–471  
  properties of, 556  
  space travel near, 561–564  
  stellar transits of, 564  
  supermassive, 569, 601–603, 624, 652  
  tidal forces of, 561–562  
“Black smoker,” 723  
Blackbody, 68  
Blackbody (Planck) curve, 68, 69, 70–71, 73,  
  428–429, 683, 684  
Blazar, 627  
Blue giants, 436  
Blue stragglers, 507, 508  
Blue supergiants, 436  
Blueshifted radiation, 74, 75, 588  
Bode, Johann, 320  
Bohr, Niels, 84, 86  
Bohr model, 84  
Boltzmann, Ludwig, 72  
Borealis Basin, 242  
Boson, 695  
Bradley, James, 42  
Brahe, Tycho, 44, 530  
Brightness, 110  
  absolute, 425. *See also* Luminosity  
  apparent, 425–426  
Brown dwarfs, 368, 370, 378, 478, 479, 512  
Brown oval, 272, 273  
Bruno, Giordano, 42  
Bug (NGC 6302) Nebula, 495, 496  
“Bullet cluster” (1E 0657-56 galaxy cluster),  
  661, 662  
Bunsen, Robert, 83  
Burst counterparts, 553  
Butterfly (NGC 6302) planetary nebula,  
  495, 496

## C

C ring, 298, 300  
Caldera, volcanic, 225  
Calendar, 34–35  
Callisto (moon), 42, 279, 280–281, 285–287,  
  307, 308, 309  
Caloris Basin, 207–208  
Calypso (moon), 314  
Canada–France–Hawaii telescope, 107  
“Canals,” Martian, 132  
Canis Major (Great Dog), 14  
Cannon, Annie, 584  
Capture theory, 210  
Caracol temple (Mexico), 35  
Carbon, life and, 722, 724  
Carbon atom, 90  
Carbon burning, 534–535  
Carbon core, of Sun-like star, 501–502  
Carbon dioxide  
  in Earth's atmosphere, 166, 167, 168, 258  
  on Mars, 252, 258–259  
  in Venus's atmosphere, 230, 233, 258  
Carbon fusion, 534–535  
Carbon-12, 534

- Carbon-14, 172  
Carbonaceous (C-type) asteroids, 341  
Carbonaceous meteorites, 362  
Carbonate rock, 246  
Carbon-detonation (Type I) supernovae, 527–528  
  energy emission, 537  
Caribbean plate, 175  
Carina Nebula, 448, 449, 490  
“Cartwheel” galaxy, 643  
Cassegrain, Guillaume, 103  
Cassegrain focus, 103  
Cassegrain telescope, 103  
Cassini, Giovanni Domenico, 298  
Cassini Division, 298, 301, 311  
*Cassini Equinox*, 306  
Cassini mission, 147, 265, 269, 291, 293, 295–296, 298, 299, 300, 301, 304, 305, 307, 308, 309, 311, 312  
  description of, 142, 147, 306  
*Cassini-Huygens* probe, 147, 314, 722  
Cassiopeia A (Cas A), 542, 543  
  supernova remnant, 126  
Castor double star, 114  
Cataclysmic variables, 581. *See also* Nova/ novae; Supernova/supernovae  
Catalyst, 164  
Catastrophic theories, of planet formation, 369  
Cat’s Eye Nebula (NGC 6543), 505  
Celestial equator, 12  
Celestial mechanics, 142  
Celestial poles, 12–13  
Celestial sphere, 12–13  
Cells, primitive, 719  
Celsius system, 69  
Centaur rocket, 205  
Centaurus A galaxy, 626, 634, 642  
Center for High Angular Resolution Astronomy (CHARA), 120–121  
Center of mass, 53  
Cepheid variable stars, 581–582, 583, 613, 615, 621  
Ceres (asteroid), 136, 340, 341, 356, 357  
  properties of, 138  
Cerro Paranal Observatory, Chile, 106, 107  
*Chandra X-Ray Observatory (CXO)*, 126–127, 490, 533, 537, 543, 546, 547, 551, 569, 600–601, 626, 653  
Chandrasekhar, Subramanyan, 126, 527  
Chandrasekhar mass, 527  
*Chandrayan-1* orbiter, 199  
*Change* program, 199  
Charge-coupled devices (CCDs), 109  
Charged particles, 63–64  
  interactions between, 411–412  
Charon (moon), 355  
Chemical bonds, 91  
Chemical evolution, 717–719  
Chemosynthesis, 724  
Chicxulub, Yucatán Peninsula, 351  
Chinese astronomy, 35–36  
Chlorine, 164  
Chlorofluorocarbons (CFCs), 164  
Chromatic aberration, 102  
Chromosphere, 390, 391, 398–399  
Chryse Planitia, 241, 242, 253  
Circinus galaxy, 625  
*Clementine* satellite, 199, 204–205  
Cleopatra crater, 222–223, 227  
Climate  
  greenhouse effect and, 167  
  solar activity and, 409  
Climate change, on Mars, 252–253  
Close encounters, collision hypothesis and, 369, 369n  
Closed universe, 676  
Cloud(s)  
  dark dust, 150, 459–461  
  interstellar, 368, 369, 463–465, 472–477, 478, 480–481  
  organic compounds in, 464n  
  on Jupiter, 267, 269, 270–272, 276  
  on Neptune, 323  
  Oort, 353  
  on Saturn, 294, 295  
  on Uranus, 319  
Clusters  
  of galaxies, 7, 617–619, 640–641  
  of galaxy clusters, 656–657  
  star. *See* Star clusters  
CNO cycle, 534  
Cobalt-56, 535, 537, 538  
Cobalt-59, 536  
Cobalt-60, 536  
*COBE (Cosmic Background Explorer)*, 684, 707–709  
Cocoon nebula, 482–483  
Coformation (sister) theory, 210  
Cold dark matter, 706  
Cold Jupiters, 374, 379–381  
Collecting area, 105  
Collision hypothesis, 369  
Collisional broadening, 95  
Collisions  
  among galaxies, 120, 124, 643–645, 646  
  of galaxy clusters, 661–662  
  of comets, 274–275  
  fragmentation and, 150  
  interstellar, 508  
  protostellar, 489  
Color-magnitude diagrams, 434, 435  
Colors of stars, 428–429  
Coma (“halo”), 346  
Coma Cluster, 608, 618, 657  
Comet(s), 136, 143, 144, 155–156, 345–353.  
  *See also specific comets*  
  appearance and structure of, 346–347  
  collision of, 274–275  
  with Jupiter, 274  
  fragments of, 359–360  
  interstellar origin of life and, 719, 720  
  nucleus of, 346  
  orbits of, 345, 352–353  
  properties of, 138, 347–348  
  space missions to, 348–351  
  tail of, 345, 346, 347  
Comparative planetology, 137  
*Compton Gamma-Ray Observatory (CGRO)*, 127, 552, 601  
Condensation nuclei, 148  
Condensation theory, 148, 150–152, 362  
  and collisions, 369  
  dust’s role in, 368  
  as evolutionary theory, 369  
  extrasolar orbits and, 379  
  jovian planets and, 152–154  
  of planet formation, 148, 150–152, 368–369  
  solar system formation and, 148, 150, 368–369  
Conjunction, 37–38, 218  
Conservation of angular momentum, 145, 149  
Conservation of mass and energy, law of, 411  
Constant(s)  
  cosmological, 679, 681  
  Hubble’s, 620–621  
  Newton’s, 51  
  Planck’s, 85  
  solar, 390  
  Stefan-Boltzmann, 72  
Constellation, 10–12  
Constructive interference, 67  
Contact binary system, 517  
Continental drift, 173–175  
Continuous spectra, 80, 83, 90  
Convection, 163–164  
  and Jupiter’s atmospheric bands, 268–270  
  and plate drift, 178–179  
  solar, 395–396  
Convection cells, 164  
Convection zone, 390, 391  
Co-orbital satellites, 313–314  
Copernican principle, 43  
Copernican revolution, 32–55  
  ancient astronomy and, 34–36  
  and birth of modern astronomy, 41–42  
  and birth of modern science, 32–55  
  defined, 40  
  dimensions of solar system, 47–49  
  foundations of, 40  
  geocentric universe and, 36–39  
  heliocentric model of solar system, 39–40  
  laws of planetary motion, 44–47  
  Newton’s laws and, 49–52  
Copernicus, Nicholas, 39–41, 42, 55  
Copernicus crater, 202, 247, 248  
Coplanar motion, of planetary orbits, 379  
Coprates canal, 243  
Coral deposits, 184  
Cordelia (moon), 334  
Cordillera Mountains, 202  
Core, solar, 390, 391  
Core hydrogen burning, 496  
Core of Earth, 162, 169, 170  
Core-accretion theory, 152–153  
Core-collapse (Type II) supernova, 526, 527–528  
Core-dominated radio galaxy, 626, 627  
Corona (volcanic structure), 226  
Corona, solar, 20, 390, 391, 399–400, 407–410  
Coronal holes, 407–408  
Coronal mass ejection, 406, 407  
CoRoT (extrasolar planet), 377, 378  
CoRoT mission, 372–373  
Corpuscular (particle) theory of light, 67  
*Cosmic Background Explorer (COBE)*, 684, 707–709  
*Cosmic Background Imager*, 708  
Cosmic density parameter, 676, 678  
Cosmic distance ladder, 621  
Cosmic distance scale, 24, 582–583  
  extending, 437–439, 583, 615–617  
Cosmic evolution, 716–721  
Cosmic history  
  epochs in, 693  
  evolutionary phases in, 716  
Cosmic inflation, 701–702  
  flatness problem and, 702–704  
  horizon problem and, 702–704  
  as theory, 701–702, 704  
Cosmic microwave background, 683–685, 690, 699  
  cosmic structure and, 707–709  
  isotropy of, 700, 705  
  map of, 707, 708  
Cosmic rays, 601  
Cosmological constant, 679, 681  
Cosmological distances, 620  
Cosmological principle, 669, 672  
Cosmological redshift, 619, 620, 673  
  relativistic, 622–623  
Cosmology, 36, 666–685, 668  
  Big Bang theory. *See* Big Bang theory  
  cosmic microwave background, 683–685, 690, 699, 705, 707–709  
  cosmological principle, 669, 672  
  cosmological redshift, 619, 620, 622–623, 673  
  dark energy and, 680–683  
  defined, 668  
  expansion of universe, 670–673  
  dark energy and, 679–680  
  future of universe, 674–675  
  geometry of space, 675–677  
  Olbers’s paradox, 670  
  relativity and, 675–676  
Cosmos  
  density of, 673–674, 676, 678, 680, 690, 698–699  
  large-scale structure of, 638–662  
Coudé focus, 103  
CP1919 pulsar, 545  
Crab Nebula, 520, 521, 528, 529, 547  
Crab pulsar, 547, 548  
Crater(s)  
  lunar, 188–189, 192, 193, 194, 201–203  
  on Mars, 236–237, 244  
  on Mercury, 194, 195, 206  
  on Saturn’s moons, 308, 311, 312  
  on Venus, 227  
Crater-chain pattern, 206  
Crescent Moon, 18, 19  
Critical density, 673–674, 676  
Critical universe, 676  
Crust of Earth, 162  
C-type asteroid, 341, 362  
Cultural evolution, 721  
*Curiosity* (spacecraft), 236–237  
*Curiosity* rover, 133, 722  
Current sheet, 277, 278  
Curtis, Heber, 575, 586  
Curved space, 560–561, 676–677  
*CXO (Chandra X-Ray Observatory)*, 126–127, 490, 533, 537, 543, 546, 547, 551, 569, 600–601, 626, 653  
Cygnus (constellation), 367, 565  
Cygnus A galaxy, 626, 627, 629  
Cygnus loop supernova remnant, 125  
Cygnus X-1, 565–566  
D ring, 298, 301  
DA14 (asteroid), 343  
Dactyl (moon), 341, 342  
Daphnis (moonlet), 301  
Dark dust cloud, 150  
  obscuration of visible light by, 459–460  
Dark dust clouds, 459–461  
Dark energy, 679–680, 690, 710  
  and cosmology, 680–683  
  repulsive effect of, 679–680  
Dark halo, 596  
Dark matter, 596–598, 699, 710  
  acoustic oscillations and, 709–710  
  galaxies and, 638–662  
  mapping, 661–662  
  microwave background and, 709  
  stellar, 699  
  search for, 598–599  
  in universe, 640–642, 678, 705–707  
Dark nebula, 453  
Dark-energy era, 693  
Dark-matter halo, 640, 641  
Daughter nuclei, 172  
Daughter (fission) theory, 210  
Davy crater, 206  
*Dawn* mission, 147, 341, 342–343  
Death, of Sun-like star, 502–509  
Debris, interplanetary, 379  
Deceleration, cosmic, 678–679  
Decoupling, 699, 700  
*Deep Impact* (spacecraft), 351, 352  
Deep space, 666, 667  
Deferent, 38  
Deimos (moon), 239, 260  
Delta, 246  
Delta Cephei, 581  
Delta ring (Uranus), 333  
Dense interstellar cloud. *See* Dark dust cloud  
Density, 139. *See also specific bodies*  
  cosmic density parameter, 676, 678  
  critical, 673–674  
  of Earth’s interior, 162, 170  
  of emission nebula, 457  
  infinite, 563–564  
  of interstellar medium, 452  
  of Moon, 208  
  of solar system objects, 138  
  of Sun, 391, 394  
  uncompressed, 141  
  of universe, 673–674, 676, 678, 680–681, 690, 698–699  
Density waves, spiral, 300, 592–594  
Deoxyribonucleic acid (DNA), 718  
Descartes, René, 145  
Destructive interference, 67  
Detached binary system, 517  
Detectors, 109–111, 692  
Deuterium, 411, 412, 696, 697, 698–699  
Deuterium bottleneck, 697  
Deuterium fusion, 479  
Deuterons, 411, 412, 697  
*Dialogue Concerning the Two Chief World Systems* (Galileo), 42  
Diameter, angular, 28  
Differential force (tidal force), 183  
  of black hole, 561–562  
  exerted by Sun, 200–201  
Differential rotation, 267  
Differentiation  
  of Earth, 170–172  
  of solar system, 150–152  
Diffraction, 106–108  
Diffraction grating, 80  
Diffraction property of light, 67  
Diffraction-limited resolution, 107  
Dinosaurs, extinction of, 344, 350–351  
Dione (moon), 303, 310, 311, 314  
Direct (prograde) motion, 37  
*Discovery* (space shuttle), 104  
Disk galaxies, 579  
Distance  
  cosmological, 620  
  look-back time and, 623  
  measurement of, 24–27, 28  
  techniques for, ladder of, 437–438, 583, 621  
Distance scale, cosmic, 582–583  
  extending, 437–439, 583, 615–617  
Disturbance, 61  
Diurnal motion, 13  
DNA (deoxyribonucleic acid), 718  
Doppler, Christian, 73

- Doppler effect, 73–75, 93–95  
 detecting extrasolar planets with, 371  
 Doppler equation, 73, 75  
 Doppler shift, 92–93  
 Double cluster (h and chi Persei), 513, 514  
 Double-line spectroscopic binary, 440, 441  
 Double-pulsar binary system, 568  
 Drake equation, 724–725  
 Dust  
 dark dust clouds, 150, 459–461  
 interplanetary, 143  
 interstellar. *See* Interstellar dust  
 lunar (regolith), 203–204  
 Rayleigh scattering and, 165  
 Dust grain, 450  
 Dust lanes, 454, 455  
 Dust tails, 346  
 Dwarf(s), 433–434  
 black, 507  
 brown, 368, 370, 378, 478, 479  
 red, 433  
 white. *See* White dwarf  
 Dwarf elliptical galaxies, 611, 612, 640  
 Dwarf irregular galaxies, 612, 640  
 Dwarf planets, 144, 357  
 Dynamo theory, 182  
 Dysnomia (moon), 356
- E**
- E ring, 298, 301, 302, 312  
 Eagle Nebula (M16), 122, 123, 454, 456  
 Early universe, 688–710  
 Big Bang and, 690–692  
 evolution of, 693–697  
 inflation of, 700–704  
 nuclei and atom formation, 697–700  
 particle production in, 691–692  
 structure in, 705–707  
 Earth, 6, 160–185  
 atmosphere of, 162–168, 231–232, 258  
 core of, 162, 169, 170  
 density of, 139  
 differentiation of, 170–172  
 distance to Moon, 190  
 escape speed of, 196  
 exoplanet closest to, 376  
 gravity of, 209  
 hydrosphere, 162  
 interior of, 168–173, 209  
 core, 162, 169, 170  
 crust, 162, 169–170  
 density, 170  
 mantle, 162, 169, 170  
 modeling, 169–170  
 life on, 720–721  
 magnetic field of, 64, 176–177  
 reversals in, 177–178  
 magnetosphere of, 162, 180–182  
 measuring, 27  
 motion through the cosmos, 684  
 nearest stars to, 423–424  
 orbital motion of, 14–18  
 properties of, 138  
 radar map of, 223  
 radio radiation from, 730–731  
 revolution around Sun, 14  
 rotation of, 12, 183–184  
 solar-terrestrial relations, 409  
 structure of, 162  
 surface activity, 173–180  
 surface heating of, 166–168  
 tides of, 182–184  
 velocity of, 685  
 water on, origins of, 185  
 Earth-crossing (Apollo) asteroid, 340, 343–344  
 Earth–Moon system, formation of, 370  
 Earthquake, 168–169  
 Earths (extrasolar planets), 374  
 searching for, 381–382  
 super, 373–374, 375, 377  
 Eccentricity, 46  
 of ellipse, 45  
 Eclipse(s), 19–24  
 annular, 21–22  
 geometry of, 23  
 lunar, 19–22  
 Pluto–Charon, 354, 355  
 solar, 20–22, 23, 398–400, 410  
 tracks of, 24  
 Eclipse seasons, 22–24  
 Eclipse year, 24  
 Eclipsing binary, 441  
 Ecliptic, 14  
 Eddington, Arthur, 393, 566, 586  
 Eddington crater, 202  
 Effective temperature of Sun, 390  
 Einstein, Albert, 51, 85, 88, 411, 416, 679  
 cosmological constant and, 681  
 theories of relativity, 74n, 75, 557–561, 566–567, 675–676  
 Einstein Cross, 660  
 Einstein Observatory (spacecraft), 126  
 Einstein's elevator, 558, 560  
 Ejecta, fluidized, 247, 248  
 Ejecta blanket, 202  
 Electric field, 63  
 Electrical charge, 63  
 Electromagnetic force, 413, 694, 695  
 Electromagnetic radiation, 60. *See also* Radiation  
 Electromagnetic spectrum, 65–68  
 Electromagnetic wave, 64–65  
 Electromagnetism, 64  
 Electron, 63  
 production of, 691, 692  
 spin of, 462  
 Electron cloud, 85  
 Electron degeneracy, 555, 556  
 Electron degeneracy pressure, 500, 500n  
 Electron transitions, 457  
 molecular, 91  
 Electron volt (eV), 86  
 Electroweak force, 413, 694, 695  
 Element(s), 90. *See also specific elements*  
 emission spectra of, 81  
 “fifth,” of universe (quintessence), 680, 680n  
 formation of, 530–538  
 abundance of matter, 531–533  
 carbon burning and helium capture, 534–535  
 heaviest elements, 536  
 hydrogen and helium burning, 533–534  
 iron formation, 535–536  
 neutron capture, 536  
 stellar nucleosynthesis, 536–538  
 types of matter, 530–531  
 primordial, 531  
 Ellipse, 45  
 Elliptical galaxies, 611–612, 614, 632  
 Elongation, maximum, 191  
 Emission lines, 80–81  
 Emission nebulae, 89, 90, 91, 453–458  
 Carina, 490  
 “forbidden” lines of, 457–458  
 interstellar shock waves generated by, 484–485  
 observations of, 453–456  
 properties of, 457  
 spectra, 456–457  
 Emission spectrum, 81  
 molecular, 91  
 Enceladus (moon), 301, 302, 303, 310, 311–312, 722  
 Encke gap, 298  
 Endeavour (space shuttle), 104  
 Endurance crater, 254  
 Energy  
 of active galaxies  
 emission, 633–634  
 production, 630–632  
 black hole and production of, 561–562  
 dark, 679–683, 690, 710  
 galaxy energy spectra, 623–624  
 generated by proton–proton chain, 412–414, 416  
 Jupiter's internal energy source, 275–276  
 kinetic, 88  
 law of conservation of mass and, 149  
 mass-energy equivalence, 410–411  
 nuclear binding, 536  
 quantized orbital energies, 85  
 solar energy production, 410, 416  
 vacuum, 702  
 Energy flux, 72, 425  
 Energy levels, atomic, 84  
 Energy transport, in solar interior, 394–396  
 Envelope, stellar, 503  
 ejected, 504  
 Epicycles, 38  
 Epimetheus (moon), 313  
 Epochs in cosmic history, 693.  
*See also specific epochs*  
 Epsilon ring (Uranus), 333  
 Equilibrium  
 hydrostatic, 392–393, 496  
 thermal, 691, 692  
 Equinox, 16, 17  
 Equivalence principle, 560  
 Eratosthenes, 27  
 Eris, 356, 357–358  
 Eros (asteroid), 342  
 Escape speed, 54–55, 196–197  
 of black hole, 556  
 calculation of, 196–197  
 critical density and, 673, 674  
 of Earth, 196  
 of Mercury, 191, 197  
 of Moon, 191, 197  
 Eskimo Nebula, 505  
 Eta Carinae, 490  
 Eta ring (Uranus), 333  
 Euclid, 28, 677  
 Euclidean geometry, 677, 682  
 Eurasian plate, 175, 177  
 Europa (moon), 42, 279, 280, 283–285, 722  
 European Southern Observatory, 98–99, 106, 112, 113  
 European Space Agency, 395, 424, 708, 722  
 European Very Large Telescope, 368  
 EUVE (*Extreme Ultraviolet Explorer*), 125  
 Event horizon, 557–558, 562–563  
 singularities and, 563–564  
 “Evil Eye” galaxy (M64), 640  
 Evolution  
 biological, 721  
 chemical, 717–719  
 cosmic, 716–721  
 cultural, 721  
 Evolutionary theory, condensation theory as  
 example of, 369  
 Evolutionary tracks, 475  
 in binary-star systems, 515–517  
 high-mass, 509–510, 512  
 prestellar, 477  
 protostellar, 475–476  
 Excited state, 85, 89  
 Extinction, 450  
 Extrasolar planets (exoplanets), 368, 373–377  
 closest to Earth, 376  
 composition of, 375, 377  
 discovery of, 137, 370–373  
 hot Jupiters and super-Earths, 373–374  
 identification of, 376  
 orbits, 371, 372, 373, 374–375, 379–380  
 properties of, 373–378  
 Extraterrestrial intelligence, 729–732  
 Extraterrestrial life, case for, 717  
 Extreme Ultraviolet Explorer (EUVE), 125  
 Extremely Large Telescope (ELT), 98–99  
 Extremophiles, 723  
 Eyepiece, 101
- F**
- F ring, 298, 302, 303–304  
 Fahrenheit temperature scale, 69  
 Failed star, 478. *See also* Brown dwarfs  
 False-color technique, 113, 120, 121, 122, 123  
 Fermi, Enrico, 127  
 Fermi Gamma-Ray Space Telescope, 127–128  
 51 Pegasi star, 371, 372  
 Filters, telescope, 110, 428  
 Fireballs, 360  
 primeval, 670–671, 693, 696  
 Firmament, 40  
 Fission, 525  
 Fission (daughter) theory, 210  
 Flares, solar, 406, 407  
 Flashes, helium-shell, 503  
 Flatness problem, 700–701  
 inflation and, 702–704  
 Fleming, Williamina, 584  
 Fluidized ejecta, 247, 248  
 Fluorescence, 88  
 FM radio signal, 164, 730  
 Focal length, 100  
 Focus  
 of ellipse, 45  
 of telescope, 100  
 Fomalhaut (star), 124, 150  
 “Forbidden” lines of emission nebulae, 457–458  
 Force(s), 49  
 fundamental, 413, 694, 695  
 gravitational, 51, 52, 53  
 Formaldehyde, 463, 464–465, 464n  
 Fornax A, 606, 607  
 47 Tucanae, 513–514, 515, 551, 597  
 “Fossil evidence,” galaxy formation and, 646  
 Fossil record, 167, 719, 720, 721, 722  
 Fragmentation, 150  
 Fraunhofer, Joseph von, 82  
 Freeze-out, 692, 695–696  
 Frequency, 61–62  
 photon energy and, 85  
 Full Moon, 18, 19, 20, 192, 194  
 Full-spectrum astronomy, 128  
 Fundamental forces, 413, 694, 695  
 Fusion  
 carbon, 534–535  
 deuterium, 411, 479  
 helium, 499–501, 534  
 nuclear, 410–411  
 proton, 411–412, 534
- G**
- G ring, 298, 301, 302, 304  
 GALA project, 424  
 Galactic activity, 599–601  
 Galactic bulge, 578, 588n  
 properties of, 589  
 Galactic cannibalism, 648–649  
 Galactic center, 422, 585, 599–603, 726  
 orbits near, 588–589, 601  
 Galactic disk, 578, 729  
 interstellar gas in, 591–592  
 orbital motion in, 588–589  
 properties of, 589  
 spiral structure and, 592–593  
 Galactic epoch, 696, 705  
 Galactic gas, 586  
 Galactic habitable zone, 726–727  
 Galactic halo, 578, 583n, 586–587  
 properties of, 589  
 Galactic nucleus, 601. *See also* Active galaxies/active galactic nuclei  
 spiral galaxies and, 608  
 supermassive black holes in, 569, 601–603, 624  
 Galactic plane, 450, 454, 588  
 Galactic rotation, 592–593, 596, 616  
 Galactic rotation curve, 596  
 Galactic “tuning fork,” 614  
 Galactic year, 588  
 Galaxy(ies), 7, 606–634. *See also* Active galaxies/active galactic nuclei; Milky Way Galaxy  
 black holes in, 569, 652–656  
 colliding, 120, 124, 643–645, 646  
 and dark matter, 638–662  
 defined, 578  
 disk, 579  
 distribution in space, 615–618  
 formation and evolution of, 645–650, 655, 705–706  
 “Great Debate” over, 574–575, 585–586, 609  
 Herschel's model of, 580  
 Hubble classification scheme, 608–614  
 elliptical galaxies, 611–612, 614  
 irregular galaxies, 612–613, 614  
 spiral galaxies, 608–611, 614  
 “tuning fork” diagram, 614  
 Hubble's law of, 619–621  
 interactions between, 646–647, 650  
 irregular, 640  
 large-scale distribution of, 656–657  
 life in, 724–729  
 masses of, 640–641  
 normal, 622, 655–656  
 properties of, 614  
 radio, 569, 625–628  
 receding, 672  
 redshift survey of, 657–658, 668  
 rotation curves of, 640  
 spectra of, 619–620  
 spiral. *See* Milky Way Galaxy; Spiral galaxies  
 starburst, 623–624, 644  
 surveys of, 657–658, 668–669  
 visible, 621n



- Galaxy clusters, 7, 617–619, 640–641  
clusters of, 656–657  
collisions of, 643–645  
distant, 618–619  
lensing, 661  
mass of, 641  
Perseus, 651  
X-ray emission, 641, 642
- Galaxy Evolution Explorer (GALEX)* satellite, 125
- Helix Nebula visualization by, 59
- Galaxyrise Over Alien Planet* (painting), 714, 715
- Gale crater, 236–237
- Galilean moons, 42, 266, 279–287, 305  
Callisto, 279, 280–281, 285–287  
Europa, 279, 280, 283–285  
Ganymede, 279, 280–281, 285–287  
Io, 279, 280, 281–283
- Galileo Galilei, 2, 41–42, 45, 49, 50, 136, 298, 321, 401, 529
- Galileo* missions, 147, 271, 272, 274, 275, 277, 279, 280, 281–282, 283, 284, 285, 286, 306, 341–342
- Galileo Regio, 285
- Galle, Johann, 321
- Galle ring (Neptune), 334
- Gamma rays, 60, 66, 412–414, 692
- Gamma ring (Uranus), 333
- Gamma-ray astronomy, 127–128
- Gamma-ray bursts, 552–555  
counterpart of, 553  
models of, 554
- Gamma-ray pulsars, 548
- Gamma-ray radiation, 65
- Ganymede (moon), 42, 279, 280–281, 285–287, 722
- Gaposhkin, Sergei, 386
- Gas  
greenhouse, 166  
interstellar, 450, 452, 455, 456, 457, 458, 591–592  
intracluster, 641–642
- Gas turbulence, 94
- Gas-exchange experiment, 250
- Gaspra (asteroid), 341–342
- Geminga pulsar, 548
- Gemini North telescope, 106, 107
- Gemini South telescope, 106
- Gene(s), 718  
of virus, 717
- General relativity, 558, 560, 675–676, 690  
tests of, 566–567
- Geocentric universe, 36–39
- Geomagnetic activity at Earth, 409
- Geometric scaling, 25
- Geometry  
of eclipses, 23  
Euclidean, 677, 682  
measuring distances with, 24–27, 28  
Riemannian, 677  
of space, 675–677
- Geysers, 311, 312  
nitrogen, 331
- Giacobini-Zinner (comet), 347
- Giant elliptical galaxy, 611
- Giant planets, 152–154. *See also* Jovian planets  
migration of, 154
- Giants, 433–434  
blue, 433, 436  
mass loss from, 511  
red, 433, 434, 499
- Gibbous phase of Moon, 18, 19
- Giotto* (spacecraft), 348, 349
- GJ 1214B (extrasolar planet), 377  
mass of, 377
- Glashow, Sheldon, 694
- Gliese 229, 479, 623
- Global warming, 167. *See also* Greenhouse effect
- Global clusters, 487–488, 507, 580–581, 583n, 585  
in Andromeda galaxy, 569  
M4, 460  
M9, 420, 421
- Globule (interstellar cloud), 451
- Globules, interstellar, 719, 720
- Glouons, 695
- Glycine, 464n
- GÖNG (Global Oscillations Network Group) project, 393
- GRAIL (Gravity Recovery and Interior Laboratory)*, 199, 209, 212
- Grand Canyon, Martian (Mariner Valley), 242–243
- Grand Unified Theories (GUTs), 694, 695, 704
- Granite, 170
- Granulation, solar, 396–397
- Gravitational competition, 471
- Gravitational constant, 51
- Gravitational deformation, 182–183
- Gravitational force, 52, 413
- Gravitational instability theory, 153
- Gravitational lensing, 598, 659–660, 659n  
galaxy clusters, 661  
microlensing, 659
- Gravitational radiation, 554, 568
- Gravitational redshift, 562–563
- Gravitational slingshot (gravity assist), 142, 147, 306
- Gravitons, 694, 695
- Gravity, 694  
atmosphere and, 196–197  
of Earth, 209  
expansion of universe slowed by, 673–674  
in general relativity, 558, 560  
heat and, 471–472  
on Jupiter, 283  
magnetism and, 471  
on Mercury, 191  
on Moon, 191  
Newton's law of, 49, 51–52  
particles and, 695  
quantum, 693, 695  
rotation and, 471  
solar, 52  
tides and, 182–183
- Gravity Recovery and Interior Laboratory (GRAIL)*, 199, 209, 212
- Gravity wave, 568
- GRB 080319B, 553
- Great Dark Spot, 325–326
- Great Red Spot, 264, 265, 268, 269, 272, 273, 274
- Great Wall, 657–658, 668
- Greenhouse effect, 166, 168  
on Earth, 231–232  
global warming and, 167  
“reverse runaway” on Mars, 256–257  
runaway, 162, 231–232  
on Venus, 231–232
- Greenhouse gas, 166, 233
- Ground state, 84
- G-type star, evolution of, 502–503
- Gula Mons, 225
- GUT epoch, 695, 696, 701, 702, 706
- GUTs (Grand Unified Theories), 694, 695, 704
- H**
- Habitable zones, 381  
galactic, 726–727  
stellar, 725–726
- Hadley, Mount, 206
- Hadley Rille, 206
- Hale, George Ellery, 400, 401
- Hale-Bopp (comet), 347, 719, 720  
properties of, 138
- Half-life, 172
- Hall, Asaph, 132, 260
- Halley, Edmund, 49, 347
- Halley's comet, 346, 347, 349, 720  
nucleus of, 348  
orbit of, 349
- Halo stars, 588, 588n
- Harvard College Observatory, 584
- Houghton crater (Arctic), 249
- Haumea (Kuiper belt object), 356, 357
- Hayabusa* (spacecraft), 135
- Hayashi, C., 476
- Hayashi track, 476, 482
- Haystack radio telescope, 117
- HD 154345 (extrasolar planet), 380
- HD 209458 (star), 372, 373
- HDE 226868 (blue supergiant), 565, 567
- Head-tail radio galaxies, 641–642
- Heat (thermal energy), 69  
atmosphere and, 166–168  
gravity and, 471–472  
solar system formation and, 148
- Heating, black hole, 561–562
- Helene (moon), 314
- Heliocentric model, 39–40
- Helioseismology, 393–394
- Helium, 90, 696  
absorption lines in stars, 430–431  
emission spectrum of, 81  
formation in early universe, 697–698  
proton-proton chain and, 412–414
- Helium burning, 499
- Helium capture, 535
- Helium core, of Sun-like star, 497, 499, 501
- Helium flash, 499–501
- Helium fusion, 499–501, 534
- Helium precipitation, 297
- Helium white dwarf, 506, 517
- Helium-shell burning, 501
- Helium-shell flashes, 503
- Helix Nebula, 58, 59
- Hellas Basin, 241, 242, 246
- Hellas Planitia, 242
- Helmholtz, Hermann von, 475
- Henize Nebula, 387
- Herbig-Haro 46, 71
- Herbig-Haro objects, 483–484
- Herschel, William, 122, 320, 322, 580
- Herschel crater, 311
- Herschel Space Observatory*, 122, 123, 469
- Hertz (Hz), 62
- Hertz, Heinrich, 62
- Hertzsprung, Ejnar, 434, 584
- Hertzsprung-Russell (H-R) diagram, 434–437  
cluster evolution on, 512–513, 514–515  
development of, 584  
of globular clusters, 487–488, 507, 509  
high-mass evolutionary tracks, 510  
horizontal branch of, 500  
instability strip on, 582  
of newborn star, 476  
of open clusters, 486, 487, 512–513  
prestellar evolutionary tracks, 477  
of protostar's physical properties, 474–475  
red-giant branch of, 499  
red-giant region of, 436, 437  
of stars, 434–437  
and stellar mass, 442  
of white dwarf, 505
- Hewish, Antony, 545
- HH1 (bipolar jet), 483–484
- HH2 (bipolar jet), 483–484
- HH30 (star system), 483
- HI regions, 456
- Hierarchical merging, 645–646, 647
- High Energy Transient Explorer 2*, 555
- High-energy telescope, 125–128
- Higher (superior) planet, 37
- Highlands, lunar, 192–193, 194, 203
- High-mass stars  
end of, 524–526  
evolution of, 509–510, 512  
mass loss from, 511
- High-resolution telescopes, 111–114
- HI1 regions, 456. *See also* Emission nebula
- Himalayas, 176, 177
- Hipparchus, 426
- Hipparcos* satellite, 11, 424, 436, 437, 438, 567, 588
- History, learning astronomy from, 508
- Holden Crater, 247
- Holes, coronal, 407–408
- Hollows, 207
- Homo sapiens*, 721
- Homogeneity, cosmic, 669, 672
- Hooke, Robert, 272
- Horizon problem, 700–701  
inflation and, 702–704
- Horizontal branch, 500
- Horsehead Nebula, 460, 461
- Hot dark matter, 706
- Hot Jupiters (extrasolar planets), 373–375, 379–381
- Hot longitudes, 200, 208
- Hot springs, 722–723
- Hot stars, 482–483
- Hourglass, 456
- H-R diagram. *See* Hertzsprung-Russell (H-R) diagram
- HST. See* Hubble Space Telescope (HST)
- Hubble, Edwin, 104, 571, 608, 620, 679
- Hubble classification scheme, 608–614  
elliptical galaxies, 611–612, 614  
galaxy formation and, 645  
irregular galaxies, 612–613, 614  
spiral galaxies, 608–611, 614
- Hubble Deep Field, 647
- Hubble diagrams, 620
- Hubble flow, 620, 672
- Hubble sequence, 613–614, 649–650
- Hubble Space Telescope (HST)*, 104–105, 109, 110, 112, 120, 122, 124, 125, 240, 266, 273, 274, 294, 295, 297, 324, 326, 356, 367, 421, 432, 455, 469, 470, 485, 490, 506, 543, 546, 547, 597, 607, 629, 631, 646, 647, 651, 653, 654, 667
- Hubble Ultra Deep Field (UDF), 575, 666, 667
- Hubble's constant, 620–621, 670–671, 672, 690
- Hubble's law, 619–621, 628, 657, 670–671  
distance measurements and, 621
- Hulse, Russell, 568
- Huygens, Christian, 67, 298, 305
- Huygens* probe, 308
- Hyades cluster, 513, 514
- Hydra (moon), 354
- Hydrocarbon, 307, 308
- Hydrogen, 85, 86, 698  
absorption lines in stars, 431  
atomic, absorption lines of, 658–659  
emission spectrum of, 81, 83, 84  
energies of excited states of, 85, 89  
heavy. *See* Deuterium  
spectrum of, 80–81, 87–88
- Hydrogen 21-cm emission, 462–463
- Hydrogen envelope, 346
- Hydrogen (Balmer) series, 86
- Hydrogen-burning reaction, proton-proton, 533
- Hydrogen-shell burning stage, 497, 498
- Hydrosphere, 162, 182
- Hydrostatic equilibrium, 392–393  
of main-sequence star, 496
- Hydrothermal vents, 722–723
- Hyperbola, 55
- Hyperion (moon), 303, 304, 313
- Hypernova, 554, 555
- Hypothesis, 9
- I**
- Iapetus (moon), 303, 310, 312–313
- IC 694 (galaxy), 644
- IC 1396 (interstellar cloud), 482
- IC 2163 (galaxy), 643
- Ice, lunar, 204–205
- Ida (asteroid), 341–342
- Igneous rocks, 179
- Image processing, 110
- Images, 100–101  
formation of, 102
- Impact crater  
on Mars, 244, 248, 249  
on Venus, 227
- Impact theory, 210–211
- Inclination, 14, 16
- Index of Prohibited Books*, 41
- Indian plate, 175, 177
- Inertia, 50
- Inferior conjunction, 37–38, 218
- Inferior orbit, 37
- Inferior (lower) planet, 37
- Infinite density, 563–564
- Inflation  
cosmic. *See* Cosmic inflation  
epoch of, 701–702
- Infrared Astronomy Satellite (IRAS)*, 482
- Infrared radiation, 58, 65, 66
- Infrared telescopes, 121–123
- Inhomogeneities, growth of, 705
- Inner core, 169
- Instability strip, 582
- Intelligence  
development of, 721  
extraterrestrial, 729–732
- Intensity, 68
- Intercloud medium, 458
- Inter crater plains, on Mercury, 194
- Interference property of light, 67
- Interferometry, 118–121, 432  
optical, 120–121
- Intermediate-mass black hole, 569, 570
- International Astronomical Union, 357
- International Ultraviolet Explorer (IUE)*, 125
- Interplanetary debris, 379
- Interplanetary matter, 143–144
- Interstellar “bubbles,” 458

- Interstellar clouds  
 contraction of, evidence for, 478, 480  
 evolution of, 472–477  
 fragmentation of, 473  
 evidence for, 480–481  
 in planet formation, 368, 369  
 in star formation, 473–474
- Interstellar collisions, 508
- Interstellar dust, 148, 150, 450, 452  
 origin of life and, 719  
 particle shapes, 452–453
- Interstellar magnetic field, 471, 473
- Interstellar medium, 448–465  
 composition of, 452  
 dark dust clouds, 150  
 defined, 450  
 density of, 452  
 emission nebulae, 453–458  
 gas and dust, 148, 150, 450  
 molecules of, 463–465  
 21-centimeter radiation, 462–463
- Interstellar message, 730
- Interstellar space, first living cells produced in, 719–720
- Intracluster gas, 641–642
- Intrinsic variable, 581
- Inverse-square law, 52, 425–426, 430, 582
- Io (moon), 42, 279, 280, 281–283
- Io plasma torus, 282–283
- Ion, 84
- Ion tail, 346, 347
- Ionized atom, 84, 455–456
- Ionosphere, 68, 163, 164
- IRAS (*Infrared Astronomy Satellite*), 482
- Iridium, 351
- Iron, 90  
 formation of, 535–536
- Iron core, collapse in high-mass star, 525
- Iron meteorites, 362
- Irregular galaxies, 612–613, 614, 640
- Ishtar Terra, 222, 224
- Isidis Planitia, 242
- Islamic astronomy, 36
- Isolated spirals, 595
- Isotopes, 412  
 element formation and, 530
- Isotropy  
 cosmic, 669, 672  
 of cosmic microwave background, 700, 705
- Itokawa (asteroid), 134–135, 338, 339
- IUE (*International Ultraviolet Explorer*), 125
- J**
- James Webb Space Telescope (JWST)*, 122
- Jansky, Karl, 114–115
- Janus (moon), 313
- Jets  
 active galaxy, 630, 631, 632, 634  
 bipolar, 483–484  
 quasar, 629, 630  
 radio galaxy, 626–628
- Joule (J), 72
- Jovian planets, 140–141, 143. *See also* Jupiter;  
 Neptune; Saturn; Uranus  
 formation of, 152–154  
 interiors of, 327  
 magnetic fields of, 327  
 moons of, 154  
 and planetary debris, 152–156  
 relative sizes of, 322  
 ring systems of, 299, 300
- Joyce, James, 694
- Jumbled (weird) terrain, 208
- Juno (asteroid), 340
- Juno mission, 147
- Jupiter, 264–287  
 atmosphere of, 268–272  
 as failed star, 478  
 internal structure of, 274–277, 327  
 magnetic field and magnetosphere of, 277–278, 282–283, 327  
 mass and radius of, 266–267  
 moons of, 42, 279–287  
 orbit of, 266, 340, 344–345, 357  
 properties of, 138, 266–268  
 rings of, 287, 300  
 rotation rate of, 267–268  
 Shoemaker-Levy 9 comet, collision with, 351  
 significance to life on Earth, 276  
 spacecraft exploration of, 146–147  
 starlike composition of, 276  
 Trojan asteroids orbiting, 340, 344  
 views of, 266, 267  
 weather on, 272–274
- Jupiter-like planets, 379–380
- Jupiters (extrasolar planets), 373–374,  
 379–381  
 cold, 374, 379–381  
 hot, 373–375, 379–381
- K**
- Keck telescopes, 103, 104, 106, 107, 111, 113,  
 120, 319, 569
- Kelvin, Lord, 69, 475
- Kelvin temperature scale, 69
- Kelvin–Helmholtz contraction phase, 475,  
 478
- Kennedy, John F., 198
- Kepler (extrasolar planets), 377
- Kepler, Johannes, 44–45, 49, 55, 530
- Kepler mission, 372–373, 374  
 for extrasolar planets, 82, 379, 381
- Kepler supernova, 530
- Kepler's laws, 45–47, 52–53, 567
- Kinetic energy, 88
- Kirchhoff, Gustav, 83
- Kirchhoff's laws, 83, 84, 88–90
- Kirkwood, Daniel, 345
- Kirkwood gaps, 345
- Kitt Peak National Observatory (KPNO),  
 105, 111, 506
- Kuiper, Gerard, 328, 330, 352
- Kuiper belt, 139–140, 143–144, 352  
 orbit of, 357
- Kuiper-belt objects (KBOs), 136, 353–357
- L**
- L977 (dark dust cloud), 459, 460
- Labeled-release experiment, 250
- Lagoon Nebula (M8), 453, 454, 456
- Lagrange, Joseph Louis, 314, 344
- Lagrangian points, 314, 344, 345, 395,  
 516–517
- Lakshmi Planum, 222, 224
- Language, development of, 721
- Laplace, Pierre Simon de, 145
- Large Hadron Collider, 689
- Large Magellanic Cloud (LMC), 612, 613
- Las Campanas Observatory (Chile), 658
- Laser altimeter, 240
- Laser Interferometer Gravity-Wave  
 Observatory (LIGO)*, 568
- Lassell, William, 328, 330
- Lassell ring (Neptune), 334
- Lava, 179
- Lava domes, 225–226
- Lava flows, 224, 225
- LCROSS (*Lunar CRater Observation and  
 Sensing Satellite*), 205
- Leap year, 23
- Leavitt, Henrietta, 583, 584
- Lens/lensing, gravitational, 598, 659–660,  
 659n  
 galaxy cluster, 661  
 microlensing, 659
- Lenticular galaxies, 612–613
- Lepton epoch, 696
- Leptons  
 particles of, 694  
 quarks and, 696
- Leverrier, Urbain, 321
- Leverrier ring (Neptune), 334
- Lick Observatory, 114, 299
- Life  
 on Earth, 720–721  
 in Galaxy, 724–729  
 interstellar origin, evidence of, 719–720  
 on Mars, 250–251  
 search for extraterrestrial intelligence,  
 729–732  
 in solar system, 722–724  
 in universe, 714–732  
 working definition of, 716–717
- Light. *See also* Radiation  
 diffraction property of, 67  
 gravitational deflection of, 564, 565, 566  
 interference property of, 67  
 particle and wave property of, 85, 87  
 particle (corpuscular) theory of, 67  
 polarization of, 67, 453  
 radiation and, 63  
 scattering of, 165  
 speed of, 64–65, 557, 558  
 visible, 60, 62–63  
 dark matter and, 641  
 obscuration of, 459–460
- Light curves  
 from carbon-detonation (Type I) super-  
 novae, 537–538  
 eclipsing binary, 441  
 supernova, 527
- Light echo, 511
- Light pollution, 111
- Lighthouse model, of neutron-star emission,  
 545–547
- Light-years, 6–7
- LIGO (*Laser Interferometric  
 Gravity-Wave Observatory*), 568
- Lin, C. C., 594
- Line broadening, 93–95
- LMC X-3, 567, 569
- Lobachevsky, Nikolai Ivanovich, 677
- Local Bubble, 458
- Local Group, 656  
 contents of, 617, 618
- Local Supercluster, 656–657
- Logarithmic scale, 65
- Loki volcano, 281
- Long-period comets, 352–353
- Look-back time, 622–623
- Lorentz contraction, 559
- Lowell, Percival, 132–133, 353, 619
- Lowell Crater, 25
- Lower (inferior) planet, 37
- Low-mass star, evolution of.  
*See* Under Stellar evolution
- LRO (*Lunar Reconnaissance Orbiter*), 199,  
 205
- Luminosity(ies)  
 active galaxy, 630  
 gamma-ray burst distances and,  
 552–553  
 in Hertzsprung–Russell diagrams,  
 434–437  
 and mass and radius, 442–443  
 of nova, 522, 523  
 period–luminosity relationship and, 583  
 of protostar, 474  
 radius–luminosity–temperature  
 relationship, 432–434  
 of stars, 425–427, 430, 433  
 classification, 438–439  
 of Sun, 390–392, 409, 410, 433  
 of supernovae, 615–616, 616n  
 of variable star, 582–583
- Luna missions, 198
- Lunar CRater Observation and Sensing  
 Satellite (LCROSS), 205
- Lunar eclipse, 9, 19–22
- Lunar Orbiter (spacecraft), 198
- Lunar phases, 18–19
- Lunar Prospector satellite, 199, 205, 208, 209,  
 248, 259
- Lunar Reconnaissance Orbiter (LRO), 199,  
 205
- Lunar tides, 183, 184
- Lyman, Theodore, 86
- Lyman series, 86
- Lyman-alpha forest, 658–659
- M**
- M2-9, 505
- M4 globular cluster, 460, 506
- M8 (Lagoon) nebula, 453, 454, 456, 599
- M9 star cluster, 420, 421
- M16 (Eagle) nebula, 454, 456
- M17 nebula, 91, 454
- M20 nebula, 453, 454, 455  
 environment of, 464  
 molecules near, 465  
 star formation in, evidence of, 478, 480
- M32 galaxy, 617
- M33 galaxy, 617
- M49 galaxy, 611
- M51 (Whirlpool) galaxy, 649
- M64 (“Evil Eye”) galaxy, 640
- M80 globular cluster, 507
- M81 spiral galaxy, 125
- M82 galaxy, 125, 569
- M84 galaxy, 611
- M86 galaxy, 618, 627
- M87 (Virgo A) galaxy, 618, 628, 629, 632,  
 634  
 disk, 632
- M100 galaxy, 123, 615
- M101 galaxy, 105, 579
- M104 (Sombrero) galaxy, 610
- M110 galaxy, 611
- MACHOs (Massive Compact Halo Objects),  
 597
- Magellan, Ferdinand, 612
- Magellan mission, 147, 206, 221, 222, 223,  
 224, 226, 227, 309
- Magellanic Clouds, 612–613, 648
- Magma, 179
- Magnetic field lines, 180
- Magnetic fields, 64. *See also* under specific  
 planets  
 of active galaxy jets, 631, 633  
 of Earth, 176–177, 182  
 reversals in, 177–178  
 interstellar, 471, 473  
 of Mars, 259  
 of Mercury, 209  
 near Galactic center, 600  
 of Neptune, 326, 327  
 of newborn neutron stars, 545  
 spectral-line broadening and, 95  
 of Sun, 401–403  
 of terrestrial planets, 141, 143  
 of Uranus, 326, 327  
 of Venus, 232–233
- Magnetic poles, 180
- Magnetism, 64  
 gravity and, 471
- Magnetopause, 181, 277–278
- Magnetosphere, 162, 180–182. *See also* under  
 specific planets
- Magnitude scale, 426–427, 430
- Main sequence, 435–436  
 defined, 435  
 leaving, 496  
 in star formation, 476–477  
 well-know stars, main properties of,  
 444  
 zero-age (ZAMS), 477–478
- Main-sequence turnoff, 513
- Major axis, 45
- Makemake (Kuiper belt object), 356, 357
- Manicouagan Reservoir, 360
- Mantle of Earth, 162, 169, 170
- Mare Imbrium, 192, 193
- Mare Nectaris, 193
- Mare Nubium, 192, 193
- Mare Orientale, 227
- Maria, lunar, 192, 193, 203, 206, 212
- Mariner missions, 146, 192, 194, 206, 207,  
 232, 240, 244, 256, 259
- Mariner Valley, 241, 242–243
- Mars, 132–133, 236–261  
 atmosphere of, 256–259  
 climate change on, 252–253  
 craters on, 236–237  
 exploration of, 236–237  
 Gusev crater, 137  
 internal structure of, 259–260  
 life on, 250–251  
 long-distance observations of, 239–240  
 moons of, 239, 260–261  
 orbit of, 238–239  
 plate tectonics on, 259  
 properties of, 138, 239  
 search for life on, 222  
 spacecraft exploration of, 147  
 spacecraft landing sites on, 241  
 study of, 132–133  
 surface of, 132, 133, 240–244  
 “canals,” 132, 133  
 “Grand Canyon,” 242–243  
 “gullies” in cliffs and crater walls,  
 248, 249  
 impact cratering, 244  
 polar caps, 239–240, 249, 252, 257  
 topography of, 240–242  
 viewed by Martian landers,  
 253–256  
 volcanism, 243–244  
 water, 244–249
- Mars Exploration Rover, 147
- Mars Express mission, 133, 147, 245, 247,  
 248, 252, 722
- Mars Global Surveyor, 147, 240, 241, 242,  
 243, 245, 246, 248, 249, 259, 722

- Mars Odyssey* mission, 147, 248, 254  
*Mars Pathfinder* mission, 147, 254  
*Mars Phoenix* lander, 147  
*Mars Reconnaissance Observer*, 245, 248  
*Mars Reconnaissance Orbiter*, 147, 242  
*Mars Science Laboratory*, 147  
 Mars-crossing (Amar) asteroid, 340  
 Mass(es), 50  
   angular momentum and, 149  
   in binary-star systems, 441–442  
   measuring, 443  
   black hole, 652, 653  
   center of, 53  
   Chandrasekhar, 527  
   of emission nebula, 457  
   of galaxies, 640–641  
     Milky Way Galaxy, 595–599  
   and law of conservation of mass and energy, 149  
   mass loss from giant stars, 511  
   of the Moon, 191  
   of planets, 138  
     Jupiter, 266–267  
     Mars, 239  
     Mercury, 191  
   of solar system objects, 138  
   of stars, 440–444  
     distribution, 442  
     measuring, 443  
   of Sun, 54  
   turnoff, 513  
 Mass function, 438  
 Mass-energy equivalence, 410–411  
 Massive Compact Halo Objects (MACHOs), 597  
 Mass–luminosity relations, 442–443  
 Mass–radius relations, 442–443  
 Mass-transferring binary (semidetached binary) system, 522–523  
 Mathematical models  
   of star rotations, 591  
   of Sun, 392  
 Mathilde (asteroid), 341, 342  
 Matter, 690. *See also* Dark matter  
   basic components of, 690  
   interplanetary, 143–144  
   living vs. nonliving, 716–717  
   normal, 699, 705–706  
   radiation–matter decoupling, 699, 700  
 Matter era, 693, 699  
 Matter oscillations, 709–710  
 Matter-dominated universe, 690–691  
 Mauna Kea, Hawaii, 103, 106, 107, 111  
 Maunder minimum, 405, 409  
 Maury, Antonia, 584  
 Maximum elongation, 191  
 Maxwell, James Clerk, 223, 299  
 Maxwell Montes range, 222, 223  
 Mayall telescope, 105  
 Mead crater, 227  
 Mean solar day, 17  
 Measurement of distance, 24–27  
 Mediocrity, assumptions of, 717  
 Mercury, 188–213  
   atmosphere of, 191–192  
   evolutionary history of, 212–213  
   interior of, 209  
   *Mariner 10* flybys of, 192, 194, 206, 207  
   mass of, 138, 191  
   orbit of, 190–191  
   phases of, 191  
   properties of, 138, 191–192  
   rotation of, 196–199  
   solar transit of, 48  
   spacecraft exploration of, 146  
   surface of, 194–195, 206–208  
   tidal forces on, 199–200  
 Mercury (element), emission spectrum of, 81  
 Merger/merging  
   between galaxies, 645–646  
   types of, 648–649, 650  
   neutron-star model, 554  
   of Sun-like stars, 508  
 Mesosphere, 163  
 Message, interstellar, 730  
*Messenger* probe, 146, 147, 192, 194, 206, 207, 209, 212  
 Messier, Charles, 453, 453n  
 Messier 2 cluster, 71  
 Metamorphic rocks, 179  
 Meteor showers, 359, 359n, 360  
 Meteor trails, 358  
 Meteorites, 359  
   interstellar origin of life and, 719, 720  
   properties of, 361–362  
   samples of, 362  
   source of, 135  
 Meteoroid swarm, 359  
 Meteoroids, 143, 201–202, 358–362  
 Meteors, 136, 201, 358  
 Methane, 324  
 Metis (moon), 287  
 Michelson, A. A., 557, 559  
 Michelson–Morley experiment, 557, 559  
 Microlensing, 659  
 Micrometeoroid swarms, 588–589  
 Micrometeoroids, 202, 204, 359  
 Microquasar, 550  
 Microsphere, protein-like, 718–719  
 Microwave background, cosmic, 683–685, 690, 699  
   and cosmic structure, 707–709  
   isotropy of, 700, 705  
   map of, 707, 708  
 Mid-Atlantic Ridge, 176–177  
 Midocean ridges, 176  
 Migration, planet, 154  
 Milky Way Galaxy, 2, 4, 117–118, 128, 450, 453, 453n, 576–603  
   condensation theory and, 148  
   formation of, 589–590  
   Galactic center, 422, 585, 599–603  
   mass of, 595–599  
   measuring, 579–586  
   orbital motion of, 588–589  
   plane of, 450, 454, 578  
   radio maps of, 591  
   size and shape of, 583, 583n, 585  
   spatial distribution of stars in, 586–587  
   spiral structure of, 580–581, 586–589, 591–592  
     origins, 595  
   star counts for, 580  
   stellar populations in, 587  
   tidal streams in, 588–589, 649  
   weighing, 596  
 Miller, Stanley, 718  
 Millisecond pulsar, 550–551  
 Mimas (moon), 302–303, 310, 311  
 Minor planet, 340. *See also* Asteroid  
 Minute, arc, 13  
 Miranda (moon), 328, 329–330, 370  
 MN (asteroid), 343  
 MN4 (asteroid), 344  
 Model(s)  
   heliocentric, 39–40  
   Ptolemaic, 38–39  
   theoretical, 8  
 Model universes, 674  
 Modeling, 25  
 Molecular clouds, 463  
   complexes, 465  
   organic compounds in, 464n  
 Molecular emission, 91, 463  
 Molecular spectral lines, 91–92  
 Molecular spectrum, 92  
 Molecular speed, 196  
 Molecular tracer, 464–465  
 Molecules, 91–92  
   interstellar, 463–465  
 Momentum  
   angular, 145, 149  
   linear, 149  
 Month  
   sidereal, 18–19  
   synodic, 19  
 Moon(s). *See also* Galilean moons; *specific moons*  
   of jovian planets, 154  
   of Jupiter, 279–287  
   of Mars, 260–261  
   of Neptune, 328, 329, 330–332  
   of Saturn, 303, 304–314  
   of terrestrial planets, 143  
   of Uranus, 328–329  
 Moon, the, 188–213, 329  
   angular diameter of, 191  
   angular size of, 13  
   atmosphere of, 191–192  
   craters on, 192–193, 194, 201–203  
   density of, 208  
   distance to Earth, 190  
   evolutionary history of, 211–212  
   explorations of, 198–199  
   interior of, 208–209  
   mass of, 191  
   motion of, 18–24  
   nodes of orbit, 22  
   orbit of, 190  
   origin of, 210–211  
   properties of, 138, 191–192  
   rotation of, 195–196  
   surface composition of, 203–206  
   surface features of, 192–194  
   synodic period of, 22  
   tidal bulge in, 196  
   tides and, 182–183  
   volcanism on, 205–206  
 Moonlets of Saturn, 301  
 Morley, E. W., 553, 559  
 Motion(s)  
   apparent, 39  
   atomic, 472  
   coplanar, 379  
   diurnal, 13  
   orbital, of Milky Way Galaxy, 588–589  
   prograde (direct), 37  
   proper, 424, 601  
   radial, 74  
   retrograde, 37, 40, 41  
   stellar, 424–425  
   transverse, 74, 74n  
   unbound, 54  
 Motion, laws of  
   Newton's, 49–51  
   planetary, 44–47  
   relativity and, 557–560  
 Mount Everest, 176, 177  
 Mount Hadley, 206  
 Mount Haleakala, Hawaii, 114  
 MRC 1138-262 (“Spiderweb” galaxy), 648  
 M-type asteroid, 341  
 Multiple-planet systems, 373, 379  
 Multiple-star systems, 440–441.  
   *See also* Binary-star systems  
 Multiple-wavelength astronomy, 128  
 Muon particles, 694  
 Murchison meteorite, 720  
  
**N**  
 N49 (supernova remnant), 538  
 Naked singularity, 564  
 Nakhla (meteorite), 251  
 Nanometer (nm), 62–63  
 Nasmyth focus, 103  
 National Astronomical Observatory of Japan, 106  
 National Radio Astronomy Observatory, 115  
 Natural selection, 728  
 Nazca plate, 175  
 Neap tides, 183  
*Near Earth Asteroid Rendezvous (NEAR)*, 342  
 Near-miss events, asteroids and, 343–344  
 NEAR-Shoemaker, 342  
 Nebula/nebulae, 145, 148, 453, 453n. *See also specific nebulae*  
   clusters and, 488  
   cocoon, 482–483  
   emission. *See* Emission nebulae  
   planetary, 502–504  
   reflection, 454, 455  
   solar, 148, 150–152, 368  
   early, temperature in, 151  
   spiral, 580–581, 586  
   structure of, 455  
 Nebular contraction, 145, 148  
 Nebular spectrum, 456–457  
 Nebular/condensation theory, 148, 369–370.  
   *See also* Condensation theory  
 Negative curvature, 676, 677  
 Neon, emission spectrum of, 81  
 Neon-20, formation of, 535  
 Neon-oxygen white dwarf, 507  
 Neptune  
   atmosphere of, 325–326  
   discovery of, 320–321  
   Great Dark Spot of, 325–326  
   internal structure of, 326, 327  
   magnetic field and magnetosphere of, 326, 327  
   mass and density of, 322  
   moons of, 328, 329, 330–332  
   orbit of, 357  
   Pluto and, 353–354  
   properties of, 138, 322–323  
   rings of, 300, 334  
   rotation rate of, 323  
   spacecraft exploration of, 146  
 Neptunes, 374  
 Neptunes (extrasolar planets), 373, 374, 379  
   cold, 374  
 Nereid (moon), 321, 328, 329, 330  
 Neutrino detectors, 414–415  
 Neutrino oscillations, 415  
 Neutrino telescope, 415  
 Neutrinos, 411–412, 414–415  
   standard solar model and, 414–415  
 Neutron capture, 536  
 Neutron degeneracy pressure, 544, 555, 556  
 Neutron stars, 544–545, 634  
   distinguishing black hole from, 569–571  
   properties of, 544–545  
   and pulsars, 545–548  
   rotation of, 544  
   as stellar remnant, 544  
 Neutronization, 525  
 Neutrons, 90  
 Neutron-star binary, 548–552  
*New Horizons* mission, 142, 147, 356  
 New Moon, 18, 19, 20  
 New Technology Telescope (NTT), 112, 113  
 Newborn stars, H–R diagrams for, 476  
   star cluster, 514  
 Newton (N), 50  
 Newton, Isaac, 49, 55, 62, 67, 103  
 Newtonian mechanics, 49, 52–55  
   escape speed and, 673, 674  
 Newtonian telescope, 103, 136  
 Newton's constant, 51  
 Newton's laws, 49–52, 597, 601  
 NGC 1201, 612  
 NGC 1252 galaxy, 576, 577  
 NGC 1265 galaxy, 642  
 NGC 1300 galaxy, 610  
 NGC 1316, 602, 603  
 NGC 1365 galaxy, 610  
 NGC 1566, 593  
 NGC 1569 galaxy, 613  
 NGC 2207 galaxy, 643  
 NGC 2808, 509  
 NGC 2859, 612  
 NGC 2997 galaxy, 609  
 NGC 3603 star cluster, 486  
 NGC 3690 galaxy, 644  
 NGC 4038/4039 (Antennae) galaxies, 644  
 NGC 4214 galaxy, 485  
 NGC 4258, 652  
 NGC 4261 galaxy, 631, 632  
 NGC 4449 galaxy, 613  
 NGC 4565 galaxy, 579  
 NGC 4603, 616  
 NGC 6240, 653, 654  
 NGC 6302 planetary nebula, 495, 496  
 NGC 6520, 150  
 NGC 6543 (Cat's Eye Nebula), 505  
 NGC 6872 galaxy, 610  
 NGC 6934 star cluster, 114  
 NGC 7635, 538  
 NGC 7742 galaxy, 624  
 Nickel meteorites, 362  
 Nickel-56, 535, 537, 538  
 Night sky, 15  
 Nitrogen geysers, 331  
 Nix (moon), 354  
 Noise, background, 110  
 Nonstellar radiation, 624, 625n, 630  
 Nonthermal radiation, 633–634  
 Normal galaxies, 622, 655–656  
   black hole at center of, 652, 655  
 Normal matter, 699, 705–706  
 North American plate, 175, 177  
 North pole, magnetic, 180  
 North star (Polaris; Pole star), 10, 12  
 Northern Lights (aurora borealis), 181, 182  
 Northern sky, 12  
 Nova Cygni, 524  
 Nova Herculis, 522  
 Nova Persei, 524  
 Nova/novae, 522–523  
   recurrent, 522  
   and supernovae, 526–527, 526n  
 Nuclear binding energy, 536  
 Nuclear epoch, 696, 698  
 Nuclear force, 411–412, 413  
 Nuclear fusion, 410–411  
   proton–proton chains and, 412–414, 416

- Nucleosynthesis  
 primordial, 697, 699  
 stellar, 531, 533–534, 536–538
- Nucleotide bases, 718
- Nucleus/nuclei  
 atomic, 84, 172  
 of comet, 346  
 condensation, 148  
 formation, 697–700  
 galactic. *See* Galactic nucleus  
 active. *See* Active galaxies/active galactic nuclei  
 Halley's comet, 348  
 supermassive black holes in, 624
- O**
- OB associations, 487
- Oberon (moon), 328, 329
- Observatories, ancient, 35
- Occam's razor, 8
- Occultation, stellar, 332
- Olbers, Heinrich, 670
- Olbers's paradox, 670
- Old star cluster, H–R diagram of, 515
- Olympus Mons, 243, 244
- Omega Centauri, 487, 488
- Omega Nebula (M17), 91
- IE 0657-56 galaxy cluster (“bullet cluster”), 661, 662
- 1862III (Swift-Tuttle) (comet), 359
- Oort, Jan, 353
- Oort cloud, 155, 353
- Opacity, 66–67, 68, 582
- Open star clusters, 486, 487, 488, 491  
 evolution of, 512–513
- Open universe, 676
- Ophelia (moon), 334
- Ophiuchus constellation, 421
- Ophiuchus dark dust cloud, 460
- Opportunity* rover, 254, 255, 722
- Opposition, 38
- Optical doubles, 441
- Optical interferometry, 120–121
- Optical telescope, 100
- Optics  
 active, 112  
 adaptive, 112–114
- Orbit(s), 37–38  
 of comets, 345, 352–353  
 of Earth, 14–18  
 extrasolar, 371, 372, 373, 374–375, 379–380  
 of Moon, 21, 190  
 near Galactic center, 588–589, 601  
 planetary, 46, 140. *See also specific planets*  
 stellar, in Milky Way Galaxy, 589  
 synchronous, 195  
 of two bodies, 53
- Orbital motion, of Milky Way Galaxy, 588–589
- Orbital period, 138
- Orbitals, 84
- Oriente Basin, 202, 203
- Orion constellation, 2, 3, 10, 11, 428, 432, 510
- Orion molecular cloud, 480–481, 483, 484
- Orion Nebula (M42), 118, 121, 122, 458, 468, 469, 481, 489
- young stars in, 489
- Oscillations  
 acoustic, 709–710  
 matter, 709–710  
 neutrino, 415  
 solar, 393
- Outer core, 169
- Outflow channels, Martian, 245–246
- Outgassing, 168, 231–232, 257
- Ovda Regio, 224
- Oxygen-16, formation of, 534–535
- Ozone layer, 164, 166
- P**
- Pacific plate, 175, 177
- Pair production, 691–692
- Pallas (asteroid), 340, 341
- Pan (moonlet), 301
- Pandora (moon), 304
- Pangaea, 179
- Parallax, 24–26  
 spectroscopic, 437–438, 437n  
 stellar, 39, 42, 422–423
- Paranal Observatory, Chile, 106, 107
- Parsecs, 423
- Partial eclipses  
 lunar, 20  
 solar, 20, 21
- Particle detectors, 692
- Particle nature of radiation, 85, 87
- Particle (corpuscular) theory of light, 67
- Particle–antiparticle annihilation, 691–692
- Particle–antiparticle pair, 691–692
- Particles  
 alpha, 499  
 charged, 63–64  
 interactions, 411–412  
 fundamental force and, 694, 695  
 threshold temperature of, 691
- Pauli, Wolfgang, 500
- Pauli exclusion principle, 500
- Payne-Gaposchkin, Cecilia, 386–387, 584
- Pencil-Beam Survey, 668, 669
- Penumbra, 21, 401
- Penzias, Arno, 683–684
- Perihelion, 46, 190, 200
- Period, 47  
 orbital, 138  
 rotation, 138, 139  
 wave, 61
- Period-luminosity relationship, 583
- Permafrost, 247
- Perseid shower, 359
- Perseus galaxy cluster, 651
- PG0052 + 251 (quasar), 654
- Phaeton (asteroid), 359n
- Phase(s)  
 lunar, 18–19  
 of Mercury, 191  
 in star formation, 480  
 T Tauri, 153, 476, 478  
 of Venus, 42, 43
- Philippine plate, 175
- Philosophiae Naturalis Principia Mathematica (The Mathematical Principles of Natural Philosophy)* (Newton), 49
- PHL 938 quasar, 658
- Phobos (moon), 239, 260, 261
- Phoebe (moon), 313
- Phoenix* mission, 147, 255
- Pholus (Kuiper belt object), 356
- Photodisintegration, 525, 535–536
- Photoelectric effect, 88
- Photoevaporation, 455
- Photometers, 111
- Photometry, 110–111, 429
- Photon, 85, 88  
 particle–antiparticle pair produced by, 691–692  
 redshift of, 673
- Photons, 695
- Photosphere, 390, 391, 474, 700
- Photosynthesis, 724
- Piazzi, Giuseppe, 340
- Pickering, E. C., 584
- Pioneer* missions, 146–147, 277, 303, 729–730, 732  
*Pioneer 10* spacecraft, plaque mounted on, 730  
 to Saturn, 293  
*Pioneer Venus* (spacecraft), 220, 221, 222, 223, 226, 228, 229, 232
- Pistol Star, 387
- Pixels, 109
- Plains, on Mercury  
 intercrater, 194  
 smooth, 194
- Planck, Max, 68, 85, 695
- Planck (blackbody) curve, 69, 70–71, 428–429, 683, 684
- Planck epoch, 695, 701, 702
- Planck* mission, 699, 708, 709
- Planck's constant, 85
- Planet(s), 36–37. *See also specific planets*  
 accreting, 151  
 alignment of, 140  
 binary-star, 727  
 condensation theory of formation of, 148, 150–152  
 dwarf, 144, 357  
 Earth-like, searching for, 725–727  
 extrasolar (exoplanet), 137, 368, 370–378  
 features attributed to random events, 370  
 formation of  
 condensation theory, 368–369  
 modeling, 368–369  
 giant, 152–154  
 habitable, 725–728  
 IAU definition of, 357  
 jovian, 140–141, 143. *See also* Jupiter;  
 Neptune; Saturn; Uranus  
 Jupiter-like, 379–380  
 mass of, 138  
 minor, 340. *See also* Asteroids  
 observations of, 36–37  
 orbits of, 46, 140  
 properties of, 138  
 pulsar, 551–552  
 rotation period, 138, 139  
 sinking, 380–381  
 spacecraft exploration of, 142  
 timeline for, 146–147  
 terrestrial, 140–141, 143. *See also* Earth;  
 Mars; Mercury; Venus
- Planet migration, 154
- Planetary debris, jovian planets and, 152–156
- Planetary irregularities, 369–370
- Planetary motion, 37  
 Kepler's laws of, 45–47, 52–53  
 laws of, 44–47  
 and Newton's laws, 52  
 types of. *See* Motion
- Planetary nebula, 502–504
- Planetary radar, 200
- Planetary system  
 discovery, 136–137  
 formation of. *See* Solar system  
 habitable planets per, 725–727  
 stars with, 725
- Planetary transit, 372–373, 374, 375, 377, 378
- Planetesimals, 150  
 ejection to form Oort cloud and Kuiper belt, 155  
 in solar system formation, 368, 369
- Planetoids. *See* Asteroids
- Planetology, comparative, 137
- Plasma torus, Io, 278, 282–283
- Plate tectonics  
 on Earth, 174–180  
 on Ganymede, 286  
 on Mars, 242, 259  
 on Venus, 223–224, 225, 228, 233
- Plates, global, 174, 175
- Plato, 38
- Pleiades (Seven Sisters), 2, 3, 486, 487
- Plum crater, 188–189
- Plutinos, 354
- Pluto (KBO), 370  
 discovery of, 132, 353–354  
 Kuiper belt and, 356  
 orbit of, 357  
 properties of, 138, 354–356  
 reclassification as dwarf planet, 143–144, 357–358
- Pluto–Charon system, formation of, 370
- Plutoids, 357
- Plutonium 241, 172, 173
- Polar caps, Martian, 239–240, 249, 252, 257
- Polar vortex, at Venus's poles, 230
- Polaris (North star; Pole star), 10, 12
- Polarity of sunspot, 402
- Polarization of light, 67  
 of starlight by dust, 453, 453n
- Pole star (North star; Polaris), 10, 12
- Poles  
 celestial, 12–13  
 magnetic, 180
- Pollution, light, 111
- Polycyclic aromatic hydrocarbons, 250, 251
- Population I and II stars, 587
- Positive curvature, 676, 677
- Positron, 411
- Precession, 17–18
- Pressure waves, 168–169
- Primary atmosphere, 168
- Primary mirror, 100
- Primary waves (P-waves), 168–169
- Prime focus, 100
- Primeval fireball, 670–671, 693, 696
- Primordial elements, 531
- Primordial intracluster gas, 642
- Primordial nucleosynthesis, 697, 699
- “Primordial soup,” 718
- Principia* (Newton), 49
- Prism, 62
- Procyon, 78–79
- Procyon B, 434, 436
- Progenitor of supernova, 526, 533
- Prograde (direct) motion, 37
- Prometheus (moon), 304
- Prometheus (volcano), 282
- Promethium, 531
- Prominences, solar, 405–406
- Proper motions, 424, 601
- Protein-like microspheres, 718–719
- Proteins, 718
- Proteus (moon), 328, 329, 330
- Proton fusion, 411–412, 534
- Proton–proton chain, 412–414, 416, 533, 697n
- Protons, 63  
 subatomic, smashing together, 688, 689
- Protoplanets, 150, 153–154
- Protostars, 474–476  
 evidence of, 481–483  
 evolution of, 476
- Protostellar collisions, 489
- Protostellar disk, 475. *See also* Solar nebula
- Protostellar winds, 482, 483–484
- Protosun, 150
- Proxima Centauri (Alpha Centauri C), 376, 423, 435, 444
- Ptolemaic model, 38–39, 55
- Ptolemy, 38–39
- Pulsar planet, 551–552
- Pulsar radiation, 545
- Pulsar wind, 546, 547
- Pulsars, 545–548  
 binary, 568  
 gamma-ray, 548  
 millisecond, 550–551  
 neutron stars and, 545–549
- Pulsating variable stars, 581, 582
- P-waves (primary waves), 168–169
- P/Wild 2 (comet), 350, 352
- Pyrolytic-release experiment, 250
- Q**
- QSO 1422 + 2309 quasar, 659
- Quantized orbital energy, 85
- Quantum fluctuation, 702, 705, 707
- Quantum gravity, 564, 693, 695
- Quantum mechanics, 85, 88
- Quaasar (Kuiper belt object), 356, 357
- Quark epoch, 696
- Quarks, 694  
 and leptons, 696
- Quarter Moon, 18, 19
- Quasar(s), 624, 628–630  
 absorption lines, 658–659  
 discovery of, 628  
 first, 683  
 “host” galaxies of, 653, 654  
 and measuring plate motion, 175  
 properties of, 628–629  
 twin, 659–660  
 typical, 630
- Quasar epoch, 652–654, 683
- Quasar feedback, 654
- Quasar jets, 629, 630
- Quasar “mirages,” 659–660
- Quasi-stellar radio sources. *See* Quasar(s)
- Quiescent prominences, solar, 405
- Quintessence, 680, 680n
- R**
- R136 star cluster, 110, 113, 470
- Radar, 48–49  
 planetary, 200
- Radar maps, of Venus and Earth, 223
- Radar ranging, 190
- Radial motion, 74
- Radial velocity, 92–93  
 of star, 371
- Radian, 28
- Radiant, 359, 360
- Radiation, 58–75. *See also* Radio radiation  
 background, ripples in, 707–709  
 defined, 60  
 diffraction property of, 67  
 distribution of, 68, 69  
 Doppler effect, 73–75  
 electromagnetic, 60  
 emitted by active galaxies, 622–624  
 gravitational, 554, 568

- Radiation (*continued*)  
infrared, 60  
interference property of, 67  
laws of, 70–73  
light and, 63  
nonstellar, 624, 625n, 630  
nonthermal, 633–634  
as particles, 85, 87  
from Seyfert galaxies, 624–625  
spectrum. *See* Spectrum/spectra, of Sun  
synchrotron, 633–634  
thermal, 68–73  
21-centimeter, 462–463  
types of, 65, 66  
ultraviolet, 163, 164  
in universe, 690–691  
wave motion of, 60–62, 67  
wave theory, 65
- Radiation darkening, 329  
Radiation era, 693, 696  
Radiation zone, 390, 391  
Radiation-dominated universe, 691  
Radiation-matter decoupling, 699, 700  
Radio astronomy, 114–118  
Radio communication, search for  
extraterrestrial intelligence using,  
730–731  
Radio emission by interstellar gas, 462–463  
Radio galaxies, 569, 624, 625–628  
core-dominated, 626, 627  
head-tail, 641–642  
Radio lobes, 625–626, 633  
Radio maps, of Milky Way, 591  
Radio radiation, 60, 65, 66, 730–731  
pulsar, 545, 546  
Radio telescopes, 114–117, 730  
interferometer, 118–120  
Radioactive dating, 172–173  
Radioactive elements, 531  
Radioactivity, 171, 172  
Radius  
angular momentum and, 149  
and mass and luminosity, 442–443  
of planet, determining, 138. *See also*  
*specific planets*  
of solar system objects, 138  
Radius–luminosity–temperature  
relationship, 432–434  
Radome, 117  
Random events, planetary features  
attributed to, 370  
Ranger series, 198  
Rapid mass transfer, 517  
Rayleigh, Lord, 165  
Rayleigh scattering, 165  
RCW 38, 538  
Receding galaxies, 672  
Recession, universal, 619–620  
Recession velocity, 74, 620, 671  
Recombination, 699  
Reconnection process, 406  
Recurrent nova, 522, 523  
Recycling, stellar, 538–539  
Red dwarfs, 433  
missing, 597  
Red giants, 434, 499  
Red River, 245  
Red Spot, 264, 265, 268, 269, 272, 273, 274  
Red Spot Junior, 273, 274  
Red supergiants, 434, 509–510  
Reddening, 451  
Red-giant branch, 498–499  
Red-giant instability, 503  
Red-giant region, 436, 437  
Red-giant wind, 511  
Redshift(s), 553  
cosmological, 619, 620, 673  
gravitational, 562–563  
relativistic, 622–623  
Redshift galaxy survey, 657–658, 668  
Redshifted radiation, 74, 75, 588  
Reflecting telescopes, 100–103  
Reflection, 101  
Reflection nebula, 454, 455  
Refracting telescopes, 100–105, 136  
Refraction, 62, 100, 101  
Regolith, 203–204  
Reinhold crater, 202  
Reionization, of universe, 696  
Relativistic fireball, gamma-ray bursts as,  
554  
Relativistic redshifts, 622–623  
Relativity  
Einstein's theory of, 74, 75, 557–561,  
675–676, 690  
tests of, 566–567  
general, 558, 560, 566–567  
special, 557–558, 559
- Remnants  
acoustic, 710  
stellar, properties of, 556. *See also*  
Supernova remnant
- Residual caps, 249, 252  
Resolution  
angular, 106–108  
diffraction-limited, 107  
Resonance, 198–200  
orbital, Trojan asteroids and, 344  
of ring particles of Saturn, 301–304  
spin-orbit, 198–199
- Retrograde motion, 37, 40, 41  
Retrograde rotation, 219–220  
Revolution of Earth around Sun, 14  
Rhea (moon), 303, 310, 311  
Rho Ophiuchi, 460  
Riemann, Georg Friedrich, 677  
Riemannian geometry, 677  
Rigel, 428, 432, 439, 510, 512  
Rille, 206  
Ring Nebula, 503–504  
Ring system  
of Jupiter, 287, 300  
of Neptune, 300, 334  
of Saturn, 290, 291, 292, 293, 298–304,  
333  
of Uranus, 300, 332–334  
Ringlets of Saturn, 300, 301  
Roche, Edouard, 300, 516  
Roche limit  
Neptune, 334  
Saturn, 299–300  
Roche lobes, 516–517  
Rock cycle, 179  
Rocks, lunar, 204  
Röntgen, Wilhelm, 126  
ROSAT X-ray satellite, 126  
Rosbud Nebula, 387  
Rosse, Earl of, 136  
Rotation  
differential, 267  
of Earth, 12, 183–184  
Galactic, 592–593, 596, 616  
gravity and, 471  
of Mercury, 196–199  
molecular, 91  
rate of, 149  
solar, 390  
spectral-line broadening and, 94  
sunspot, 402–403  
of terrestrial planets, 141, 219
- Rotation curve  
Galactic, 596  
mass measurement of spiral galaxies  
and, 640
- Rotation period  
of Moon, 195–196  
of planets, 138, 139  
solar, 390  
of solar system objects, 138
- Rotational flattening, 268  
r-process, 536  
RR Lyrae variable stars, 581–582  
Runaway greenhouse effect, 168, 231–232  
Runoff channels, Martian, 245  
Russell, Henry Norris, 434, 584
- S**  
S106 (nebula), 366, 367  
Sagittarius A, 600–601  
Sagittarius dwarf galaxy, 648, 649  
Salam, Abdus, 694  
San Andreas Fault, 175, 176, 177  
Saros cycle, 23  
Satellite(s). *See also specific satellites*  
co-orbital, 313–314  
shepherd, 301–304, 333–334  
of solar system objects, 138
- Saturn, 3, 290–314  
atmosphere of, 292–296  
interior structure of, 296–297, 327  
internal heating of, 296–297, 327  
magnetic field and magnetosphere of,  
297, 327  
moons of, 303, 304–314  
orbit of, 292, 293  
properties of, 138  
rings of, 292, 293, 298–304, 313, 333  
rotation of, 292  
spacecraft exploration of, 146–147  
storms on, 295–296  
weather on, 294–296
- SB0 galaxies, 612  
Scalar fields, 701–702  
Scarps, 206–207  
Scattering of sunlight, Rayleigh, 165  
Schiaparelli, Giovanni, 132, 133, 196–197  
Schmitt, Harrison, 198  
Schwarzschild, Karl, 557  
Schwarzschild radius, 557  
Scientific method, 8–10  
active galaxies and, 656  
stars and, 514–515
- Scientific notation, 7  
Scorpius–Centaurus association, 458  
Scotia plate, 175  
SDO (*Solar Dynamics Observatory*), 389, 395  
SDSS (*Sloan Digital Sky Survey*), 649, 651,  
658, 668, 710  
Seafloor spreading, 176, 178  
Search for extraterrestrial intelligence  
(SETI), 729, 731–732  
Seasonal caps, 249, 252  
Seasons and seasonal changes, 14–17  
Second, arc, 13  
Secondary atmosphere, 168  
Secondary mirror, 103  
Secondary waves (S-waves), 168–169  
Sedimentary rock, 179  
Sedna (Kuiper belt object), 356, 357  
Seeing, 111  
Seeing disk, 111, 112  
Seismic waves, 168–169  
Seismograph, 168  
Self-propagating star formation, 594–595  
Semidetached binary (mass-transferring  
binary) system, 522–523  
Semimajor axis, 45, 46  
Sensitivity, telescope size and, 106  
SETI (search for extraterrestrial  
intelligence), 729, 731–732  
Seven Sisters (Pleiades), 2, 3, 486, 487  
Seyfert, Carl, 624  
Seyfert galaxies, 624–625  
Sgr A\*, 600, 601–602  
Shadow zones, 169  
Shapley, Harlow, 574, 583, 584, 585  
Shear waves, 168–169  
Shepherd moon, 303  
Shepherd satellites, 301–304, 333–334  
Shield volcanoes, 225  
Shock waves, star formation and, 484–485,  
594  
Shoemaker–Levy 9 (comet), 274–275, 351  
Shooting stars. *See* Meteors  
Short-period comet, 352  
Shu, Frank, 594  
SI (Système Internationale) metric system, 7  
Sidereal day, 14  
Sidereal month, 18–19  
Sidereal orbital period, 138  
Sidereal year, 17  
*Sidereus Nuncius (The Starry Messenger)*  
(Galileo), 42  
Sif Mons, 225  
Silicate meteorite, 362  
Silicon, 724  
Single-line spectroscopic binary, 440–442  
Singularity, 563–564, 693  
naked, 564  
Sinking planets, 380–381  
Sirius A, 432, 443, 444, 505,  
506, 508  
Sirius B, 433, 443, 505, 506, 508  
Sister (coformation) theory, 210  
*Skylab* space station, 407  
Slingshot, gravitational, 142, 147, 306  
Slipher, Vesto M., 132, 619  
*Sloan Digital Sky Survey (SDSS)*, 649, 651,  
658, 668, 710  
Sloan Great Wall, 668. *See also* Great Wall  
Small Magellanic Cloud, 612  
Smog, 122  
Smooth plains, on Mercury, 194  
SN1987A, 526, 532–533  
SNO (Sudbury Neutrino Observatory)  
detector, 415  
SO galaxies, 612  
Sodium, emission spectrum of, 81, 82–83
- Sojourner* rover, 254  
*Solar and Heliospheric Observatory (SOHO)*,  
393, 406, 408  
description of, 395  
Solar constant, 390  
Solar cycle, 403–405, 409  
Solar day, 13  
mean, 17  
*Solar Dynamics Observatory (SDO)*, 389, 395  
Solar eclipse, 20–22, 23, 398–400, 410  
Solar energy production, 410, 416  
Solar flares, 406, 407  
Solar magnetism, 395, 400–405  
Solar maximum, 404  
Solar minimum, 404  
Solar nebula, 148, 151, 152, 368, 475  
early, temperature in, 151  
temperature profile of, 150–152  
Solar neutrino problem, 414  
Solar neutrinos, 412, 414–415  
Solar pressure waves, 393  
Solar prominences, 405–406  
Solar rotation, 394  
Solar system, 36, 134–156. *See also* Sun;  
*specific planets*  
condensation theory of. *See* Condensa-  
tion theory  
debris. *See* Asteroids; Comet(s); Kuiper-  
belt objects (KBOs); Meteorites;  
Meteoroids; Meteors  
differentiation of, 150–152  
dimensions of, 47–49  
formation of, 144–152, 154  
condensation theory,  
368–369  
modeling, 368–369. *See also* Con-  
densation theory  
geocentric model of, 36–39  
heliocentric model of, 39–41  
interplanetary matter, 143–144  
inventory of, 136–137  
jovian planets in, 140–141, 143, 152–154  
layout of, 139–140  
life in, 722–724  
measuring planets, 138–139  
objects in, properties of, 138  
and other planetary systems compared,  
379–382  
planets beyond. *See* Extrasolar planets  
properties of, 144–145  
regularities and irregularities of,  
369–370  
spacecraft exploration of, 142  
timeline, 146–147  
terrestrial planets in, 140–141, 143  
Solar tides, 184  
Solar transit, 48  
Solar wind, 143, 180, 390, 391, 400  
Venus and, 232–233
- Solstice, 15–16  
Sombrero galaxy (M104), 610  
Sound waves, cosmic, 707, 709  
South American plate, 175  
South pole, magnetic, 180  
South Pole–Aitken Basin, 194  
Southern lights (aurora australis), 181, 182
- Space  
curvature of, 560–561, 676–677  
geometry of, 675–677  
Space warping, 561  
Space-based astronomy, 121–128  
Spacecraft exploration  
of Moon, 198–199  
of solar system, 134–135, 142  
Spacetime, 558  
Special relativity, 557–558, 559  
Spectral class, stellar, 431–432, 431n  
Spectral lines, molecular, 463–464  
Spectral window, 66, 68  
Spectral-line analysis, 92–95  
Spectrograph (spectrometer), 80  
Spectroscope, 80  
Spectroscopic binary, 440–441  
Spectroscopic parallax, 437–438, 437n  
Spectroscopic radio astronomy, 591  
Spectroscopic thermometer, 92  
Spectroscopy, 78–95, 111  
Spectrum/spectra, 62, 80–81, 111  
absorption, 460–461  
absorption lines, 82–83  
atomic, 87  
blackbody (Planck) curve, 68, 69,  
428–429, 683, 684

- continuous, 80, 81  
 electromagnetic, 65–68  
 emission lines, 80–81  
 formation of, 87–91  
 galaxy, 619–620  
 galaxy energy, 623–624  
 helium, 81  
 hydrogen, 80, 81, 83, 84, 87–88  
 identifying starlight, 83  
 Kirchhoff's laws, 83, 84, 88–90  
 line broadening, 93–95  
 mercury, 81  
 molecular, 92, 463–464  
 of more complex elements, 90–91  
 nebular, 456–457  
 neon, 81  
 nonthermal, 72  
 sodium, 81, 82–83  
 of spiral nebulae, 586  
 of stars, 83, 429–432  
 of Sun, 65–68, 82, 397–398  
 of 3C 273 quasar, 629  
 velocity information from spectral lines, 92–93  
 of white light, 62
- Speed, 46  
 escape. *See* Escape speed  
 of light, 64–65, 557, 558  
 molecular, 196  
 vs. velocity, 50
- Spica B, 444  
 Spicules, solar, 399  
 Spiderweb (MRC 1138-262) galaxy, 644  
 Spin-orbit resonance, 198–199  
 Spiral arms, 592–595  
 Spiral density waves, 300, 592–594  
 Spiral galaxies, 580–581, 608–611, 614. *See also* Milky Way Galaxy  
 barred, 609–611  
 M81, 125  
 M82, 125  
 M100, 123  
 M101, 105  
 origins of spiral structure in, 595
- Spiral nebulae, 580–581  
 brightness and spectra of, 586  
 rotation of, 586
- Spirit* robot, 136  
*Spirit* rover, 137, 147, 245, 254  
*Spirit/Opportunity* lander, 133  
 Spitzer, Lyman, Jr., 121  
*Spitzer Space Telescope (SST)*, 121–123, 150, 312, 377, 479, 480, 489, 543
- Spring tides, 183  
 s-process, 536, 537  
*Sputnik*, 198–199  
 SS 433, 549–550, 549n
- Stadium (unit of measurement), 27  
 Standard candles, 615–616, 621  
 Standard Solar Model, 391, 392  
 neutrino physics and, 414–415
- Star(s), 420–444. *See also* Stellar entries; Sun  
 binary. *See* Binary-star systems  
 brightness of, 425–427  
 absolute, 425  
 apparent, 425–426, 581, 582, 586, 598  
 classification of, 438–439  
 variations in, 439  
 clusters of. *See* Star cluster  
 colors of, 428–429  
 defined, 390  
 distances to, 422–424  
 halo, 588, 588n  
 Hertzsprung–Russell (H–R) diagram, 434–437  
 lifetimes of, 444, 496  
 luminosity of, 425–427, 430, 433  
 classes, 438–439  
 main-sequence, key properties of, 444  
 mass of, 440–442  
 distribution, 442  
 measuring, 443  
 and other stellar properties, 442–444  
 of Milky Way, spatial distribution of, 586–587  
 motion of, 424–425  
 neutron. *See* Neutron stars  
 newborn, H–R diagram for, 476  
 orbits of, in Milky Way Galaxy, 589  
 with planetary systems, 725  
 scientific method and, 514–515  
 sizes of, 432–434  
 solar-type. *See* Sun-like stars  
 in Southern sky, 376  
 spectra of, 78–79, 83  
 classification, 431–432, 431n  
 spectroscopic parallax, 437–438, 437n  
 Sun-like. *See* Sun-like stars  
 temperatures of, 428–429  
 variable, 387, 432, 581–582, 583
- Star clusters, 468, 486–491  
 associations, 486–487  
 blue stragglers in, 507, 508  
 environment of, 488–489  
 globular, 487–488, 507, 580–581, 585  
 lifetimes of, 491  
 nebulae and, 488  
 newborn, 486, 514  
 observing stellar evolution in, 512–515  
 open, 486, 487
- Star formation, 468–491. *See also* Stellar evolution; Stellar explosions  
 characteristic stages of, 472–473  
 different masses and, 477–478  
 earliest stars and, 683  
 failed stars and, 478  
 generations of, 485  
 gravitational competition and, 471  
 modeling, 472  
 of nebulae and star clusters, 486–491  
 observational evidence of, 478–484  
 phases in, 480  
 rate of, 725  
 regions of, 470–472  
 self-propagating, 594–595  
 shock waves and, 484–485, 594
- Star Wars program (Strategic Defense Initiative), 113  
 Starburst galaxies, 623–624, 644  
*Stardust* mission, 350–351
- Starlight  
 aberration of, 42  
 spectral information from, 95
- Stefan, Josef, 70  
 Stefan-Boltzmann constant, 72  
 Stefan-Boltzmann equation, 72  
 Stefan-Boltzmann law, 72, 433  
 Stefan's law, 70, 72  
 Stellar epoch, 697
- Stellar evolution, 494–517. *See also* Star formation  
 of binary-star systems, 515–517  
 cycle of, 538–539  
 end points for stars of different masses, 512  
 of high-mass stars, 509–510, 512  
 leaving main sequence, 496  
 stages in, characteristics of, 472–473, 498  
 in star clusters, observing, 512–515  
 of Sun-like stars, 496–502  
 carbon core, 501–502  
 death of, 502–509  
 helium core, 497, 499, 501  
 helium flash, 499–501  
 helium fusion, 499–501  
 planetary nebulae, 502–504  
 red-giant branch, 498–499  
 subgiant branch, 497–498  
 white dwarf, 504–507  
 theory of, 514–515  
 reality vs., 507–509
- Stellar explosions, 520–539. *See also* Element(s)  
 end of high-mass star, 524–526  
 novae, 522–523  
 supernovae, 525–526
- Stellar habitable zone, 725–726  
 Stellar nucleosynthesis, 531, 533–534  
 observational evidence for, 536–538
- Stellar occultation, 332  
 Stellar parallax, 25, 39, 42, 422–423  
 Stellar populations, 587  
 Stellar recycling, 538–539  
 Stellar remnants, 556. *See also* Supernova remnant  
 Stellar transits, of black holes, 564  
 Stellar wind, 511  
 Stickney crater, 260  
 Stonehenge, 34  
 Storms, on Saturn, 295–296
- Strategic Defense Initiative (Star Wars program), 113  
 Stratosphere, 163
- Stray asteroids, 360–361  
 Strings/string theory, 694  
 Strong nuclear force, 411, 413, 694, 695  
 Structure in early universe, 705–707  
 S-type asteroid, 341, 362  
 Subaru telescope, 106, 107, 113, 367, 569  
 Subatomic particles, 597  
 Subduction zones, 176  
 Subgiant branch, 497–498  
 on H–R diagram, 499  
 Sublimation, 346  
 “Substellar” object. *See* Brown dwarf  
 Sudbury Neutrino Observatory (SNO) detector, 415  
 “Sugar-scoop” antenna, 683  
 Sulfur-32, formation of, 535  
 Sulfuric acid in Venus's atmosphere, 230  
 Summer solstice, 15–16
- Sun, 6, 388–416, 444. *See also* Solar entries  
 absorption lines of, 397–398  
 active, 405–409  
 active regions of, 405–406  
 angular size of, 13  
 atmosphere of, 397–400  
 CNO cycle in, 534  
 composition of, 398  
 changes in (theoretical), 497  
 convection and, 395–396  
 corona of, 399–400, 407–410  
 density of, 391, 394  
 energy production of, 410, 416  
 and granulation, 396–397  
 interaction with planets, 52  
 interior of, 392–397  
 luminosity of, 390–392, 409, 410, 433, 435  
 magnetism of, 395, 400–405  
 neutrinos of, 414–415  
 nuclear fusion in, 411–412  
 oscillations, 393  
 physical properties of, 138, 390–392  
 Procyon star, 78–79  
 radiation of, 166, 394–396  
 rotation of, 394, 403  
 solar-terrestrial relations, 409  
 spectrum of, 82, 397–398  
 stars like the. *See* Sun-like stars  
 stellar classification for, 431, 432, 439  
 structure of, 390, 391  
 temperature of, 391, 394, 399–400, 401  
 tidal forces exerted by, 200–201  
 weight of, 54  
 X-ray observations of, 407, 408
- Sun Dagger (Chaco Canyon, New Mexico), 35
- Sun-like stars  
 evolution of, 496–502  
 formation of, 472–477  
 merger of, 508  
 zero-age main sequence (ZAMS), 477–478
- Sunspot cycle, 403–404, 405  
 Sunspot minimum, 405  
 Sunspots, 42, 401–405, 634  
 rotation of, 402–403
- Super Kamiokande telescope, 415  
 Superclusters, 656–657, 678  
 Super-Earths, 373–374, 375  
 and Earth comparisons, 377  
 “Superforce,” 694, 695
- Supergiants, 434  
 mass loss from, 511  
 red, 509–510
- Supergranulation, 397  
 Superior conjunction, 38, 218  
 Superior orbit, 37  
 Superior (higher) planet, 37  
 Supermassive black holes, 569, 624, 652  
 in galactic nuclei, 569, 601–603, 624  
 Supernova remnants, 125, 528–530  
 Cassiopeia A (Cas A), 126  
 Crab, 529  
 in Crab Nebula, 520, 521  
 Cygnus, 125  
 near Galactic center, 600  
 Vela, 528–529, 530
- Supernova/supernovae, 526–530  
 ancient Chinese data on, 35–36  
 explosion, 525–526  
 interstellar shock waves generated by, 468, 469  
 progenitor of, 526, 533  
 remnants. *See* Supernova remnants
- Type I (carbon-detonation), 527–528, 679  
 Type II (core-collapse), 526, 527–528
- Supersymmetric partners, 694  
 Supersymmetry, 694  
*Surveyor* missions, 198  
 S-waves (secondary waves), 168–169  
*Swift* mission, 553  
 Swift-Tuttle (1862III) (comet), 359  
 Synchronous orbit, 195  
 Synchrotron radiation, 633–634  
 Synodic month, 19  
 Syntaxis (Ptolemy), 38–39  
 Système Internationale (SI) metric system, 7
- T**
- T association, 487  
 T Tauri phase, 153, 476, 478  
 T Tauri stars, 153, 487  
 Tail of comet, 345, 346, 347  
 Tau particles, 694  
 Taylor, Joseph, 568  
 Technetium, 531, 537  
 Technetium-99, 537  
 Technological civilization(s), 716, 721, 728–729  
 average lifetime of, 728–729  
 number in Galaxy, 729
- Tectonic fractures, 242, 311  
 Telescopes, 41, 98–129  
 defined, 100  
 focus of, 100  
 full-spectrum, 128  
 high-energy, 125–128  
 high-resolution, 111–114  
 image acquisition by, 109  
 image processing in, 110  
 infrared, 121–123  
 interferometry, 118–121  
 invention of, 136  
 light-gathering power of, 105–106  
 neutrino, 415  
 optical, 100  
 photometry, 110–111  
 radio, 114–117, 730  
 interferometer, 118–120  
 reflecting, 100–103  
 refracting, 100–105, 136  
 resolving power of, 105, 106–108  
 size of, 105–108  
 spectroscopy with, 111  
 ultraviolet, 123
- Teleso (moon), 314  
 Tempel 1 (comet), 352  
 Temperature, 68–73  
 absolute, 70  
 differentiation of solar system and, 150–152  
 in early solar nebula, 151  
 of Earth's interior, 170  
 of emission nebula, 457  
 habitable zone of “comfortable,” 725–726  
 in Hertzsprung–Russell diagrams, 434, 435  
 Kelvin scale, 69  
 line intensity and, 92  
 on Mars, 256  
 of stars, 428–429  
 of Sun, 390, 391, 394, 399–400, 401  
 threshold, of particle, 691
- Terminator line, on Moon, 192, 193  
 Terrestrial planets, 140–141, 143. *See also* Earth; Mars; Mercury; Venus  
 composition of, 139  
 rotational properties of, 138, 219  
 surface conditions of, 141
- Terzan 2 (globular cluster), 549  
 Tethys (moon), 303, 310, 311, 314  
 Tharsis bulge, 241, 242, 244  
 Theoretical model, 8  
 Theory, 8–9. *See also specific theories*  
 “Theory of Everything,” 693  
 Thermal broadening, 94  
 Thermal energy (heat), 69  
 atmosphere and, 166–168  
 gravity and, 471–472  
 solar system formation and, 148  
 Thermal equilibrium, 691, 692  
 Thermal motions of atoms, 94  
 Thermal radiation, 68–73  
 Thermometer, spectroscopic, 92

- Thick disk component of Milky Way, 587  
*Third Cambridge Catalog*, 628  
 Thorium-232, 172, 173  
 3C 48 quasar, 624–625  
 3C 84 galaxy, 625  
 3C 175 quasar, 630  
 3C 273 quasar, 628–629  
   spectrum of, 629  
 3C296 galaxy, 569  
 Threshold temperature of particle, 691  
 Thuban, 18  
 Tidal bulge, 183, 184  
   in Moon, 196  
 Tidal force (differential force), 183, 199–200  
   of black holes, 561–562  
   exerted by Sun, 200–201  
 Tidal stability limit (Roche limit), 299  
 Tidal streams, 588–589, 649  
 Tides, 182–184  
   lunar, 183, 184  
   solar, 184  
 Time delay, quasar mirage and, 660  
 Time dilation, 563  
 Titan (moon), 297, 303, 306  
   atmosphere of, 305, 307–308  
   surface and interior structure of, 308–310, 722  
 Titania (moon), 328, 329  
 Tombaugh, Clyde, 132, 354  
 Total eclipse  
   lunar, 20  
   solar, 20, 21  
 Toutatis (asteroid), 343, 344  
 Tracer, molecular, 464–465  
*Transition Region and Coronal Explorer (TRACE)* satellite, 402, 406  
 Transition zone, 390, 391, 399–400  
 Transits  
   planetary, 372–373, 374, 375, 377, 378  
   solar, 48  
   stellar, 564  
 Trans-Neptunian object (TNO), 353, 356–357  
 Transverse Doppler shift, 74n  
 Transverse motion, 74, 74n  
 Transverse velocity of star, 424–425  
 Trapezium (stars), 487, 489  
 Triangulation, 24–26  
 Trifid Nebula, 454  
 Trigonometry, 24  
 Triple-alpha process, 499  
 Triple-alpha reaction, 501, 534  
 Triton (moon), 321, 328, 330–332, 370  
 Trojan asteroids, 340, 344  
 Tropical year, 17  
 Troposphere, 163  
 Tully-Fisher relation, 616–617, 621, 640  
 Tunguska event of 1908, 351, 361  
 “Tuning fork” diagram (of Hubble), 614  
 Turbulence  
   atmospheric, 111, 112  
   clear-air, 164  
   gas, 94  
 Turnoff mass, 513  
 21-centimeter radiation, 462–463  
 0024 + 1654 galaxy cluster, 661  
 Twin quasars, 659–660  
 2M1207 (planetary system), 368  
 Tycho’s supernova, 530  
 Type I (carbon–detonation) supernova, 527–528, 679  
   energy emission, 537  
 Type II (core-collapse) supernova, 527–528
- U**  
 UDF (Hubble Ultra Deep Field), 575, 666, 667  
*Uhuru* satellite, 565  
 Ultra Deep Field, 666, 667. *See also* Hubble Ultra Deep Field (UDF)  
 Ultraviolet astronomy, 123–128  
 Ultraviolet radiation, 60, 65, 66, 163, 164  
 Ultraviolet telescopes, 123  
*Ulysses* (spacecraft), 408  
 Umbra, 21, 23, 24, 401  
 Umbriel (moon), 328, 329  
 Unbound motion, 54  
 Universal recession, 619–620  
 Universe, 6. *See also* Early universe; Life  
   acceleration of, 678–679  
   age of, 671, 682–683  
   birth of, 670–671, 693, 695  
   closed, 676  
   composition of, 680–681, 682, 690  
   cosmological constant and, 679, 681  
   critical, 676  
   dark matter in, 640–642  
   density of, 673–674, 676, 678, 680  
   early, 688–710  
   evolution of, 693–697  
   expansion of, 670–673  
     critical density and, 673–674  
     dark energy and, 679–680  
   fate of, 673–675  
   flat, 700–701  
   future of, 674–675  
   geocentric, 36–39  
   geometry of, 682  
   large-scale structure in, 651, 656–658, 668–669, 705–707  
   life in, 714–732  
   matter-dominated, 690–691  
   model, 674  
   open, 676  
   radiation in, 690–691  
   radiation-dominated, 691  
   reionization of, 696  
   relativity and, 681  
   size and scale of, 6–7  
 Ununpentium, 530  
 Upper mantle, 170, 176  
 Upsilon Andromedae (Sun-like star), 371, 372  
 Uraniborg observatory, 44  
 Uranium-235, 172, 173  
 Uranium-238, 172, 173  
 Uranus, 318  
   atmosphere of, 324–325  
   discovery of, 320  
   internal structure of, 326, 327  
   magnetic field and magnetosphere of, 326, 327  
   mass and density of, 322  
   moons of, 328–329  
   properties of, 138, 322–323, 324  
   rings of, 300, 332–334  
   rotation of, 322–323, 324, 370  
   seasons on, 322–323  
   spacecraft exploration of, 146  
 Urey, Harold, 718  
 Urey-Miller experiment, 718, 719, 728  
*U.S. Mars Reconnaissance Orbiter*, 133  
 U.S. National Observatory, 111  
 Utopia region (Mars), 253
- V**  
 V838 Monocerotis, 511  
 Vacuum energy, 679, 702  
 Valhalla basin, 286  
 Valhalla ridges, 286  
 Valles Marineris (Mariner Valley), 241, 242–243  
 Valley networks, 245  
 Van Allen belts, 180–182, 409, 634  
 Variable stars, 387, 432, 581–582, 583  
 Vega (star), 431, 432, 444  
*Vega 2*, 348  
 Vela series of satellites, 127  
 Vela supernova remnant, 528–529, 530  
 Velocity, 50  
   of Earth, 685  
   measuring, with Doppler effect, 75, 92–93  
   radial, 92–93  
   recessional, 73–74, 620, 671  
   transverse, 424–425  
   wave, 62  
*Venera* missions, 146, 221, 222, 224, 226, 228, 233  
 Venus, 216–233  
   atmosphere of, 220–221, 228–232, 258  
   brightness of, 218  
   density of, 219  
   distance from Earth to, 48–49  
   internal structure of, 232–233  
   long-distance observations of, 220–221  
   magnetic field of, 232–233  
   mass of, 138, 219  
   orbit of, 218  
   phases of, 42, 43  
   properties of, 138, 219–220  
   radar map of, 223  
   radius of, 219  
   rotation rate of, 219–220, 370  
   spacecraft exploration of, 146–147  
   surface of, 221–228  
*Venus Express* mission, 147, 220, 221, 227, 229  
 Vernal equinox, 17  
 Very Large Telescope (VLT), 106, 107, 113, 120, 461, 555, 607  
 Very Long Baseline Interferometry (VLBI), 120, 652  
 Vesta (asteroid), 340, 341, 342–343  
 Vibration, molecular, 91  
*Viking* missions, 146, 239, 240, 241, 243, 248, 250, 253–254, 256, 259, 260, 722  
 Virgo A (M87) galaxy, 618, 628, 629, 632, 634  
 Virgo Cluster, 618, 621, 627, 631, 632, 656–657  
 Virgo Supercluster, 656–657  
 Virus, 717  
 Visible galaxy, 621n  
 Visible light, 60, 62–63  
   obscurance of, 459–460  
 Visible spectrum, 62  
 Visual binary, 440, 441  
 VLA interferometer, 119  
 Voids, 657, 658, 668, 669  
 Volcanoes/volcanism, 170, 173, 174  
   on Enceladus, 311, 312  
   on Io, 281–283  
   lunar, 205–206  
   on Mars, 243–244  
   on Mercury, 206  
   shield volcanoes, 225  
   on Venus, 225–228  
   on Vesta, 341  
*Voyager* missions, 146, 147  
   description of, 142  
   interstellar probes, 729–730  
   Jupiter, 268, 269, 271, 272, 274, 275, 277, 279, 280, 281, 282, 283–284, 287  
   Neptune, 321, 323, 325–326, 330  
   Saturn, 292, 293, 297, 299, 300–301, 302, 304, 305, 307, 313  
   Uranus, 320, 322, 324, 325, 326, 328, 329, 332–333, 334
- W**  
 Wabar (meteorite), 361  
 “Walls,” 658  
   Great Wall, 657–658  
   Sloan Great Wall, 668  
 Waning Moon, 18, 19  
 Warm longitudes, 200  
 Water  
   on Earth, origins of, 185  
   on Europa, 283–285  
   life and, 731–732  
   on Mars, 244–249  
 Water hole, 731–732  
 Water wave, 61  
 Watt, 72  
 Wave frequency, 65, 66  
 Wave motion of radiation, 60–62  
 Wave period, 61  
 Wave theory of radiation, 65, 67  
 Wave velocity, 62  
 Wavelength, 61, 65, 66  
   angular resolution of telescope and, 106–108  
 Waxing Moon, 18, 19  
 Weak nuclear force, 412, 413, 695
- Weakly Interacting Massive Particles (WIMPs), 597  
 Weather  
   on Earth, 14–17  
   on Jupiter, 272–274  
   on Mars, 256  
   on Saturn, 294–296  
   on Uranus and Neptune, 324–326  
 Wegener, Alfred, 174  
 Weight, 49  
 Weinberg, Steven, 694  
 Weird (jumbled) terrain, 208  
 Wheeler, John Archibald, 560  
 Whirlpool (M51) Galaxy, 649  
 White dwarf region, 435, 436  
 White dwarfs, 434, 436, 504–507.  
   *See also* Nova/novae  
   distant, 506  
   helium, 506  
   on H-R diagram, 505, 507  
   life after death for, 522–523  
   neon-oxygen, 507  
 White ovals, 272  
 Wien, Wilhelm, 70  
 Wien’s law, 70, 72, 73, 166, 525  
*Wilkinson Microwave Anisotropy Probe (WMAP)*, 699, 707–709  
 Wilson, Robert, 683–684  
 WIMPs (Weakly Interacting Massive Particles), 597  
 Wind(s)  
   on Jupiter, 269–270  
   on Neptune, 326  
   protostellar, 482, 483–484  
   pulsar, 546, 547  
   on Saturn, 294, 295  
   solar, 143, 180, 390, 391, 400  
   stellar, 511  
   on Uranus, 325  
 Winter solstice, 17  
 Wispy terrain, 311  
*WMAP (Wilkinson Microwave Anisotropy Probe)*, 699, 707–709  
 Wollaston, William, 82  
 Wr 124 (variable star), 387  
 WW Cygni (star), 581
- X**  
*XMM-Newton* satellite, 127  
 X-ray bursters, 548–549  
   millisecond pulsars and, 550  
 X-ray observations, of Sun, 407, 408  
 X-ray radiation, 60, 66, 67  
 X-ray (high-energy) telescopes, 125–128, 407  
 X-rays, detection methods for, 125–126
- Y**  
 Year  
   eclipse, 22–24  
   Galactic, 588  
   leap, 23  
   sidereal, 17  
   tropical, 17  
 Yerkes observatory, 102  
*Yohkoh* X-ray solar observatory, 407  
 Young, Thomas, 67  
 Young star cluster, H-R diagram of, 513, 514  
 Yuty crater, 247, 248
- Z**  
 Zeeman effect, 95, 401  
 Zero-age main sequence (ZAMS), 477–478  
 Zodiac, 14, 15  
 Zonal flow  
   on Jupiter, 269–270  
   on Saturn, 294, 295  
 “Zone of influence,” in binary-star system, 516  
 Zones, in jovian atmospheres, 268–270


# Star Charts

Have you ever become lost in an unfamiliar city or state? Chances are you used two things to get around: a map and some signposts. In much the same way, these two items can help you find your way around the night sky in any season. Fortunately, in addition to the seasonal Star Charts on the following pages, the sky provides us with two major signposts. Each seasonal description will talk about the Big Dipper—a group of seven bright stars that dominates the constellation Ursa Major the Great Bear. Meanwhile, the


constellation Orion the Hunter plays a key role in finding your way around the sky from late autumn until early spring.

Each chart portrays the sky as seen from near 35° north latitude at the times shown at the top of the page. Located just outside the chart are the four directions: north, south, east, and west. To find stars above your horizon, hold the map overhead and orient it so a direction label matches the direction you're facing. The stars above the map's horizon now match what's in the sky.


## WINTER




## SPRING



## SUMMER



## AUTUMN



Reproduced by permission. © 2007, Astronomy Magazine, Kalmbach Publishing Co.



## Exploring the Winter Sky

Winter finds the Big Dipper climbing the northeastern sky, with the three stars of its handle pointing toward the horizon and the four stars of its bowl standing highest. The entire sky rotates around a point near Polaris, a 2nd-magnitude star found by extending a line from the uppermost pair of stars in the bowl across the sky to the left of the Dipper. Polaris also performs two other valuable functions: The altitude of the star above the horizon equals your latitude north of the equator, and a straight line dropped from the star to the horizon.


Turn around with your back to the Dipper and you'll be facing the diamond-studded winter sky. The second great signpost in the sky, Orion the Hunter, is central to the brilliant scene. Three closely spaced, 2nd-magnitude stars form a straight line that represents the unmistakable belt of Orion. Extending the imaginary line joining these stars to the upper right leads to Taurus the Bull and its orangish 1st-magnitude star, Aldebaran. Reverse the direction of your gaze to the belt's lower left and you cannot miss Sirius the Dog Star—brightest in all the heavens at magnitude  $-1.5$ .

Now move perpendicular to the belt from its westernmost star, Mintaka, and find at the upper left of Orion the red supergiant star Betelgeuse. Nearly a thousand times the Sun's diameter, Betelgeuse marks one shoulder of Orion. Continuing this line brings you to a pair of bright stars, Castor and Pollux. Two lines of fainter stars extend from this pair back toward Orion—these represent Gemini the Twins. At the northeastern corner of this constellation lies the beautiful open star cluster M35. Head south of the belt instead and your gaze will fall on the blue supergiant star Rigel, Orion's other luminary.

Above Orion and nearly overhead on winter evenings is brilliant Capella, in Auriga the Charioteer. Extending a line through the shoulders of Orion to the east leads you to Procyon in Canis Minor the Little Dog. Once you have these principal stars mastered, using the chart to discover the fainter constellations will be a whole lot easier. Take your time, and enjoy the journey. Before leaving Orion, however, aim your binoculars at the line of stars below the belt. The fuzzy "star" in the middle is actually the glorious Orion Nebula (M42), a stellar nursery illuminated by bright, newly formed stars.

# WINTER

2 a.m. on December 1; midnight on January 1; 10 p.m. on February 1



Reproduced by permission. © 2007, Astronomy Magazine, Kalmbach Publishing Co.

## Exploring the Spring Sky

The Big Dipper, our signpost in the sky, swings high overhead during the spring and lies just north of the center of the chart. This season of rejuvenation encourages us to move outdoors with the milder temperatures, and with the new season a new set of stars beckons us.

Follow the arc of stars outlining the handle of the Dipper away from the bowl and you will land on brilliant Arcturus. This orangish star dominates the spring sky in the kite-shaped constellation Boötes the Herdsman. Well to the west of Boötes lies Leo the Lion. You can find its brightest star, Regulus, by using the pointers of the Dipper in reverse. Regulus lies at the base of a group of stars shaped like a sickle or backward question mark, which represents the head of the lion.

Midway between Regulus and Pollux in Gemini, which is now sinking in the west, is the diminutive group Cancer the Crab. Centered in this group is a hazy patch of light that binoculars reveal as the Beehive star cluster (M44).

To the southeast of Leo lies the realm of the galaxies and the constellation Virgo the Maiden. Virgo's brightest star, Spica, shines at magnitude 1.0.


During springtime, the Milky Way lies level with the horizon, and it's easy to visualize that we are looking out of the plane of our Galaxy. In the direction of Virgo, Leo, Coma Berenices, and Ursa Major lie thousands of galaxies whose light is unhindered by intervening dust in our own Galaxy. However, all these galaxies are elusive to the untrained eye and require binoculars or a telescope to be seen.

Boötes lies on the eastern border of this galaxy haven. Midway between Arcturus and Vega, the bright "summer" star rising in the northeast, is a region where no star shines brighter than 2nd magnitude. A semicircle of stars represents Corona Borealis the Northern Crown, and adjacent to it is a large region that houses Hercules the Strongman, the fifth-biggest constellation in the sky. It is here we can find the northern sky's brightest globular star cluster, M13. A naked-eye object from a dark site, it looks spectacular when viewed through a telescope.

Returning to Ursa Major, check the second-to-last star in the Dipper's handle. Most people will see it as double, while binoculars show this easily. The pair is called Mizar and Alcor, and they lie just  $0.2^\circ$  apart. A telescope reveals Mizar itself to be a double. Its companion star shines at magnitude 4.0 and lies 14 arc seconds away.

# SPRING

1 a.m. on March 1; 11 p.m. on April 1; 9 p.m. on May 1. Add one hour for daylight-saving time



Reproduced by permission. © 2007, Astronomy Magazine, Kalmbach Publishing Co.

## Exploring the Summer Sky

The richness of the summer sky is exemplified by the splendor of the Milky Way. Stretching from the northern horizon in Perseus, through the cross-shaped constellation Cygnus overhead, and down to Sagittarius in the south, the Milky Way is packed with riches. These riches include star clusters, nebulae, double stars, and variable stars.

Let's start with the Big Dipper, our perennial signpost, which now lies in the northwest with its handle still pointing toward Arcturus. High overhead, and the first star to appear after sunset, is Vega in Lyra the Harp. Vega forms one corner of the summer triangle, a conspicuous asterism of three stars. Near Vega lies the famous double-double, Epsilon ( $\epsilon$ ) Lyrae. Two 5th-magnitude stars lie just over 3 arc minutes apart and can be split when viewed through binoculars. Each of these two stars is also double, but you need a telescope to split them.

To the east of Vega lies the triangle's second star: Deneb in Cygnus the Swan (some see a cross in this pattern). Deneb marks the tail of this graceful bird, the cross represents its outstretched wings, and the base of the cross denotes its head, which is marked by the incomparable double star Albireo. Albireo matches a 3rd-magnitude yellow star and a

5th-magnitude blue star and offers the finest color contrast anywhere in the sky. Deneb is a supergiant star that pumps out as much light as 60,000 Suns. Also notice that the Milky Way splits into two parts in Cygnus, a giant rift caused by interstellar dust blocking starlight from beyond.


Altair, the third star of the summer triangle and the one farthest south, is the second brightest of the three. Lying 17 light-years away, it's the brightest star in the constellation Aquila the Eagle.

Frequently overlooked to the north of Deneb lies the constellation Cepheus the King. Shaped rather like a bishop's hat, the southern corner of Cepheus is marked by a compact triangle of stars that includes Delta ( $\Delta$ ) Cephei. This famous star is the prototype of the Cepheid variable stars used to determine the distances to some of the nearer galaxies. It varies regularly from magnitude 3.6 to 4.3 and back again with a 5.37-day period.

Hugging the southern horizon, the constellations Sagittarius the Archer and Scorpius the Scorpion lie in the thickest part of the Milky Way. Scorpius's brightest star, Antares, is a red supergiant star whose name means "rival of Mars" and derives from its similarity to the planet in both color and brightness.

# SUMMER

1 a.m. on June 1; 11 p.m. on July 1; 9 p.m. on August 1. Add one hour for daylight-saving time



Reproduced by permission. © 2007, Astronomy Magazine, Kalmbach Publishing Co.

## Exploring the Autumn Sky

The cool nights of autumn are here to remind us the chill of winter is not far off. Along with the cool air, the brilliant stars of the summer triangle descend in the west to be replaced with a rather bland-looking region of sky. But don't let initial appearances deceive you. Hidden in the fall sky are gems equal to summertime.

The Big Dipper swings low this season, and for parts of the Southern United States it actually sets. Cassiopeia the Queen, a group of five bright stars in the shape of a "W" or "M," reaches its highest point overhead, the same spot the Big Dipper reached six months ago. To the east of Cassiopeia, Perseus the Hero rises high. Nestled between these two groups is the wondrous Double Cluster—NGC 869 and NGC 884—a fantastic sight in binoculars or a low-power telescope.

Our view to the south of the Milky Way is a window out of the plane of our Galaxy in the opposite direction to that visible in spring. This allows us to look at the Local Group of galaxies. Due south of Cassiopeia is the Andromeda Galaxy (M31), a 4th-magnitude smudge of light that passes directly overhead


around 9 p.m. in mid-November. Farther south, between Andromeda and Triangulum, lies M33, a sprawling face-on spiral galaxy best seen in binoculars or a rich-field telescope.

The Great Square of Pegasus passes just south of the zenith. Four 2nd- and 3rd-magnitude stars form the square, but few stars can be seen inside of it. If you draw a line between the two stars on the west side of the square and extend it southward, you'll find 1st-magnitude Fomalhaut in Piscis Austrinus the Southern Fish. Fomalhaut is the solitary bright star low in the south. Using the eastern side of the square as a pointer to the south brings you to Diphda in the large, faint constellation of Cetus the Whale.

To the east of the Square lies the Pleiades star cluster (M45) in Taurus, which reminds us of the forthcoming winter. By late evening in October and early evening in December, Taurus and Orion have both cleared the horizon and Gemini is rising in the northeast. In concert with the reappearance of winter constellations, the view to the northwest finds summertime's Cygnus and Lyra about to set. The autumn season is a great transition period, both on Earth and in the sky, and a fine time to experience the subtleties of these constellations.

# AUTUMN

1 a.m. on September 1; 11 p.m. on October 1; 9 p.m. on November 1. Add one hour for daylight-saving time




Reproduced by permission. © 2007, Astronomy Magazine, Kalmbach Publishing Co.




*This page intentionally left blank*

# The Entire Electromagnetic Spectrum



# The Distance Scale



## Some Useful Constants and Physical Measurements\*

astronomical unit	$1 \text{ A.U.} = 1.496 \times 10^8 \text{ km}$ ( $1.5 \times 10^8 \text{ km}$ )
light year	$1 \text{ ly} = 9.46 \times 10^{12} \text{ km}$ ( $10^{13} \text{ km}$ ; 6 trillion miles)
parsec	$1 \text{ pc} = 3.09 \times 10^{13} \text{ km} = 3.3 \text{ ly}$
speed of light	$c = 299,792,458 \text{ km/s}$ ( $3 \times 10^5 \text{ km/s}$ )
Stefan-Boltzmann constant	$\sigma$ [Greek sigma] = $5.67 \times 10^{-8} \text{ W/m}^2 \cdot \text{K}^4$
Planck's constant	$h = 6.63 \times 10^{-34} \text{ Js}$
gravitational constant	$G = 6.67 \times 10^{-11} \text{ Nm}^2/\text{kg}^2$
mass of Earth	$M_{\oplus} = 5.97 \times 10^{24} \text{ kg}$ ( $6 \times 10^{24} \text{ kg}$ ; about 6000 billion billion tons)
radius of Earth	$R_{\oplus} = 6378 \text{ km}$ (6500 km)
mass of the Sun	$M_{\odot} = 1.99 \times 10^{30} \text{ kg}$ ( $2 \times 10^{30} \text{ kg}$ )
radius of the Sun	$R_{\odot} = 6.96 \times 10^5 \text{ km}$ ( $7 \times 10^5 \text{ km}$ )
luminosity of the Sun	$L_{\odot} = 3.90 \times 10^{26} \text{ W}$
effective temperature of the Sun	$T_{\odot} = 5778 \text{ K}$ (5800 K)
Hubble constant	$H_0 \approx 70 \text{ km/s/Mpc}$
mass of the electron	$m_e = 9.11 \times 10^{-31} \text{ kg}$
mass of the proton	$m_p = 1.67 \times 10^{-27} \text{ kg}$

\*The rounded-off values used in the text are shown in parentheses.

### Conversions between English and Metric Units

1 inch	=	2.54 centimeters (cm)	1 mile	=	1.609 kilometers (km)
1 foot (ft)	=	0.3048 meters (m)	1 pound (lb)	=	453.6 grams (g) or 0.4536 kilograms (kg) (on Earth)

## Main-Sequence Stellar Properties by Spectral Class

Spectral Class	Typical Surface Temperature	Color	Mass*	Luminosity*	Lifetime*	Familiar Examples
	(K)		( $M_{\odot}$ )	( $L_{\odot}$ )	( $10^6 \text{ yr}$ )	
O	>30,000	Electric Blue	>20	>100,000	<2	Mintaka (O9)
B	20,000	Blue	7	500	140	Spica (B2)
A	10,000	White	3	60	500	Vega (A0) Sirius (A1)
F	7,000	Yellow - white	1.5	7	2000	Procyon (F5)
G	6,000	Yellow	1.0	1.0	10,000	Sun (G2) Alpha Centauri (G2)
K	4,000	Orange	0.8	0.3	30,000	Epsilon Eridani (K2)
M	3,000	Red	0.1	0.00006	16,000,000	Proxima Centauri (M5) Barnard's Star (M5)

\*Approximate values for stars of solar composition

*This page intentionally left blank*



THE  
**BIG**  
PICTURE

**The Big Picture** Your students deserve the latest content and the best technology in their Introduction to Astronomy course.

**The Big Question** Who but Chaisson and McMillan would you trust to deliver what your students deserve?



# IMPROVED PEDAGOGY


**The Big Picture** There is something philosophically intriguing about the idea that the deaths of some stars cause the birth of others. Build up, break down, change . . . dust to dust is a scientific concept. Many of the elements composing our world and ourselves were created in the violent explosions of long-gone stars. It sounds rather poetic that we are made of mostly stardust, but it happens to be true.

**THE BIG PICTURE** feature on each chapter-opening spread explains how chapter content fits in with an overall perspective on introductory astronomy, helping students see how each chapter is connected to a broad understanding of the universe.

**The Big Question** Despite widespread observations of supernovae and their scattered debris, researchers still don't know exactly how these massive stars actually manage to explode. Although many times the mass of the Sun, these stars defy gravity by ripping themselves apart. That they blow their whole being to smithereens is well established, but how do they do it, reversing their catastrophic inward collapse to become outwardly exploding stars?

**THE BIG QUESTION** mirrors The Big Picture and piques student interest at the end of each chapter. It identifies a critical unanswered question relating to the chapter content, reinforcing that astronomy is still an evolving science.

**ANNOTATED ART** now appears in close to half of all the figures in *Astronomy Today*, 8th edition. Informed by education research, these blue annotations function much like the professor's voice, interpreting and explaining the art for the student.



## Radiation

INFORMATION FROM THE COSMOS

Astronomical objects are more than just things of beauty in the night sky. Planets, stars, and galaxies are of vital significance if we are to fully understand our place in the larger scheme of things in the universe. Each object is a source of information about the material aspects of our universe—its state of motion, its temperature, its chemical composition, and even its past history.

Much of this information comes to us in the form of light. When we look at the stars, the light we see actually began its journey to Earth decades, centuries—even millennia—ago. The faint rays from the most distant galaxies have taken billions of years to reach us. The stars and galaxies in the night sky show us the far away and the long ago. In this chapter, we begin our study of how astronomers extract information from the light emitted by astronomical objects. These basic concepts of radiation are central to modern astronomy.

**The Big Picture** Our human eyes actually see only a small part of the universe—we literally see optical, or visible, light. There is a much bigger picture to be sensed beyond visible light—invisible radiation such as heat or radio waves or X-rays. Many different kinds of radiation are constantly traveling through space, and detailed study of this wider range of visible and invisible information is the main way that astronomers study stars and other distant objects well beyond Earth.

**LEFT:** In about 5 billion years, the Sun will begin running out of fuel. Its hydrogen gas will become depleted at its center, causing the bulk of our old and decrepit star to slowly disperse into space. We can actually watch such an amazing event by observing other stars that are now dying. This stunning image captures the Helix Nebula about 650 light-years away—but not in visible light. Instead, it shows invisible radiation emitted by the former star: infrared radiation (mostly yellow in this image) captured by the Spitzer Space Telescope and ultraviolet radiation (mostly black) by the galaxy evolution Explorer satellite enable close inspection of the remarkable process of stellar death. (NASA, Caltech)

### Learning Outcomes

Studying this chapter will enable you to

1. Outline the basic properties of wave motion.
2. Tell how electromagnetic radiation transfers energy and information through interstellar space.
3. Describe the major regions of the electromagnetic spectrum and explain how Earth's atmosphere affects our ability to make astronomical observations at different wavelengths.
4. Explain what is meant by the term "blackbody radiation" and describe the basic properties of such radiation.
5. Tell how we can determine the temperature of an object by observing the radiation it emits.
6. Describe how the relative motion between a source of radiation and its observer can change the perceived wavelength of the radiation, and explain the importance of this phenomenon to astronomy.

MasteringAstronomy® Visit the MasteringAstronomy Study Area for quizzes, animations, videos, interactive figures, and self-guided tutorials.

59

## Chapter Review

### SUMMARY

1. **Electromagnetic radiation** (p. 60) travels through space in the form of a **wave** (p. 60). A wave is characterized by its **period** (p. 61), the length of time taken for one complete cycle; its **wavelength** (p. 61), the distance between successive wave crests; and its **amplitude** (p. 61), which measures the size of the disturbance associated with the wave. A wave's **frequency** (p. 61) is the number of wave crests that pass a given point in one second.
2. Any electrically charged object is surrounded by an **electric field** (p. 63) that determines the force the object exerts on other charged objects. When a charged particle moves, information about its motion is transmitted via the particle's changing **electric and magnetic fields** (pp. 63, 64). The information travels at the **speed of light** (p. 64) as an electromagnetic wave. **Diffraction** (p. 67) and **interference** (p. 67) are properties of radiation that mark it as a wave phenomenon.
3. The color of visible light is simply a measure of its wavelength—and light has a longer wavelength than blue light. The entire **electromagnetic spectrum** (p. 65) consists of (in order of increasing frequency) **radio waves**, **infrared radiation**, **visible light**, **ultraviolet radiation**, **X-rays**, and **gamma rays** (p. 66). Only radio waves, some infrared wavelengths, and visible light can penetrate the atmosphere and reach the ground from space.
4. The **temperature** (p. 68) of an object is a measure of the speed with which its constituent particles move. The intensity of radiation emitted by an object has a characteristic distribution, called a **blackbody curve** (p. 68), which depends only on the object's temperature.
5. **Wien's law** (p. 70) tells us that the wavelength at which the object radiates most of its energy is inversely proportional to its temperature. Measuring that peak wavelength tells us the object's temperature. **Stefan's law** (p. 70) states that the total amount of energy radiated is proportional to the fourth power of the temperature.
6. Our perception of the wavelength of a beam of light can be altered by the source's velocity relative to us. This motion-induced change in the observed frequency of a wave is called the **Doppler effect** (p. 73). Any net motion of the source away from the observer causes a **redshift**—a shift to lower frequencies—in the received beam. Motion toward the observer causes a **blueshift**. The extent of the shift is directly proportional to the source's radial velocity relative to the observer.

**LEARNING OUTCOMES** have replaced the Learning Goals of previous editions and help guide students in structuring their understanding of complex topics; their connection to the end-of-chapter Summary and Review and Discussion questions are made explicit and are consistent with the AAS outcomes.




# MasteringAstronomy®

www.masteringastronomy.com


MasteringAstronomy is designed to improve results by helping students quickly master concepts. Students benefit from wrong answer specific feedback, hints, and a huge variety of educationally effective content while unrivalled gradebook diagnostics allow an instructor to pinpoint the weaknesses and misconceptions of their class.

## NEW NARRATED FIGURES

provide students with expanded audio explanations and animation of figures from the text and include pause-and-predict questions. Assignable activities written for Narrated Figures can be found in the MasteringAstronomy item library.



## MasteringAstronomy®



**INTERACTIVE FIGURES AND PHOTOS** illustrate key concepts from the book and bring important static figures to life through animation.

**NEW TO THIS EDITION** Visual activities or tutorials are available for every chapter.

## LEARNING OUTCOMES

quantify students' learning gains and allow you to share those results quickly and easily.

**TUTORIALS** are available for every major learning outcome.

## MasteringAstronomy®


Item Library


Preview items available for assignment ([How do I assign items?](#))

[Create New Item](#) [Import Items](#)

Book/Source:	Chapter:	Display By:	Learning Outcomes:	Publisher Items	My Items	TIME	DIFFICULTY (5-hardest)	USAGE STATISTICS (Roll over any colored segment)
Chaisson/McMillan, Astronomy Today, 7e	6 THE SOLAR SYSTEM	Learning Outcomes	Explain how the gravitational force between two objects depends on their masses and the distance between them.	<input checked="" type="checkbox"/> Tutorial <input checked="" type="checkbox"/> End-of-Chapter	<input checked="" type="checkbox"/> My Items	3m	1	
			Identify the information needed to compute the mass of an orbiting system using Newton's version of Kepler's third law.	<input checked="" type="checkbox"/> Coaching Activities <input checked="" type="checkbox"/> Test Bank		3m	1	
			Describe the major patterns we find among the objects orbiting the Sun.			--	--	
			Describe the nebular theory for solar system formation.			--	--	
			Explain how the nebular theory accounts for the motions of the planets and their moons.			--	--	
			Explain how the nebular theory accounts for the dependence of planetary composition on distance from the Sun.			--	--	
			Explain how the nebular theory accounts for the existence of asteroids and comets.			--	--	
			Arrange the jovian planets in order of increasing mass or radius.			--	--	
			Distinguish between comets, asteroids, meteors, and meteorites.			--	--	


The H-R diagram plots stars by luminosity (vertical axis) and temperature or spectral class (horizontal axis). The dashed diagonal lines are lines of constant radius.






### Overlay 1

Adding stars reveals how the H-R diagram organizes stellar information. Hotter (bluer) stars lie to the left; cooler (redder) stars lie to the right. Some specific stars are noted.



## Overlay 2


Patterns and groups emerge. Note especially the main sequence. Our star, the Sun, is by definition 1 solar unit in luminosity and 1 solar radius in size. It is a yellow G-type star, now on the main sequence.



### Overlay 3

The track of a 1-solar-mass star on the H–R diagram:

- 5 Protostar
- 6 Star
- 7 Main-sequence star
- 8 Subgiant branch
- 9 Helium flash
- 10 Horizontal branch
- 11 Asymptotic giant branch
- 12 Carbon core / planetary nebula
- 13 White dwarf
- 14 Black dwarf



#### Overlay 4

Some data for rich globular clusters (M80 mostly, shown at right). At the lower end the stars lie along the main sequence; just beyond G they turn off, following the subgiant, red giant, and horizontal branch stages. An age over 10 billion years is indicated by this degree of evolution (see Figure 20.17). Because globular cluster stars contain far fewer heavy elements than the Sun, those stars tend to be hotter and brighter for the same mass.

